

# Pacific Journal of Mathematics

## **STIEFEL'S THEOREM AND TORAL ACTIONS**

JANEY ANTONIO DACCACH AND ARTHUR GABRIEL WASSERMAN

## STIEFEL'S THEOREM AND TORAL ACTIONS

J. A. DACCACH AND A. G. WASSERMAN

The second Stiefel-Whitney class of an orientable manifold admitting an effective codimension three toral action is shown to be Poincaré dual to the fixed point sets of circle subgroups and cyclic subgroups of even order.

**0. Introduction.** Stiefel [5] proved that the second Stiefel-Whitney class of an orientable 3-manifold is zero. A simple generalization of this theorem is that the second Stiefel-Whitney class of an orientable  $(n + 3)$ -manifold,  $M$ , vanishes if  $M$  admits a free action of  $T^n$ , the  $n$ -torus. If  $T^n$  acts almost freely and effectively on an orientable manifold  $M^{n+3}$ , that is, if all isotropy subgroups are finite, then  $w_2(M)$  need not vanish; in fact  $w_2(M)$  is Poincaré dual to the fixed point sets of cyclic subgroups of even order.

In this note, we consider an arbitrary effective action of  $T^n$  on an orientable manifold  $M^{n+3}$  and show that the Poincaré dual of  $w_2(M)$  is represented by the fixed point sets of circle subgroups and cyclic subgroups of even order.

In §1 we establish some notation and give a precise statement of the theorem. In §2 we reduce the proof of the theorem to the case of compact manifolds having only cyclic or circle isotropy subgroups. In §3 we study the cases of only cyclic isotropy subgroups or only circle isotropy subgroups and in §4 we prove the theorem by reducing to the two special cases of §3.

**1. Statement of Theorem.** If  $M$  is a manifold with a smooth action of  $T^n$  and  $H$  is any subgroup of  $T^n$  then  $E(M, H) = \{x \in M \mid T_x^n = H\}$  where  $T_x^n$  is the isotropy subgroup at  $x$ ;  $E(M, H)$  is open in  $F(M, H) = \{x \in M \mid h(x) = x \text{ for all } h \in H\}$ , the fixed point set of  $H$  [3]. If  $X$  is any component of  $F(M, H)$  then  $X$  is a closed submanifold of  $M$  and  $X \cap E(M, H)$  is either empty or dense in the  $T^n/H$  space  $X$  by [3]; hence  $E(M, H)$  is a union of disjoint closed submanifolds of  $M$ .

If  $A^a$  is a closed submanifold of a manifold  $B^b$ , the Poincaré dual of  $A$ ,  $D(A) \in H^{b-a}(B; Z_2) \simeq \text{Hom}(H_{b-a}(B; Z_2); Z_2)$ , is defined by  $\langle D(A), Z \rangle = \text{number of points in } A \pitchfork Z \pmod{2}$  where  $\pitchfork$  indicates the intersection of  $A$  with the cycle  $Z$  in general position [4].

If  $T^n$  acts smoothly and effectively on an orientable manifold  $M^{n+3}$ , then for any cyclic subgroup  $H \subset T^n$ ,  $\overline{E(M, H)}$  is a codimension two submanifold since  $H$  must act linearly, effectively, and in an orientation preserving manner on the three dimensional slice. If  $H$  is isomorphic to  $S^1$ ,  $\overline{E(M, H)}$  will in general have components of codimension two and four; let  $E_2(M, H)$  denote the components of codimension two. Finally, for  $T^n$  acting smoothly and effectively on an orientable manifold  $M^{n+3}$ , we define  $A(M) \in H^2(M; Z_2)$  by

$$A(M) = \sum_{H \simeq S^1} D(\overline{E_2(M, H)}) + \sum_{H \simeq Z_{2m}} D(\overline{E(M, H)}).$$

**THEOREM.** *If  $T^n$  acts smoothly and effectively on an orientable  $(n + 3)$ -manifold,  $M^{n+3}$ , then  $A(M) = w_2(M)$ , the second Stiefel-Whitney class of  $M$ .*

The remainder of the paper is devoted to the proof of this theorem.

**2. First reduction.**

**LEMMA 1.** *It is sufficient to prove the theorem for compact manifolds without boundary.*

*Proof.* We wish to prove that  $A(M) = w_2(M)$  for an arbitrary orientable  $(n + 3)$ -manifold  $M$  with effective smooth  $T^n$  action. If  $A(M) - w_2(M) \neq 0 \in H^2(M; Z_2)$  then there is a singular cycle  $Z \in H_2(M, Z_2)$  with  $\langle A(M) - w_2(M), Z \rangle \neq 0$ . But any such cycle is contained in a compact, invariant, submanifold with boundary  $X$  that is  $Z = i_*Z'$  for  $Z' \in H_2(X; Z_2)$ ,  $i: X \hookrightarrow M$ . Hence

$$\begin{aligned} \langle A(M) - w_2(M), Z \rangle &= \langle A(M) - w_2(M), i_*Z' \rangle \\ &= \langle i^*A(M) - i^*w_2(M), Z' \rangle = \langle A(X) - w_2(X), Z' \rangle \end{aligned}$$

and so it is sufficient to prove the theorem for compact manifolds, possibly with boundary. Finally, let  $Y = X \cup_{\partial X} X$  be the double of  $X$  and  $j: X \hookrightarrow Y$ . Then

$$\begin{aligned} \langle w_2(Y) - A(Y), j_*Z' \rangle &= \langle j^*A(Y) - j^*w_2(Y), Z' \rangle \\ &= \langle A(X) - w_2(X), Z' \rangle \end{aligned}$$

so it is sufficient to prove the theorem for  $Y$ , that is, for compact manifolds without boundary. □

If  $T^n$  acts effectively on the orientable manifold  $M^{n+3}$ , then  $T_x^n$ , the isotropy subgroup at  $x$ , can only be isomorphic to one of  $\{e\}$ ,  $Z_m$ ,  $S^1$ ,  $S^1 \times Z_m$ ,  $T^2$  or  $T^3$  because the isotropy subgroup must act linearly, effectively, and in an orientation preserving manner on the slice at  $x$ .

**LEMMA 2.** *It is sufficient to prove the theorem for compact manifold  $M$  without boundary, such that  $E(M, H) = \phi$  for  $H$  isomorphic to  $T^3$ ,  $T^2$  or  $S^1 \times Z_m$  and such that  $E(M, H) = E_2(M, H)$  for  $H$  isomorphic to  $S^1$ .*

*Proof.* Let  $H_1, \dots, H_r$  be the isotropy subgroups isomorphic to  $T^3$ ,  $T^2$  or  $S^1 \times Z_m$  and suppose that  $H_1$  is isomorphic to  $T^3$ . Then  $H_1$  is a maximal isotropy subgroup and hence  $E(M, H_1) = F(M, H_1)$  is a closed submanifold of  $M$  of dimension  $n - 3$  since  $T^3$  must act effectively on fibre of the normal bundle of  $F(M, H_1)$  in  $M$ . Let  $U$  be a closed, invariant tubular neighborhood of  $E(M, H_1)$ ,  $\partial U$  the boundary of  $U$ . Then we have  $H^2(M, M - U; Z_2) \simeq H^2(U, \partial U; Z_2)$  by excision and  $H^2(U, \partial U; Z_2) \simeq H_{n+1}(U; Z_2) \simeq H_{n+1}(F(M, H_1); Z_2) = 0$  by Poincaré duality, and hence we have in the exact sequence

$$H^2(M, M - U; Z_2) \rightarrow H^2(M; Z_2) \xrightarrow{i_*} H^2(M - U; Z_2)$$

that  $i_*$  is injective and hence it is sufficient to prove  $i^*(A(M) - w_2(M)) = 0$ , that is  $A(M - U) = w_2(M - U)$ .

By doubling  $M - U$  as in Lemma 1 we see that it is sufficient to consider compact manifolds without boundary having isotropy subgroups  $H_2, \dots, H_r$ . Repeating this argument a finite number of times removes all isotropy subgroups isomorphic to  $T^3$ . If  $H_s$  is isomorphic to  $T^2$  and if  $E(M, H) = \phi$  for  $H$  isomorphic to  $T^3$  then  $E(M, H_s) = F(M, H_s)$  is a closed invariant submanifold of codimension four and hence  $H_{n+1}(F(M, H_s); Z_2) = 0$  as before and we may repeat the argument. Continuing in this manner, we remove next  $F(M, H)$  for  $H$  isomorphic to  $S^1 \times Z_m$  since codimension  $F(M, H)$  is four in this case and finally we remove the components of  $F(M, H)$ ,  $H$  isomorphic to  $S^1$ , whose codimension is four.  $\square$

### 3. Two special cases.

**PROPOSITION 3.** *Let  $T^n$  act effectively and smoothly on the closed orientable manifold  $M^{n+3}$  with all isotropy subgroups finite. Then  $w_2(M) = A(M)$ .*

LEMMA 4. *With the same hypothesis of Proposition 3,  $M^{n+3}/T^n$  is a closed orientable 3-manifold.*

*Proof.* Let  $\pi: M^{n+3} \rightarrow M^{n+3}/T^n$  denote the orbit map. To give a chart  $(U_i, \phi_i)$  in  $M^{n+3}/T^n$  it is sufficient to give a smooth invariant map  $\bar{\phi}_i: \pi^{-1}(U_i) \rightarrow \mathbf{R}^3$ . If  $x \in M$  with  $T_x^n = e$  then the slice at  $x$ ,  $S_x$  is diffeomorphic to  $\mathbf{R}^3$  [1], and we take  $U = T^n \times S$ ,  $\bar{\phi}: T^n \times S \rightarrow S \simeq \mathbf{R}^3$ . If  $T_x^n = Z_m$  then  $S_x \simeq C \times \mathbf{R}$  with  $g \in Z_m$  acting linearly on  $C \times \mathbf{R}$  via  $g(z, t) = (\xi z, t)$  for  $\xi$  an  $m$ th root of unity. Define  $\bar{\Phi}: T^n \times_{Z_m} S_x \rightarrow C \times \mathbf{R}$  via  $\bar{\phi}(X, z, t) = (z^m, t)$ .

To prove that  $M/T^n$  is orientable we note that  $T^n$  acts freely on  $M - \bigcup_{H \neq e} F(M, H)$  so that  $(M - \bigcup_{H \neq e} F(M, H))/T^n$  is orientable and  $\bigcup_{H \neq e} F(M, H)/T^n$  consists of isolated curves in  $M/T^n$  so  $M/T^n$  is orientable also. □

*Proof of Proposition 3.* Let  $x \in H_2(M; Z_2)$  and let  $x$  be represented by a submanifold  $Q^2 \subset M$  [6]. We must show  $\langle w_2(M), Q \rangle = \langle A(M), Q \rangle$ . We may assume that  $Q$  intersects each fixed point set  $F(M, H)$  transversally, that  $Q \cap \bigcup_{H \neq e} F(M; H) = \{p_1, \dots, p_r\}$ , and that in a neighborhood of an intersection point  $p_i \in Q \cap F(M, H_i)$ ,  $Q$  coincides with a fiber of a tubular neighborhood. More precisely, there is a slice at  $p_i$ ,  $S_{p_i} \simeq C \times \mathbf{R}$  and a neighborhood  $U_{p_i}$  of  $p_i$  in  $Q$  with  $U_{p_i} \subset C \times 0$ . To compute  $\langle w_2(M), Q \rangle$  we split  $T(M)/Q$  as  $\theta^{n+1} \oplus \eta^2$  where  $\theta^{n+1}$  is a trivial bundle and then  $\langle w_2(M), Q \rangle = \langle w_2(\eta^2), Q \rangle = \langle \chi(\eta^2), Q \rangle =$  number of zeroes of a generic section of  $\eta^2 \bmod 2$ , where  $\chi$  denotes the Euler class. First note that we have a splitting  $T(M) = \theta^n \oplus \xi^3$  where  $\theta^n$ , the bundle of tangents to the orbits, is trivial since all isotropy subgroups are finite. We must now split  $\xi^3/Q = \eta^2 \oplus \theta^1$ . We have  $d\pi: T(M) \rightarrow T(M/T^n)$  is an epimorphism with kernel  $\theta^n$  off the fixed point sets; therefore  $d\pi: \xi^3 \rightarrow T(M/T^n)$  is an isomorphism off the fixed point sets, and at a fixed point  $x$  of  $H$ ,  $d\pi_x$  has rank 1. Since every orientable 3-manifold is parallelizable by Stiefel's theorem [5], we can find vector fields  $X_1, X_2, X_3$  which are linearly independent at every point of  $M/T^n$ . Around each point  $p_i$  we can choose a slice  $S_{p_i} \simeq C \times \mathbf{R}$  with coordinate functions  $(z, t)$  such that  $\pi(S_{p_i})$  is a coordinate chart of  $M/T^n$  at  $\pi(p_i)$  with coordinate functions  $(w, t)$  where  $\pi(z, t) = (z^m, t)$ . So we can consider in  $\pi(S_{p_i})$  the vector fields  $\partial/\partial x, \partial/\partial y, \partial/\partial t$  where  $w = (x, y)$ . Since, up to permutation of the indices, the frame  $X_1|_{\pi(S_{p_i})}, X_2|_{\pi(S_{p_i})}, X_3|_{\pi(S_{p_i})}$  is homotopic to the frame  $\partial/\partial x, \partial/\partial y, \partial/\partial t$  we can modify the vector fields  $X_1, X_2, X_3$  to  $\tilde{X}_1, \tilde{X}_2, \tilde{X}_3$  so that in the neighborhood  $\pi(S_{p_i})$

we have  $\tilde{X}_1 = \partial/\partial x$ ,  $\tilde{X}_2 = \partial/\partial y$ ,  $\tilde{X}_3 = \partial/\partial t$ . Note that the vector field  $Y_1$  on  $S_{p_i}$  given by  $Y_1 = \partial/\partial t$  satisfies  $d\pi(y_1) = \tilde{X}_3$ . Then on  $Q - P_i$  we have a well defined nonzero vector field  $Y_2$  that corresponds via  $d\pi$  to the vector field  $\tilde{X}_3$ , since  $d\pi$  is an isomorphism off the fixed point set. Restricting the vector field  $Y_1$  to  $U_i$  we have that  $Y_1 = Y_2$  in  $U_i - \{P_i\}$ . So  $Y_2$  extends to a nonzero vector field  $Y$  in  $Q$ . Let us use this vector field to split  $\xi^3|_Q \simeq \eta^2 \oplus \theta$ . Now  $Z = d\pi^{-1}(\tilde{X}_1)$  is a section of  $\eta^2|_{Q - \{P_i\}}$ . Denoting by  $\chi_i$  the index of  $Z$  at  $P_i$  we have that  $\chi(\eta^2) \bmod 2 = \sum_{i=1}^r \chi_i \bmod 2$ . The map  $\pi$  restricted to  $U_i$  is given by  $\pi(z) = z^{m_i}$  and  $d\pi(Z) = \partial/\partial x$  for  $Z|_{U_i - p_i}$ . Writing this in a matrix form we have

$$m_i r^{m_i} \begin{pmatrix} \cos(m_i - 1)\theta & -\sin(m_i - 1)\theta \\ \sin(m_i - 1)\theta & \cos(m_i - 1)\theta \end{pmatrix} \begin{pmatrix} z_1^i \\ z_2^i \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

where  $z = re^{i\theta}$  and  $Z = (z_1^i, z_2^i)$  so in  $U_i$  the vector  $Z$  can be written in the form  $Z = (1/m_i r^{m_i})(\cos(m_i - 1)\theta, -\sin(m_i - 1)\theta)$  which shows that the degree of  $Z$  at  $p_i$  is  $1 - m_i$ . Then  $\chi(\eta^2) \bmod 2$  is the number of  $p_i$ 's mod 2 such that  $m_i$  is even. □

LEMMA 5. Let  $\pi: E^{n+3} \rightarrow B^{n+1}$  be a smooth  $T^n$  vector bundle  $\nu$  such that

- (i)  $F(E^{n+3}, H) = E(E^{n+3}, H) = B^{n+1}$  for some  $H \simeq S^1$ ,
- (ii)  $E$  is an orientable manifold,
- (iii)  $T^n$  acts effectively on  $E^{n+1}$ .

Then  $w_2(E^{n+3}) = A(E^{n+3}) = D(B^{n+1})$ .

*Proof.* Since  $H$  acts effectively on the fibers of  $E$  by (iii),  $T_x^n = e$  for  $x \notin B$  and hence  $A(E^{n+3}) = D(B^{n+1})$ . We have  $T(E) = \pi^*(T(B) \oplus \nu) = \pi^*(T(B) \oplus \nu)$ . Since  $H \simeq S^1$  acts effectively on  $E$ ,  $E$  has a complex structure and is hence orientable and thus  $B$  is orientable and  $w_2(E) = \pi^*(w_2(B) + w_2(\nu))$ . Since  $T^n/H \simeq T^{n-1}$  acts freely on  $B$ ,  $B/T^{n-1}$  is an orientable 2-manifold and thus  $w_2(B/T^{n-1}) = 0$ . From the fibration  $p: B \rightarrow B/T^{n-1}$  we have that  $T(B) = p^*T(B|T^{n-1}) \oplus T_F$  where  $T_F$ , the tangent bundle along the fibers is a trivial bundle [1] and hence  $w_2(B) = p^*w_2(B|T^{n-1}) = 0$ . Thus  $w_2(E) = \pi^*w_2(\nu)$ . Now any  $z \in h_2(E; \mathbb{Z}_2)$  can be written as  $z = i_*z'$ ,  $z' \in H_2(B; \mathbb{Z}_2)$  and  $z'$  can be represented by a 2-manifold  $Q \subset B$ . Then

$$\langle A(E), z \rangle = \langle D(B), z \rangle = Q \cap B \bmod 2$$

and

$$\begin{aligned} \langle w_2(E), z \rangle &= \langle \pi^* w_2(\nu), i_* z' \rangle = \langle w_2(\nu), \pi_* i_* z' \rangle \\ &= \langle w_2(\nu), z' \rangle = \langle w_2(\nu|Q), Q \rangle = \langle \chi(\nu|Q), Q \rangle \\ &= \text{number of zeroes mod 2 of a generic section } s, \end{aligned}$$

where  $\chi$  denotes the Euler class. But this generic section can be used to put  $Q$  in general position with respect to  $B$  and the number of points in the intersection  $s(Q) \cap B$  is just the number of zeroes of the section.  $\square$

#### 4. Proof of Theorem.

DEFINITION. A  $T^n$  manifold  $M^{n+3}$  is said to be a nice  $T^n$  manifold if

- (i)  $M$  is closed and orientable,
- (ii) every isotropy subgroup is cyclic or isomorphic to  $S^1$ ,
- (iii) for  $H$  isomorphic to  $S^1$ , such that  $E(M, H) \neq \emptyset$ ,  $E(M, H)$  has codimension two,
- (iv) for every component  $F$  of  $E(M, H)$ ,  $H$  isomorphic to  $S^1$ ,  $w_2(\nu(F, M)) \neq 0$  where  $\nu(F, M)$  is the normal bundle of  $F$  in  $M$ .

LEMMA 6. *The theorem is true for nice  $T^n$  manifolds.*

*Proof.* Let  $E$  be a tubular neighborhood of  $\bigcup_{H \simeq S^1} E(M, H)$  with boundary  $\partial E$ . Consider the cohomology Mayer-Vietoris sequence for  $(M, \bar{E}, M - E)$  with  $Z_2$  coefficients

$$\begin{aligned} \rightarrow H^1(\bar{E}) \oplus H^1(M - E) \xrightarrow{i_1^* + i_2^*} H^1(\partial E) \xrightarrow{\delta} H^2(M) \\ \xrightarrow{j_1^* \oplus j_2^*} H^2(E) \oplus H^2(M - E). \end{aligned}$$

By Proposition 3,  $j^* w_2(M) = w_2(M - E) = A(M - E) = j_2^* A(M)$ , and by Lemma 5  $j_1^* w_2(M) = w_2(\bar{E}) = A(\bar{E}) = j_1^* A(M)$ , hence  $w_2(M) - A(M) \in \text{image } \delta$ . To prove the lemma it suffices to show  $i_1^*: H^1(E) \rightarrow H^1(\partial E)$  is onto and hence that  $\delta = 0$ . To prove that  $i_1^*$  is onto, we consider the Gysin sequence for one component  $\pi: E_1 \rightarrow F$  of the vector bundle  $E \rightarrow \bigcup_{H \simeq S^1} E(M, H)$

$$\rightarrow H^1(E_1) \xrightarrow{i_1^*} H^1(\partial E) \xrightarrow{\beta} H^0(E_1) \xrightarrow{\alpha} H^2(E_1)$$

and note that  $H^0(E_1) \simeq Z_2$  since  $E_1$  is connected and that  $\alpha(1) = \pi^* w_2(\nu(F, E_1))$  [2], and  $\pi^* w_2(\nu(F, E_1)) \neq 0$  by condition (iv). Hence  $\alpha$  is

1-1,  $\beta$  is zero and  $i_i^* | H^1(E_i)$  is onto. Taking direct sums over the components of  $E$  yields  $i_1^*: H^1(E) \rightarrow H^1(\partial E)$  is onto.  $\square$

To complete the theorem we need one final reduction lemma.

LEMMA 7. *It is sufficient to prove the theorem for nice actions of  $T^n$ .*

*Proof.* We shall show that for any smooth effective action of  $T^n$  on an orientable manifold  $M^{n+3}$  and any  $x \in H_2(M; Z_2)$  there exists a nice action of  $T^n$  on a manifold  $\tilde{M}^{n+3}$  and an  $\tilde{x} \in H_2(\tilde{M}, Z_2)$  such that  $\langle w_2(M), x \rangle = \langle w_2(\tilde{M}), \tilde{x} \rangle$  and  $\langle A(M), x \rangle = \langle A(\tilde{M}), \tilde{x} \rangle$ . Clearly that will prove Lemma 7.

We may assume, by Lemma 2, that the action of  $T^n$  on  $M^{n+3}$  satisfies the niceness conditions (i), (ii) and (iii). Let  $x$  be represented by a closed submanifold  $Q^2 \subset M^{n+3}$  which is transverse to  $F(M, H)$  for every isotropy subgroup  $H$  isomorphic to  $S^1$ . To construct the manifold  $\tilde{M}$  we will first choose an invariant neighborhood  $U$  of  $Q^2$  and then construct  $\tilde{M}$  so that  $U$  is contained in  $\tilde{M}$ . The class  $\tilde{x}$  will then be represented by  $Q \subset U \subset \tilde{M}$ . Clearly then

$$\langle A(M), i_* Q \rangle = \langle i^* A(M), Q \rangle = \langle A(U), Q \rangle = \langle j^* A(\tilde{M}), Q \rangle$$

and

$$\begin{aligned} \langle w_2(M), i_* Q \rangle &= \langle i^* w_2(M), Q \rangle = \langle w_2(U), Q \rangle \\ &= \langle j^* w_2(\tilde{M}), Q \rangle = \langle w_2(\tilde{M}), j_* Q \rangle \end{aligned}$$

where  $i: U \hookrightarrow M, j: U \hookrightarrow \tilde{M}$  are inclusions.

Since  $Q$  is transverse to  $\bigcup_{H \simeq S^1} F(M, H)$  the intersection is finite, say  $\{P_1, \dots, P_r\}$ . Also for any isotropy subgroup  $H \simeq S^1$ ,  $F(M, H)$  is an  $(n+1)$ -dimensional oriented closed manifold with free action of  $T^n/H \simeq T^{n-1}$ , hence  $F(M, H)/T^n$  is a 2-manifold and  $\pi: \bigcup_{H \simeq S^1} F(M, H) \rightarrow \bigcup_{H \simeq S^1} F(M, H)/T^n$  is a union of principal  $T^{n-1}$  bundles. Choose neighborhoods  $V_i$  of  $\pi(P_i)$  in  $F(M, H)/T^n$  with  $V_i$  diffeomorphic to the open disc  $\mathring{D}^2$ . Then  $\pi^{-1}(V_i) \simeq T^n/H \times V_i$ . Let  $C = \bigcup_{H \simeq S^1} F(M, H) - \bigcup_{i=1}^r \pi^{-1}(V_i)$ . Note that  $C$  is compact and invariant and hence  $M - C$  is an open invariant neighborhood of  $Q$ . Let  $U$  be an open invariant neighborhood of  $Q$ ,  $Q \subset U \subset M - C$  such that  $\bar{U} \subset M - C$  and  $\bar{U}$  is a manifold with boundary and let  $M_1^{n+3} = \text{double of } \bar{U} = \bar{U} \cup_{\partial \bar{U}} \bar{U} = \partial(\bar{U} \times I)$ . Note that  $M_1$  is not a nice  $T^n$  manifold, in fact, the normal bundle of every fixed point set  $F(M_1, H)$  in  $M_1$  is trivial. But  $M_1$  does



have the property that  $H^2(F; Z_2) \neq 0$  for any component  $F$  of a fixed point set  $F(M_1, H)$ ,  $H \simeq S^1$ . (One could have a  $T^2$  action on  $M^5$  for example, with  $F(M^5, H) \simeq S^3$ ). To prove the last statement we note that

$$\bigcup_{H \simeq S^1} F(\bar{U}, H) \subset \bigcup_{H \simeq S^1} F(M - C, H) = \bigcup_{i=1}^r T^n/H_i \times V_i$$

and hence

$$\bigcup_{H \simeq S^1} F(\bar{U}, H) = \bigcup_{i=1}^r T^n/H_i \times W_i$$

for  $W_i$  a compact 2-manifold with boundary  $\subset V_i \simeq \mathring{D}^2$ . Thus

$$\bigcup_{H \simeq S^1} F(M_1, H) = \bigcup_{i=1}^r T^n/H_i \times (W_i \cup_{\partial W_i} W_i)$$

and

$$H^2(W_i \cup_{\partial W_i} W_i; Z_2) \neq 0.$$

To construct the manifold  $\tilde{M}$  we shall modify the manifold  $M_1$  by twisting the normal bundle at each component of each fixed point set  $F(M_i, H)$ ,  $H \simeq S^1$  to satisfy condition (iv) of niceness. Let  $F_1, \dots, F_r$  be the components of  $\bigcup_{H \simeq S^1} F(M_1, H)$  and let  $\{p_1, \dots, p_r\} = Q \cap \bigcup_{H \simeq S^1} F(M_1, H)$  with  $p_i \in F_i$ ,  $F_i$  a component of  $F(M_i, H_i)$ . Choose  $q_i \in F_i$  so that  $p_i$  and  $q_i$  are on different orbits. Choose a tubular neighborhood  $C_i$  of the orbit  $q_i$ , diffeomorphic to  $T^n \times_{H_i} \mathbf{C} \times D^2$ .

Let

$$M_2 = \left( M_1 - \bigcup_{i=1}^r C_i \right) \cup_f \bigcup_{i=1}^r C_i''$$

where

$$C_i' = T^n \times_{H_i} \mathbf{C}(2) \times \mathring{D}^2, \quad C_i'' = T^n \times_{H_i} \mathbf{C}(1) \times D^2,$$

where  $\mathbf{C}(r)$  denotes the closed disk of radius  $r$  in  $\mathbf{C}$ , and  $f: T^n \times_{H_i} \mathbf{C}(1) \times \partial D^2 \rightarrow T^n \times_{H_i} \mathbf{C}(2) \times \partial D^2$  is given by  $f(t, z, \theta) = (t, e^{i\theta}z, \theta)$ . Note that  $F(M_2, H_i) = F(M_1, H_i)$ . See Figure 1.




FIGURE 1

Finally  $\tilde{M}$  is constructed by choosing  $M_3 \subset M_2$  so that  $M_3$  is a manifold with boundary and  $\cup_{H \simeq S^1} F(M_2, H) \subset \text{int } M_3$ ,  $U \subset M_3$  and setting  $\tilde{M} = \text{double of } M_3$ .  $\tilde{M}$  clearly satisfies conditions (i), (ii) and (iii) of niceness and  $Q \subset U \subset \tilde{M}$ . So we need only show that condition (iv) is satisfied. To that end we note that  $\nu(F(M_1, H_i))$  is a trivial bundle and therefore has a nonvanishing section  $X$ . Then, in  $M_2$ , we have a nonvanishing section of  $\nu(F(M_2, H_i), M_2) | F(M_2, H_i) - T^n/H_i \times \mathring{D}^2$ . Recall that  $F(M_2, H_i) = T^n/H \times (W_i \cup_{\partial W_i} W_i) = T^n/H \times B^2$  where  $B^2$  is a closed 2-manifold.

We shall show that  $\langle w_2(\nu(F(M_2, H_i), H_2), B^2) \neq 0$ . The section  $X$  restricts to a section of  $\nu(F(M_2, H_i), M_2) | B^2 - \mathring{D}^2$  and we want to look at this section in local coordinates on  $D^2$ . Then  $\tilde{X}: \partial D^2 \rightarrow C$  has the form  $\tilde{X}(\theta) = e^{i\theta} X(\theta)$  where  $\deg X = 0$ ,  $X: \partial D^2 \rightarrow C - \{0\}$ , since  $X$  extends to  $D^2 \subset M_1$  and therefore  $\deg \tilde{X}(\theta) = 1$  and hence extending  $\tilde{X}$  generically gives  $\langle \chi(\nu(F(M_2, h_i)), B^2) \neq 0$  where  $\chi$  is the Euler class.  $\square$

## REFERENCES

- [1] G. Bredon, *Introduction to Compact Transformation Groups*, Academic Press, New York and London, 1972.
- [2] J. W. Milnor and J. D. Stasheff, *Characteristic Classes*, Annals of Math. Studies No. 76, Princeton University Press, 1974.
- [3] D. Montgomery, *Orbits of highest dimension*, Seminar on Transformation Groups, Annals of Math. Studies No. 46, Chapter IX, Princeton University Press, 1960.
- [4] M. Seifert and W. Threlfall, *A Textbook of Topology*, Academic Press, 1980.
- [5] E. Stiefel, *Richtungsfelder and Fernparallelismus in Mannigfaltigkeiten*. Comm. Math. Helv., **8** (1936), 3-51.
- [6] R. Thom, *Quelques propriétés globales des variétés différentiables*, Comm. Math. Helv., **28** (1954), 17-86.

Received March 17, 1982. The first named author was partially supported by FAPESP-Fundação de Amparo À Pesquisa do Estado de São Paulo-BRASIL Proc 81 | 0803-3.

THE UNIVERSITY OF MICHIGAN  
ANN ARBOR, MI 48109

AND

UNIVERSIDADE FEDERAL DE SÃO CARLOS  
SÃO CARLOS-SP. BRASIL-13560

THE UNIVERSITY OF MICHIGAN  
ANN ARBOR, MI 48109

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California

Los Angeles, CA 90024

HUGO ROSSI

University of Utah

Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS

University of California

Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics

University of Southern California

Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON

Stanford University

Stanford, CA 94305

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

(1906–1982)

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P. O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS. A NON-PROFIT CORPORATION

Copyright © 1984 by Pacific Journal of Mathematics

<b>Kenneth F. Andersen and Wo-Sang Young</b> , On the reverse weak type inequality for the Hardy maximal function and the weighted classes $L(\log L)^k$ .....	257
<b>Richard Eugene Bedient</b> , Double branched covers and pretzel knots .....	265
<b>Harold Philip Boas</b> , Holomorphic reproducing kernels in Reinhardt domains .....	273
<b>Janey Antonio Daccach and Arthur Gabriel Wasserman</b> , Stiefel's theorem and toral actions .....	293
<b>Michael Fried</b> , The nonregular analogue of Tchebotarev's theorem .....	303
<b>Stanley Joseph Gurak</b> , Minimal polynomials for circular numbers .....	313
<b>Norimichi Hirano and Wataru Takahashi</b> , Nonlinear ergodic theorems for an amenable semigroup of nonexpansive mappings in a Banach space ...	333
<b>Jim Hoste</b> , Sewn-up $r$ -link exteriors .....	347
<b>Mohammad Ahmad Khan</b> , The existence of totally dense subgroups in LCA groups .....	383
<b>Mieczysław Kula, Murray Angus Marshall and Andrzej Śladek</b> , Direct limits of finite spaces of orderings .....	391
<b>Luis Montejano Peimbert</b> , Flat Hilbert cube manifold pairs .....	407
<b>Steven C. Pinault</b> , An a priori estimate in the calculus of variations .....	427
<b>McKenzie Y. K. Wang</b> , Some remarks on the calculation of Stiefel-Whitney classes and a paper of Wu-Yi Hsiang's .....	431
<b>Brian Donald Wick</b> , The calculation of an invariant for Tor .....	445
<b>Wolfgang Wollny</b> , Contributions to Hilbert's eighteenth problem .....	451