

Pacific Journal of Mathematics

**A GALOIS-CORRESPONDENCE FOR GENERAL LOCALLY
COMPACT GROUPS**

JOACHIM BOIDOL

A GALOIS-CORRESPONDENCE FOR GENERAL LOCALLY COMPACT GROUPS

JOACHIM BOIDOL

We give a characterization in terms of \hat{G} of those parts in the unitary dual of a locally compact group G , which correspond to closed normal subgroups of G . These are exactly the sets $S \subset \hat{G}$, which have the property that for all $\pi, \rho \in S$ the support of $\pi \otimes \bar{\rho}$ is contained in S and which are closed in a topology on \hat{G} , which is in general weaker than the standard topology on \hat{G} , and which we call the L^1 -hull-kernel-topology. As an easy consequence we obtain that for $*$ -regular groups G the mapping $N \rightarrow N^\perp = \{\pi \in \hat{G} | \pi|_N = 1|_{\mathcal{N}_N}\}$ is a bijection from the set of closed normal subgroups of G onto the set of closed subsets $S \subset \hat{G}$ with the property that $\pi \otimes \bar{\rho}$ has support in S for all $\pi, \rho \in S$. This generalizes and unifies results of Pontryagin, Helgason and Hauenschild, with a considerably simplified proof. Furthermore we prove that $*$ -regular groups have the weak Frobenius property (TP 1), i.e. 1_G is weakly contained in $\pi \otimes \bar{\pi}$ for all unitary representations π of G , generalizing a result of E. Kaniuth.

Let G be a locally compact group with unitary dual \hat{G} and let \mathcal{N}_G denote the set of closed normal subgroups of G . To every $N \in \mathcal{N}_G$ corresponds a canonical subset of \hat{G} , namely the annihilator $N^\perp = \{\pi \in \hat{G} | \pi|_N = 1|_{\mathcal{N}_N}\}$ of N . By the Gelfand-Raikov theorem $N \rightarrow N^\perp$ is an injective mapping from \mathcal{N}_G into the subsets of \hat{G} and it is an important problem in harmonic analysis to describe the image of this mapping in terms of \hat{G} .

DEFINITION. A nontrivial subset S of \hat{G} is called a *subdual* of \hat{G} , if for all $\pi, \rho \in S$ the tensor product $\pi \otimes \bar{\rho}$ of π and the conjugate $\bar{\rho}$ of ρ has support in S . We denote by \mathcal{S}_G the set of closed subduals of \hat{G} .

It is clear that $N \rightarrow N^\perp$ is an injective mapping from \mathcal{N}_G into \mathcal{S}_G . Let $[H]$ be the class of locally compact groups G , for which $N \rightarrow N^\perp$ is a surjection onto \mathcal{S}_G .

As a well known consequence of the duality theorem of Pontryagin one obtains that all *abelian locally compact groups* belong to $[H]$ (see for example [6], Chap. II, §1.7). S. Helgason proved in [8], Theorem 1, that all *compact groups* belong to $[H]$. It was then W. Hauenschild, who generalized and unified these results in [7], and proved that all *Moore groups*,

i.e. all locally compact groups G , which have only finite dimensional irreducible unitary representations, belong to the class $[H]$.

On the other hand the support \hat{G}_r of the left regular representation λ_G of a locally compact group G is clearly a closed subdual of \hat{G} . If $\hat{G}_r = N^\perp$ for some $N \in \mathcal{N}_G$, then $N = \{e\}$ and $\hat{G}_r = \hat{G}$. Therefore every group G , which belongs to $[H]$, has to be amenable.

We recall that the (standard) topology on \hat{G} is induced by the Jacobson topology on the primitive ideal space $\text{Prim}(G)$ of the group C^* -algebra $C^*(G)$ of G via the mapping $\pi \rightarrow \ker_{C^*(G)} \pi$. Let $\text{Prim}_* L^1(G)$ denote the space of kernels in $L^1(G)$ of topologically irreducible $*$ -representations of $L^1(G)$ in Hilbert spaces. $\text{Prim}_* L^1(G)$ is also a topological space with the Jacobson topology and the mapping $\pi \rightarrow \ker_{L^1(G)} \pi$ defines a second topology on \hat{G} , which we call the L^1 -hull-kernel-topology. This topology is weaker than the standard one and in general both topologies are different. Both topologies coincide if and only if the canonical continuous and surjective mapping $\Psi: \text{Prim}(G) \rightarrow \text{Prim}_* L^1(G)$, given by $\Psi(I) = I \cap L^1(G)$, is a homeomorphism, i.e. if G is $*$ -regular.

DEFINITION. Let $\mathcal{S}_G^* \subset \mathcal{S}_G$ be the set of subduals of \hat{G} , which are closed in the L^1 -hull-kernel-topology.

The main result of our paper will be that \mathcal{S}_G^* is the exact image of the mapping $N \rightarrow N^\perp$ for general locally compact groups. The results of Helgason and Hauenschild will be an easy consequence. But first we need the following

PROPOSITION. For every unitary representation of G in a Hilbert space \mathcal{H}_π we have $\ker_{L^1(G)} \pi \otimes \bar{\pi} \subset \ker_{L^1(G)} 1_G$.

Proof. Let $\overline{\mathcal{H}_\pi}$ be the adjoint space of \mathcal{H}_π and denote by $\bar{\eta}$ the vector $\eta \in \mathcal{H}_\pi$ considered as element of $\overline{\mathcal{H}_\pi}$. Then $\bar{\pi}$ is the representation π considered as a representation acting in $\overline{\mathcal{H}_\pi}$. We fix a unit vector $\xi \in \mathcal{H}_\pi$ and an orthonormal basis $\{\xi_i\}_{i \in I}$ of \mathcal{H}_π . Then for all $x \in G$ we have $\langle \bar{\pi}(x)\bar{\xi}, \bar{\xi}_i \rangle = \langle \overline{\pi(x)\xi}, \bar{\xi}_i \rangle$ and we obtain for all $x \in G$

$$1 = \langle \xi, \xi \rangle = \langle \pi(x)\xi, \pi(x)\xi \rangle = \sum_{i \in I} \langle \pi(x)\xi, \xi_i \rangle \langle \bar{\pi}(x)\bar{\xi}, \bar{\xi}_i \rangle.$$

Let \mathcal{F} denote the family of all finite sums of the functions

$$\langle \pi(x)\xi, \xi_i \rangle \langle \bar{\pi}(x)\bar{\xi}, \bar{\xi}_i \rangle,$$

which are matrix-coefficients of $\pi \otimes \bar{\pi}$. If $\varphi \in \mathcal{F}$ then φ is continuous and $0 \leq \varphi \leq 1$. Furthermore $1 = \sup_{\varphi \in \mathcal{F}} \varphi$.

Assume now that $f \in \text{kern}_{L^1(G)} \pi \otimes \bar{\pi}$. Then $\int_G f(x)\varphi(x) dx = 0$ for all $\varphi \in \mathcal{F}$. Given $\varepsilon > 0$ choose a compact set $\mathcal{X} \subset G$ such that $\int_{F \setminus \mathcal{X}} |f(x)| dx \leq \varepsilon/2$. By Dini there exists a sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ in \mathcal{F} (depending on \mathcal{X}), such that $1 = \lim_{n \rightarrow \infty} \varphi_n$ uniformly on \mathcal{X} . Then

$$\begin{aligned} \left| \int_G f(x) dx \right| &= \lim_{n \rightarrow \infty} \left| \int_{\mathcal{X}} f(x)\varphi_n(x) dx + \int_{G \setminus \mathcal{X}} f(x) dx \right| \\ &= \lim_{n \rightarrow \infty} \left| \int_{G \setminus \mathcal{X}} f(x) dx - \int_{G \setminus \mathcal{X}} f(x)\varphi_n(x) dx \right| \\ &\leq 2 \int_{G \setminus \mathcal{X}} |f(x)| dx \leq \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ was arbitrary, we obtain $\int_G f(x) dx = 0$.

The following corollary generalizes a result of E. Kaniuth (see [9], Lemma 1):

COROLLARY 1. *Let G be a $*$ -regular locally compact group. Then for every unitary representation π of G $\pi \otimes \bar{\pi}$ weakly contains the trivial representation, i.e. every $*$ -regular group has the property (TP 1) of [9].*

Proof. For $*$ -regular groups $\text{kern}_{L^1(G)} \pi * \bar{\pi} \subset \text{kern}_{L^1(G)} 1_G$ implies that 1_G is weakly contained in $\pi \otimes \bar{\pi}$.

REMARK. Corollary 1 shows that a quite big class of amenable groups has the weak Frobenius property (TP 1). This supports the conjecture that all amenable groups have the property (TP 1).

THEOREM. *For every locally compact group G the mapping $N \rightarrow N^\perp$ is a bijection from \mathcal{N}_G onto \mathcal{S}_G^* .*

Proof. As we remarked above, the mapping $N \rightarrow N^\perp$ is an injection from \mathcal{N}_G into \mathcal{S}_G . If $N \in \mathcal{N}_G$, then N^\perp corresponds to the set of topological irreducible $*$ -representations of $L^1(G)$, which are trivial on the kernel of the canonical homomorphisms from $L^1(G)$ onto $L^1(G/N)$. Therefore $N^\perp \in \mathcal{S}_G^*$, and we only have to prove that every set $\mathcal{S} \in \mathcal{S}_G^*$ is of the form N^\perp for some $N \in \mathcal{G}_G$.

First observe that by the proposition every $S \in \mathcal{S}_G^*$ has the following properties:

- (i) S contains 1_G and $\pi \in S$ implies $\bar{\pi} \in S$.
- (ii) for all $\pi, \rho \in S$ the support of $\pi * \rho$ is in S .

Furthermore since $S^\perp = \{x \in G \mid \pi(x) = 1|_{\mathcal{X}_\pi} \text{ for all } \pi \in S\}$ is a closed normal subgroup of G , we can consider S as a subdual of $(G/S^\perp)^\wedge$, which separates the points of G/S^\perp and is closed in the L^1 -hull-kernel-topology.

It is therefore sufficient to prove that a set $S \in \mathcal{S}_G^*$, which separates the points of G , is equal to \hat{G} .

Let S be such a set and let \mathcal{P} be the set of unitary representations of G , which have support in S . Since S is closed in \hat{G} , S and \mathcal{P} are weakly equivalent sets of representations of G . Since S has properties (i) and (ii) above, \mathcal{P} contains the trivial representation and is closed under the tensor product and under conjugation. It follows by a Stone-Weierstraß argument (see [1], Theorem) that \mathcal{P} is L^1 -separating, i.e. if $f \in L^1(G)$ and $\pi(f) = 0$ for all $\pi \in \mathcal{P}$, then $f = 0$. But then also S is L^1 -separating, i.e. its kernel in $L^1(G)$ is the trivial ideal $\{0\}$. Since S is closed in the L^1 -hull-kernel-topology, we obtain $S = \hat{G}$.

COROLLARY 2. *A locally compact group belongs to the class $[H]$ if and only if $\mathcal{S}_G = \mathcal{S}_G^*$. Especially every $*$ -regular locally compact group belongs to $[H]$ and every locally compact group in $[H]$ is amenable.*

REMARK. Let G be a locally compact group, such that all quotients G/N are C^* -unique, i.e. $L^1(G/N)$ has a unique C^* -norm (see [5]). The same arguments as in the proof of the theorem give that G belongs to $[H]$. We do not know whether this class of groups is really bigger than the class of $*$ -regular groups.

The following is known about $*$ -regular groups:

- (A) *Every $*$ -regular group is amenable (see [2]).*
- (B) *All groups G with polynomially growing Haar measure are $*$ -regular (see [2]).*
- (C) *All semidirect product $G = H \rtimes N$ with abelian H and N are $*$ -regular (see [4]).*
- (D) *A connected group G is $*$ -regular if and only if all $I \in \text{Prim}(G)$ are polynomially induced (see [3]).*

It follows from the classification of Moore groups given by C. C. Moore in [10], that all *Moore groups* have polynomial growth and so are $*$ -regular by (B). Therefore the result of W. Hauenschild is an immediate consequence of the Corollary 2 and (B). It should be noted that the proofs of the results of Pontryagin, Helgason and Hauenschild depend explicitly or implicitly on the fact that the groups under consideration are $*$ -regular. Besides this they make use of central theorems as the Pontryagin duality theorem, the Peter-Weyl theorem or structure theorems for Moore groups, which are specific for these classes of groups.

Recently E. Kaniuth proved by quite different methods that a big class of amenable groups, including the *almost connected amenable groups*, belong to $[H]$. (Cf. E. Kaniuth, *Weak containment and tensor products of group representations*. II, Math. Ann., **270** (1985), 1–15.) There seems to be some hope that the class $[H]$ coincides with the class of amenable groups.

REFERENCES

- [1] B. A. Barnes, *A note on separating families of representations*, Proc. Amer. Math. Soc., **87** (1983), 95–98.
- [2] J. Boidol, H. Leptin, J. Schürmann and D. Vahle, *Räume primitiver Ideale von Gruppenalgebren*, Math. Ann., **236** (1978), 1–13.
- [3] J. Boidol, *Connected groups with polynomially induced dual*, J. reine angew. Math., **331** (1982), 32–46.
- [4] _____, **-regularity of some classes of solvable groups*, Math. Ann. **261** (1982), 477–481.
- [5] _____, *Group algebras with a unique C^* -norm*, Functional Anal., **56** (1984), 220–232.
- [6] N. Bourbaki, *Théories Spectrales*, Paris: Hermann 1967.
- [7] W. Haenschild, *Subhypergroups and normal subgroups*, Math. Ann., **256** (1981), 1–18.
- [8] S. Helgason, *Lacunary Fourier series on noncommutative groups*, Proc. Amer. Math. Soc., **9** (1958), 782–790.
- [9] E. Kaniuth, *Weak containment and tensor products of group representations*, Math. Z., **180** (1982), 107–117.
- [10] C. C. Moore, *Groups with finite dimensional irreducible representations*, Trans. Amer. Math. Soc., **166** (1972), 401–410.

Received March 30, 1984 and in revised form August 28, 1984.

UNIVERSITÄT BIELEFELD
UNIVERSITÄTSSTRAÙE
4800 BIELEFELD 1
W. GERMANY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN (Managing Editor)
University of California
Los Angeles, CA 90024

CHARLES R. DEPRIMA
California Institute of Technology
Pasadena, CA 91125

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

H. SAMELSON
Stanford University
Stanford, CA 94305

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906-1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1985 by Pacific Journal of Mathematics

Philip Marshall Anselone and Mike Treuden, Regular operator approximation theory	257
Giuseppe Baccella, Semiprime \aleph-QF3 rings	269
Earl Robert Berkson and Thomas Alastair Gillespie, The generalized M. Riesz theorem and transference	279
Joachim Boidol, A Galois-correspondence for general locally compact groups	289
Joseph Eugene D'Atri, Josef Dorfmeister and Yan Da Zhao, The isotropy representation for homogeneous Siegel domains	295
C. Debiève, On Banach spaces having a Radon-Nikodým dual	327
Michael Aaron Freedman, Existence of strong solutions to singular nonlinear evolution equations	331
Francisco Jose Freniche, Grothendieck locally convex spaces of continuous vector valued functions	345
Hans-Peter Künzi and Peter Fletcher, Extension properties induced by complete quasi-uniformities	357
Takaši Kusano, Charles Andrew Swanson and Hiroyuki Usami, Pairs of positive solutions of quasilinear elliptic equations in exterior domains ...	385
Angel Rafael Larotonda and Ignacio Zalduendo, Spectral sets as Banach manifolds	401
J. Martínez-Maurica and C. Pérez García, A new approach to the Kreĭn-Milman theorem	417
Christian Pommerenke, On the boundary continuity of conformal maps	423
M. V. Subba Rao, Some Rogers-Ramanujan type partition theorems	431
Stephen Edwin Wilson, Bicontactual regular maps	437
Jaap C. S. P. van der Woude, Characterizations of (H)PI extensions	453
Kichoon Yang, Deformation of submanifolds of real projective space	469
Subhashis Nag, Errata: "On the holomorphy of maps from a complex to a real manifold"	493