

# Pacific Journal of Mathematics

**PAIRS OF POSITIVE SOLUTIONS OF QUASILINEAR  
ELLIPTIC EQUATIONS IN EXTERIOR DOMAINS**

TAKAÛI KUSANO, CHARLES ANDREW SWANSON AND HIROYUKI USAMI

## PAIRS OF POSITIVE SOLUTIONS OF QUASILINEAR ELLIPTIC EQUATIONS IN EXTERIOR DOMAINS

TAKAŠI KUSANO, CHARLES A. SWANSON AND HIROYUKI USAMI

**Our main objective is to prove the existence of infinitely many pairs  $(u_1, u_2)$  of positive solutions of quasilinear elliptic differential equations**

$$(1.1) \quad \Delta u - q(|x|)u = f(x, u, \nabla u), \quad x \in \Omega_\alpha,$$

**throughout exterior domains  $\Omega_\alpha \subset \mathbf{R}^N$ ,  $N \geq 2$ , of the type**

$$\Omega_\alpha = \{x \in \mathbf{R}^N : |x| > \alpha\}, \quad \alpha > 0,$$

**where  $x = (x_1, \dots, x_N)$ ,  $\nabla u = (\partial u / \partial x_1, \dots, \partial u / \partial x_N)$ , and  $\Delta = \nabla \cdot \nabla$ . Each pair has the property that  $u_1(x)/u_2(x)$  has uniform limit zero in  $\Omega_\alpha$  as  $|x| \rightarrow \infty$ . In particular, if  $q(t) \equiv 0$  and  $N \geq 3$ ,  $u_1(x)$  has limit 0 as  $|x| \rightarrow \infty$ , and  $u_2(x)$  is bounded above and below by positive constants in  $\Omega_\alpha$ .**

**1.** The function  $q$  in (1.1) is assumed to be nonnegative and locally Hölder continuous in  $\mathbf{R}_+ = [0, \infty)$ , and  $f: \Omega_\alpha \times \mathbf{R}_+ \times \mathbf{R}^N \rightarrow \mathbf{R}$  is locally Hölder continuous in  $\Omega_\alpha \times \mathbf{R}_+ \times \mathbf{R}^N$  and satisfies a Nagumo condition. Detailed hypotheses are listed in §3.

Specific asymptotic estimates for the growth (decay) of the solutions  $u_1(x)$ ,  $u_2(x)$  as  $|x| \rightarrow \infty$  follow easily from our construction. In particular sufficient conditions are given for the quasilinear equation

$$\Delta u - |x|^{2r}u = \phi(x)u^\gamma + \psi(x)|\nabla u|^\beta, \quad x \in \Omega_\alpha,$$

for constants  $r \geq 0$ ,  $\gamma \geq 0$ ,  $0 \leq \beta \leq 2$ , to have positive solutions  $u_1(x)$ ,  $u_2(x)$  in  $\Omega_\alpha$  such that  $u_i(x)$  is bounded above and below by positive constant multiples of

$$|x|^{-\lambda} \exp\left[(-1)^i |x|^{r+1} / (r+1)\right], \quad x \in \Omega_\alpha, \quad i = 1, 2,$$

where  $\lambda = (N + r - 1)/2$ .

By a similar method we also prove the existence of infinitely many positive solutions of the boundary value problem

$$(1.2) \quad \begin{aligned} \Delta u - q(|x|)u &= f(x, u, \nabla u), & x \in \Omega_\alpha, \\ u|_{\partial\Omega_\alpha} &= 0 \end{aligned}$$

under the same hypotheses as for (1.1).

The sharpness of our results is indicated by known oscillation criteria for semilinear elliptic equations [6, 10]. For example, in the case  $\psi(x) = 0$  in equation (3.26), our conditions (3.27)–(3.30) are all known to be necessary and sufficient conditions for the existence of a positive solution of (3.26), of the type described in Corollaries 3.8–3.11, in some exterior domain [11].

Our procedure will be to construct solutions of (1.1) or (1.2) which are squeezed between subsolutions and supersolutions. The latter are obtained as spherically symmetric solutions of elliptic equations with  $f(x, u, \nabla u)$  in (1.1) replaced by radial majorants. We therefore begin by proving global existence theorems for quasilinear ordinary differential equations (2.1) below. These results have independent interest, and in fact are of quite general nature. The procedure will be to find solutions of integro-differential equations as fixed points of associated operators from closed convex subsets  $\mathcal{Y}$  of  $C^1[t_0, \infty)$  into subsets of  $\mathcal{Y}$  with compact closure.

In the case of *semilinear* equations  $\Delta u = f(x, u)$ , in which  $q(t) \equiv 0$ , global existence theorems for boundary value problems in exterior domains have been obtained by Kawano and Naito [2], Kenig and Ni [3], and Noussair and Swanson [8]. Existence of positive bounded solutions in the entire space  $\mathbf{R}^N$ ,  $N \geq 3$ , have been proved by Kawano [1], Kusano and Oharu [4], and Ni [5].

**2. Positive solutions of quasilinear ordinary differential equations.**

Existence theorems will be proved for quasilinear ordinary differential equations of the type

$$(2.1) \quad Ly = h(t, y, y'), \quad t \geq t_0 > 0,$$

where  $L$  is the linear differential operator defined by

$$(2.2) \quad Lz = \frac{1}{p_2(t)} \frac{d}{dt} \left[ \frac{1}{p_1(t)} \frac{d}{dt} \left( \frac{z}{p_0(t)} \right) \right], \quad z \in C^2[t_0, \infty).$$

**ASSUMPTIONS.**

(A<sub>1</sub>) Each  $p_i(t)$  is positive in  $[t_0, \infty)$ ,  $p_i \in C^{2-i}[t_0, \infty)$ ,  $i = 0, 1, 2$ , and  $\lim_{t \rightarrow \infty} P(t) = +\infty$ , where

$$P(t) = \int_{t_0}^t p_1(s) ds.$$

(A<sub>2</sub>)  $h: [t_0, \infty) \times \mathbf{R}_+ \times \mathbf{R} \rightarrow \mathbf{R}$ ,  $\mathbf{R}_+ = [0, \infty)$ , is continuous and satisfies

$$|h(t, y, z)| \leq H(t, |y|, |z|)$$

for all  $t \in [t_0, \infty)$ ,  $y \in \mathbf{R}_+$ ,  $z \in \mathbf{R}$ , where  $H(t, u, v)$  is continuous in  $[t_0, \infty) \times \mathbf{R}_+ \times \mathbf{R}_+$ , nondecreasing in  $u$  for each  $t, v$ , and nondecreasing in  $v$  for each  $t, u$ .

The notation below will be used throughout:

$$(2.3) \quad p(t) = p_0(t)p_1(t)/P(t) + |p'_0(t)|.$$

For  $t > t_0$ ,  $(A_1)$ ,  $(A_2)$  and (2.2) show that the linear equation  $Lz = 0$  has linearly independent positive solutions

$$z_1(t) = p_0(t), \quad z_2(t) = p_0(t)P(t),$$

which are *asymptotically ordered*, i.e.,  $\lim_{t \rightarrow \infty} z_1(t)/z_2(t) = 0$ . The objective of this section is to establish sufficient conditions for equation (2.1) to have solutions  $y_1(t)$  and  $y_2(t)$  on the entire interval  $[t_0, \infty)$  with the same asymptotic behavior as  $z_1(t)$  and  $z_2(t)$ , respectively, as  $t \rightarrow \infty$ . We also prove the existence of a positive solution  $y(t)$  of (2.1) in  $(t_0, \infty)$  satisfying the boundary condition  $y(t_0) = 0$ .

It is well known [12] that every nonoscillatory linear differential operator  $L$  of second order in  $(0, \infty)$  can be written in the factorized form (2.2), where

$$(2.4) \quad p_0 = z_1, \quad p_1 = (z_2/z_1)', \quad p_2 = 1/p_0 p_1$$

for linearly independent, asymptotically ordered solutions  $z_1(t)$  and  $z_2(t)$  of  $Lz = 0$ , provided  $t_0$  is large enough so that these solutions are positive in  $[t_0, \infty)$ . The functions (2.4) are unique up to multiplicative constants.

The following hypotheses are used in the theorem below:

$$(2.5) \quad \int_{t_0}^{\infty} p_2(t)P(t)H(t, ap_0(t), bp(t)) dt < \infty;$$

$$(2.6) \quad \int_{t_0}^{\infty} p_2(t)H(t, ap_0(t)P(t), bp(t)P(t)) dt < \infty$$

for some positive constants  $a$  and  $b$ .

**THEOREM 2.1.** *In addition to  $(A_1)$ ,  $(A_2)$  suppose that  $\lambda^{-1}H(t, \lambda u, \lambda v)$  is a nondecreasing function of  $\lambda \in (0, \infty)$  and*

$$(2.7) \quad \lim_{\lambda \rightarrow 0^+} \lambda^{-1}H(t, \lambda u, \lambda v) = 0$$

for each fixed  $(t, u, v) \in [t_0, \infty) \times \mathbf{R}_+ \times \mathbf{R}_+$ .

(i) *Condition (2.5) implies that equation (2.1) has infinitely many positive (negative) solutions  $y(t)$  in  $[t_0, \infty)$  such that  $\lim_{t \rightarrow \infty} y(t)/z_1(t)$  exists and is positive (negative, respectively).*

(ii) Condition (2.6) implies that equation (2.1) has infinitely many positive (negative) solutions  $y(t)$  in  $(t_0, \infty)$  such that  $\lim_{t \rightarrow \infty} y(t)/z_2(t)$  exists and is positive (negative, respectively).

*Proof of (i).* Let  $m = \min\{a, b\}$ . The nondecreasing hypothesis on  $\lambda^{-1}H(t, \lambda u, \lambda v)$  shows that

$$(2.8) \quad \lambda^{-1}p_2(t)P(t)H(t, \lambda p_0(t), \lambda p(t)) \leq m^{-1}p_2(t)P(t)H(t, mp_0(t), mp(t))$$

for  $0 < \lambda \leq m$  and for all  $t \in [t_0, \infty)$ . By  $(A_2)$  and (2.5), the right member of (2.8) is integrable in  $(t_0, \infty)$ , and by (2.7) the left member has limit zero as  $\lambda \rightarrow 0+$  at every  $t \in [t_0, \infty)$ . The dominated convergence theorem then implies that

$$\lim_{\lambda \rightarrow 0+} \int_{t_0}^{\infty} \lambda^{-1}p_2(t)P(t)H(t, \lambda p_0(t), \lambda p(t)) dt = 0,$$

and therefore a sufficiently small constant  $k > 0$  exists such that

$$(2.9) \quad \int_{t_0}^{\infty} p_2(t)P(t)H(t, 3kp_0(t), 3kp(t)) dt \leq k.$$

With this choice of  $k$  we consider

$$(2.10) \quad \mathcal{Y} = \{y \in C^1[t_0, \infty) : kp_0(t) \leq y(t) \leq 3kp_0(t), |y'(t)| \leq 3kp(t), t \geq t_0\},$$

where  $C^1[t_0, \infty)$  denotes the locally convex space of all continuously differentiable functions in  $[t_0, \infty)$  with the topology of uniform convergence of functions and their first derivatives on compact subintervals of  $[t_0, \infty)$ . It is clear that  $\mathcal{Y}$  is a closed convex subset of  $C^1[t_0, \infty)$ . Define the integro-differential operator  $M$  by

$$(2.11) \quad (My)(t) = p_0(t) \left[ 2k + \int_t^{\infty} \left( \int_t^s p_1(\sigma) d\sigma \right) p_2(s)h(s, y(s), y'(s)) ds \right], \quad t \geq t_0.$$

If  $y \in \mathcal{Y}$ , (2.9)–(2.11) show that

$$\left| \frac{(My)(t)}{p_0(t)} - 2k \right| \leq \int_{t_0}^{\infty} p_2(s)P(s)H(s, 3kp_0(s), 3kp(s)) ds \leq k$$

and

$$\begin{aligned} |(My)'(t)| &\leq |p'_0(t)| \left[ 2k + \int_{t_0}^\infty p_2(s)P(s)H(s, 3kp_0(s), 3kp(s)) ds \right] \\ &\quad + p_0(t)p_1(t) \int_t^\infty p_2(s)H(s, 3kp_0(s), 3kp(s)) ds \\ &\leq 3k|p'_0(t)| + \frac{p_0(t)p_1(t)}{P(t)} \int_t^\infty p_2(s)P(s)H(s, 3kp_0(s), 3kp(s)) ds \\ &\leq 3k|p'_0(t)| + kp_0(t)p_1(t)/P(t) < 3kp(t). \end{aligned}$$

Therefore  $M$  maps  $\mathcal{Y}$  into  $\mathcal{Y}$ . The continuity of  $M$  follows from (2.9) and the dominated convergence theorem. The fact that  $M\mathcal{Y}$  has compact closure can be easily checked via Ascoli's theorem. It then follows from the Schauder-Tychonoff fixed point theorem that  $M$  has a fixed point  $y \in \mathcal{Y}$ . Differentiation of the equation  $y = My$  twice shows that  $y(t)$  is a positive solution of (2.1) in  $[t_0, \infty)$ , and clearly  $\lim_{t \rightarrow \infty} y(t)/p_0(t) = 2k$  by (2.11). Since infinitely many distinct choices of  $k$  are possible in (2.9), equation (2.1) has infinitely many positive solutions  $y(t)$  such that  $\lim_{t \rightarrow \infty} y(t)/p_0(t)$  exists and is positive. A slight modification establishes the analogous statement for negative solutions.

*Proof of (ii).* It follows as in part (i) that

$$\lim_{\lambda \rightarrow 0^+} \int_{t_0}^\infty \lambda^{-1} p_2(t)H(t, \lambda p_0(t)P(t), \lambda p(t)P(t)) dt = 0,$$

from which a sufficiently small constant  $k > 0$  exists such that

$$(2.12) \quad \int_{t_0}^\infty p_2(t)H(t, 3kp_0(t)P(t), 3kp(t)P(t)) dt \leq k.$$

With  $k$  as in (2.12), we define analogues  $\tilde{\mathcal{Y}}$  and  $\tilde{M}$  of (2.10) and (2.11), respectively, by

$$(2.13) \quad \tilde{\mathcal{Y}} = \{ y \in C^1[t_0, \infty) : kp_0(t)P(t) \leq y(t) \leq 3kp_0(t)P(t), \\ |y'(t)| \leq 3kp(t)P(t), t \geq t_0 \},$$

$$(2.14) \quad (\tilde{M}y)(t) \\ = p_0(t) \left[ 2kP(t) - \int_{t_0}^t p_1(s) \int_s^\infty p_2(\sigma)h(\sigma, y(\sigma), y'(\sigma)) d\sigma ds \right], \\ t \geq t_0.$$

If  $y \in \tilde{\mathcal{Y}}$ , one finds from (2.12) and (2.14) upon interchanging the order of integration that

$$\begin{aligned} & \left| \frac{(\tilde{M}y)(t)}{p_0(t)} - 2kP(t) \right| \\ & \leq P(t) \int_{t_0}^{\infty} p_2(\sigma) H(\sigma, 3kp_0(\sigma)P(\sigma), 3kp(\sigma)P(\sigma)) d\sigma \\ & \leq kP(t), \end{aligned}$$

and

$$\begin{aligned} & |(\tilde{M}y)'(t)| \\ & \leq |p'_0(t)| \left[ 2kP(t) + P(t) \int_{t_0}^{\infty} p_2(\sigma) H(\sigma, 3kp_0(\sigma)P(\sigma), 3kp(\sigma)P(\sigma)) d\sigma \right] \\ & \quad + p_0(t)p_1(t) \left[ 2k + \int_{t_0}^{\infty} p_2(\sigma) H(\sigma, 3kp_0(\sigma)P(\sigma), 3kp(\sigma)P(\sigma)) d\sigma \right] \\ & \leq 3k|p'_0(t)|P(t) + 3kp_0(t)p_1(t) = 3kp(t)P(t), \quad t \geq t_0. \end{aligned}$$

This shows that  $\tilde{M}$  maps  $\tilde{\mathcal{Y}}$  into  $\tilde{\mathcal{Y}}$ . The remainder of the proof closely parallels that of part (i) and will be deleted.

**THEOREM 2.2.** *In addition to  $(A_1)$  and  $(A_2)$  suppose that  $\mu^{-1}H(t, \mu u, \mu v)$  is a nonincreasing function of  $\mu \in (0, \infty)$  and*

$$(2.15) \quad \lim_{\mu \rightarrow \infty} \mu^{-1}H(t, \mu u, \mu v) = 0$$

for each fixed  $(t, u, v) \in [t_0, \infty) \times \mathbf{R}_+ \times \mathbf{R}_+$ . Then both conclusions (i) and (ii) of Theorem 2.1 hold verbatim et literatim.

*Proof.* Let  $C = \max\{a, b\}$ . Then, in analogy with (2.8),

$$\mu^{-1}p_2(t)P(t)H(t, \mu p_0(t), \mu p(t)) \leq C^{-1}p_2(t)P(t)H(t, Cp_0(t), Cp(t))$$

for all  $\mu \geq C$  and  $t \in [t_0, \infty)$ . It follows from  $(A_2)$ , (2.5) and (2.15) exactly as in Theorem 2.1 that

$$\lim_{\mu \rightarrow \infty} \int_{t_0}^{\infty} \mu^{-1}p_2(t)P(t)H(t, \mu p_0(t), \mu p(t)) dt = 0,$$

and hence there exists a sufficiently large positive constant  $k$  such that (2.9) is satisfied. Proceeding exactly as in the proof of part (i) of Theorem 2.1, we obtain a positive solution  $y(t)$  of (2.1) in  $[t_0, \infty)$  with  $\lim_{t \rightarrow \infty} y(t)/p_0(t) = 2k > 0$  as a fixed point of the operator  $M$  in the set  $\mathcal{Y}$ .

To prove part (ii), it suffices to observe that

$$\lim_{\mu \rightarrow \infty} \mu^{-1} \int_{t_0}^{\infty} p_2(t) H(t, \mu p_0(t) P(t), \mu p(t) P(t)) dt = 0.$$

This implies that (2.12) holds for sufficiently *large* positive constants  $k$ , and the proof is completed as in Theorem 2.1.

The proof of Theorem 2.1 (ii) also establishes the existence of infinitely many positive solutions of the quasilinear boundary value problem

$$(2.16) \quad \begin{aligned} Ly &= h(t, y, y'), & t \geq t_0, \\ y(t_0) &= 0. \end{aligned}$$

**THEOREM 2.3.** *Suppose that  $(A_1)$  and  $(A_2)$  hold. Let  $H$  be as in either Theorem 2.1 or Theorem 2.2. Then the condition (2.6) is sufficient for the boundary value problem (2.16) to have infinitely many positive solutions in  $(t_0, \infty)$ .*

In fact, for infinitely many admissible constants  $k$  in (2.6), Theorem 2.1 (ii) or Theorem 2.2 (ii) shows that equation (2.1) has a solution  $y \in \tilde{\mathcal{Y}}$  satisfying  $y(t) = (\tilde{M}y)(t)$  for  $t \geq t_0$ . Since  $P(t_0) = 0$  by  $(A_1)$ ,  $y(t)$  is a positive solution of (2.16) in  $(t_0, \infty)$  by (2.13) and (2.14).

**3. Pairs of positive solutions of quasilinear elliptic equations.** The radial component of the linear part of equation (1.1) (i.e.,  $f$  replaced by 0) is

$$(3.1) \quad Lz = t^{1-N} \frac{d}{dt} \left( t^{N-1} \frac{dz}{dt} \right) - q(t)z = 0, \quad t \geq \alpha.$$

Since (3.1) is nonoscillatory at  $\infty$ , there exist linearly independent positive solutions  $z_1(t)$  and  $z_2(t)$  of (3.1) on some fixed interval  $[t_0, \infty)$ , relabelled as  $[\alpha, \infty)$ , such that  $\lim_{t \rightarrow \infty} z_1(t)/z_2(t) = 0$ . Then the operator  $L$  has the factorized form (2.2) in  $[\alpha, \infty)$ , where  $p_0, p_1$  and  $p_2$  are defined by (2.4).

By applying the results of §2, we intend to find sufficient conditions for equation (1.1) to have positive solutions  $u_1(x)$  and  $u_2(x)$  in the entire domain  $\Omega_\alpha$  with the same asymptotic behavior at  $\infty$  as the functions  $z_1(|x|)$  and  $z_2(|x|)$ , respectively. The standing hypotheses on equation (1.1) are listed below.

**ASSUMPTIONS.**

$(B_1)$   $q(t)$  is nonnegative and locally Hölder continuous in  $\mathbf{R}_+$ , and  $f(x, u, p)$  is locally Hölder continuous in  $\Omega_\alpha \times \mathbf{R}_+ \times \mathbf{R}^N$ .

(B<sub>2</sub>) For any bounded subdomain  $G$  of  $\Omega_\alpha$  and any constant  $C > 0$ , there corresponds a constant  $\rho(G, C)$  such that

$$|f(x, u, p)| \leq \rho(G, C)(1 + |p|^2)$$

for all  $x \in G, 0 \leq u \leq C$ , and  $p \in \mathbf{R}^N$  (Nagumo condition).

(B<sub>3</sub>) There exists a continuous function  $F: \mathbf{R}_+^3 \rightarrow \mathbf{R}_+$  such that  $F(t, u, v)$  is nondecreasing in  $u$  and  $v$  and

$$|f(x, u, p)| \leq F(|x|, |u|, |p|), \quad x \in \Omega_\alpha, u \in \mathbf{R}_+, p \in \mathbf{R}^N.$$

(B<sub>4</sub>) For each fixed  $(t, u, v) \in [\alpha, \infty) \times \mathbf{R}_+ \times \mathbf{R}_+$ ,  $\lambda^{-1}F(t, \lambda u, \lambda v)$  is a nondecreasing function of  $\lambda \in (0, \infty)$  with  $\lim_{\lambda \rightarrow 0^+} \lambda^{-1}F(t, \lambda u, \lambda v) = 0$  (superlinearity).

A companion theorem will also be proved with (B<sub>4</sub>) replaced by the sublinear hypothesis (B<sub>4</sub><sup>\*</sup>) below:

(B<sub>4</sub><sup>\*</sup>) For each fixed  $(t, u, v) \in [\alpha, \infty) \times \mathbf{R}_+ \times \mathbf{R}_+$ ,  $\mu^{-1}F(t, \mu u, \mu v)$  is a nonincreasing function of  $\mu \in (0, \infty)$  with  $\lim_{\mu \rightarrow \infty} \mu^{-1}F(t, \mu u, \mu v) = 0$ .

The existence theorems below are proved under one of the following hypotheses:

$$(3.2) \quad \int_\alpha^\infty p_2(t) \frac{z_2(t)}{z_1(t)} F(t, az_1(t), bp(t)) dt < \infty,$$

$$(3.3) \quad \int_\alpha^\infty p_2(t) F\left(t, az_2(t), \frac{bp(t)z_2(t)}{z_1(t)}\right) dt < \infty$$

for some positive constants  $a$  and  $b$ , where  $z_1(t)$  and  $z_2(t)$  are positive solutions of (3.1) as described above, and

$$p_2(t) = \left[ z_1(t) \left( \frac{z_2}{z_1} \right)'(t) \right]^{-1},$$

$$p(t) = z_1(t) \frac{d}{dt} \log \left[ \frac{z_2(t)}{z_1(t)} \right] + |z_1'(t)|.$$

These forms of  $p_2(t)$  and  $p(t)$  follow from (2.3) and (2.4).

**THEOREM 3.1.** *If (B<sub>1</sub>)–(B<sub>3</sub>) and either (B<sub>4</sub>) or (B<sub>4</sub><sup>\*</sup>) hold, then (3.2) is a sufficient condition for equation (1.1) to have infinitely many positive solutions  $u(x)$  such that  $u(x)/z_1(|x|)$  is bounded above and below by positive constants in  $\Omega_\alpha$ .*

**THEOREM 3.2.** *If (B<sub>1</sub>)–(B<sub>3</sub>) and either (B<sub>4</sub>) or (B<sub>4</sub><sup>\*</sup>) hold, then (3.3) is sufficient for equation (1.1) to have infinitely many positive solutions  $u(x)$  such that  $u(x)/z_2(|x|)$  is bounded above and below by positive constants in  $\Omega_\alpha$ .*

**THEOREM 3.3.** *If  $(B_1)$ – $(B_3)$  and either  $(B_4)$  or  $(B_4^*)$  hold, then (3.3) is sufficient for the boundary value problem (1.2) to have infinitely many positive solutions in  $\Omega_\alpha$ .*

*Proof of Theorem 3.1.* Consider the ordinary differential equations

$$(3.4) \quad Ly = F(t, y, |y'|), \quad t \geq \alpha,$$

$$(3.5) \quad LY = -F(t, Y, |Y'|), \quad t \geq \alpha,$$

where  $L$  is the linear operator given by (3.1). Since (3.4) and (3.5) have the form (2.1), Hypotheses  $(B_1)$ ,  $(B_3)$ , and  $(B_4)$  (respectively,  $(B_4^*)$ ) show that Theorem 2.1 (respectively, Theorem 2.2) is applicable. By (2.4),  $p_1 = (z_2/z_1)'$  and  $P = z_2/z_1 + \text{constant}$ . Then condition (2.5) reduces to (3.2), and it follows from Theorem 2.1 (i) or Theorem 2.2 (i) that equations (3.4) and (3.5) have positive solutions  $y(t)$  and  $Y(t)$ , respectively, in the interval  $[\alpha, \infty)$  such that

$$\lim_{t \rightarrow \infty} \frac{y(t)}{z_1(t)} = 2k, \quad \lim_{t \rightarrow \infty} \frac{Y(t)}{z_1(t)} = 2K$$

for all sufficiently small positive constants  $k$  and  $K$  (or all sufficiently large  $k$  and  $K$  in the case of hypothesis  $(B_4^*)$ ). These constants can be chosen so that

$$(3.6) \quad 0 < y(t) < Y(t) \quad \text{for all } t \geq \alpha.$$

Indeed, in the case of hypothesis  $(B_4)$ , let  $Y(t)$  be determined first satisfying

$$K \leq \frac{Y(t)}{z_1(t)} \leq 3K, \quad t \geq \alpha,$$

as in the construction of Theorem 2.1 (i). Then choose  $0 < k \leq K/3$  and an associated  $y(t) = y(t, k)$  satisfying

$$(3.7) \quad k \leq \frac{y(t)}{z_1(t)} \leq 3k \leq K \leq \frac{Y(t)}{z_1(t)}, \quad t \geq \alpha.$$

The argument establishing (3.6) is similar in the case of hypothesis  $(B_4^*)$ .

We define functions  $w$  and  $v: \Omega_\alpha \rightarrow \mathbf{R}_+$  by

$$w(x) = y(|x|), \quad v(x) = Y(|x|), \quad x \in \Omega_\alpha.$$

Then (3.1), (3.4), and (3.5) show that  $w$  and  $v$  satisfy

$$\begin{aligned} \Delta w - q(|x|)w &= F(|x|, w, |\nabla w|), \quad x \in \Omega_\alpha, \\ \Delta v - q(|x|)v &= -F(|x|, v, |\nabla v|), \quad x \in \Omega_\alpha, \end{aligned}$$

respectively, and hence by  $(B_3)$  they satisfy the elliptic inequalities

$$(3.8) \quad \Delta w - q(|x|)w \geq f(x, w, \nabla w), \quad x \in \Omega_\alpha,$$

$$(3.9) \quad \Delta v - q(|x|)v \leq f(x, v, \nabla v), \quad x \in \Omega_\alpha.$$

Since  $0 < w(x) < v(w)$  throughout  $\Omega_\alpha$  by (3.6), it follows [7, p. 125] that there exists a solution  $u(x)$  of

$$\Delta u - q(|x|)u = f(x, u, \nabla u), \quad x \in \Omega_\alpha,$$

with locally Hölder continuous second derivatives in  $\Omega_\alpha$  such that  $w(x) \leq u(x) \leq v(x)$  for all  $x \in \Omega_\alpha$ . As indicated by (3.7), this implies that

$$(3.10) \quad k \leq \frac{y(|x|)}{z_1(|x|)} \leq \frac{u(x)}{z_1(|x|)} \leq \frac{Y(|x|)}{z_1(|x|)} \leq 9k$$

throughout  $\Omega_\alpha$ .

This completes the proof of Theorem 3.1. The proofs of Theorems 3.2 and 3.3 are virtually the same by application of Theorems 2.1 (ii) and 2.2 (ii).

EXAMPLE 3.4. If  $q(t)$  is identically zero, (1.1) reduces to

$$(3.11) \quad \Delta u = f(x, u, \nabla u), \quad x \in \Omega_\alpha.$$

In this case a fundamental set of asymptotically ordered positive solutions of  $Lz = 0$  in  $[\alpha, \infty)$  is

$$(3.12) \quad \begin{aligned} z_1(t) &= 1, & z_2(t) &= \log(t/\varepsilon) & \text{if } N = 2, \\ z_1(t) &= t^{2-N}, & z_2(t) &= 1 & \text{if } N \geq 3, \end{aligned}$$

for  $0 < \varepsilon < \alpha$ . Condition (3.2) is equivalent to

$$(3.13) \quad \begin{aligned} \int_\alpha^\infty t \log t F\left(t, a, \frac{b}{t \log t}\right) dt &< \infty & \text{if } N = 2, \\ \int_\alpha^\infty t^{N-1} F(t, at^{2-N}, bt^{1-N}) dt &< \infty & \text{if } N \geq 3, \end{aligned}$$

and condition (3.3) is equivalent to

$$(3.14) \quad \begin{aligned} \int_\alpha^\infty t F\left(t, a \log t, \frac{b}{t}\right) dt &< \infty & \text{if } N = 2, \\ \int_\alpha^\infty t F\left(t, a, \frac{b}{t}\right) dt &< \infty & \text{if } N \geq 3. \end{aligned}$$

Then (3.13) implies that equation (3.11) has infinitely many positive solutions  $u(x)$  in  $\Omega_\alpha$  for any  $\alpha > 0$  such that  $u(x)/z_1(|x|)$  is bounded above and below by positive constants in  $\Omega_\alpha$ ; and in particular  $u(x) \rightarrow 0$

as  $|x| \rightarrow \infty$  in dimensions  $N \geq 3$ . Similarly (3.14) implies the conclusions of Theorems 3.2 and 3.3, and in particular implies the existence of infinitely many solutions of (3.11) in  $\Omega_\alpha$  which are uniformly bounded above and below by positive constants if  $N \geq 3, \alpha > 0$ .

EXAMPLE 3.5. Consider the equation

$$(3.15) \quad \Delta u - \rho^2|x|^{2r}u = \phi(x)u^\gamma + \psi(x)|\nabla u|^\beta, \quad x \in \Omega_\alpha,$$

where  $\beta, \gamma, \rho, r$  are constants,  $\gamma \geq 0, 0 \leq \beta \leq 2, \rho > 0, 2r$  is a nonnegative integer, and  $\phi, \psi$  are locally Hölder continuous functions in  $\Omega_\alpha$  for some  $\alpha > 0$ . Equation (3.15) is an example of (1.1) in which  $q(t) = \rho^2t^{2r}$  and  $f(x, u, p) = \phi(x)u^\gamma + \psi(x)|p|^\beta$ . Define

$$\phi^*(t) = \max_{|x|=t} |\phi(x)|, \quad \psi^*(t) = \max_{|x|=t} |\psi(x)|,$$

$$(3.16) \quad F(t, u, v) = \phi^*(t)u^\gamma + \psi^*(t)v^\beta, \quad t \geq \alpha, u \in \mathbf{R}_+, v \in \mathbf{R}_+.$$

Then  $|f(x, u, p)| \leq F(|x|, |u|, |p|)$  for all  $x \in \Omega_\alpha, u \in \mathbf{R}_+, p \in \mathbf{R}^N$ , and hypothesis  $(B_3)$  holds for this choice of  $F$ .

In the case of (3.15), equation (3.1) becomes

$$(3.17) \quad t^{1-N} \frac{d}{dt} \left( t^{N-1} \frac{dz}{dt} \right) - \rho^2 t^{2r} z = 0, \quad t \geq \alpha,$$

which possesses linearly independent solutions with the asymptotic behavior [9, p. 285]

$$z_1(t) \sim t^{-\lambda} \exp \left[ -\frac{\rho t^{r+1}}{r+1} \right], \quad z_2(t) \sim t^{-\lambda} \exp \left[ \frac{\rho t^{r+1}}{r+1} \right]$$

as  $t \rightarrow \infty$ , where  $\lambda = (N + r - 1)/2$ . Evidently these solutions are asymptotically ordered at  $\infty$ , and we assume that  $\alpha$  has been selected large enough so that they are positive in  $[\alpha, \infty)$ . Using (2.4) one easily verifies the asymptotic forms

$$p_1(t) \sim 2\rho t^r \exp \left[ \frac{2\rho t^{r+1}}{r+1} \right] \quad \text{as } t \rightarrow \infty;$$

$$p_2(t) \sim \frac{t^{\lambda-r}}{2\rho} \exp \left[ -\frac{\rho t^{r+1}}{r+1} \right] \quad \text{as } t \rightarrow \infty;$$

$$\frac{p_2(t)z_2(t)}{z_1(t)} \sim \frac{t^{\lambda-r}}{2\rho} \exp \left[ \frac{\rho t^{r+1}}{r+1} \right] \quad \text{as } t \rightarrow \infty.$$

We note from (3.16) that hypothesis  $(B_4)$  holds if

$$(3.18) \quad \gamma > 1 \quad \text{and} \quad \beta > 1$$

and hypothesis  $(B_4^*)$  holds if

$$(3.19) \quad 0 \leq \gamma < 1 \quad \text{and} \quad 0 \leq \beta < 1.$$

In the present example, condition (3.2) is satisfied if both

$$(3.20) \quad \int_{\alpha}^{\infty} t^{\lambda(1-\gamma)-r} \exp\left[\frac{\rho(1-\gamma)t^{r+1}}{r+1}\right] \phi^*(t) dt < \infty$$

and

$$(3.21) \quad \int_{\alpha}^{\infty} t^{(\lambda-r)(1-\beta)} \exp\left[\frac{\rho(1-\beta)t^{r+1}}{r+1}\right] \psi^*(t) dt < \infty,$$

and condition (3.3) holds if both

$$(3.22) \quad \int_{\alpha}^{\infty} t^{\lambda(1-\gamma)-r} \exp\left[\frac{\rho(\gamma-1)t^{r+1}}{r+1}\right] \phi^*(t) dt < \infty$$

and

$$(3.23) \quad \int_{\alpha}^{\infty} t^{(\lambda-r)(1-\beta)} \exp\left[\frac{\rho(\beta-1)t^{r+1}}{r+1}\right] \psi^*(t) dt < \infty.$$

We conclude from Theorem 3.1 under hypotheses (3.20), (3.21), and either (3.18) or (3.19) that equation (3.15) has infinitely many bounded positive solutions  $u(x)$  in  $\Omega_{\alpha}$  such that

$$c_1|x|^{-\lambda} \exp\left[-\frac{\rho|x|^{r+1}}{r+1}\right] \leq u(x) \leq c_2|x|^{-\lambda} \exp\left[-\frac{\rho|x|^{r+1}}{r+1}\right],$$

$x \in \Omega_{\alpha}$ , for some positive constants  $c_1$  and  $c_2$ . Furthermore, Theorem 3.2 shows that (3.22), (3.23), and either (3.18) or (3.19) imply the existence of infinitely many unbounded positive solutions  $u(x)$  of (3.15) in  $\Omega_{\alpha}$  such that

$$c_3|x|^{-\lambda} \exp\left[\frac{\rho|x|^{r+1}}{r+1}\right] \leq u(x) \leq c_4|x|^{-\lambda} \exp\left[\frac{\rho|x|^{r+1}}{r+1}\right], \quad x \in \Omega_{\alpha},$$

for some positive constants  $c_3$  and  $c_4$ . Theorem 3.3 shows that the same conditions are sufficient for the existence of positive solutions of (3.15) in  $\Omega_{\alpha}$  satisfying the boundary condition  $u|_{\partial\Omega_{\alpha}} = 0$ .

Example 3.5 can be extended to mixed sublinear-superlinear equations (3.15), in which  $\gamma > 1$  and  $0 \leq \beta < 1$ , via the corollary below. More generally we consider equations (1.1) with the structure

$$(3.24) \quad \Delta u - q(|x|)u = f_1(x, u, \nabla u) - f_2(x, u, \nabla u), \quad x \in \Omega_{\alpha},$$

where each  $f_i$  satisfies hypotheses  $(B_1)$ ,  $(B_2)$ ,  $f_1$  satisfies  $(B_4)$ ,  $(f_2)$  satisfies  $(B_4^*)$ , but  $(B_3)$  is replaced by  $(B_3^*)$  below:

$(B_3^*)$  There exist continuous functions  $F_i: \mathbf{R}_+^3 \rightarrow \mathbf{R}_+$ ,  $i = 1, 2$ , such that each  $F_i(t, u, v)$  is nondecreasing in  $u$  and  $v$  and

$$(3.25) \quad \begin{aligned} 0 &\leq f_1(x, u, p) \leq F_1(|x|, u, |p|) \\ 0 &\leq -f_2(x, u, p) \leq F_2(|x|, u, |p|) \end{aligned}$$

for all  $x \in \Omega_{\alpha}$ ,  $u \in \mathbf{R}_+$ ,  $p \in \mathbf{R}^N$ .

**COROLLARY 3.6.** *Suppose that  $q$  and each  $f_i$  satisfies  $(B_1)$ ,  $(B_2)$ ,  $(B_3^*)$ , and (3.2), and that  $f_1, f_2$  satisfy  $(B_4)$ ,  $(B_4^*)$ , respectively, with  $F$  replaced by  $F_1, F_2$ , respectively. Then equation (3.24) has infinitely many positive solutions  $u(x)$  such that  $u(x)/z_1(|x|)$  is bounded above and below by positive constants throughout  $\Omega_\alpha$ .*

*Proof.* Consider the elliptic equations

$$\begin{aligned} \Delta w - q(|x|)w &= f_1(x, w, \nabla w) \\ \Delta v - q(|x|)v &= -f_2(x, v, \nabla v) \end{aligned}$$

in  $\Omega_\alpha$ . By Theorem 3.1, these equations have solutions  $v, w \in C_{loc}^{2+\lambda}(\Omega_\alpha)$  such that

$$\begin{aligned} k_1 z_1(|x|) &\leq w(x) \leq k_2 z_1(|x|), \quad x \in \Omega_\alpha, \\ k_3 z_1(|x|) &\leq v(x) \leq k_4 z_1(|x|), \quad x \in \Omega_\alpha, \end{aligned}$$

for all sufficiently small positive constants  $k_1, k_2 > k_1$  and sufficiently large positive constants  $k_3, k_4 > k_3$ . Since each  $f_i \geq 0$ ,  $w$  and  $v$  satisfy the elliptic differential inequalities

$$\begin{aligned} \Delta w - q(|x|)w &\geq f_1(x, w, \nabla w) - f_2(x, w, \nabla w), \\ \Delta v - q(|x|)v &\leq f_1(x, v, \nabla v) - f_2(x, v, \nabla v), \end{aligned}$$

respectively, in  $\Omega_\alpha$ . We can choose  $k_3$  large enough so that  $0 < w(x) \leq v(x)$  throughout  $\Omega_\alpha$ , and it follows [7, p. 125] as in Theorem 3.1 that (3.24) has a solution  $u \in C_{loc}^{2+\lambda}(\Omega_\alpha)$  satisfying  $w(x) \leq u(x) \leq v(x)$  throughout  $\Omega_\alpha$ .

**COROLLARY 3.7.** *Let  $q$  and each  $f_i$  be as in Corollary 3.6 except that (3.2) is replaced by (3.3). Then equation (3.24) has infinitely many positive solutions  $u(x)$  such that  $u(x)/z_2(|x|)$  is bounded above and below by positive constants throughout  $\Omega_\alpha$ .*

The proof, based on Theorem 3.2, is similar to that of Corollary 3.6.

These corollaries give new global existence theorems even for semilinear equations of the type

$$(3.26) \quad \Delta u = \phi(x)u^\gamma - \psi(x)u^\beta, \quad x \in \Omega_\alpha,$$

where  $\gamma > 1$ ,  $0 \leq \beta < 1$ , and  $\phi$  and  $\psi$  are nonnegative locally Hölder continuous functions in  $\Omega_\alpha$ ,  $\alpha > 0$ . Corollaries 3.6 and 3.7 and Example 3.4 imply the following additional corollaries.

COROLLARY 3.8. Equation (3.26) has a positive solution which is bounded and bounded away from zero in an arbitrary exterior domain  $\Omega \subset \mathbf{R}^2$  if both

$$(3.27) \quad \int_{\alpha}^{\infty} t \log t \phi^*(t) dt < \infty \quad \text{and} \quad \int_{\alpha}^{\infty} t \log t \psi^*(t) dt < \infty$$

for some  $\alpha > 0$ .

For example, we can choose

$$\alpha = \inf\{|x| : x \in \partial\Omega\}$$

to obtain  $\Omega \subset \Omega_{\alpha}$ , and apply the earlier results to  $\Omega_{\alpha}$ .

COROLLARY 3.9. Equation (3.26) has a positive solution which is bounded above and below by positive constant multiples of  $\log(|x|/\varepsilon)$  in an arbitrary domain  $\Omega_{\alpha} \subset \mathbf{R}^2$ ,  $0 < \varepsilon < \alpha$ , if both

$$(3.28) \quad \int_{\varepsilon}^{\infty} t(\log t)^{\gamma} \phi^*(t) dt < \infty \quad \text{and} \quad \int_{\varepsilon}^{\infty} t(\log t)^{\beta} \psi^*(t) dt < \infty.$$

COROLLARY 3.10. Equation (3.26) has a positive solution which is bounded and bounded away from zero in an arbitrary exterior domain  $\Omega \subset \mathbf{R}^N$ ,  $N \geq 3$ , if both

$$(3.29) \quad \int_{\alpha}^{\infty} t \phi^*(t) dt < \infty \quad \text{and} \quad \int_{\alpha}^{\infty} t \psi^*(t) dt < \infty.$$

COROLLARY 3.11. Equation (3.26) has a positive solution which is bounded above and below by positive constant multiples of  $|x|^{2-N}$  in an arbitrary exterior domain  $\Omega \subset \mathbf{R}^N$ ,  $N \geq 3$ , if both

$$(3.30) \quad \int_{\alpha}^{\infty} t^{\sigma} \phi^*(t) dt < \infty \quad \text{and} \quad \int_{\alpha}^{\infty} t^{\rho} \psi^*(t) dt < \infty,$$

where  $\sigma = (N - 1) - \gamma(N - 2)$ ,  $\rho = (N - 1) - \beta(N - 2)$ .

Theorems 3.1–3.3 also apply to quasilinear equations of the type

$$\Delta u - q(|x|)u = \phi(x)u^{\gamma} + u^{\beta} \sum_{i,j=1}^N \psi_{ij}(x) \frac{\partial u}{\partial x_i} \frac{\partial u}{\partial x_j},$$

where  $\phi$  and each  $\psi_{ij}$  are locally Hölder continuous in  $\Omega_{\alpha}$ , and the matrix  $(\psi_{ij}(x))$  is symmetric and positive semidefinite in  $\Omega_{\alpha}$ .

## REFERENCES

- [1] N. Kawano, *On bounded entire solutions of semilinear elliptic equations*, Hiroshima Math. J., **14** (1984), 125–158.
- [2] N. Kawano and M. Naito, *Positive solutions of semilinear second order elliptic equations in exterior domains*, Hiroshima Math. J., **12** (1982), 141–149.
- [3] C. E. Kenig and W-M. Ni, *An exterior Dirichlet problem with applications to some nonlinear equations arising in geometry*, Amer. J. Math., **106** (1984), 689–702.
- [4] Takaši Kusano and Shinnosuke Oharu, *Bounded entire solutions of second order semilinear elliptic equations with application to a parabolic initial value problem*, Indiana Univ. Math. J., **34** (1985), 85–95.
- [5] W-M. Ni, *On the elliptic equation  $\Delta u + K(x)u^{(N+2)/(N-2)} = 0$ , its generalizations, and applications in geometry*, Indiana Univ. Math. J., **31** (1982), 493–529.
- [6] E. S. Noussair and C. A. Swanson, *Positive solutions of semilinear Schrödinger equations in exterior domains*, Indiana Univ. Math. J., **28** (1979), 993–1003.
- [7] ———, *Positive solutions of quasilinear elliptic equations in exterior domains*, J. Math. Anal. Appl., **75** (1980), 121–133.
- [8] ———, *Global positive solutions of semilinear elliptic equations*, Canad. J. Math., **35** (1983), 839–861.
- [9] L. Sirovich, *Techniques of Asymptotic Analysis*, Springer-Verlag, New York-Heidelberg-Berlin, 1971.
- [10] C. A. Swanson, *Criteria for oscillatory sublinear Schrödinger equations*, Pacific J. Math., **104** (1983), 483–493.
- [11] ———, *Extremal positive solutions of semilinear Schrödinger equations*, Canad. Math. Bull., **26** (1983), 171–178.
- [12] W. F. Trench, *Canonical forms and principal systems for general disconjugate equations*, Trans. Amer. Math. Soc., **189** (1974), 319–327.

Received February 24, 1984. The first author was supported in part by Grant-in-Aid for Scientific Research (No. 58460004), Ministry of Education (Japan). The second author gratefully acknowledges support from NSERC (Canada) under Grant A-3105.

HIROSHIMA UNIVERSITY  
HIROSHIMA, JAPAN 730

AND

UNIVERSITY OF BRITISH COLUMBIA  
VANCOUVER, B. C.  
CANADA V6T 1Y4



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN (Managing Editor)  
University of California  
Los Angeles, CA 90024

CHARLES R. DEPRIMA  
California Institute of Technology  
Pasadena, CA 91125

R. FINN  
Stanford University  
Stanford, CA 94305

HERMANN FLASCHKA  
University of Arizona  
Tucson, AZ 85721

RAMESH A. GANGOLLI  
University of Washington  
Seattle, WA 98195

ROBION KIRBY  
University of California  
Berkeley, CA 94720

C. C. MOORE  
University of California  
Berkeley, CA 94720

H. SAMELSON  
Stanford University  
Stanford, CA 94305

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS    E. F. BECKENBACH    B. H. NEUMANN    F. WOLF    K. YOSHIDA  
(1906–1982)

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1985 by Pacific Journal of Mathematics

# Pacific Journal of Mathematics

Vol. 120, No. 2

October, 1985

<b>Philip Marshall Anselone and Mike Treuden</b> , Regular operator approximation theory .....	257
<b>Giuseppe Baccella</b> , Semiprime $\aleph$ -QF3 rings .....	269
<b>Earl Robert Berkson and Thomas Alastair Gillespie</b> , The generalized M. Riesz theorem and transference .....	279
<b>Joachim Boidol</b> , A Galois-correspondence for general locally compact groups .....	289
<b>Joseph Eugene D'Atri, Josef Dorfmeister and Yan Da Zhao</b> , The isotropy representation for homogeneous Siegel domains .....	295
<b>C. Debiève</b> , On Banach spaces having a Radon-Nikodým dual .....	327
<b>Michael Aaron Freedman</b> , Existence of strong solutions to singular nonlinear evolution equations .....	331
<b>Francisco Jose Freniche</b> , Grothendieck locally convex spaces of continuous vector valued functions .....	345
<b>Hans-Peter Künzi and Peter Fletcher</b> , Extension properties induced by complete quasi-uniformities .....	357
<b>Takaši Kusano, Charles Andrew Swanson and Hiroyuki Usami</b> , Pairs of positive solutions of quasilinear elliptic equations in exterior domains ...	385
<b>Angel Rafael Larotonda and Ignacio Zalduendo</b> , Spectral sets as Banach manifolds .....	401
<b>J. Martínez-Maurica and C. Pérez García</b> , A new approach to the Kreĭn-Milman theorem .....	417
<b>Christian Pommerenke</b> , On the boundary continuity of conformal maps ....	423
<b>M. V. Subba Rao</b> , Some Rogers-Ramanujan type partition theorems .....	431
<b>Stephen Edwin Wilson</b> , Bicontactual regular maps .....	437
<b>Jaap C. S. P. van der Woude</b> , Characterizations of (H)PI extensions .....	453
<b>Kichoon Yang</b> , Deformation of submanifolds of real projective space .....	469
<b>Subhashis Nag</b> , Errata: "On the holomorphy of maps from a complex to a real manifold" .....	493