

Pacific Journal of Mathematics

**GROUPS OF KNOTS IN HOMOLOGY 3-SPHERES THAT ARE
NOT CLASSICAL KNOT GROUPS**

MAGNHILD LIEN

GROUPS OF KNOTS IN HOMOLOGY 3-SPHERES THAT ARE NOT CLASSICAL KNOT GROUPS

MAGNHILD LIEN

In this paper we attempt to enlarge classical knot groups K by adding a root to a meridian of K . Thus if K is a classical knot group with a meridian μ , then the groups we study are of the form $G = K *_{\mu=t^q} \langle t \rangle$.

This group can always be realized as the group of a knotted 3-sphere in S^5 . By using explicit geometric constructions we also show that such a group G is a 2-knot group and the group of a knot in a homology 3-sphere. Finally, we show that G is not realizable by any knot in S^3 .

1. Introduction. In [R₁] J. Ratcliffe gave an example of a group Γ that is the group of a fibered knot in a homology 3-sphere which cannot be realized as the group of a classical knot. Let K be the group of the trefoil knot and let $\mu \in K$ represent a meridian. As seen in [R₁] Γ can be expressed as a free product with amalgamation $K *_{\mu=t^2} \langle t \rangle$.

In this paper we generalize the result in [R₁]. We study groups G obtained from classical knot groups K by forming an amalgamated free product of K with Z . More specifically if K is a classical knot group with meridian μ , then $G = K *_{\mu=t^q} \langle t \rangle$. Note that if $q = 1$ then $G = K$. Hence we assume $q > 1$. One natural question to ask about G is if G is the group of a knot in any dimension. We show that G is the group of a knotted 2-sphere in S^4 . Furthermore we show that G can be realized as the group of a knot in a homology 3-sphere. However, G is not a classical knot group.

I would like to thank the referee for his suggestion of how to construct the 2-sphere in §3. This made the third section a lot simpler than it was in the first version of this paper.

2. Preliminaries. In this paper we work in the smooth category. S^n and B^n denote the standard n -sphere and n -ball. If N and M are manifolds and $f: M \rightarrow N$ is a map then both of the induced homomorphisms $\pi_1(M) \rightarrow \pi_1(N)$ or $H_1(M) \rightarrow H_1(N)$ will be denoted by f_* . Homeomorphism between spaces and isomorphism between groups are denoted by \cong . An n -dimensional knot is the image of a smooth

embedding Σ^n of S^n into S^{n+2} or R^{n+2} . By the knot group we mean $\pi_1(S^{n+2} - \Sigma^n)$. For $n = 1$, we call these groups classical knot groups.

We define the deficiency of a group presentation with n generators and m relators to be the integer $n - m$. The following well known proposition is due to Kervaire [K].

PROPOSITION 2.1. *If a group K has a deficiency one presentation and $K/K' \cong Z$ then $H_2(K) = 0$.*

Consider an oriented knot Σ in S^3 . Remove an open neighbourhood N of Σ in S^3 to produce the knot manifold $X = S^3 - N$. The preferred meridian, longitude pair (μ, λ) of Σ are two nontrivial simple closed curves on $\text{Bd}(X)$ such that μ bounds a disk in N and λ is homologically trivial in X .

DEFINITION 2.2. A (p, q) -curve is a simple closed curve J on $\text{Bd}(X)$ that is homotopic to $\mu^p \lambda^q$ where p and q are relatively prime. We also call J a (p, q) -cable about Σ .

3. A 2-knot with group $K *_{\mu=t^q} \langle t \rangle$. Let K be a classical knot group with meridian μ . We construct a new group G by adding a q th root (via amalgamted free product) to the meridian μ of K , i.e., $G = K *_{\mu=t^q} \langle t \rangle$. The following proposition is easy to verify using Kervaire's characterization of high dimensional knot groups ([K], Theorem 1).

PROPOSITION 3.1. *The group G is a high dimensional knot group.*

Proof. Since K is a classical knot group it has a deficiency one presentation. We can thus obtain a deficiency one presentation of G from a presentation of K by adding one more generator (t) and one more relation $(\mu = t^q)$. Moreover since K is a knot group and hence satisfies the conditions of Kervaire's characterization it is straightforward to check that $G/G' \cong Z$ and that $G/\langle\langle t \rangle\rangle = 1$. By Proposition 2.1 we obtain $H_2(G) = 0$, and it follows that G can be realized as the group of a knotted 3-sphere in S^5 . □

Let Σ be a knot in S^3 with group K . We shall construct a knotted 2-sphere Σ^4 with $\pi_1(S^4 - \Sigma^2) = K *_{\mu=t^q} \langle t \rangle$. The equatorial cross-section of this 2-sphere will be a $(1, q)$ -cable about the composite knot $\Sigma\# - \Sigma^*$, where $-\Sigma^*$ is the mirror image of Σ with its orientation reversed.

DEFINITION 3.2. A knot Σ in S^3 is a slice knot if there exists a smooth disk D in B^4 such that $\text{Bd}(D) = \Sigma$.

THEOREM 3.3. *Let Σ be a knot in S^3 and let L be the $(1, q)$ -cable about $\Sigma\# - \Sigma^*$. Then L is a slice knot.*

Proof. For any knot Σ , the knot $\Sigma\# - \Sigma^*$ is a slice knot [F, M]. To construct a slice disk D in B^4 with $\text{Bd}(D) = \Sigma\# - \Sigma^*$, we do as follows. First note that $(S^3, \Sigma) = (B^3, \beta) \cup_{\partial} (B^3, B^1)$ where β is a knotted arc and (B^3, B^1) is a standard ball pair. Remove the ball pair (B^3, B^1) from (S^3, Σ) and cross (B^3, β) with the interval to obtain a disk $D = \beta \times I$ contained in B^4 . Then $\text{Bd}(D) = \Sigma\# - \Sigma^*$. Thus D is the desired disk. Let $N = D \times \text{int } B^2$ be an open neighbourhood of the slice disk and let $M = B^4 - N$. Then $D \times \text{Bd}(B^2)$ is in $\text{Bd}(M)$. We shall attach a 2-handle $B^2 \times B^2$ which contains a slice disk for the trivial $(1, q)$ -torus knot to M along $D \times \text{Bd}(B^2)$. If $B^2 \times B^2$ is attached along $S^1 \times B^2$ then let the torus knot be the $(1, q)$ -cable about the core of the solid torus $B^2 \times S^1$. Note that the attaching sphere $S^1 \times \{0\}$ represents the q th power of the meridian of the torus knot. Since the image of $\{*\} \times S^1$ under the attaching map is $\text{Bd}(D) = \Sigma\# - \Sigma^*$ it follows that the image of the boundary of the slice disk for the $(1, q)$ -torus knot is L . Thus there exists a disk in B^4 with boundary L . □

Since L is a slice knot, we can use L as the equatorial section of a knotted 2-sphere in S^4 by joining together smooth disks in B^4_+ and B^4_- bounded by L . We denote the 2-sphere obtained this way by $S(\Sigma, q)$.

THEOREM 3.4. *Let Σ be a knot in S^3 with group K and meridian μ , $q > 1$ and let $S(\Sigma, q)$ be the 2-knot described above. Then*

$$\pi_1(S^4 - S(\Sigma, q)) = K *_{\mu = t^q} \langle q \rangle.$$

Proof. If \tilde{D} is the slice disk for the $(1, q)$ -cable about $\Sigma\# - \Sigma^*$ it suffices to show that $\pi_1(B^4 - \tilde{D}) = K *_{\mu = t^q} \langle t \rangle$. Using the notation from the proof of Theorem 3.3 we have that

$$B^4 - \tilde{D} = M \cup_{S^1 \times B^2 - \text{Bd}(\tilde{D})} (B^2 \times B^2) - \tilde{D}.$$

Since M is homotopic to $B^4 - D$ which equals $(B^3 - \beta) \times I$ it follows that $\pi_1(M) = \pi_1(B^3 - \beta) = K$. Moreover, since $\text{Bd}(\tilde{D}_1)$ is the (trivial) $(1, q)$ -torus knot we get that the fundamental group of $B^2 \times B^2 - \tilde{D}$ is

infinite cyclic generated by the meridian of the torus knot. Thus the fundamental group of $B^4 - \tilde{D}$, is obtained from $\pi_1(M)$ by attaching a q th root of the original meridian μ , i.e. $\pi_1(B^4 - \tilde{D}) = K *_{\mu=t^q} \langle t \rangle$. \square

4. 3-manifolds that can realize the group $K *_{\mu=t^q} \langle t \rangle$. We now consider how close G is to being a classical knot group. As the following shows the Alexander polynomial $\Delta_G(t)$ for G is symmetric and it satisfies $\Delta_G(1) = \pm 1$.

THEOREM 4.1. *Let K be a classical knot group with Alexander polynomial $\Delta_K(t)$ and let $G = K *_{\mu=t^q} \langle t \rangle$. Then the Alexander polynomial $\Delta_G(t)$ for G satisfies $\Delta_G(t) = \Delta_K(t^q)$.*

Proof. Let $K = \langle x_0, x_1, \dots, x_n; R_1, \dots, R_n \rangle$ be a standard Wirtinger presentation for K and let $A = [\partial R_i / \partial x_j]$ be the $n \times (n + 1)$ Alexander matrix with respect to this presentation.

$$A \text{ is equivalent to } B = \begin{bmatrix} \frac{\partial R_1}{\partial X_0} & \cdots & \frac{\partial R_1}{\partial X_{n-1}} & 0 \\ & & & \vdots \\ \frac{\partial R_n}{\partial X_0} & \cdots & \frac{\partial R_n}{\partial X_{n-1}} & 0 \end{bmatrix}$$

which is obtained from A by adding the first n columns to the last column. The Alexander polynomial for K , $\Delta_K(t)$ is the generator of the principal ideal generated by the determinants of all the $n \times n$ submatrices of B . Thus $\Delta_K(t) =$ determinant of the $n \times n$ submatrix obtained from B by deleting the last column. The group G has a presentation $\langle t, x_0, \dots, x_n; R_1, \dots, R_n, x_0 t^{-q} \rangle$ and its Alexander matrix is

$$\begin{bmatrix} & & & & 0 \\ & & & & 0 \\ 1 & 0 & \cdots & 0 & k(t) \end{bmatrix}$$

where $k(t) = \partial(x_0 t^{-q}) / \partial t$. This matrix is equivalent to

$$C = \begin{bmatrix} \frac{\partial R_1}{\partial X_0}(t^q) & \cdots & \frac{\partial R_1}{\partial X_{n-1}}(t^q) & 0 & 0 \\ \frac{\partial R_n}{\partial X_0}(t^q) & & \frac{\partial R_n}{\partial X_{n-1}}(t^q) & 0 & 0 \\ 1 & 0 & \cdots & 0 & 1 & k(t) \end{bmatrix}$$

The ideal generated by all $(n + 1) \times (n + 1)$ minors of C is easily seen to be principal and its generator is $\Delta_K(t^q)$. Hence G has an Alexander polynomial $\Delta_G(t)$ and moreover $\Delta_G(t) = \Delta_K(t^q)$. \square

Since K is a classical knot group there is a knot Σ in S^3 with knot manifold X such that $\pi_1(X) = K$. Let (μ, λ) be the preferred meridian, longitude pair for $\text{Bd}(X)$, and let $A = \mu \times I$ be an annulus on $\text{Bd}(X)$. By T^3 we mean the standard solid torus $S^1 \times D^2$ in R^3 . Furthermore, let J be a $(1, q)$ -curve on the boundary of T^3 and let $B = J \times I$ be an annulus on $\text{Bd}(T^3)$. We construct a cabled 3-manifold M $[\mathbf{J}, \mathbf{M}]$ by glueing together X and T^3 along the two annuli A and B , i.e. $M = X \cup_{A=B} T^3$.

PROPOSITION 4.2. $\pi_1(M) = K *_{\mu=t^q} \langle t \rangle$.

Proof. Let $\pi_1(T^3) = \langle t \rangle$, $\pi_1(A) = \langle \mu \rangle$ and $\pi_1(B) = \langle j \rangle$. The image of μ in $\pi_1(X)$ under the homomorphism induced by the inclusion map $A \rightarrow M$ is μ , and the image of j in $\pi_1(T^3)$ under the homomorphism induced by the inclusion map $B \rightarrow T^3$ is t^q . Thus by the Van-Kampen Theorem we conclude that $\pi_1(M) = K *_{\mu=t^q} \langle t \rangle$. \square

The boundary of M is homeomorphic to $S^1 \times S^1$, and if (μ, λ) is a standard meridian, longitude pair for $\text{Bd}(X)$, then a basis for $\text{Bd}(M)$ is $\mu, \lambda t^{-1}$.

THEOREM 4.3. *If K is a classical knot group with meridian μ , then $G = K *_{\mu=t^q} \langle t \rangle$ is the group of a knot in a homology 3-sphere.*

Proof. We construct a 3-manifold M' by sewing a solid torus to M in such a way that a meridian of the solid torus is identified with the curve λt^{-1} . Then $\pi_1(M')$ is obtained from $\pi_1(M)$ by adding the relation $t = \lambda$. Since $\pi_1(M) = K *_{\mu=t^q} \langle t \rangle$ we get that $\pi_1(M') = K / \langle \langle \mu \lambda^{-q} \rangle \rangle$. It is now easy to see that $H_1^{\mu=t^q}(M') \cong 0$ and we conclude, using the usual Poincaré duality argument that M' is a homology 3-sphere.

In some recent work done by Culler, Gordon, Luecke, and Shalen, it is shown that if $(1, q)$ -surgery on a nontrivial knot yields a simply connected manifold then $|q| = 0$ or 1 [C, G, L, S]. If a knot Σ has group K and $m, l \in K$ is the meridian and the longitude, then the fundamental group of the surgery manifold $\Sigma(1, q)$ is $K / \langle \langle ml^q \rangle \rangle$. We thus have the following proposition.

PROPOSITION 4.3. *If K is a classical knot group and $q > 1$ then $K/\langle\langle ml^q \rangle\rangle \neq 1$.*

THEOREM 4.4. *Let K be a classical knot group with meridian μ and let $G = K *_{\mu=t^q} \langle t \rangle$, $q > 1$. Then G is not a classical knot group.*

Proof. Let $M = X \cup_A T^3$ be the cabled manifold we constructed in the paragraph preceding Proposition 4.2, then $\pi_1(M) \cong G$. Suppose there exists a knot L in S^3 with $\pi_1(S^3 - L) = G$. We shall eventually show that L must be a cable knot about some core L' and that the surgery manifold $L'(1, q)$ is simply connected which contradicts Proposition 4.3.

The knot manifold X is aspherical [P]. Moreover, M as the union of aspherical spaces sewn together along an incompressible subspace is aspherical [W₂]. Let N be the knot manifold for L . Since $\pi_1(M)$ is isomorphic to $\pi_1(N)$ and M and N are aspherical spaces, there exists a homotopy equivalence $f: N \rightarrow M$ that induces the group isomorphism.

The annulus A is bicollared in M . Furthermore, $\pi_1(A) \rightarrow \pi_1(M)$ is injective since $\pi_1(A) \rightarrow \pi_1(X)$ and $\pi_1(A) \rightarrow \pi_1(T^3)$ are injective; also $\pi_2(A) = \pi_2(M) = \pi_3(M) = \pi_2(M - A) = 0$ since M , X and T^3 are aspherical spaces. Hence $\ker(\pi_j(A) \rightarrow \pi_j(M)) = 0$, $j = 1, 2$. By Lemma 1.1 in [W₁] there exists a map g that is homotopic to f such that:

1. g is transverse with respect to A , i.e. there exists a neighbourhood $g^{-1}(A) \times I$ of $g^{-1}(A)$ so that $g(x, y) = (g(x), y)$ for every $x \in g^{-1}(A)$ and $y \in I$.
2. $g^{-1}(A)$ is an orientable compact 2-manifold and $g^{-1}(A) \cap \text{Bd}(M) = \text{Bd}(g^{-1}(A))$.
3. If F is a component of $g^{-1}(A)$ then $\ker(\pi_j(F) \rightarrow \pi_j(N)) = 0$, $j = 1, 2$.

Choose a map g that minimizes the number of components of $g^{-1}(A)$. We shall show that for such a map g , $g^{-1}(A)$ is just one annulus F and that F separates N into a solid torus V and a knot manifold Y having the same group as X . We use techniques similar to those used in [F, W] and [S] to prove the following assertions. (Some proofs are omitted since they are the same as proofs in [F, W] and [S].)

Claim 4.4.1. $g^{-1}(A)$ is nonempty.

Claim 4.4.2. Let F be a component of $g^{-1}(A)$. Then F is an essential annulus.

We thus have $g^{-1}(A) = F_1 \cup \dots \cup F_K$, $k \geq 1$. Since each component of $g^{-1}(A)$ is an essential annulus we get that the core L of $S^3 - \text{Int}(N)$ is either a composite knot or a cable knot ([W₃], Lemma 1.1).

Claim 4.4.3. L is a cable knot.

Since $g^{-1}(A) = F_1 \cup \dots \cup F_K$ and L is a (p', q') -cable about a knot L' we have for each i , $1 \leq i \leq k$, $N = Y_i \cup_{F_i} V_i$ where V_i is a solid torus and Y_i is a knot manifold. Moreover a boundary component of $\text{Bd}(F_i)$ is a (p', q') -curve on $S^3 - \text{Int}(Y_i)$.

Claim 4.4.4. $\pi_1(N)$ has trivial center.

Proof of 4.4.4. Since $\pi_1(N) = K *_{\mu=t^q} \langle t \rangle$, the center of $\pi_1(N) = C(K) \cap \langle \mu \rangle$ ([M, K, S], Cor. 4.5). If K is not a torus knot group then $C(K) = 1$ and consequently $\pi_1(N)$ has no center. On the other hand if K is a torus knot group then $C(K) = \langle \mu^{p^q \lambda} \rangle$, but $\langle \mu^{p^q \lambda} \rangle \cap \langle \mu \rangle = 1$ in K and consequently in $K *_{\mu=t^q} \langle t \rangle$. □

Claim 4.4.5. The annuli F_1, \dots, F_K are parallel in N .

Proof of 4.4.5. N is prime since it is a cable knot manifold. Moreover N is irreducible. Since $\pi_1(N)$ has trivial center, N cannot be a Siefert fiber space with decomposition surface a disk with 3 singular fibers as such a space has fundamental group with nontrivial center [J]. By Lemma 2.4 in [J, M] there exists a unique (up to ambient isotopy) essential annulus embedded in N . Hence each annulus F_i is parallel to F_1 . □

Claim 4.4.6. For each i , $1 \leq i \leq k$, $g|_{F_i}$ is homotopic to a homeomorphism.

Proof of 4.4.6. It suffices to show that $g_*: H_1(F_i) \rightarrow H_1(A)$ is an isomorphism. Since $H_1(F_i) \cong \pi_1(F_i)$ and $H_1(A) \cong \pi_1(A)$, this implies that $g_*: \pi_1(F_i) \rightarrow \pi_1(A)$ is an isomorphism. Hence $g|_{F_i}$ is a homotopy equivalence and is therefore homotopic to a homeomorphism.

Let f_i be a generator for $H_1(F_i)$ and let a be a generator for $H_1(A)$. Then $g_*(f_i) = a^r$. We wish to show that $r = \pm 1$. We use a homotopy inverse h of g ($h: M \rightarrow N$). As done earlier in the proof we can modify h such that $h^{-1}(F_i)$ is a collection of essential annuli, B_1, \dots, B_n . Since M is irreducible ([J], Lemma 3.1) and M is not a twisted I -bundle over the Klein bottle (if M is a twisted I -bundle over the Klein bottle then $\pi_1(M)$

abelianizes to $Z \oplus Z_2$) there exists an isotopy h_t ($0 \leq t \leq 1$) such that $h_1(\text{Bd}(B_i)) \cap \text{Bd}(A) \neq \emptyset$. Hence $\text{Bd}(B_i)$ is homologous to $\text{Bd}(A)$ in $H_1(M)$, i.e. the generator a for $H_1(A)$ is homologous to a generator b_i for $H_1(B_i)$. Furthermore, $h_*(b_i) = f_i^s$. Since h is a homotopy inverse of g we get $g_*(h_*(b_i)) = b_i$, i.e. $a^{sr} = b_i$. Since a is homologous to b_i in $H_1(M)$ we obtain $a^{sr} = a$ in $H_1(M)$. Hence $sr = 1$, and it follows that $r = \pm 1$. \square

Claim 4.4.7. $p'q' = q$.

Proof of 4.4.7. $z_{p'q'} \cong H_1(N)/H_1(F_i) \cong H_1(M)/H_1(A) \cong Z_q$. \square

Recall that $g^{-1}(A) = F_1 \cup \dots \cup F_k$.

Claim 4.4.8. k is odd.

Proof of 4.4.8. Since the annuli F_i , $1 \leq i \leq k$ are all parallel V_k contains a core v_1 of V_1 . Let α be a path in V_1 from F_1 to v_1 , then $\pi_1(Y_1)$ and $\alpha v_1 \alpha^{-1}$ generate $\pi_1(N)$. If k is even then g maps $\text{Int}(Y_1)$ and v_1 into the same component of $M - A$. If Y_1 and v_1 are both mapped into $X - A$, then $\pi_1(M)/\pi_1(X) \cong 1$ which contradicts the fact that $H_1(M)/H_1(X) \cong Z_q$. On the other hand Y_1 cannot be mapped into T^3 , because that would imply that $g_*: \pi_1(Y_1) \rightarrow \pi_1(T^3)$ is 1-1 which contradicts the fact that Y_1 is a knot manifold. \square

Claim 4.4.9. $k = 1$.

The proof of 4.4.9 is the same as the proof of Claim 7 in [S].

Since $k = 1$, we have that $N = Y \cup_F V$, where Y is the knot manifold for the knot L' , V is a solid torus, and a boundary component of $\text{Bd}(F)$ is a (p', q') -curve of the boundary of $S^3 - \text{Int}(Y)$. Moreover we have a homotopy equivalence $g: Y \cup_F V \rightarrow X \cup_A T^3$ such that $g|_F$ is a homeomorphism. We saw in the proof of Claim 4.4.8 that $g(Y)$ is not contained in T^3 . Hence $g(Y) \subseteq X$ and $g(V) \subseteq T^3$ and $g_*(\pi_1(Y)) \subseteq \pi_1(X)$ and $g_*(\pi_1(V)) \subseteq \pi_1(T^3)$. Let s be a generator for $\pi_1(V)$ and let m, l be a meridian, longitude pair for $\pi_1(Y)$, then $\pi_1(N) = \pi_1(Y) *_{\pi_1(F)} \pi_1(V) = \pi_1(Y) *_{m^{p'}l^{q'} = s^{q'}, \langle s \rangle} \langle s \rangle$. Recall that $\pi_1(M) = \pi_1(x) *_{\pi_1(A)} \pi_1(T^3)$ where $\pi_1(x) = K$ and $\pi_1(T^3) = \langle t \rangle$. Since g_* is an isomorphism and $g_*(\pi_1(F)) = \pi_1(A)$ we have the following $g_*(\pi_1(N)) = g_*(\pi_1(Y)) *_{\pi_1(A)} g_*(\pi_1(V)) = K *_{\pi_1(A)} \langle t \rangle$. By Proposition 2.5 in [B] we conclude that $g_*|_{\pi_1(Y)}: \pi_1(Y) \rightarrow K$ and $g_*|_{\pi_1(V)}: \pi_1(V) \rightarrow \pi_1(T^3)$ is an isomorphism. Therefore, since

$q = p'q'$ we have the following: $g_*((m^{p'l^{q'}})^{p'}) = g_*(s^{q'p'}) = g_*(s^q) = t^{\pm q} = \mu^{\pm 1}$. Hence $\pi_1(Y)/\langle\langle(m^{p'l^{q'}})^{p'}\rangle\rangle \cong K/\langle\langle\mu^{\pm 1}\rangle\rangle = 1$. Since $\pi_1(Y)/\langle\langle(m^{p'l^{q'}})^{p'}\rangle\rangle$ abelianizes to $Z_{(p')^2}$ this implies that $p' = \pm 1$ which in turn implies that $q' = \pm q$. Hence $\pi_1(Y)/\langle\langle ml^q\rangle\rangle = 1$ which contradicts Proposition 4.3. \square

REFERENCES

- [B] E. M. Brown, *Unknotting in $M^2 \times I$* , Trans. Amer. Math. Soc., **123** (1966), 480–505.
- [C, G, L, S] M. Culler, C. McA. Gordon, J. Luecke, and P. Shalen, *Dehn surgery on knots*, Bull. Amer. Math. Soc., **13**, no. 1 (1985), 43–45.
- [F, W] C. D. Feustel, and W. Whitten, *Groups and complements of knots*, Canad. J. Math., **30**, no. 6 (1978), 1284–1295.
- [F] R. H. Fox, *A quick trip through knot theory*, in Topology of 3-Manifolds and Related Topics, M. K. Fort, Ed., Prentice Hall (1966), 120–167.
- [F, M] R. H. Fox and J. W. Milnor, *Singularities of 2-spheres in 4-space and cobordism of knots*, Osaka J. Math., **3** (1966), 257–267.
- [J] W. Jaco, *Lectures on three-manifold topology*, Conference Board of the Mathematical Sciences by the Amer. Math. Soc. no. 43, (1977).
- [J, M] W. Jaco and R. Meyers, *An algebraic determination of closed orientable 3-manifolds*, Trans. Amer. Math. Soc., **253** (1979), 149–170.
- [K] M. Kervaire, *On higher dimensional knots*, in Differential and Combinatorial Topology, S. S. Cairns, Ed., Princeton University Press.
- [M, K, S] W. Magnus, A. Karrass and D. Solitar, *Combinatorial Group Theory*, (2nd revised ed.), Dover Pub. Inc., New York (1966).
- [P] C. D. Papakyriakopoulos, *On Dehn's Lemma and the asphericity of knots*, Proc. Nat. Acad. Sci., **43** (1957), 169–172.
- [R₁] J. G. Ratcliffe, *A fibered knot in a homology 3-sphere whose group is non-classical*, Contemporary Math., **20** (1983).
- [R₂] D. Rolfsen, *Knots and Links*, Publish or Perish Inc. (1976).
- [S] J. K. Simon, *An algebraic classification of knots in S^3* , Ann. of Math., (2) **97** (1973), 1–13.
- [W₁] F. Waldhausen, *Gruppen mit Zentrum und 3-dimensionale Mannigfaltigkeiten*, Topology, **6** (1967), 505–517.
- [W₂] J. H. C. Whitehead, *On the asphericity of regions in a 3-sphere*, Fund. Math., **32** (1939), 259–270.
- [W₃] W. Whitten, *Algebraic and geometric characterizations of knots*, Invent. Math., **26** (1974), 259–270.

Received June 1, 1986 and in revised form September 8, 1986.

COLLEGE OF CHARLESTON
CHARLESTON, SC 29424

Current address: Department of Mathematics
California State University, Northridge
Northridge, CA 91330

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

K. Adachi , Continuation of bounded holomorphic functions from certain subvarieties to weakly pseudoconvex domains	1
Erazm Jerzy Behr , Enveloping algebras of Lie superalgebras	9
Dong M. Chung , Scale-invariant measurability in abstract Wiener spaces	27
Peter Gerard Dodds and Bernardus de Pagter , Algebras of unbounded scalar-type spectral operators	41
Wu-Yi Hsiang and Hsueh-Ling Huynh , Generalized rotational hypersurfaces of constant mean curvature in the Euclidean spaces. II	75
Harvey Bayard Keynes and M. Sears , Time changes for \mathbf{R}^n flows and suspensions	97
Frances Kirwan, Ronnie Lee and Steven Howard Weintraub , Quotients of the complex ball by discrete groups	115
Magnhild Lien , Groups of knots in homology 3-spheres that are not classical knot groups	143
Juan Carlos Migliore , Liaison of a union of skew lines in \mathbf{P}^4	153
Jesper M. Møller , Spaces of sections of Eilenberg-Mac Lane fibrations	171
Daniel Ullman , A generalization of a theorem of Atkinson to noninvariant measures	187
Kohhei Yamaguchi , Operations which detect \mathcal{P}^1 in odd primary connective K -theory	195