

Pacific Journal of Mathematics

**A SEMILINEAR WAVE EQUATION WITH NONMONOTONE
NONLINEARITY**

ALFONSO CASTRO AND SUMALEE UNSURANGSIE

A SEMILINEAR WAVE EQUATION WITH NONMONOTONE NONLINEARITY

ALFONSO CASTRO AND SUMALEE UNSURANGSIE

We prove that a semilinear wave equation in which the range of the derivative of the nonlinearity includes an eigenvalue of infinite multiplicity has a solution. The solution is obtained through an iteration scheme which provides a priori estimates.

1. Introduction. Here we study the nonlinear wave equation

$$(1.1) \quad u_{tt} - u_{xx} + \lambda u = cq(x, t) + r(x, t) + g(u), \quad (x, t) \in [0, \pi] \times \mathbf{R},$$

$$(1.2) \quad u(0, t) = u(\pi, t) = 0, \quad u(x, t) = u(x, t + 2\pi), \\ (x, t) \in [0, \pi] \times \mathbf{R},$$

where $\lambda \in \mathbf{R} - \{k^2 - j^2: k = 1, 2, 3, \dots, j = 0, 1, 2, \dots\}$ and g is a function of class C^1 such that

$$(1.3) \quad \lim_{|u| \rightarrow \infty} g'(u) = 0.$$

A main difficulty in studying (1.1)–(1.2) arises when $-\lambda \in g'(\mathbf{R})$. This causes compactness arguments to fail because 0 is an eigenvalue of $u_{tt} - u_{xx}$, (1.2) of infinite multiplicity. Recent studies on (1.1)–(1.2) either: (i) assume that $g(u) - \lambda u$ is monotone (see [B-N], [R], [W]), or (ii) assume enough symmetry on g, q , and r so that the kernel of $u_{tt} - u_{xx}$, (1.2) reduces to $\{0\}$ (see [Co]), or (iii) use dichotomy on whether the Palais-Smale condition holds proving existence for values of $cq+r$ which cannot be given explicitly (see [H], [W]). Our main result (Theorem A below) does not fall in any of the above three classes.

Let $\Omega = [0, \pi] \times [0, 2\pi]$. Let H^1 , L^2 , and L^∞ denote the Sobolev spaces $H^1(\Omega)$, $L^2(\Omega)$, and $L^\infty(\Omega)$ respectively. We let $\|\cdot\|_1$, $\|\cdot\|$, and $\|\cdot\|_\infty$ denote the norms in H^1 , L^2 , and L^∞ respectively. Let

$$N = \left\{ u \in L^2: u = \sum_{k=1}^{\infty} (a_k \sin(kx) \sin(kt) + b_k \sin(kx) \cos(kt)) \right\}.$$

Let $N^\perp \subseteq L^2$ denote the orthogonal complement to N in L^2 . Let P denote the orthogonal projection onto N and Q the orthogonal projection

onto N^\perp . Let φ be a solution to

$$(1.4) \quad \begin{aligned} \varphi_{tt} - \varphi_{xx} + \lambda\varphi &= q(x, t), & (x, t) \in [0, \pi] \times [0, 2\pi], \\ \varphi(0, t) = \varphi(\pi, t) &= 0, & \varphi(x, t) = \varphi(x, t + 2\pi). \end{aligned}$$

Throughout the rest of this paper we will assume that

$$(1.5) \quad m\{(x, t) \in \gamma; |\varphi(x, t)| < \delta\} \rightarrow 0 \quad \text{as } \delta \rightarrow 0$$

uniformly in γ , where γ is any characteristic of the operator $\partial_{tt} - \partial_{xx}$. Our main result is

THEOREM A. *If $(P(r))_t \in L^2$, $\varphi_t \in L^\infty$, and (1.5) holds, then there exists c_0 such that for $|c| > c_0$ the problem (1.1)–(1.2) has a weak solution $u \in H^1 \cap L^\infty$.*

Our proof is based on an iteration argument that resembles the proof of the inverse function theorem. We give (see §3) estimates in $H^1 \cap L^\infty$ which show the convergence of the scheme. Even though our arguments and main result suggest applicability of the standard inverse function theorem, our hypotheses are not enough to guarantee it.

2. Preliminaries and notations. A direct calculation shows that if $u \in N$, then $u_{tt} - u_{xx} = 0$ in the sense of distributions. Let $Au = u_{tt} - u_{xx}$. We recall (see [B-N]) that A subject to condition (1.2) is selfadjoint, the range of A is closed in L^2 , and $R(A) = N^\perp$. The eigenvalues of A subject to (1.2) form the set $\{k^2 - j^2: k = 1, 2, 3, \dots, j = 0, 1, 2, \dots\}$. The corresponding eigenfunctions are $\sin(kx) \sin(jt)$ and $\sin(kx) \cos(jt)$. The operator A^{-1} is compact from N^\perp into N^\perp , and there exists a real number c such that

$$\begin{aligned} \|A^{-1}f\|_\infty &\leq c\|f\| & \text{for all } f \in R(A), \\ \|A^{-1}f\|_1 &\leq c\|f\| & \text{for all } f \in R(A). \end{aligned}$$

Using Fourier series it is easy to show that if $u \in H^1, v \in N \cap H^1$ then

$$(2.1) \quad \iint (P(u))_t v_t = \iint u_t v_t,$$

$$(2.2) \quad \int_0^{2\pi} (v)^2(x, s) ds \leq \int \int_\Omega (v)^2 dx dt \quad \text{for all } x \in [0, \pi].$$

Let A_1 denote the operator defined by

$$A_1 \left(\sum_{\substack{k=1 \\ j=0 \\ k \neq j}}^{\infty} (a_{kj} \sin(kx) \sin(jt) + b_{kj} \sin(kx) \cos(jt)) \right) \\ = \sum_{\substack{k=1 \\ j=0 \\ k \neq j}}^{\infty} \left[\frac{a_{kj}}{k^2 - j^2 + \lambda} \sin(kx) \sin(jt) + \frac{b_{kj}}{k^2 - j^2 + \lambda} \sin(kx) \cos(jt) \right].$$

It is easy to see that if $f \in L^2$, then $w = A_1(f)$ is a weak solution to $w_{tt} - w_{xx} + \lambda w = f$. An elementary Fourier series argument shows that if $w = A_1(Q(f))$, then

$$(2.3) \quad \int_0^{2\pi} ((w_x)^2(x, t) + (w_t)^2(x, t)) dt \leq b_0^2 \|Q(f)\|^2$$

for all $x \in [0, \pi]$.

where $b_0 = \max\{(2/\pi)(k^2 + j^2)/(k^2 - j^2 + \lambda)^2 : k \neq j, k = 1, 2, 3, \dots, j = 1, 2, 3, \dots\}$. In particular

$$(2.4) \quad \|A_1(Q(f))\|_1 \leq b_0 \pi^{1/2} \|Q(f)\|.$$

Also it is easy to show that

$$(2.5) \quad \|A(Q(f))\|_{\infty} \leq b_1 \|Q(f)\|$$

where

$$b_1 = \left(\sum_{\substack{k=1 \\ j=0 \\ k \neq j}}^{\infty} \frac{2}{(k^2 - j^2 + \lambda)^2} \right)^{1/2}$$

We can rewrite (1.1) as the following:

$$(2.6) \quad u_{tt} - u_{xx} + \lambda u = (c/R)Rq(x, t) + r(x, t) + g(u)$$

where

$$R = \max\{2(2)^{1/2}dK\lambda, 2(2)^{1/2}, 16d^4 \|g'\|_{\infty}^3 \lambda, 16 \|g'\|_{\infty}^3 / \lambda\} / (\|\varphi_t\|_{\infty} + 2^{1/2}),$$

$K = b_0 \pi^{1/2} + b_1$ and $d > 0$ is a constant such that $\|u\|_{L^4} \leq d \|u\|_1$ and $\|u\| \leq d \|u\|_1$ for all $u \in H^1$ (the existence of d follows the fact that

the embeddings $H^1 \rightarrow L^4$ and $H^1 \rightarrow L^2$ are continuous (see [A]). Let $w = Ru/c$ and $\beta = R/c$. Then (2.6) becomes

$$(2.7) \quad w_{tt} - w_{xx} + \lambda w = Rq(x, t) + \beta(r(x, t) + g(w/\beta)).$$

Inductively we define $w_0 = 0$, w_1 as the solution to

$$(2.8) \quad \begin{aligned} (w_1)_{tt} - (w_1)_{xx} + \lambda w_1 &= \beta(r(x, t) + g(R\varphi/\beta)), \\ w_1(0, t) = w_1(\pi, t) &= 0, \quad w_1(x, t) = w_1(x, t + 2\pi), \end{aligned}$$

and w_{n+1} as the solution to

$$(2.9) \quad \begin{aligned} (w_{n+1})_{tt} - (w_{n+1})_{xx} + \lambda w_{n+1} &= \beta(r(x, t) + g((R\varphi + w_n)/\beta)), \\ w_{n+1}(0, t) = w_{n+1}(\pi, t) &= 0, \quad w_{n+1}(x, t) = w_{n+1}(x, t + 2n). \end{aligned}$$

3. Estimates. For the sake of simplicity we will assume throughout the rest of this paper that $\lambda > 0$, and $c > 0$. The case $\lambda < 0$, or $c < 0$ requires only obvious modifications.

LEMMA 3.1. *Let $\{w_n\}_n$ be defined by (2.8) and (2.9). Under the assumption of Theorem A, there exists $\beta_1 > 0$ such that if $\beta \in (0, \beta_1)$ then for all $n = 1, 2, 3, \dots$ we have*

$$(3.1) \quad \|w_n\|_1 + \|w_n\|_\infty \leq 1.$$

Proof. Since for each characteristic γ , $m\{(x, t) \in \gamma : |\varphi(x, t)| < \delta\} \rightarrow 0$ as $\delta \rightarrow 0$ uniformly in γ then there exists $\delta_0 > 0$ such that if $0 < \delta < \delta_0$ then

$$(3.2) \quad \begin{aligned} m\{(x, t) \in \gamma : |\varphi(x, t)| < \delta\} \\ < (\lambda/(32\pi R(\|g'\|_\infty + 1)(\|\varphi_t\|_\infty + 2^{1/2})))^2 \end{aligned}$$

for any characteristic γ . Because of (1.3) there exists $L \geq 0$ such that for all $u \in \mathbf{R}$

$$(3.3) \quad |g(u)| \leq (|u|/64\pi KR(\|\varphi\|_\infty + 1)) + L,$$

and there exists $M > 0$ such that if $|u| > M$ then

$$(3.4) \quad |g'(u)| < \lambda/(64\pi^2 R(\|\varphi_t\|_\infty + 2^{1/2})).$$

Now we define β_1 by

$$(3.5) \quad \beta_1 = \min\{1/(32(M + (L + 1)\pi K + K\|r\|)), \\ \lambda/(32\pi(\|r_t\| + 1)), R\delta_0/M\}.$$

Next we prove (3.1) by induction. First we observe that

$$\begin{aligned}
 (3.6) \quad \|g(R\varphi/\beta)\|^2 &= \iint_{\Omega} (g(R\varphi/\beta))^2 \\
 &\leq 2 \iint_{\Omega} L^2 + 2 \left(\iint_{\Omega} (R\varphi/\beta)^2 \right) / (64\pi KR(\|\varphi\|_{\infty} + 1))^2 \\
 &\leq 4\pi^2 L^2 + (1/32K\beta)^2.
 \end{aligned}$$

We write $w_1 = v_1 + z_1$ with $v_1 \in N$ and $z_1 \in N^{\perp}$. From (2.4), (2.5), and (2.7) it follows that if $\beta \in (0, \beta_1)$ then

$$(3.7) \quad \|z_1\|_1 + \|z_1\|_{\infty} \leq 1/4.$$

Projecting (2.8) into N , differentiating with respect to t , multiplying by $(v_1)_t$, and integrating over Ω we have

$$\begin{aligned}
 \lambda \iint_{\Omega} ((v_1)_t)^2 dx dt &= \beta \iint_{\Omega} (P(r + g(R\varphi/\beta)))_t (v_1)_t dx dt \\
 &= \beta \iint_{\Omega} (r + g(R\varphi/\beta))_t (v_1)_t dx dt \\
 &= \beta \iint_{\Omega} r_t (v_1)_t + \iint_{\Omega} g'(R\varphi/\beta)(R\varphi_t)(v_1)_t \\
 &= \beta \|r_t\| \cdot \|(v_1)_t\| + R\|\varphi_t\|_{\infty} \|(v_1)_t\| \left(\iint_{\Omega} (g'(R\varphi/\beta))^2 \right)^{1/2}.
 \end{aligned}$$

Therefore

$$\|(v_1)_t\| \leq \left(\beta \|r_t\| + R\|\varphi_t\|_{\infty} \left(\iint_{\Omega} (g'(R\varphi/\beta))^2 \right)^{1/2} \right) / \lambda.$$

In order to estimate $\iint_{\Omega} (g'(R\varphi/\beta))^2$ we define $s_{\beta} = \{(x, t) : |R\varphi(x, t)| \leq M\beta\}$ and $c_{\beta} = \Omega - s_{\beta}$. Since $M\beta/R < \delta_0$ we have

$$m(s_{\beta}) < (\lambda/(32\pi R(\|g'\|_{\infty} + 1)(\|\varphi_t\|_{\infty} + 2^{1/2})))^2$$

(see (3.2) and (3.5)); then

$$\begin{aligned}
 \left(\iint_{\Omega} (g'(R\varphi/\beta))^2 \right)^{1/2} &= \left[\iint_{s_{\beta}} (g'(R\varphi/\beta))^2 + \iint_{c_{\beta}} (g'(R\varphi/\beta))^2 \right]^{1/2} \\
 &\leq [\|g'\|_{\infty}^2 m(s_{\beta}) + 2\pi^2 \lambda^2 / (64\pi^2 R(\|\varphi_t\|_{\infty} + 2^{1/2}))^2]^{1/2} \\
 &\leq [2\lambda^2 / (32R(\|\varphi_t\|_{\infty} + 2^{1/2}))^2]^{1/2} \\
 &\leq \lambda / (16R(\|\varphi_t\|_{\infty} + 2^{1/2})).
 \end{aligned}$$

Hence if $\beta \in (0, \beta_1)$ then $\|(v_1)_t\| \leq 1/8$. Since $\|v_t\| = \|v_x\|$ for all $v \in N \cap H^1$, we have $\|v_1\|_1 \leq 1/4$. Because of (2.2) we have

$$\begin{aligned} |v_1(x, t)| &= \left| \int_0^t (v_1)_t(x, s) ds \right| \leq \left(\int_0^t ((v_1)_t)^2(x, s) ds \right)^{1/2} \left(\int_0^t ds \right)^{1/2} \\ &\leq (2\pi)^{1/2} \|(v_1)_t\| \leq (2\pi)^{1/2}/8 \leq 1/3. \end{aligned}$$

Therefore

$$(3.8) \quad \|v_1\|_1 + \|v_1\|_\infty \leq 7/12.$$

Combining (3.7) and (3.8) we have

$$\|w_1\|_1 + \|w_1\|_\infty \leq 1.$$

Suppose now that $\|w_n\|_1 + \|w_n\|_\infty \leq 1$. We write $w_{n+1} = v_{n+1} + z_{n+1}$ with $v_{n+1} \in N$ and $z_{n+1} \in N^\perp$. Again from (2.4), (2.5) and (2.7) we have

$$\|z_{n+1}\|_1 + \|z_{n+1}\|_\infty \leq \beta K(\|r\| + \|g((R\varphi + w_n)/\beta)\|).$$

In order to estimate $\|g((R\varphi + w_n)/\beta)\|$ we observe that

$$\begin{aligned} &\iint_{\Omega} (g((R\varphi + w_n)/\beta))^2 \\ &\leq 2 \iint_{\Omega} L^2 + 2 \left(\iint_{\Omega} ((R\varphi + w_n)/\beta)^2 \right) / (64\pi RK(\|\varphi\|_\infty + 1))^2 \\ &\leq 4\pi^2 L^2 + 8\pi^2 [(R\|\varphi\|_\infty + 1) / (64\pi RK(\|\varphi\|_\infty + 1))]^2 / \beta^2. \end{aligned}$$

Therefore if $\beta \in (0, \beta_1)$ then

$$(3.9) \quad \|z_{n+1}\|_1 + \|z_{n+1}\|_\infty \leq 1/4.$$

Now projecting (2.9) into N , differentiating with respect to t , multiplying by $(v_{n+1})_t$ and integrating over Ω we have

$$\begin{aligned} (3.10) \quad &\lambda \iint_{\Omega} ((v_{n+1})_t)^2 \\ &= \beta \left(\iint_{\Omega} (r_t)(v_{n+1})_t \right. \\ &\quad \left. + \iint_{\Omega} g'((R\varphi + w_n)/\beta)((R\varphi + w_n)/\beta)_t (v_{n+1})_t \right) \\ &\leq \beta \|r_t\| \cdot \|(v_{n+1})_t\| \\ &\quad + (2\pi R\|\varphi_t\|_\infty + 2^{1/2}) \left[\iint_{\Omega} (g'((R\varphi + w_n)/\beta))^2 ((v_{n+1})_t)^2 \right]^{1/2}. \end{aligned}$$

Now we consider

$$(3.11) \quad \begin{aligned} I &= \iint_{\Omega} (g'((R\varphi + w_n)/\beta))^2 ((v_{n+1})_t)^2 \\ &= \int_0^{2\pi} \int_0^\pi (g'((R\varphi + w_n)/\beta))^2 ((v_{n+1})_t)^2 dx dt. \end{aligned}$$

Without loss of generality we can assume that $v_{n+1} = h(x - t)$ or $(v_{n+1})_t = -h'(x - t)$. Because the integrand in (3.11) is 2π periodic in t we have

$$I = \int_0^\pi \int_x^{2\pi+x} (g'((R\varphi + w_n)(x, t)/\beta))^2 (h'(x - t))^2 dt dx.$$

By defining $\eta = x$, $\zeta = -x + t$, $\gamma_\zeta = \{(s, s + \zeta) : s \in [0, \pi]\}$ and $A_\beta = \{(x, t) \in \Omega : |R\varphi(x, t)| \leq M\beta + 1\}$ we have

$$(3.12) \quad \begin{aligned} I &= \int_0^{2\pi} \int_0^\pi (g'((R\varphi(\eta, \eta + \zeta) + w_n(\eta, \eta + \zeta))/\beta))^2 (h'(-\zeta))^2 d\eta d\zeta \\ &= \int_0^{2\pi} (h'(-\zeta))^2 \left(\int_0^\pi (g'((R\varphi(\eta, \eta + \zeta) + w_n(\eta, \eta + \zeta))/\beta))^2 d\eta \right) d\zeta \\ &= \int_0^{2\pi} (h'(-\zeta))^2 \left[\int_{\gamma_\zeta \cap A_\beta} (g'((R\varphi(\eta, \eta + \zeta) + w_n(\eta, \eta + \zeta))/\beta))^2 d\eta \right. \\ &\quad \left. + \int_{\gamma_\zeta - (\gamma_\zeta \cap A_\beta)} (g'((R\varphi(\eta, \eta + \zeta) + w_n(\eta, \eta + \zeta))/\beta))^2 d\eta \right] d\zeta \\ &\leq \|(v_{n+1})_t\|^2 [(\|g'\|_\infty \lambda / (32\pi(\|g'\|_\infty + 1)))^2 \\ &\quad + (\lambda\pi^{1/2}/64\pi^2)^2] / (2R(\|\varphi_t\|_\infty + 2^{1/2}))^2. \end{aligned}$$

Hence if $\beta \in (0, \beta_1)$ then

$$\|(v_{n+1})_t\| \leq 1/8.$$

Imitating the argument in (3.8) we have

$$(3.13) \quad \|v_{n+1}\|_1 + \|v_{n+1}\|_\infty \leq 7/12.$$

Combining (3.9) and (3.13) we have

$$\|w_{n+1}\|_1 + \|w_{n+1}\|_\infty \leq 1,$$

which proves the lemma.

LEMMA 3.2. *If $\{a_n\}$ is a sequence of nonnegative real numbers such that*

$$a_{n+1} \leq \tau(a_n + a_{n-1}), \quad n = 2, 3, 4, \dots,$$

then

$$a_{n+1} \leq c_{n+1}k\tau^{[n/2]},$$

where $k = \max\{a_1, a_2\}$, and c_n is the n th Fibonacci number of the sequence defined by $c_{n+1} = c_n + c_{n-1}$, $c_2 = 2$ and $c_1 = 0$, and $[x]$ denotes the largest integer less than or equal to x . In particular the series $\sum a_n$ converges if τ is small enough.

Proof. We prove the lemma by induction. For $n = 2$ we have

$$a_3 \leq \tau(a_2 + a_1) \leq 2\tau k = (c_2 + c_1)\tau k = c_3k\tau^{[2/2]}.$$

Suppose that

$$a_n \leq c_nk\tau^{[(n-1)/2]}.$$

If n is even we have

$$\begin{aligned} a_{n+1} &\leq \tau(a_n + a_{n-1}) \leq \tau(c_nk\tau^{[(n-1)/2]} + c_{n-1}k\tau^{[(n-2)/2]}) \\ &\leq \tau k(c_n\tau^{(n-2)/2} + c_{n-1}\tau^{(n-2)/2}) \\ &\leq (c_n + c_{n-1})k\tau^{n/2} \leq c_{n+1}k\tau^{[n/2]}. \end{aligned}$$

Similarly if n is odd we have

$$\begin{aligned} a_{n+1} &\leq \tau(c_nk\tau^{(n-1)/2} + c_{n-1}k\tau^{(n-3)/2}) \\ &= c_nk\tau^{(n-1)/2} + c_{n-1}k\tau^{(n-1)/2} \leq (c_n + c_{n-1})k\tau^{[n/2]}, \end{aligned}$$

which proves Lemma 3.2.

4. Proof of Theorem A. For $n = 1, 2, 3, \dots$ we write $w_n = v_n + z_n$ where $v_n \in N$ and $z_n \in N^\perp$. Let $n > 2$. From the Sobolev imbedding theorem and (2.4) and (2.5) we have

$$\begin{aligned} (4.1) \quad \|z_{n+1} - z_n\|^2 &\leq d^2 \|z_{n+1} - z_n\|_1^2 \\ &\leq (dK\beta)^2 \iint_{\Omega} (g((R\varphi + w_n)/\beta) - g((R\varphi + w_{n-1})/\beta))^2 \\ &= (dK)^2 \iint_{\Omega} (g'((R\varphi + \zeta)/\beta))^2 (w_{n-1} - w_n)^2 \\ &\leq 2(dK)^2 \left[\iint_{\Omega} (g'((R\varphi + \zeta)/\beta))^2 (v_n - v_{n-1})^2 \right. \\ &\qquad \qquad \qquad \left. + \iint_{\Omega} (g'((R\varphi + \zeta)/\beta))^2 (z_n - z_{n-1})^2 \right] \end{aligned}$$

where $\zeta \in [w_n(x, t), w_{n-1}(x, t)] \cup [w_{n-1}(x, t), w_n(x, t)]$. Since v_n and $v_{n-1} \in N$, imitating the argument in (3.11)–(3.12) we see that

$$\begin{aligned} &\iint_{A_\beta} (g'((R\varphi + \zeta)/\beta))^2 (v_n - v_{n-1})^2 \\ &\leq (\lambda \|v_n - v_{n-1}\| / 32\pi R (\|\varphi_t\|_\infty + 2^{1/2}))^2. \end{aligned}$$

Thus

$$\begin{aligned}
 (4.2) \quad & 2(dK)^2 \iint_{\Omega} (g'((R\varphi + \zeta)/\beta))^2 (v_n - v_{n-1})^2 \\
 & \leq 2(dK)^2 \left[\iint_{A_\beta} (g'((R\varphi + \zeta)/\beta))^2 (v_n - v_{n-1})^2 \right. \\
 & \quad \left. + \iint_{\Omega - A_\beta} (g'((R\varphi + \zeta)/\beta))^2 (v_n - v_{n-1})^2 \right] \\
 & \leq 2(dK)^2 [(\lambda/32\pi R(\|\varphi_t\|_\infty + 2^{1/2}))^2 \\
 & \quad + (\lambda/64\pi R(\|\varphi_t\|_\infty + 2^{1/2}))^2] \|v_n - v_{n-1}\|^2 \\
 & \leq 2(dK)^2 (\lambda/16R(\|\varphi_t\|_\infty + 2^{1/2}))^2 \|v_n - v_{n-1}\|^2.
 \end{aligned}$$

where we also have used that $|(R\varphi + \zeta)/\beta| > M$ for $(x, t) \in \Omega - A_\beta$ (see (3.4)). On the other hand we have

$$\begin{aligned}
 (4.3) \quad & 2(dK)^2 \iint_{\Omega} (g'((R\varphi + \zeta)/\beta))^2 (z_n - z_{n-1})^2 \\
 & \leq 2(dK)^2 \left[\iint_{A_\beta} (g'((R\varphi + \zeta)/\beta))^2 (z_n - z_{n-1})^2 \right. \\
 & \quad \left. + \iint_{\Omega - A_\beta} (g'((R\varphi + \zeta)/\beta))^2 (z_n - z_{n-1})^2 \right] \\
 & \leq 2(dK)^2 \left[\|g'\|_\infty^2 \iint_{A_\beta} (z_n - z_{n-1})^2 \right. \\
 & \quad \left. + (\lambda/64\pi^2 R(\|\varphi_t\|_\infty + 2^{1/2}))^2 \|z_n - z_{n-1}\|^2 \right] \\
 & \leq 2(dK)^2 \left[\|g'\|_\infty^2 \iint_{\Omega} (\chi_{A_\beta} (z_n - z_{n-1}))^2 \right. \\
 & \quad \left. + (\lambda/64\pi^2 R(\|\varphi_t\|_\infty + 2^{1/2}))^2 \|z_n - z_{n-1}\|^2 \right] \\
 & \leq 2(dK)^2 \left[\|g'\|_\infty^2 \left(\iint_{\Omega} (\chi_{A_\beta})^2 \right)^{1/2} \left(\iint_{\Omega} (z_n - z_{n-1})^4 \right)^{1/2} \right. \\
 & \quad \left. + (\lambda/64\pi^2 R(\|\varphi_t\|_\infty + 2^{1/2}))^2 \|z_n - z_{n-1}\|^2 \right] \\
 & \leq 2(dK)^2 [\|g'\|_\infty^2 \lambda d^2 \|z_n - z_{n-1}\|^2 / 32\pi R(\|g'\|_\infty + 1)(\|\varphi_t\|_\infty + 2^{1/2}) \\
 & \quad + (\lambda/64\pi^2 R(\|\varphi_t\|_\infty + 2^{1/2}))^2 \|z_n - z_{n-1}\|^2],
 \end{aligned}$$

where we have used that by the Sobolev imbedding theorem (see [A]) $\|z_n - z_{n-1}\|_{L^4} \leq d\|z_n - z_{n-1}\|_1$. Also since g' is bounded,

$$\|\beta(g(\varphi + w_n)/\beta) - g((\varphi + w_{n-1})/\beta)\| \leq \|g'\|_\infty \|w_{n-1} - w_{n-2}\|_1.$$

This, (4.2) and (4.3) give

$$\begin{aligned} (4.4) \quad & 2(dK)^2 \iint_{\Omega} (g'((R\varphi + \zeta)/\beta))^2 (z_n - z_{n-1})^2 \\ & \leq 2(dK)^2 \frac{\|g'\|_\infty^4 \lambda(dK)^2 \|w_{n-1} - w_{n-2}\|^2}{32\pi R(\|g'\|_\infty + 1)(\|\varphi_t\|_\infty + 2^{1/2})} \\ & \leq 2(dK)^2 [\|g'\|_\infty^3 \lambda(dK)^2 \|w_{n-1} - w_{n-2}\|^2 / 32R(\|\varphi_t\|_\infty + 2^{1/2}) \\ & \quad + (\lambda/64\pi(\|\varphi_t\|_\infty + 2^{1/2}))^2 \|z_n - z_{n-1}\|^2]. \end{aligned}$$

Combining (4.2) and (4.4) we have

$$\begin{aligned} (4.5) \quad \|z_{n+1} - z_n\| & \leq 2^{1/2} dK \lambda \|w_n - w_{n-1}\| \\ & \quad + 2^{1/2} (dK)^2 \|g'\|_\infty^{3/2} \|w_{n-1} - w_{n-2}\| \\ & \quad \times (\lambda/32R(\|\varphi_t\|_\infty + 2^{1/2}))^{1/2}. \end{aligned}$$

Also

$$\begin{aligned} \lambda^2 \|v_{n+1} - v_n\|^2 & \leq \iint_{\Omega} (g'((\varphi + \zeta)/\beta))^2 (w_n - w_{n-1})^2 \\ & \leq 2 \iint_{\Omega} (g'((\varphi + \zeta)/\beta))^2 [(v_n - v_{n-1})^2 + (z_n - z_{n-1})^2]. \end{aligned}$$

Using now (4.2) and (4.4) we have

$$\begin{aligned} \lambda^2 \|v_{n+1} - v_n\|^2 & \leq 2(\lambda/16R(\|\varphi_t\|_\infty + 2^{1/2}))^2 \|v_n - v_{n-1}\|^2 \\ & \quad + 2[\|g'\|_\infty^3 \lambda(dK)^2 \|w_{n-1} - w_{n-2}\|^2 / 32R(\|\varphi_t\|_\infty + 2^{1/2}) \\ & \quad + (\lambda\|z_n - z_{n-1}\|/64R(\|\varphi_t\|_\infty + 2^{1/2}))^2]. \end{aligned}$$

Hence

$$\begin{aligned} (4.6) \quad \|v_{n+1} - v_n\| & \leq 2^{1/2} \|w_n - w_{n-1}\| / 8R(\|\varphi_t\|_\infty + 2^{1/2}) \\ & \quad + 2^{1/2} \|g'\|_\infty^{3/2} \|w_{n-1} - w_{n-2}\| / (32\lambda R(\|\varphi_t\|_\infty + 2^{1/2}))^{1/2}. \end{aligned}$$

Combining (4.5) and (4.6) and using the definition of R we have

$$\|w_{n+1} - w_n\| \leq (\|w_n - w_{n-1}\| + \|w_{n-1} - w_{n-2}\|) / 8.$$

Hence by Lemma 3.2 we have

$$\|w_{n+1} - w_n\| \leq k2^n (1/8)^{[n/2]}, \quad n = 2, 3, 4, \dots$$

Hence $\sum_{n=3}^{\infty} \|w_n - w_{n-1}\|$ converges. Thus the sequence $\{w_n = (w_n - w_{n-1}) + (w_{n-1} - w_{n-2}) + \cdots + (w_2 - w_1) + (w_1 - w_0)\}$ converges in L^2 to some $w \in L^2$. Since $\{w_n\}$ is bounded in $H^1 \cap L^\infty$ we see that w also belongs to $H^1 \cap L^\infty$. Hence by (2.11) we see that $w + R\phi$ is a solution to (1.2), (2.7). Hence $u = c(w + R\phi)$ is a solution to (1.1)–(1.2) which proves the Theorem.

REMARK. Double checking the proofs it is easily seen that Theorem A also holds when the limits in (1.3) are allowed to be in some interval of the form $(-s, s)$, with s depending on the distance from λ to $\{k^2 - j^2: k = 1, 2, 3, \dots, j = 0, 1, 2, 3, \dots\}$.

REFERENCES

- [A] R. Adams, *Sobolev Spaces*, Academic Press 1975.
- [B] H. Brezis, *Periodic solutions of nonlinear vibrating strings and duality principles*, Bull. Amer. Math. Soc., **8**, No. 3 (1983), 409–426.
- [B-N] H. Brezis and L. Nirenberg, *Forced vibrations for a nonlinear wave equation*, Comm. Pure Appl. Math., **31** (1978), 1–30.
- [C-W] D. Costa and M. Willem, *Multiple critical points of invariant functionals and applications*, to appear in Nonlinear Analysis TMA.
- [Co] J. M. Coron, *Periodic solutions of a nonlinear wave equation without assumption of monotonicity*, Math. Anal., **262** (1983), 273–285.
- [H] H. Hofer, *On the range of a wave operator with nonmonotone nonlinearity*, Math. Nachr., **106** (1982), 327–340.
- [R] P. H. Rabinowitz, *Free vibrations for a semilinear wave equation*, Comm. Pure Appl. Math., **31** (1978), 165–194.
- [W] M. Willem, *Density of the range of potential operators*, Proc. Amer. Math. Soc., **83** (1981), 341–344.

Received December 15, 1986.

NORTH TEXAS STATE UNIVERSITY
DENTON, TX 76203-5116

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906-1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1988 by Pacific Journal of Mathematics

Jeffery Marc Bergen and Luisa Carini, A note on derivations with power central values on a Lie ideal	209
Alfonso Castro and Sumalee Unsurangsi, A semilinear wave equation with nonmonotone nonlinearity	215
Marius Dadarlat, On homomorphisms of matrix algebras of continuous functions	227
A. Didierjean, Quelques classes de cobordisme non orienté refusant de se fibrer sur des sphères	233
Edward George Effros and Zhong-Jin Ruan, On matricially normed spaces	243
Sherif El-Helaly and Taqdir Husain, Orthogonal bases are Schauder bases and a characterization of Φ -algebras	265
Edward Richard Fadell and Peter N-S Wong, On deforming G -maps to be fixed point free	277
Jean-Jacques Gervais, Stability of unfoldings in the context of equivariant contact-equivalence	283
Douglas Martin Grenier, Fundamental domains for the general linear group	293
Ronald Scott Irving and Brad Shelton, Loewy series and simple projective modules in the category $\mathcal{O}_{\mathcal{S}}$	319
Russell Allan Johnson, On the Sato-Segal-Wilson solutions of the K-dV equation	343
Thomas Alan Keagy and William F. Ford, Acceleration by subsequence transformations	357
Min Ho Lee, Mixed cusp forms and holomorphic forms on elliptic varieties	363
Charles Livingston, Indecomposable surfaces in 4-space	371
Geoffrey Lynn Price, Shifts of integer index on the hyperfinite II_1 factor	379
Andrzej Śladek, Witt rings of complete skew fields	391