

Pacific Journal of Mathematics

REMARKS ON THE DIFFERENTIAL ENVELOPES OF ASSOCIATIVE ALGEBRAS

R. COQUEREAUX AND D. KASTLER

REMARKS ON THE DIFFERENTIAL ENVELOPES OF ASSOCIATIVE ALGEBRAS

R. COQUEREAUX AND D. KASTLER

Dedicated to the memory of Henry Dye

We study the relationship between the two types of differential envelopes of $\mathbf{Z}/2$ -graded associative complex algebras; and describe the differential envelopes of a $\mathbf{Z}/2$ -graded algebra as a simple modification of those of the associated ungraded algebra.

In this note we describe some aspects of the two existing notions of differential envelope of a $\mathbf{Z}/2$ -graded complex (or, for that matter real) associative algebra. Both notions are basic in the non commutative geometry under current development.

In §1, which fixes notation, we recall the definitions and main properties of the two types of differential envelope (= universal differential algebra) one attaches to a $\mathbf{Z}/2$ -graded algebra A . The first, $\tilde{\Omega}(A) = \mathbf{C}\tilde{\mathbf{1}} \oplus \Omega(A)$, pertains to general algebras (unital or not), and has a differential which vanishes nowhere on A (in particular, if A is unital with unit $\mathbf{1}$, $d\mathbf{1} \neq 0$). The second, ΩA , is defined only for unital algebras, and has a differential vanishing on the unit. $\tilde{\Omega}(A)$ can accordingly be directly defined in terms of tensors over A , whilst the construction of ΩA uses either tensors over an A -bimodule, or a subfamily of tensors over A . Both $\tilde{\Omega}(A)$ (for that matter $\Omega(A)$) and ΩA are universal objects (= tied up with functors), the first in the general and the second in the unital category.

In §2 we discuss the relationship between the two notions, each of which is definable in terms of the other. On the one hand one has $\tilde{\Omega}(A) = \Omega\tilde{A}$, $\tilde{A} = \mathbf{C}\tilde{\mathbf{1}} \oplus A$ the augmented ($\mathbf{Z}/2$ -graded) algebra. On the other hand there is an injective multiplicative linear map of ΩA into the right ideal $A\tilde{\Omega}(A)$ of $\tilde{\Omega}(A)$, allowing one to consider ΩA as a linear subspace of tensors over A (vanishing under the “consecutive diagonal mappings”, and multiplying under “concatenation”).

In §3 we show how the differential envelopes of a $\mathbf{Z}/2$ -graded algebra A can be constructed in terms of those of the associated ungraded algebra (merely through a simple modification of the latter’s product).

This result is important from the algorithmic point of view, since it leads to a reformulation yielding formulae in which the “grading signs” occurs in a more transparent way (important in particular with respect to the algorithmics of $\mathbf{Z}/2$ -graded cyclic cohomology).

1. The two types of differential envelopes of an associative complex algebra. For the results quoted in this section we refer to [1] through [5]. In the sequel $A = A^0 \oplus A^1$ throughout denotes a $\mathbf{Z}/2$ -graded complex algebra¹, with set $A^\cdot = A^0 \cup A^1$ of homogeneous elements², and grading denoted by ∂ ($\partial a = 0$ resp. $1 \pmod{2}$ for $a \in A^0$, resp. $a \in A^1$). \tilde{A} denotes the augmented $\mathbf{Z}/2$ -graded algebra³, with unit $\tilde{\mathbf{1}}$ (irrespective of whether A possesses a unit of its own or not—if this is the case, this unit is denoted by $\mathbf{1}$).

1.1. *The first kind of differential envelope: the differential algebras $(\tilde{\Omega}(A), d)$ and $(\Omega(A), d)$.* Let A be a $\mathbf{Z}/2$ -graded algebra (not necessarily unital). We define, as an \mathbf{N} -graded vector space

$$(1.1) \quad \tilde{\Omega}(A) = \sum_{n \in \mathbf{N}}^{\oplus} \tilde{\Omega}(A)^n$$

where

$$(1.2) \quad \tilde{\Omega}(A)^n = \tilde{A} \otimes A^{\otimes n}, \quad n \in \mathbf{N}$$

with $\tilde{\Omega}(A)^0 = \tilde{A}$. The *intrinsic grading* ∂_0 is defined by setting, for $a_0 \otimes a_1 \otimes \dots \otimes a_n \in \tilde{\Omega}(A)^n$:

$$(1.3) \quad \partial_0(a_0 \otimes a_1 \otimes \dots \otimes a_n) = \sum_{k=0}^n \partial a_k,$$

the *total grading* ∂ is then the sum of the intrinsic and the \mathbf{N} -grading.

$\tilde{\Omega}(A)^n$ is made into a ∂_0 -graded unital \tilde{A} -bimodule, by setting, for $a_0, x = \lambda \tilde{\mathbf{1}} + \underline{x} \in \tilde{A}$, $\lambda \in \mathbf{C}$, $a_1, \dots, a_n \in A^\cdot$:

$$(1.4) \quad x(a_0 \otimes a_1 \otimes \dots \otimes a_n) = xa_0 \otimes a_1 \otimes \dots \otimes a_n$$

¹“Algebra” always means “complex associative algebra” (whereby “complex” could throughout be replaced by “real”, changing \mathbf{C} to \mathbf{R}).

²Generally, we denote by E^\cdot the set $E^0 \cup E^1$ of homogeneous elements of any $\mathbf{Z}/2$ -graded vector space $E = E^0 \oplus E^1$.

³ $\tilde{A} = \mathbf{C} \oplus A$ with product $(\lambda, a)(\mu, b) = (\lambda\mu, \lambda b + \mu a + ab)$, $\lambda, \mu \in \mathbf{C}$, $a, b \in A$; and $\mathbf{Z}/2$ -grading $\tilde{A}^0 = \mathbf{C} \oplus A^0$, $\tilde{A}^1 = 0 \oplus A^1$. The “added unit” is $\tilde{\mathbf{1}} = (1, 0)$ (thus $\tilde{A} = \mathbf{C}\tilde{\mathbf{1}} \oplus A$). If A is unital with unit $\mathbf{1}$, it appears as the ideal $(0, A)$ of \tilde{A} generated by the idempotent $(0, \mathbf{1})$.

and⁴

$$(1.5) \quad (a_0 \otimes a_1 \otimes \cdots \otimes a_n)x \\ = \sum_{i=0}^{n-1} (-1)^{n+i+\sum_{k=i+1}^n \partial a_k} a_0 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_n \otimes \underline{x} \\ + a_0 \otimes a_1 \otimes \cdots \otimes a_n x.$$

Defining then the product of $a_0 \otimes a_1 \otimes \cdots \otimes a_n \in \tilde{\Omega}(A)^n$ and $b_0 \otimes b_1 \otimes \cdots \otimes b_m \in \tilde{\Omega}(A)^m$ as

$$(1.6) \quad (a_0 \otimes a_1 \otimes \cdots \otimes a_n)(b_0 \otimes b_1 \otimes \cdots \otimes b_m) \\ = \{(a_0 \otimes a_1 \otimes \cdots \otimes a_n)b_0\} \otimes b_1 \otimes \cdots \otimes b_m,$$

$\tilde{\Omega}(A)$ becomes an associative complex algebra, graded for both $\mathbf{Z}/2$ -gradings ∂_0 and ∂ , and for the \mathbf{N} -grading.

Setting now, for $a_0 = \lambda \tilde{\mathbf{1}} + \underline{a}_0 \in \tilde{A}$, $\underline{a}_0, a_1, \dots, a_n \in A$

$$(1.7) \quad d(a_0 \otimes a_1 \otimes \cdots \otimes a_n) = \tilde{\mathbf{1}} \otimes \underline{a}_0 \otimes a_1 \otimes \cdots \otimes a_n$$

specifies d as a linear operator of $\tilde{\Omega}(A)$, of \mathbf{N} - and ∂ -grade 1, and of vanishing square, which fulfills the graded derivation property

$$(1.8) \quad \begin{cases} d(\omega\psi) = (d\omega)\psi + (-1)^{\partial\omega}\omega d\psi, \\ \omega, \psi \in \tilde{\Omega}(A), \omega \text{ of total grade } \partial\omega, \end{cases}$$

thereby making $\tilde{\Omega}(A)$ a ∂ -graded (in fact bigraded⁵ differential algebra): elements of $\tilde{\Omega}(A)$ are then reinterpreted as follows:

$$(1.9) \quad a_0 \otimes a_1 \otimes \cdots \otimes a_n = a_0 da_1 \cdots da_n.$$

It is useful to note that

$$(1.10) \quad \tilde{\Omega}(A) = \mathbf{C}\tilde{\mathbf{1}} \oplus \Omega(A)$$

where $\Omega(A)$, the direct sum of the

$$(1.11) \quad \Omega(A)^n = \begin{cases} A, & n = 0, \\ \tilde{\Omega}(A)^n, & n \geq 1 \end{cases}$$

is an ideal of $\tilde{\Omega}(A)$ decomposing in turn as

$$(1.12) \quad \Omega(A) = d\Omega(A) \oplus A\Omega(A)$$

⁴This formula becomes transparent by identifying the l.h.s. with $(a_0 d_1 \cdots da_n)x$ (cf. (1.9) below), and transforming the latter expression using the graded derivation property of d (cf. (1.8) and (1.9) below).

⁵a $\mathbf{Z}/2$ -graded differential algebra $(\Xi = \Xi^+ \oplus \Xi^-, \delta)$ is a *bigraded differential algebra* whenever the algebra Ξ has \mathbf{N} -grading $\Xi = \sum_{n \in \mathbf{N}}^{\oplus} \Xi^n$ such that (i) $\Xi^n = \Xi^n \cap \Xi^+ \oplus \Xi^n \cap \Xi^-$, $n \in \mathbf{N}$ and $\delta(\Xi^n) \subset \Xi^{n+1}$, $n \in \mathbf{N}$.

with $A\Omega(A)$ a right ideal of $\Omega(A)$ (generated by the unit $\mathbf{1}$ of A for a unital A).

$\Omega(A)$ is a universal object, which could more transparently (but less explicitly) be defined as the quotient of the free algebra of the set $\{a, da; a \in A\}$ through the ideal generated by⁶

$$(1.13) \quad \begin{cases} \lambda \cdot a \dot{+} \mu \cdot b \dot{-} (\lambda a + \mu b), \\ \lambda \cdot da \dot{+} \mu \cdot db \dot{-} d(\lambda a + \mu b), \\ a \cdot b \dot{-} ab, \\ da \cdot b \dot{+} (-1)^{\partial a} a \cdot db, \end{cases} \quad \begin{cases} a, b \in A, \\ \lambda, \mu \in \mathbf{C}, \\ a \in A^*, b \in A. \end{cases}$$

As a consequence⁷, given a $\mathbf{Z}/2$ -graded complex algebra B , and a pair (α, Δ) of a homomorphism⁸ $\alpha: A \rightarrow B$ and a graded α -derivation $\Delta: A \rightarrow B$ (i.e. a grade one linear map fulfilling

$$(1.14) \quad \Delta(ab) = (\Delta a)\alpha(b) + (-1)^{\partial a}\alpha(a)\Delta b, \quad \begin{cases} a \in A, \\ b \in A, \end{cases}$$

there is a unique homomorphism⁸ $\theta: \Omega(A) \rightarrow B$ such that

$$(1.15) \quad \begin{cases} \theta(a) = \alpha(a), \\ \theta(da) = \Delta a, \end{cases} \quad , a \in A,$$

specifically given by

$$(1.16) \quad \begin{cases} \theta(a_0 da_1 \cdots da_n) = \alpha(a_0)\Delta a_1 \cdots \Delta a_n, \\ \theta(da_1 \cdots da_n) = \Delta a_1 \cdots \Delta a_n. \end{cases}$$

In fact we have a bijection between such pairs (α, Δ) and homomorphisms: $\Omega(A) \rightarrow B$ (of grade 0):

$$(1.17) \quad \begin{array}{ccc} \Omega(A) & & \\ \uparrow & \searrow \theta & \\ A & \xrightarrow[\Delta]{\alpha} & B \end{array}$$

This applies in particular (making $\Delta = \delta \circ \alpha$) to homomorphisms $\alpha: A \rightarrow B$ into $\mathbf{Z}/2$ -graded algebras B possessing a graded derivation δ . And the further specialization $B = \Xi$, (Ξ, δ) a $\mathbf{Z}/2$ -graded differential

⁶Operations within the free algebra are indicated by a dot.

⁷And also by direct proof, using the original definition (1.12) of $\Omega(A)$ —this showing the identity of the original definition and the definition as a quotient.

⁸Of $\mathbf{Z}/2$ -graded complex algebras (in particular, of grade zero).

algebra⁹, then yields a functor $\Omega()$ from the category of $\mathbf{Z}/2$ -graded algebras to that of $\mathbf{Z}/2$ -graded differential algebras:

$$(1.18) \quad \begin{array}{ccc} \Omega(A) & & \\ \uparrow & \searrow \Omega(\alpha) & \\ A & \xrightarrow{\alpha} & (\Xi, \delta) \end{array}$$

The special case where (Ξ, δ) is bigraded, and α ranges in the $\mathbf{N} = 0$ part of Ξ , then furnishes a functor from the category of $\mathbf{Z}/2$ -graded algebras to that of bigraded differential algebras.

1.2. *The second kind of differentiable envelope: the differentiable algebra $\Omega A, A$ unital.* Now let A be a unital $\mathbf{Z}/2$ -graded algebra with unit $\mathbf{1}$.

Defining as follows a linear map $m: A \otimes A \rightarrow A$:¹⁰

$$(1.19) \quad m \left(\sum_{i=1}^k a_i \otimes b_i \right) = \sum_{i=1}^k (-1)^{\partial b_i} a_i b_i, \\ a_i \in A, \quad b_i \in A^*, \quad i = 1, \dots, k$$

as well as linear maps $\phi^\perp, \phi: A \otimes A \rightarrow A \otimes A$:

$$(1.20) \quad \begin{cases} \phi^\perp(T) = m(T) \otimes \mathbf{1}, & T \in A \otimes A, \\ \phi = \text{id} - \phi^\perp, \end{cases}$$

we get supplementary idempotents ϕ, ϕ^\perp , such that

$$(1.21) \quad \begin{cases} J = \text{Im } \phi = \text{Ker } \phi^\perp = \text{Ker } m, \\ K = \text{Ker } \phi = \text{Im } \phi^\perp = A \otimes \mathbf{1}. \end{cases}$$

Moreover, if one sets

$$(1.22) \quad \begin{cases} x(a \otimes b) = xa \otimes b, \\ (a \otimes b)x = a \otimes bx, \end{cases} \quad a, b, x \in A,$$

J becomes a graded A -bimodule: therefore the direct sum

$$(1.23) \quad \Omega A = \sum_{n \in \mathbf{N}} \Omega A^n, \quad \begin{cases} \Omega A^0 = A, \\ \Omega A^n = J \otimes_A J \otimes_A \dots \otimes_A J (n \text{ factors}) \end{cases}$$

is a unital algebra with unit $\mathbf{1}$ (for the product \otimes_A), \mathbf{N} -graded and $\mathbf{Z}/2$ -graded for both the *intrinsic grading* ∂_0 (stemming from the grading of $J^{\otimes n}$ as a tensor product) and the *total grading*, sum of the intrinsic and the \mathbf{N} -grading. Further, setting

$$(1.24) \quad da = \mathbf{1} \otimes a - (-1)^{\partial a} a \otimes \mathbf{1}, \quad a \in A,$$

⁹I.e. δ is of grade one and $\delta^2 = 0$.

¹⁰In the case of an algebra of functions m is the "diagonal map" assigning to functions of two variables their values on the diagonal.

specifies a graded derivation $d: A \rightarrow J$:

$$(1.25) \quad d(ab) = (da)b \oplus (-1)^{\partial a}adb, \quad a \in A', \quad b \in A,$$

vanishing on the unit $\mathbf{1}$ and fulfilling

$$(1.26) \quad adb = a \otimes b - (-1)^{\partial b}ab \otimes \mathbf{1} = \phi(a \otimes b), \quad a, b \in A.$$

Since d uniquely extends to a differential d of ΩA (i.e. a ∂ -graded derivation of vanishing square), moreover of \mathbf{N} -grade 1, ΩA becomes a $\mathbf{Z}/2$ -graded (in fact a bigraded) differential algebra.

As was the case above for $\Omega(A)$, ΩA is a universal object—but now “in the unital category”. ΩA could be defined as the quotient of the free algebra of the set $\{a, da; a \in A\}$ through the ideal generated by¹¹

$$(1.27) \quad \begin{cases} \lambda \cdot a \dot{+} \mu \cdot b \dot{-} (\lambda a + \mu b), \\ \lambda \cdot da \dot{+} \mu \cdot db \dot{-} d(\lambda a + \mu b), \\ a \cdot b \dot{-} ab, \\ da \cdot b \dot{+} (-1)^{\partial a}a \cdot db \dot{-} d(ab), \\ d\mathbf{1} \end{cases} \quad \begin{cases} \lambda, \mu \in \mathbf{C}, \\ a, b \in A, \\ a \in A', \quad b \in A. \end{cases}$$

Moreover, we have universality properties analogous to those of $\Omega(A)$, (but within the unital category): to each $\mathbf{Z}/2$ -graded algebra B , and pair (α, Δ) of a homomorphism⁸: $A \rightarrow B$ and a *unital* graded derivation α -derivation Δ (i.e.

$$(1.28) \quad \Delta \mathbf{1} = 0),$$

there is a homomorphism $\theta: \Omega A \rightarrow B$ lifting α , namely

$$(1.29) \quad \begin{cases} \theta(a_0 da_1 \cdots da_n) = \alpha(a_0)\Delta a_1 \cdots \Delta a_n, \\ a_0, a_1, \dots, a_n \in A. \end{cases}$$

If B and α are unital, (1.28) is automatic and θ is unital. In fact, there is a bijection between such pairs (α, Δ) , and homomorphisms: $\Omega A \rightarrow B$:

$$(1.30) \quad \begin{array}{ccc} & \Omega A & \\ & \uparrow \quad \searrow \theta & \\ & A & \xrightarrow[\Delta]{\alpha} B \end{array}$$

¹¹As was the case for $\Omega(A)$, the identity of the two definitions results from the fact that both lead to the same universally property, cf. (1.31) below. Notice, on the other hand, that the second definition implies a linear bijection: $a_0 da_1 \cdots da_n \leftrightarrow a_0 \otimes \hat{a}_1 \otimes \cdots \otimes \hat{a}_n, \hat{a}_i = a_i \pmod{\mathbf{C}\mathbf{1}}$.

This applies (with $\Delta = \delta \circ \alpha$) to *unital* homomorphism⁸ $\alpha: A \rightarrow B$, B equipped with a unital graded derivation δ ($\delta \mathbf{1}_B = 0$). The specialization: $B = \Xi$, (Ξ, δ) a unital $\mathbf{Z}/2$ graded differential algebra (i.e. $\delta \mathbf{1} = 0$) yields a functor from the category of unital algebras (with unital morphisms) to that of unital $\mathbf{Z}/2$ -graded differential algebras (with differential vanishing on the unit, and unital morphisms).

$$(1.31) \quad \begin{array}{ccc} & \Omega A & \\ & \uparrow & \searrow \Omega\alpha \\ & A & \xrightarrow{\alpha} (\Xi, \delta) \end{array}$$

Ω induces in turn a functor from the category of unital $\mathbf{Z}/2$ -graded algebras to that of unital bigraded differential algebras, taking bigraded differential algebras (Ξ, δ) and homomorphisms α with range within the $n = 0$ part of Ξ .

2. Relationship between the two types of differential envelopes.

2.1. *The isomorphism $\tilde{\Omega}(A) \cong \Omega\tilde{A}$.* Let A be an arbitrary $\mathbf{Z}/2$ -graded algebra (unital or not): the differential envelope $(\tilde{\Omega}(A), d)$ of the first kind is in fact isomorphic (as a bigraded differential algebra) to the differential envelope $(\Omega\tilde{A}, d)$ of the second kind pertaining to the augmentation \tilde{A} of A ¹².

This fact easily follows from the above-mentioned universality properties. On the one hand (1.18) (with $\alpha = i: A \hookrightarrow \tilde{A}$, and $(\Xi, \delta) = (\Omega\tilde{A}, d)$) yields a homomorphism $\Omega(i): \Omega(A) \rightarrow \Omega\tilde{A}$ with unique extension to $\tilde{\Omega}(A)$ as a unital homomorphism. On the other hand (1.31) (with A replaced by \tilde{A} , $\alpha = \text{id}: \tilde{A} \rightarrow \tilde{A}$ and $(\Xi, \delta) = (\tilde{\Omega}(A), d)$) yields a unital homomorphism $\tilde{\Omega}\text{id} = \Omega\tilde{A} \rightarrow \tilde{\Omega}(A)$, inverse of the preceding.

As is to be expected (1.21) combined with the fact $\tilde{A} = \mathbf{C}\tilde{\mathbf{1}} \oplus A$ implies that we have:

$$(2.1) \quad \tilde{\Omega}(A)^1 = \tilde{A} \otimes \tilde{A} \pmod{\tilde{A} \otimes \tilde{\mathbf{1}} \cong \tilde{A} \otimes A.}$$

2.2. *ΩA as consisting of tensors over A .* A being now a unital $\mathbf{Z}/2$ -graded algebra with unit $\mathbf{1}$, we will conversely construct the differential envelope ΩA of the second type in terms of that of the first kind—in fact as a subalgebra of the right ideal $A\Omega(A) = \mathbf{1}\Omega(A)$ of $\Omega(A)$,

¹²Hence we could consider the notation $\tilde{\Omega}(A)$ as superfluous, keeping the symbols $\Omega\tilde{A}$ and $\Omega(A)$ with $\Omega\tilde{A} = \mathbf{C}\tilde{\mathbf{1}} + \Omega(A)$.

with the product reducing to “concatenation”, and under a modified differential. We noted earlier (cf. footnote 11) the linear isomorphism

$$(2.2) \quad \Omega A \cong \sum_{n \in \mathbb{N}}^{\oplus} A \otimes (A/\mathbf{C1})^{\otimes n} \cong \sum_{n \in \mathbb{N}}^{\oplus} A^{\otimes n} / J_n$$

with

$$(2.3) \quad J_n = A \otimes \mathbf{1} \otimes A \otimes \cdots \otimes A + A \otimes A \otimes \mathbf{1} \otimes A \otimes \cdots \otimes A + \cdots + A \otimes A \otimes \cdots \otimes \mathbf{1}.$$

This amounts in fact to quotienting $\Omega(A)$ by an ideal, the corresponding exact sequence of algebras being split, hence yielding a supplementary subalgebra of $\Omega(A)$ isomorphic to ΩA .

We begin by describing these objects.

LEMMA. *With a $\mathbb{Z}/2$ -graded unital, we denote by Δ the linear operator of $A\Omega(A)$ obtained by applying d and projecting back in $A\Omega(A)$:*

$$(2.4) \quad \Delta\omega = \mathbf{1}d\omega, \quad \omega \in A\Omega(A);$$

and consider the direct sums

$$(2.5) \quad K_* = \sum_{n=1}^{\infty} \oplus K_n$$

where K_n is the linear closure of elements $a_0 da_1 \cdots da_n$ with $a_0, a_1, \dots, a_n \in A$ and $a_k = \mathbf{1}$ for at least one $k \geq 1$; and

$$(2.6) \quad J_* = \sum_{n=1}^{\infty} \oplus J_n$$

where $J_n \subset A\Omega(A)^n$ equals

$$(2.7) \quad J_n = \bigcap_{i=0}^{n-1} \text{Ker } m_i^n$$

the linear “consecutive diagonal maps” m_i^n being given as follows: for a_0, a_1, \dots, a_n

$$(2.8) \quad \begin{cases} m_0^n(a_0 da_1 \cdots da_n) = (-1)^{\partial a_1} a_0 a_1 da_2 \cdots da_n, \\ m_i^n(a_0 da_1 \cdots da_n) = (-1)^{\partial a_{i+1}} a_0 da_1 \cdots d(a_i a_{i+1}) \cdots da_n, i \geq 0. \end{cases}$$

We then have that

(i) K_* is the (graded) ideal of $\Omega(A)$ intersection of $A\Omega(A)$ and the ideal $\overline{K_*}$ of $\Omega(A)$ generated by $d\mathbf{1}$.

(ii) Δ is a “differential modulo K_* ”, i.e. one has for $\omega, \psi \in A\Omega(A)$, ω of grade $\partial\omega$

$$(2.9) \quad \Delta(\omega\psi) - (\Delta\omega)\psi - (-1)^{\partial\omega}\omega\Delta\psi \in K_*$$

$$(2.10) \quad \Delta^2\psi \in K_*$$

(iii) J_* is a subalgebra of $A\Omega(A)$ on which the product reduces to “concatenation” \times specified as follows¹³: for $a_0da_1 \cdots da_n \in J_n$, $b_0db_1, \dots, db_m \in J_m$:

$$(2.11) \quad (a_0da_1 \cdots da_n) \times (b_0db_1 \cdots db_m) \\ = a_0da_1 \cdots da_{n-1}d(a_nb_0)db_1 \cdots db_m$$

Note that the signs $(-1)^{\partial a_1}$, resp. $(-1)(-1)^{\partial a_{i+1}}$ amount to applying the linear operators ε_1 , resp. ε_{i+1} where $\varepsilon_k = \text{id} \otimes \cdots \otimes \text{id} \otimes \varepsilon \text{id} \otimes \cdots \otimes \text{id}$ with the $(k+1)$ th factor ε is the grading involution of A (we here use the identification (1.9)). As a consequence, leaving out these signs in the definition of m_n^n , resp. m_i^n would not alter the definition of J_n .

Proof. (i). To prove that K_* is an ideal it suffices to check that, given $\omega = a_0da_1 \cdots da_n$, $a_0, a_1, \dots, a_n \in A$, $a_k = \mathbf{1}$ for $k \geq 1$, and $\psi = b_0db_1 \cdots db_m$, $b_0, b_1, \dots, b_m \in A$, one has $\psi\omega \in J_*$ and $\omega\psi \in J_*$. Now $\psi\omega \in J_*$ because “reordering”¹⁴ ψb_0 does not affect the factor $da_k = d\mathbf{1}$. To check that $\omega\psi \in J_*$ it obviously suffices to check that $\omega b_0 \in J_*$: this however follows from the fact that in reordering ωb_0 , the two terms not displaying a factor $d\mathbf{1}$ exactly cancel:

$$(2.12) \quad \sum (-1)^{n+k-1+\sum_{i=k}^n \partial a_i} a_0da_1 \cdots d(a_{k-1}\mathbf{1})da_{k+1} \cdots da_ndb_0 \\ + \sum (-1)^{n+k+\sum_{i=k}^n \partial a_i} a_0da_1 \cdots d(a_{k-1})d(\mathbf{1}a_{k+1})da_ndb_0 = 0.$$

We now claim that \bar{K}_* is the linear closure of monomials $a_0da_1 \cdots da_n$, $a_0 \in \tilde{A}$, $a_1, \dots, a_n \in A$ with $a_k = \mathbf{1}$ for at least one $k \geq 1$. Indeed, reordering $\omega(d\mathbf{1})\psi$, $\omega, \psi \in \Omega(A)$, $\psi = xd\psi'$, $x \in A$, amounts to rearranging $\omega(d\mathbf{1})x$ whereby the factor $d\mathbf{1}$ always persists for the same reason as before. Now $K_* \subset A\Omega(A) \cap \bar{K}_*$ is obvious. And the opposite inclusion follows from the above description of \bar{K}_* and the fact that $\omega \in A\Omega(A)$ implies $\omega = \mathbf{1}\omega$.

(ii) Check of (2.9): the l.h.s. equals

$$(2.13) \quad \mathbf{1}d(\omega\psi) - \mathbf{1}(d\omega)\psi - (-1)^{\partial\omega}\omega\mathbf{1}d\psi = (-1)^{\partial\omega}\omega(d\mathbf{1})\psi$$

thus belongs to $\bar{K}_* \cap A\Omega(A) = K_*$.

¹³In fact the product of $\Omega(A)$ reduces to the “concatenation product” \times more generally for the first factor within J_* and the second factor any element in $A\Omega(A)$. Note, on the other hand, that I_1 and K_1 coincide via (1.9) with the respective I and K introduced above, cf. (1.21).

¹⁴For shortness we use the word “reordering” to mean: writing as a sum of “monomials”.

Check of (2.10): we have

$$(2.14) \quad \Delta^2 \psi = \mathbf{1}d(\mathbf{1}d\psi) = \mathbf{1}(d\mathbf{1})d\psi$$

also contained in $\overline{K}_* \cap A\Omega(A) = K_*$.

(iii) Let $\omega = a_0 da_1 \cdots da_n = a_0 \otimes a_1 \otimes \cdots \otimes a_n \in J_n$ and $\psi = b_0 db_1 \cdots db_m = a_0 \otimes b_1 \otimes \cdots \otimes b_m \in A\Omega(A)^m$. Applying (1.6) and (1.5) with $x = b_0$ (where all but the last term vanish owing to (2.7), (2.8)) we see that the product $\omega\psi$ reduces to (2.11), whose r.h.s. clearly vanishes under all maps m_i^{n+m} for $i = 1, 2, \dots, n - 1$ and, if $\psi \in J_*$, also for $i = n, \dots, n + m - 1$.

PROPOSITION. *Let A be a unital $\mathbf{Z}/2$ -graded complex algebra, with Δ, K_* and J_* as in the last lemma. We have a split short exact sequence*

$$(2.15) \quad 0 \rightarrow K_* \hookrightarrow A\Omega(A) \begin{array}{c} \xrightarrow{\pi} \\ \xleftarrow{\theta} \end{array} \Omega A \rightarrow 0$$

where¹⁵ π is the homomorphism of algebras, restriction to $A\Omega(A)$ of the homomorphism π in the diagram (cf. (1.17))

$$(2.16) \quad \begin{array}{ccc} & \Omega(A) & \\ & \uparrow \searrow \pi & \\ & A & \begin{array}{c} \xrightarrow{\text{id}} \\ \xrightarrow{d} \end{array} \Omega A \end{array}$$

i.e.

$$(2.16a) \quad \pi(a_0 da_1 \cdots da_n) = a_0 \underline{d}a_1 \cdots \underline{d}a_n.$$

The lift θ of $\pi(\theta \circ \pi = \text{id}_{\Omega A})$ is the homomorphism of algebras defined by the diagram (cf. (1.30)).

$$(2.17) \quad \begin{array}{ccc} & \Omega A & \\ & \uparrow \searrow \theta & \\ & A & \begin{array}{c} \xrightarrow{\text{id}} \\ \xrightarrow{D} \end{array} \Omega(A) \end{array}$$

i.e.

$$(2.17a) \quad \theta(a_0 \underline{d}a_1 \cdots \underline{d}a_n) = a_0 D a_1 \cdots D a_n$$

¹⁵In the remainder of this section we will denote by \underline{d} the differential of ΩA to distinguish it from the differential d of $\Omega(A)$.

where

$$(2.18) \quad Da = \mathbf{1}da\mathbf{1} = \mathbf{1}da - (-1)^{\partial a}ad\mathbf{1}.$$

One has moreover

$$(2.19) \quad \text{Im } \theta = J_* \quad (\theta(\Omega(A)^n) = J_n);$$

consequently $\phi_* = \theta \circ \pi$ and $\phi_*^\perp = \text{id} - \phi_*$ (resp. their restriction ϕ_n and ϕ_n^\perp to $A\Omega(A)^n$) are the projections in $A\Omega(A)$ (resp. $A\Omega(A)^n$) corresponding to the direct sum decomposition

$$(2.20) \quad \begin{cases} A\Omega(A) = J_* \oplus K_* \\ (\text{resp. } A\Omega(A)^n = J_n \oplus K_n). \end{cases}$$

Finally we have the intertwining properties

$$(2.21) \quad \pi \circ \Delta = \underline{d} \circ \pi,$$

$$(2.22) \quad \theta \circ \underline{d} = D \circ \theta,$$

$$(2.23) \quad \phi_* \circ \Delta = D \circ \theta.$$

It follows in particular that if one defines

$$(2.24) \quad D(a_0Da_1 \cdots Da_n) = Da_0Da_1 \cdots Da_n.$$

(J_*, D) , with the concatenation product and the N, ∂ and ∂_0 gradings inherited from $A\Omega(A)$, becomes a bigraded differential algebra isomorphic to $(\Omega A, \underline{d})$.

Proof. The homomorphism π being onto gives rise to a short exact sequence. The definition (2.17) of the homomorphism θ is justified by the fact that D is a graded derivation, vanishing on the unit, as one immediately checks:

$$(2.25) \quad \begin{cases} D(ab) = (Da)b + (-1)^{\partial a}aDb, & a \in A, \quad b \in A, \\ D\mathbf{1} = 0. \end{cases}$$

We note that the r.h.s. of (2.17a) is unchanged if one replaces throughout D and D' where

$$(2.26) \quad D'a = da - (-1)^{\partial a}ad\mathbf{1} = (da)\mathbf{1};$$

indeed both procedures lead to the expression

$$(2.27) \quad a_0(da_1)\mathbf{1}(da_2)\mathbf{1} \cdots \mathbf{1}(da_n)\mathbf{1},$$

the use of D' showing that $\phi_n^\perp(a_0da_1 \cdots da_n)$ is a sum of terms of the type

$$(2.28) \quad \pm a_0 \cdot \text{product of factors } da_i \text{ and factors } a_k d\mathbf{1}$$

(in arbitrary order, with at least one factor of the type $a_k d\mathbf{1}$, two such factors being never contiguous).

We now show the inclusion (hence equalities)

$$(2.29) \quad K_n \subset \text{Ker } \phi_n \subset \text{Im } \phi_n^\perp \subset K_n.$$

The first inclusion follows from $K_n \subset \text{Ker } \pi$, immediate from (2.16a) since $d\mathbf{1} = 0$; the second is obvious the third follows from the noted fact that $\text{Im } \phi_n^\perp$ is a sum of terms of the type (2.28), where the last factor $d\mathbf{1}$ to the right persists whilst “reordering”. Since $\phi_n^2 - \phi_n = \phi_n \phi_n^\perp$, we established that ϕ_n and ϕ_n^\perp are idempotent.

We next show that $\text{Im } \phi_n \subset J_n$ by induction. Equality holds for $n = 1$. Induction step: for a monomial $\omega \in J_n$ and $a_{n+1} \in A$:

$$(2.30) \quad \omega D' a_{n+1} = \omega \times D a_{n+1} = \omega \times (\mathbf{1} d a_n - (-1)^{\partial a_{n+1}} a_{n+1} d \mathbf{1})$$

obviously vanishes under the maps m_i , $i = 1, 2, \dots, n$, but also under m_{n+1} , since

$$(2.31) \quad m(x d a_n - (-1)^{\partial a_{n+1}} x a_{n+1} d \mathbf{1}) = 0.$$

The converse inequality $\text{Ker } \phi_n^\perp \supset J_n$ follows from the fact that for

$$(2.32) \quad \omega = \sum_{i=1}^k a_0^i \cdots d a_1^i \cdots d a_n^i \in J_n$$

each of the contributions to $\phi_n^\perp \omega$ corresponding to one given type of product (2.28) individually vanishes: indeed we obtain (2.28) from $a_0 d a_1 \cdots d a_n$ by replacing successively factors $d a_k$ of increasing k by factors $a_k d \mathbf{1}$: however, the first replacement applied to (2.20) yields zero owing to vanishing of ω under all contiguous diagonal maps.

Since (2.21) and (2.22) (hence (2.23)) immediately follow from (2.16a), (2.17a) and (2.24), we completed our proof.

To summarise we have found a variety of equivalent descriptions of the bigraded differential algebra $(\Omega A, \underline{d})$:

- as the tensorial algebra of the A -bimodule J under the product \otimes_A and the differential extending (1.24);
- as linear combinations of expressions

$$(2.33) \quad a_0 \underline{d} a_1 \cdots \underline{d} a_n \cong a_0 \otimes \hat{a} \otimes \cdots \otimes \hat{a}_n, \quad a_0, a_1, a_n \in A,$$

where $\hat{a} = a \pmod{\mathbf{C}\mathbf{1}}$, and with multiplication dictated by the fact that \underline{d} is a differential vanishing on the unit;

- as the quotient of $A\Omega(A)$ by its ideal K_* , and Δ as a “differential modulo K_* ”, cf. (2.9), (2.10);

- most explicitly, as the subalgebra $J_* \subset A\Omega(A)$, with multiplication, the “concatenation product” \times , and differential D , cf. (2.6), (2.7), (2.11), (2.18), (2.14).

3. The $\mathbf{Z}/2$ -graded case in terms of the ungraded case. The following proposition contains two statements pertaining to the two kinds of differential envelope¹⁶.

3.1. PROPOSITION. *Let $A = A^0 \oplus A^1$ be a unital $\mathbf{Z}/2$ -graded algebra (resp. a $\mathbf{Z}/2$ -graded algebra), with \underline{A} the corresponding ungraded algebra¹⁷.*

And consider in parallel the differential envelopes¹⁸ $(\Omega A, d)$ (resp. $(\tilde{\Omega}(A), d)$) generated by monomials

$$(3.1) \quad a_0 da_1 \cdots da_n, \quad \begin{cases} a_0 \in A \text{ (resp. } \in \tilde{A}), \\ a_1, \dots, a_n \in A, \end{cases}$$

and $(\Omega \underline{A}, d)$ (resp. $(\tilde{\Omega}(\underline{A}), d)$) generated by tensors

$$(3.2) \quad a_0 \otimes a_1 \otimes \cdots \otimes a_n, \quad \begin{cases} a_0 \in A \text{ (resp. } \in \tilde{A}), \\ a_1, \dots, a_n \in A \text{ mod } \mathbf{C1} \text{ (resp. } \in A), \end{cases}$$

with the product of $\Omega \underline{A}$ (resp. $\tilde{\Omega}(\underline{A})$) denoted by \odot (hence we have, for $x \in A$ (resp. $x = \lambda \tilde{\mathbf{1}} + \underline{x} \in \tilde{A}$, $\lambda \in \mathbf{C}$))

$$(3.3) \quad \begin{cases} x \odot (a_0 \otimes a_1 \otimes \cdots \otimes a_n) = xa_0 \otimes a_1 \otimes \cdots \otimes a_n, \\ (a_0 \otimes a_1 \otimes \cdots \otimes a_n) \odot x \\ = \sum_{i=0}^n (-1)^{n+i} a_0 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_n \otimes \{x \text{ (resp. } \underline{x})\} \\ + a_0 \otimes a_1 \otimes \cdots \otimes a_{n-1} \otimes a_n x, \end{cases}$$

where the products in the r.h.s. have to be taken mod $\mathbf{C1}$ in the case of $\Omega \underline{A}$.

If we equip $\Omega \underline{A}$ (resp. $\tilde{\Omega}(\underline{A})$) with the new bilinear product given by

$$(3.4) \quad S \circ T = (-1)^{m\partial_0 S} S \odot T,$$

where $S, T \in \Omega \underline{A}$ (resp. $\tilde{\Omega}(\underline{A})$) are respectively of intrinsic grade $\partial_0 S$ and of \mathbf{N} -grade m , we then have that $(\Omega A, d)$ and $(\Omega \underline{A}, d)$ (resp. $(\tilde{\Omega}(A), d)$

¹⁶The second statement follows from the first via $\tilde{\Omega}(A) = \Omega \tilde{A}$.

¹⁷I.e. $\underline{A} = A$ as an algebra, with the trivial grading $A^0 = A, A^1 = \{0\}$.

¹⁸ ΩA and $\Omega \underline{A}$ (resp. $\tilde{\Omega}(A)$ and $\tilde{\Omega}(\underline{A})$) coincide as \mathbf{N} -, ∂_0 -, and ∂ -graded vector spaces.

and $(\tilde{\Omega}(\underline{A}), d)$ are isomorphic bigraded differential algebras¹⁹, with the isomorphism given by

$$(3.5) \quad \begin{cases} i(a_0 da_1 \cdots da_n) = (-1)^n \sum_{k=0}^n \partial a_k + \sum_{k \text{ odd}} \partial a_k a_0 \otimes a_1 \otimes \cdots \otimes a_n, \\ a_0 da_1 \cdots da_n \in \Omega A \text{ (resp. } \tilde{\Omega}(A)), \\ a_0 \in A^\cdot \text{ (resp. } \tilde{A}^\cdot, a_1, \dots, a_n \in A^\cdot). \end{cases}$$

REMARK. With $\omega \in \Omega A$ (resp. $\in \tilde{\Omega}(A)$) the “monomial”

$$(3.6) \quad \begin{cases} \omega = a_0 da_1 \cdots da_n, \\ a_0 \in A \text{ (resp. } \in \tilde{A}^\cdot), a_1, \dots, a_n \in A^\cdot \end{cases}$$

and defining

$$(3.7) \quad \partial^\pm \omega = \sum_{\substack{k \text{ even} \\ \text{odd}}} \partial a_k, \quad \Delta \omega = n \partial_0 \omega + \partial^- \omega,$$

the sign factor $(-1)^{\Delta \omega}$ appearing r.h.s. of (3.5) corresponds to “extraction of the n (grade one) symbols d towards the left” whilst effecting the passage

$$(3.8) \quad a_0 da_1 \cdots da_n \rightarrow a_0 \otimes a_1 \otimes \cdots \otimes a_n.$$

In “ $\mathbb{Z}/2$ -spirit”, the map i hence appears as more natural than the former identification (1.9). Note also that $\Delta \omega$ is coherently defined for monomials in ΩA , since the intrinsic grade passes to the quotient $\hat{a}_k = a_k \text{ mod } \mathbf{C1}$, $a_k \in A$. Caution: $\Delta \omega$ is only defined for monomials, and does not make sense in general for homogeneous elements. We note that we have, for ω as in (3.6)

$$(3.9) \quad \Delta \omega = \begin{cases} \partial^- \omega & \text{for } n \text{ even,} \\ \partial^+ \omega & \text{for } n \text{ odd,} \end{cases}$$

$$(3.10) \quad \Delta(xd\omega) = \Delta \omega + (n + 1) dx, \quad x \in A^\cdot \text{ (resp. } x \in \tilde{A}^\cdot),$$

$$(3.11) \quad \begin{cases} \Delta(\omega d\psi) = \Delta \omega + \Delta \psi + (m + 1) \partial_0 \psi, \\ \psi = b_0 db_1 \cdots db_m \in \Omega \underline{A} \text{ (resp. } \tilde{\Omega} \underline{A}). \end{cases}$$

The claim in Proposition 3.1 that $(\Omega \underline{A}, d)$ (resp. $(\tilde{\Omega}(\underline{A}), d)$) are bigraded differential algebras results by taking $(\Omega, d) = (\Omega \underline{A}, d)$ (resp. $(\tilde{\Omega}(\underline{A}), d)$) in the following general fact.

¹⁹ $\Omega \underline{A}$ (resp. $\tilde{\Omega}(\underline{A})$) are equipped with the new product (3.4); the differential d and the gradings ∂, ∂_0 , remaining unchanged. Note that (3.5) defines i coherently as a linear map since the sign factor r.h.s. amounts to applying the linear operators $\{\prod_{k=0}^n \varepsilon_k\}^n \prod_{k \text{ odd}} \varepsilon_k$ the operator $\text{id} \otimes \cdots \otimes \text{id} \otimes \varepsilon \otimes \text{id} \otimes \cdots \otimes \text{id}$, where the k th factor ε is the grading involution of A or $A \text{ mod } \mathbf{C1}$ (resp. of \tilde{A} or A).

LEMMA. Let $\Omega = \sum_{n \in \mathbf{N}}^{\oplus} (\Omega^n, d)$ be an \mathbf{N} -graded differential algebra, with product \odot , and assume the existence of an intrinsic $\mathbf{Z}/2$ -grading ∂_0 commuting with the \mathbf{N} -grading

$$(3.12) \quad \begin{cases} \Omega^n = \Omega^{n(+)} + \Omega^{n(-)} \\ \text{with } \begin{cases} \Omega^{n(\pm)} \odot \Omega^{m(+)} = \Omega^{n+m(\pm)}, \\ \Omega^{n(\pm)} \odot \Omega^{m(-)} = \Omega^{n+m(\mp)}, \end{cases} \end{cases} \quad n, m \in \mathbf{N},$$

and with the differential:

$$(3.13) \quad d\Omega^{n(\pm)} \subset \Omega^{n+1(\pm)}, \quad n \in \mathbf{N}.$$

Then (Ω, d) becomes a bigraded differential algebra (with total grading ∂ the sum of the intrinsic and the \mathbf{N} -grading) under the new product \cdot specified as follows:

$$(3.14) \quad \begin{cases} \omega \cdot \psi = (-1)^{m\partial_0\omega} \omega \odot \psi, \\ \omega, \psi \in \Omega, \omega \text{ of intrinsic grade } \partial_0\omega, \psi \text{ of } \mathbf{N}\text{-grade } m. \end{cases}$$

To prove this lemma, we have to check that the product \cdot is associative and makes d a ∂ -graded derivative. Now let $\omega, \psi, \theta \in \Omega$, with ω, ψ of respective intrinsic grades $\partial_0\omega, \partial_0\psi$, and ψ, θ of respective \mathbf{N} -grades m, p . We have on the one hand

$$(3.15) \quad \begin{aligned} (\omega \cdot \psi) \cdot \theta &= (-1)^{m\partial_0\omega} (\omega \odot \psi) \cdot \theta \\ &= (-1)^{m\partial_0\omega + p(\partial_0\omega + \partial_0\psi)} \omega \odot \psi \odot \theta \\ &= (-1)^{(m+p)\partial_0\omega} \omega \odot (\psi \cdot \theta) = \omega \cdot (\psi \cdot \theta) \end{aligned}$$

and on the other hand

$$(3.16) \quad \begin{aligned} d(\omega \cdot \psi) &= (-1)^{m\partial_0\omega} d(\omega \odot \psi) \\ &= (-1)^{m\partial_0\omega} [(d\omega) \odot \psi] + (-1)^n \omega \odot d\psi \\ &= d\omega \cdot \psi + (-1)^{n+\partial_0\omega} \omega \cdot d\psi. \end{aligned}$$

The remainder of Proposition 3.1 now follows from the fact that the map i (obviously linear bijective) turns the product of ΩA (resp. $\tilde{\Omega}(A)$) into the new product of $\Omega \underline{A}$ (resp. $\tilde{\Omega}(\underline{A})$): we have

$$(3.17) \quad i(\omega\psi) = i(\omega) \cdot i(\psi), \quad \omega, \psi \in \Omega A \text{ (resp. } \in \tilde{\Omega}(A))$$

which it suffices to check for ω and ψ “monomials”²⁰ We first do

²⁰The following argument applies to both the case of ΩA and that of $\tilde{\Omega}(A)$. In what follows $n(\omega)$ denotes the \mathbf{N} -grade of ω .

this for $\psi = d\psi'$, ψ' a monomial: we have, with $T(a_0 da_1 \cdots da_n) = a_0 \otimes \cdots \otimes a_n$ and taking account of (3.11),

$$\begin{aligned}
 (3.18) \quad i(\omega d\psi') &= (-1)^{\Delta(\omega d\psi')} T(\omega d\psi') \\
 &= (-1)^{\Delta(\omega d\psi')} T(\omega) \otimes T(\psi') \\
 &= (-1)^{\Delta\omega + \Delta(d\psi') + n(\partial\psi')\partial_0\omega} T(\omega) \odot T(\psi') \\
 &= (-1)^{n(d\psi')\partial_0\omega} i(\omega) \odot i(d\psi') = i(\omega) \cdot i(d\psi').
 \end{aligned}$$

It remains to check (3.17) for $\psi = bd\psi'$, $b \in A'$; ψ' a monomial, however, this boils down to the simpler case of $\psi = b$: the knowledge of (3.17) in that case indeed allows one to infer from (3.18) and the associativity of the product \cdot that one has

$$\begin{aligned}
 (3.19) \quad i(\omega bd\psi') &= i(\omega b) \cdot i(d\psi') = i(\omega) \cdot i(b) \cdot i(d\psi') \\
 &= i(\omega) \cdot i(bd\psi').
 \end{aligned}$$

The remaining proof of (3.17) for $\psi = b \in A'$ will now proceed by induction w.r.t. the \mathbf{N} -grade of ω ²¹.

For $x, y \in A'$ (resp. $x \in \tilde{A}'$, $y \in A$) we have

$$\begin{aligned}
 (3.20) \quad i[y(dx)b] &= i[yd(xb) - (-1)^{\partial x} yx db] \\
 &= (-1)^{\partial y} y \otimes xb - (-1)^{\partial x + \partial(yx)} yx \otimes b \\
 &= (-1)^{\partial y} y \odot (\mathbf{1} \otimes x) \odot b \\
 & \quad (\text{resp. } = (-1)^{\partial y} y \odot (\tilde{\mathbf{1}} \otimes x) \odot b \\
 & = i(y dx) \odot b = i(y dx) \cdot i(b)).
 \end{aligned}$$

We now effect the induction step: assuming (2.19) to hold for ω and $\psi = b$, we check that it holds also for ω replaced by ωdx , $x \in A'$, and $\psi = b$, we have, by what precedes

$$\begin{aligned}
 (3.21) \quad i[\omega(dx)b] &= i[\omega d(xb) - (-1)^{\partial x} \omega x db] \\
 &= i(\omega) \cdot i(d(xb)) - (-1)^{\partial x} i(\omega x) \cdot i(db) \\
 &= i(\omega) \cdot [i(d(xb)) - (-1)^{\partial x} i(xdb)] \\
 &= i(\omega) \cdot i[d(xb)] = i(\omega) \cdot i(dx) \cdot i(b) \\
 &= i(\omega dx) \cdot i(b)
 \end{aligned}$$

completing our proof.

3.2. We shall now briefly show how the replacement of $(\tilde{\Omega}(A), d)$ by $(\tilde{\Omega}(\underline{A}), d)$ equipped with the product (2.5) simplifies the formulae

²¹One could also perform a direct verification based on (1.5) rewritten using the identification (1.9).

of $\mathbf{Z}/2$ -graded cyclic cohomology²². We recall that the latter can be formulated using operators $\beta', \beta, \alpha, \gamma, \rho, \lambda, \Sigma$ on $\Omega(A)$ defined as follows [4], [5]²³: for $\omega \in \Omega$ of total grade $\partial\omega$, and $x \in A^0, a \in A$:

$$(3.22) \quad \begin{cases} \beta'(\omega dx) = (-1)^{\partial\omega} \omega x, \\ \beta'(a) = 0, \quad \beta'(da) = a, \end{cases}$$

$$(3.23) \quad \begin{cases} \alpha(\omega dx) = (-1)^{(1+\partial x)\partial\omega} x\omega, \\ \alpha(a) = 0, \quad \alpha(da) = a, \end{cases}$$

$$(3.24) \quad \beta = \beta' - \alpha,$$

$$(3.25) \quad \begin{cases} \rho(\omega dx) = (-1)^{(1+\partial x)\partial\omega} (dx)\omega, \\ \rho(a) = a, \quad \rho(da) = a, \end{cases}$$

$$(3.26) \quad \gamma = \text{id} - \rho,$$

$$(3.27) \quad \begin{cases} \lambda(\omega dx) = (-1)^{(1+\partial x)(1+\partial\omega)} x d\omega, \\ \lambda(a) = a, \quad \lambda(da) = 0, \end{cases}$$

and, for $a_0 = s\mathbf{1} + \underline{a}_0 \in \tilde{A}, \underline{a}_0, a_1, \dots, a_n \in A, s \in \mathbf{C}$,

$$(3.28) \quad \begin{aligned} \Sigma(a_0 da_1 \cdots da_{n+2}) &= (-1)^{\partial a_1} a_0 a_1 a_2 da_3 \cdots da_{n+2} \\ &+ \sum_{i=2}^{n+1} (-1)^{\partial a_i} (a_0 da_1 \cdots da_{i-1}) a_i a_{i+1} da_{i+2} \cdots da_{n+2} \\ &+ (-1)^{n+1} \sum_{k=1}^{n+1} \partial a_k s (a_1 da_2 \cdots da_{n+1}) a_{n+2}. \end{aligned}$$

If one replaces $(\Omega(A), d)$ by $(\Omega(\underline{A}), d)$ the above operators are replaced by their images which we denote by corresponding underlined symbols:

$$(3.29) \quad \underline{0} = i \circ 0 \circ i^{-1}$$

and we have then, for $T \in \Omega(\underline{A})^n$ of intrinsic grade $\partial_0 T$ and $x \in A$ of grade ∂x

$$(3.30) \quad \underline{\beta}'(T \odot dx) = (-1)^n T \odot x,$$

$$(3.31) \quad \underline{\alpha}(T \odot dx) = (-1)^{n+\partial x \partial_0 T} x \odot T,$$

$$(3.32) \quad \underline{\rho}(T \odot dx) = (-1)^{n+\partial x \partial_0 T} dx \odot T,$$

$$(3.33) \quad \underline{\lambda}(T \odot dx) = (-1)^{n+1+\partial x \partial_0 T} x \odot dT.$$

²²Cf. [1] formula (a'), (b'), p. 60, and [5] p. 47a.

²³With $\Omega(A)^*$ the algebraic dual of $\Omega(A)$, the cyclic cochains, resp. cyclic cocycles are the $\varphi \in \Omega(A)^*$ fulfilling $\varphi \circ \lambda = \varphi (\leftrightarrow \varphi \circ \gamma = \varphi \circ d = 0)$, resp. $\varphi \circ \beta = \varphi \circ d = 0 (\leftrightarrow \varphi$ is a closed graded trace of $\Omega(A))$. The operator Σ is proportional to the transpose of Connes's operator $S: S_\varphi = 2\pi\varphi \circ \Sigma$.

For $a_0 = s\mathbf{1} + \underline{a}_0 \in \tilde{A}^0$, $\underline{a}_0, a_1, \dots, a_{n+2} \in A^0$, $s \in \mathbf{C}$, we have then

$$(3.4) \quad \begin{aligned} \underline{\beta}'(a_0 \otimes a_1 \otimes \cdots \otimes a_{n+1}) \\ = \sum_{i=1}^n (-1)^i \underline{a}_0 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_{n+1}, \end{aligned}$$

$$(3.5) \quad \begin{aligned} \underline{\alpha}(a_0 \otimes a_1 \otimes \cdots \otimes a_{n+1}) \\ = (-1)^{n+\partial a_{n+1} + \sum_{k=0}^n \partial a_k} a_{n+1} \otimes a_0 \otimes \cdots \otimes a_n, \end{aligned}$$

$$(3.36) \quad \begin{aligned} \underline{\lambda}(a_0 \otimes a_1 \otimes \cdots \otimes a_{n+1}) \\ = (-1)^{n+1+\partial a_{n+1}} \sum_{k=1}^n \partial a_k a_{n+1} \otimes \underline{a}_0 \otimes a_1 \otimes \cdots \otimes a_n, \end{aligned}$$

and

$$(3.37) \quad \begin{aligned} \underline{\Sigma}(a_0 \otimes a_1 \otimes \cdots \otimes a_{n+2}) \\ = \sum_{i=1}^{n+1} (a_0 \otimes a_1 \otimes \cdots \otimes a_{i-1}) \odot a_i a_{i+1} \otimes a_{i+2} \otimes \cdots \otimes a_{n+2} \\ + (-1)^{n+1} s(a_1 \otimes a_2 \otimes \cdots \otimes a_{n+1}) \odot a_{n+2}. \end{aligned}$$

As we see the formulae (3.34) and (3.37) are free of sign factors depending upon the intrinsic grading—a notable simplification w.r.t. the formulae relative to β' and Σ (cf. (1.5) and (3.28)).

Acknowledgments. The authors express hearty thanks to Professor M. Berger for the hospitality of IHES, where part of this work was completed.

REFERENCES

- [1] A. Connes, *Non commutative differential geometry I. The Chern Character in K-homology II. De Rham Cohomology and Non commutative Algebra*. Pub. Math. IHES 257 (1985).
- [2] M. Karoubi, *Connections, courbures et classes caractéristiques en K-théorie algébrique*, Canad. Math. Soc. Proc., Vol. 2, part 2, Part 1. 19 (1982). *Homologie cyclique et K-théorie*, Astérisque no. 149, Société Mathématique de France (1987).
- [3] W. Arveson, *The harmonic analysis of automorphism groups*, Operator algebras and applications. Proc. Symposia Pure Math. 38 part I 199 (1982).
- [4] D. Kastler, *Z/2-graded cohomology within the differential envelope*, Contemporary Mathematics 62, 221 (1987).

- [5] ———, *Cyclic Cohomology within the Differential Envelope; An Introduction to Alain Connes' Non Commutative Differential Geometry*. Travaux en cours. Hermann, Paris (1987).

Received February 26, 1988. Research of the first author was supported in part by the U.S. Dept. of Energy under Grant no. DE-FG02-84ER40158 with Harvard University.

CENTRE DE PHYSIQUE THEORIQUE
CNRS-LUMINY, CASE 907
F-13288, MARSEILLE CEDEX 09 (FRANCE)

Current address: R. Coquereaux
Gordon McKay Laboratory
Harvard University
Cambridge, MA 02138
USA

