

# Pacific Journal of Mathematics

**THE  $n$ -DIMENSIONAL ANALOGUE OF THE CATENARY:  
EXISTENCE AND NONEXISTENCE**

ULRICH DIERKES AND GERHARD HUISKEN

# THE $n$ -DIMENSIONAL ANALOGUE OF THE CATENARY: EXISTENCE AND NON-EXISTENCE

U. DIERKES AND G. HUISKEN

We study “heavy”  $n$ -dimensional surfaces suspended from some prescribed  $(n - 1)$ -dimensional boundary data. This leads to a mean curvature type equation with a non-monotone right hand side. We show that the equation has no solution if the boundary data are too small, and, using a fixed point argument, that the problem always has a smooth solution for sufficiently large boundary data.

Consider a material surface  $M$  of constant mass density which is suspended from an  $(n - 1)$ -dimensional surface  $\Gamma$  in  $\mathbb{R}^n \times \mathbb{R}^+$ ,  $\mathbb{R}^+ = \{t > 0\}$ , and hangs under its own weight. If  $M$  is given as graph of a regular function  $u: \Omega \rightarrow \mathbb{R}^+$  on a domain  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , then  $u$  provides an equilibrium for the potential energy  $\mathcal{E}$  under gravitational forces,

$$\mathcal{E}(u) = \int_{\Omega} u \sqrt{1 + |Du|^2}.$$

Thus  $u$  solves the Dirichlet problem

$$(1) \quad \operatorname{div} \left\{ \frac{u \cdot Du}{\sqrt{1 + |Du|^2}} \right\} = \sqrt{1 + |Du|^2} \quad \text{in } \Omega,$$

$$u = \varphi \quad \text{on } \partial\Omega$$

The corresponding variational problem

$$(2) \quad \int_{\Omega} u \sqrt{1 + |Du|^2} + \frac{1}{2} \int_{\partial\Omega} |u^2 - \varphi^2| d\mathcal{H}_{n-1} \rightarrow \min$$

in the class

$$BV_2^+(\Omega) := \{u \in L_2(\Omega) : u \geq 0, u^2 \in BV(\Omega)\}$$

has been solved by Bemelmans and Dierkes in [BD]. It was shown in [BD, Theorem 7] that the coincidence set  $\{u = 0\}$  of a minimizer  $u$  is non-empty provided that

$$(3) \quad |\varphi|_{\infty, \partial\Omega} < \frac{|\Omega|}{\mathcal{H}_{n-1}(\partial\Omega)},$$

where  $|\Omega|$  denotes the Lebesgue measure of  $\Omega$  and  $\mathcal{H}_n$  denotes  $n$ -dimensional Hausdorff measure.

We want to show here that (1) has *no* solution in case (3) holds, whereas (1) has *always* a solution for sufficiently large boundary data. More precisely we prove the following existence–non-existence result.

**THEOREM.** *Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be a bounded domain of class  $C^{2,\alpha}$ ,  $\alpha > 0$ , with non-negative (inward) mean curvature. Suppose  $\varphi \in C^{2,\alpha}(\overline{\Omega})$  satisfies*

$$(4) \quad k_0 := \inf_{\partial\Omega} \varphi \geq c(n) \left(1 + \sqrt{2^{n+1}}\right)^2 |\Omega|^{1/n},$$

where  $c(n) = n^{-1} \omega_n^{-1/n}$  is the isoperimetric constant. Then the Dirichlet problem (1) has a global regular solution  $u \in C^{2,\alpha}(\overline{\Omega})$ . Moreover, if  $u \in C^{0,1}(\overline{\Omega})$  is a weak positive solution of (1) with Lipschitz constant  $L$ , then we have

$$(5) \quad h := \sup_{\partial\Omega} \varphi \geq (1 + L^{-2})^{1/2} \frac{|A|}{\mathcal{H}_{n-1}(\partial A)}$$

for every Caccioppoli set  $A \subset \Omega$ .

Since  $c(n)$  is the isoperimetric constant, we have

$$c(n)|\Omega|^{1/n} \geq \frac{|\Omega|}{\mathcal{H}_{n-1}(\partial\Omega)}$$

and therefore it is an interesting question whether our existence result remains true if we replace (4) with an inequality of the form

$$k_0 \geq \text{const.} \frac{|\Omega|}{\mathcal{H}_{n-1}(\partial\Omega)}.$$

The proof of the theorem is based on a priori bounds for solutions to the related problem

$$\Delta u - \frac{D_i u D_j u}{1 + |Du|^2} D_i D_j u = f^{-1},$$

which enable us to apply a fixed point argument. Notice that the operator

$$\Delta - \frac{D_i u D_j u}{1 + |Du|^2} D_i D_j = (1 + |Du|^2) \cdot \Delta_M$$

where  $\Delta_M$  is the Laplace-Beltrami operator on  $M = \text{graph } u$ .

Let us make some comments on the literature. For two dimensional parametric surfaces in  $\mathbb{R}^3$  the existence problem has been investigated

by Böhme, Hildebrandt and Tausch [BHT]. To our knowledge the first existence result for the Dirichlet problem (1), in case  $n = 2$ , is due to Dierkes [D1]. The variational problem (2) is solved in [BD]. It is shown in [D2] that minima  $u$  of (2) are regular up to the boundary provided only the boundary is mean curvature convex. A non-existence result of a different type has been proved by J. C. C. Nitsche in [N].

*Proof.* We consider regular solutions  $u_f \in C^{2,\alpha}(\overline{\Omega})$  of the related problem

$$(6) \quad \begin{aligned} \sqrt{1 + |Du|^2} \operatorname{div} \frac{Du}{\sqrt{1 + |Du|^2}} &= f^{-1} && \text{in } \Omega, \\ u &= \varphi && \text{on } \partial\Omega, \end{aligned}$$

where  $f \in C^{1,\alpha}(\overline{\Omega})$  and  $0 < d \leq f$ . As a first step we establish a priori bounds for  $\sup_{\Omega} u$  and  $\inf_{\Omega} u$ .

**LEMMA.** *Let  $u_f \in C^{2,\alpha}(\overline{\Omega})$  be a solution to the Dirichlet problem (6). If*

$$f \geq d \geq \left(1 + \sqrt{2^{n+1}}\right) c(n) |\Omega|^{1/n}$$

and

$$k_0 = \inf_{\partial\Omega} \varphi \geq \left(1 + \sqrt{2^{n+1}}\right)^2 c(n) |\Omega|^{1/n},$$

then we have  $h \geq u_f \geq d$ .

*Proof of the Lemma.* The first inequality follows immediately from the maximum principle since  $f$  is positive. To prove the second relation we chose  $\delta \geq -k_0$  and put  $w = \min(u + \delta, 0)$ ,  $A(\delta) = \{x \in \Omega : u < -\delta\}$ . Multiplying (6) with  $w$ , integrating by parts and using  $w|_{\partial\Omega} = 0$ , we obtain

$$\begin{aligned} \int_{\Omega} \frac{|Dw|^2}{\sqrt{1 + |Dw|^2}} &= \int_{A(\delta)} \frac{|w|}{f \sqrt{1 + |Du|^2}}, \quad \text{hence} \\ \int_{\Omega} |Dw| &\leq |A(\delta)| + d^{-1} \int_{A(\delta)} |w|. \end{aligned}$$

We use Sobolev's inequality on the left and Hölder's inequality on the right hand side and get with  $c(n) = n^{-1} \omega_n^{-1/n}$

$$|w|_{n/n-1} \cdot \{c^{-1}(n) - d^{-1} |\Omega|^{1/n}\} \leq |A(\delta)|,$$

where  $|w|_{n/n-1}$  stands for the  $L_{n/n-1}$ -norm of  $w$ . Another application of Hölder's inequality yields

$$(\delta_1 - \delta_2)|A(\delta_1)| \leq \left\{ \frac{c(n)d}{d - c(n)|\Omega|^{1/n}} \right\} |A(\delta_2)|^{1+1/n}$$

for all  $\delta_1 \geq \delta_2 \geq -k_0$ . In view of a well-known lemma due to Stampacchia, [St, Lemma 4.1], this is easily seen to imply

$$|A(-k_0 + 2^{n+1} \cdot c_1 |A(-k_0)|^{1/n})| = 0, \quad \text{where}$$

$$c_1 = \frac{c(n)d}{d - c(n)|\Omega|^{1/n}}.$$

Clearly this means that

$$u \geq k_0 - \frac{2^{n+1}dc(n)|\Omega|^{1/n}}{d - c(n)|\Omega|^{1/n}}.$$

Since  $k_0 \geq (1 + \sqrt{2^{n+1}})d$  and  $d \geq (1 + \sqrt{2^{n+1}})c(n)|\Omega|^{1/n}$  we finally obtain  $u \geq d$ .  $\square$

To derive a gradient estimate at the boundary, we rewrite (6) into

$$(7) \quad (1 + |Du|^2)\Delta u - D_i u D_j u D_i D_j u = f^{-1}(1 + |Du|^2).$$

We can then apply the results of Serrin [Se1], see also [GT, Chapter 14.3]. Equation (7) satisfies the structure condition (14.41) in [GT] and the RHS is  $\mathcal{O}(|Du|^2)$ . So we obtain a gradient estimate on the boundary which is independent of  $|Df|$ :

$$\sup_{\partial\Omega} |Du_f| \leq c_2 = c_2(n, \Omega, h, |\varphi|_{2,\Omega}),$$

provided only that  $\partial\Omega$  has non-negative (inward) mean curvature.

It is not possible to derive interior gradient estimates independent of  $|Df|$ , but we can prove

$$(8) \quad \sup_{\Omega} |Du_f| \leq \max \left\{ 2, \frac{1}{4} \sup_{\Omega} |Df|, 2e^{4(hd^{-1}-1)} \sup_{\partial\Omega} |Du_f| \right\},$$

which will be sufficient for our fixed point argument. Estimate (8) can be obtained from a careful analysis of the structure conditions in [GT, Chapter 15]. Here we present a selfcontained proof, using the geometric nature of equation (6). For a similar procedure we refer to [K].

In the following let  $v = (1 + |Du|^2)^{1/2}$  and denote by  $H$  and  $\Delta$  the mean curvature and the Laplace-Beltrami operator on  $M = \text{graph } u$  respectively. Then equation (6) takes the form

$$(9) \quad v^2 \Delta u = f^{-1} \Leftrightarrow H = f^{-1} v^{-1}.$$

Let  $\tau_1, \tau_2, \dots, \tau_n, \nu$  be an adapted local orthonormal frame on  $M$ , such that  $\nu$  is the upper unit normal and

$$\nabla_i \nu = -h_{il} \tau_l, \quad \nabla_i \tau_j = h_{ij} \nu,$$

where  $\nabla_i$  is the tangential derivative with respect to  $\tau_i$  and  $h_{il}$  is the second fundamental form. Then we get for  $v = (1 + |Du|^2)^{1/2} = \langle \nu, e_{n+1} \rangle^{-1}$  the Jacobi-Codazzi equation

$$\begin{aligned} \Delta v &= \nabla_i \nabla_i \langle \nu, e_{n+1} \rangle^{-1} = \nabla_i (v^2 \langle h_{il} \tau_l, e_{n+1} \rangle) \\ &= |A|^2 v + 2v^{-1} |\nabla v|^2 + v^2 \langle \nabla H, e_{n+1} \rangle, \end{aligned}$$

where  $|A|^2 = h_{il} h^{il}$ . Now (9) implies

$$(10) \quad \Delta v = |A|^2 v + 2v^{-1} |\nabla v|^2 - f^{-2} v \langle \nabla f, e_{n+1} \rangle - f^{-1} \langle \nabla v, e_{n+1} \rangle.$$

If we now extend all functions from  $M$  to  $\mathbb{R}^{n+1}$  by

$$f(\hat{x}, x_{n+1}) = f(\hat{x})$$

such that

$$(11) \quad \begin{aligned} \nabla f &= Df - \nu \langle Df, \nu \rangle, D_{n+1} f = 0 \quad \text{and} \\ \langle \nabla f, e_{n+1} \rangle &= -v^{-1} \langle Df, \nu \rangle \end{aligned}$$

then we derive from (10) and (11)

$$(12) \quad \Delta v \geq 2v^{-1} |\nabla v|^2 - f^{-1} \langle \nabla v, e_{n+1} \rangle - f^{-2} |Df|.$$

Next we compute for  $\alpha > 0$  and  $g = e^{\alpha u} \cdot v$  the inequality

$$\begin{aligned} \Delta g &\geq e^{\alpha u} \{ 2v^{-1} |\nabla v|^2 - f^{-1} \langle \nabla v, e_{n+1} \rangle - f^{-2} |Df| \\ &\quad + 2\alpha \nabla_i v \nabla_i u + \alpha v \Delta u + v \alpha^2 |\nabla u|^2 \}. \end{aligned}$$

Using again the equation (9) and

$$\nabla_i g = \nabla_i v e^{\alpha u} + \alpha v e^{\alpha u} \nabla_i u$$

we obtain

$$\begin{aligned} \Delta g &\geq 2v^{-1} \nabla_i v \nabla_i g - f^{-1} \langle \nabla g, e_{n+1} \rangle + \alpha f^{-1} e^{\alpha u} v \langle \nabla u, e_{n+1} \rangle \\ &\quad - f^{-2} |Df| e^{\alpha u} + \{ v^{-1} \alpha f^{-1} + v \alpha^2 |\nabla u|^2 \} e^{\alpha u}. \end{aligned}$$

In view of relation (11) we finally conclude

$$\Delta g \geq 2v^{-1} \nabla_i v \nabla_i g - f^{-1} \langle \nabla g, e_{n+1} \rangle + \{ \alpha^2 |\nabla u|^2 - \alpha f^{-1} - v^{-1} f^{-2} |Df| \} g$$

Now let again  $d \leq f \leq h$  and choose  $\alpha = 4d^{-1}$ . Then, since

$$|\nabla u|^2 = \frac{|Du|^2}{1 + |Du|^2} \geq \frac{1}{2} \quad \text{for } |Du| \geq 1,$$

we see that  $g$  cannot have an interior maximum if

$$v \geq \max \left\{ 2, \frac{1}{4} \sup_{\Omega} |Df| \right\}.$$

Therefore we get the estimate

$$\sup_{\Omega} v \leq \max \left\{ 2, \frac{1}{4} \sup_{\Omega} |Df|, e^{4(hd^{-1}-1)} \sup_{\partial\Omega} v \right\}$$

yielding (8).

To prove existence of a solution to equation (1) we now define the set

$$\mathcal{M} := \left\{ f \in C^{1,\alpha}(\overline{\Omega}) : d \leq f \leq h, \sup_{\Omega} |Df| \leq M \right\}$$

for  $M > 0$  large and consider the operator

$$\begin{aligned} T: \mathcal{M} &\rightarrow C^{1,\alpha}(\overline{\Omega}), \\ f &\rightarrow u_f. \end{aligned}$$

In view of our estimates for  $u_f$  and  $|Du_f|$  we may then choose  $M$  so large that

$$T(\mathcal{M}) \subset \mathcal{M}.$$

Moreover, standard theory ensures that  $T$  is continuous and  $T(\mathcal{M})$  is precompact. So we can apply Schauder's fixed point theorem, see e.g. ([GT], Cor. 11.2) to obtain the existence of a regular  $u \in C^{2,\alpha}(\overline{\Omega})$  satisfying (1).

To prove the necessary conditions (5) we proceed similarly as in [G]. To this end let  $A \Subset \Omega$  have finite perimeter  $\mathbb{M}(\partial A)$ . There exists a sequence of positive functions  $\varphi_k \in C_c^1(\Omega)$  such that  $\varphi_k \rightarrow \varphi_A$  in  $L_1(\Omega)$ , and

$$\int_{\Omega} |D\varphi_k| \rightarrow \mathbb{M}(\partial A),$$

where  $\varphi_A$  denotes the characteristic function of the set  $A$ .

We test (1) with  $\varphi_k$  and integrate,

$$(13) \quad \int_{\Omega} \left\{ \frac{u Du D\varphi_k}{\sqrt{1 + |Du|^2}} + \varphi_k \sqrt{1 + |Du|^2} \right\} dx = 0.$$

Now, since  $u \in \text{Lip}(\overline{\Omega})$  it follows from standard regularity theory that  $u \in C^\infty(\Omega)$  and therefore

$$\text{div} \frac{Du}{\sqrt{1 + |Du|^2}} \geq 0 \quad \text{on } \Omega, \text{ whence } u \leq h.$$

Using this in (13) we get

$$\int_{\Omega} \varphi_k dx \leq \frac{h \cdot L}{\sqrt{1 + L^2}} \int_{\Omega} |D\varphi_k|$$

and, letting  $k \rightarrow \infty$ ,

$$|A| \leq \frac{h \cdot L}{\sqrt{1 + L^2}} \mathbb{M}(\partial A), \quad \text{or}$$

$$h \geq \{1 + L^{-2}\}^{1/2} \frac{|A|}{\mathbb{M}(\partial A)}.$$

The general case follows by an approximation argument, using the fact that

$$\mathbb{M}(\partial[A \cap \Omega_\varepsilon]) \rightarrow \mathbb{M}(\partial A) \quad \text{as } \varepsilon \rightarrow 0,$$

where

$$\Omega_\varepsilon := \{x \in \Omega : \text{dist}(x, \partial\Omega) > \varepsilon\}.$$

This completes the proof of the theorem.

**REMARK.** With the same method we could as well deal with the integral

$$\int_{\Omega} u^\gamma \sqrt{1 + |Du|^2}, \quad \gamma > 0,$$

the Euler equation of which is given by

$$\text{div} \frac{Du}{\sqrt{1 + |Du|^2}} = \frac{\gamma}{u \sqrt{1 + |Du|^2}}.$$

Clearly, in this case the constants appearing in the theorem would depend on  $\gamma$  too, however we shall not dwell on this.

#### REFERENCES

- [BD] J. Bemelmans and U. Dierkes, *On a singular variational integral with linear growth I: existence and regularity of minimizers*, Arch. Rat. Mech. Anal., **100** (1987), 83–103.
- [BHT] R. Böhme, S. Hildebrandt and E. Tausch, *The two dimensional analogue of the catenary*, Pacific J. Math., **88** (1980), 247–278.



- [D1] U. Dierkes, *A geometric maximum principle, Plateau's problem for surfaces of prescribed mean curvature, and the two-dimensional analogue of the catenary*, in: *Partial Differential Equations and Calculus of Variations*, pp. 116–141, Springer Lecture Notes in Mathematics **1357** (1988).
- [D2] ———, *Boundary regularity for solutions of a singular variational problem with linear growth*, *Arch. Rat. Mech. Anal.*, **105** (1989), 286–298.
- [G] M. Giaquinta, *On the Dirichlet problem for surfaces of prescribed mean curvature*, *Manuscripta Math.*, **12** (1974), 73–86.
- [GT] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, Berlin-Heidelberg-New York, 1977.
- [K] N. Korevaar, *An easy proof of the interior gradient bound for solutions to the prescribed mean curvature equation*, *Proc. Symp. Pure Math.*, **45** (1986), 81–89.
- [LU] O. A. Ladyzhenskaya and N. N. Ural'tseva, *Local estimates for the gradients of solutions of non-uniformly elliptic and parabolic equations*, *CPA M*, **23** (1970), 677–703.
- [N] J. C. C. Nitsche, *A non-existence theorem for the two-dimensional analogue of the catenary*, *Analysis*, **6** No. 2/3 (1986), 143–156.
- [SE1] J. Serrin, *The problem of Dirichlet for quasilinear elliptic differential equations with many independent variables*, *Phil. Trans. Roy. Soc. London, Ser. A*, **264** (1969), 413–496.
- [SE2] *Gradient estimates for solutions of nonlinear elliptic and parabolic equations*, in: *Contributions to nonlinear functional analysis*, p. 565–602. Academic Press, New York 1971.
- [St] G. Stampacchia, *Equations elliptiques du second ordre à coefficients discontinus*, Les Presse de l'Université, Montréal 1966.

Received January 15, 1988.

UNIVERSITÄT DES SAARLANDES  
D-66 SAARBRÜCKEN  
FEDERAL REPUBLIC OF GERMANY

AND

AUSTRALIAN NATIONAL UNIVERSITY  
GPO BOX 4  
CANBERRA ACT 2601, AUSTRALIA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093

R. FINN  
Stanford University  
Stanford, CA 94305

HERMANN FLASCHKA  
University of Arizona  
Tucson, AZ 85721

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305

ROBION KIRBY  
University of California  
Berkeley, CA 94720

C. C. MOORE  
University of California  
Berkeley, CA 94720

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH  
(1906-1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

<b>Yusuf Abu-Muhanna and Abdallah Khalil Lyzzaik, The boundary behaviour of harmonic univalent maps</b> .....	1
<b>Lawrence Jay Corwin, Allen Moy and Paul J. Sally, Jr., Degrees and formal degrees for division algebras and <math>GL_n</math> over a <math>p</math>-adic field</b> .....	21
<b>Ulrich Dierkes and Gerhard Huisken, The <math>n</math>-dimensional analogue of the catenary: existence and nonexistence</b> .....	47
<b>Peter Gerard Dodds, C. B. Huijsmans and Bernardus de Pagter, Characterizations of conditional expectation-type operators</b> .....	55
<b>Douglas Dokken and Robert Ellis, The Poisson flow associated with a measure</b> .....	79
<b>Larry J. Santoni, Horrocks' question for monomially graded modules</b> .....	105
<b>Zbigniew Slodkowski, Pseudoconvex classes of functions. II. Affine pseudoconvex classes on <math>R^N</math></b> .....	125
<b>Dean Ellis Smith, On the Cohen-Macaulay property in commutative algebra and simplicial topology</b> .....	165
<b>Michał Szurek and Jarosław Wisniewski, Fano bundles over <math>P^3</math> and <math>Q_3</math></b> ...	197