

Pacific Journal of Mathematics

KAPLANSKY'S THEOREM AND BANACH PI-ALGEBRAS

VLADIMÍR MÜLLER

KAPLANSKY'S THEOREM AND BANACH PI-ALGEBRAS

VLADIMÍR MÜLLER

By the theorem of Kaplansky a bounded operator in a Banach space is algebraic if and only if it is locally algebraic. We prove a generalization of this theorem. As a corollary we obtain the analogous result for finite (or countable) families of operators. Further we prove that a Banach algebra is PI (i.e. it satisfies a polynomial identity) if and only if it is locally PI.

Let T be a bounded operator on a Banach space X . The classical theorem of Kaplansky [5] states that T is algebraic (i.e. $p(T) = 0$ for some polynomial $p \neq 0$) if and only if it is locally algebraic (i.e. for every $x \in X$ there exists a non-zero polynomial p_x such that $p_x(T)x = 0$). In this paper we prove (Theorem 1) a generalized version of this theorem. As its corollaries it is possible to obtain the original theorem of Kaplansky, the theorem of Sinclair [9] and also new analogical results for finite or countable families of operators.

In the second part of the paper we deal with Banach PI-algebras (i.e. Banach algebras satisfying a polynomial identity). PI-rings and PI-algebras were studied intensely from the algebraic point of view, see e.g. [4], [8]. On the other hand Banach PI-algebras are much less known even though they form a very interesting class of Banach algebras. They are a natural generalization of commutative Banach algebras and it is possible to develop the complete analogy of the Gelfand theory, see [6].

In this paper we prove a theorem of Kaplansky's type for Banach PI-algebras. This result is closely related to earlier results of Grabiner [2] and Dixon [1].

The author wishes to thank Professor B. Silbermann for calling his attention to the interesting field of Banach PI-algebras and fruitful discussions about it.

Let n be a positive integer. We denote by $\mathcal{P}^{(n)}$ the set of all complex polynomials in n non-commutative indeterminates i.e. the free algebra over \mathbb{C} with n generators and with the unit element. Similarly we denote by $\mathcal{P}^{(\infty)} = \bigcup_{n=1}^{\infty} \mathcal{P}^{(n)}$ the set of all complex polynomials with countably many indeterminates.

Let X and Y be Banach spaces. Then $B(X, Y)$ denotes the set of all bounded operators from X to Y ; we write shortly $B(X)$ instead of $B(X, X)$.

Let X be a Banach space, $1 \leq n < \infty$ and let $T_1, \dots, T_n \in B(X)$. We say that the n -tuple (T_1, \dots, T_n) is algebraic if $p(T_1, \dots, T_n) = 0$ for some $p \in \mathcal{P}^{(n)}$, $p \neq 0$. We say that (T_1, \dots, T_n) is locally algebraic if, for every $x \in X$, there exists a non-zero polynomial $p_x \in \mathcal{P}^{(n)}$ such that $p_x(T_1, \dots, T_n)x = 0$.

These definitions can be used also for an infinite sequence $\{T_i\}_{i=1}^{\infty}$ of bounded operators on X (for $p \in \mathcal{P}^{(n)} \subset \mathcal{P}^{(\infty)}$ we have $p(T_1, T_2, \dots) = p(T_1, \dots, T_n)$). Equivalently, the sequence $\{T_i\}_{i=1}^{\infty}$ is locally algebraic if, for every $x \in X$ there exist n and $0 \neq p \in \mathcal{P}^{(n)}$ such that $p(T_1, \dots, T_n)x = 0$.

We start with the following generalization of Kaplansky's theorem.

THEOREM 1. *Let M be a linear space of countable (infinite) dimension, let Y, Z be Banach spaces and let $R: M \rightarrow B(Y, Z)$ be a linear mapping with the property that for every $y \in Y$ there exists $m \in M$, $m \neq 0$ such that $R(m)y = 0$. Then there exists $m \in M$, $m \neq 0$ such that $R(m)$ is a finite-dimensional operator.*

Proof. Let e_1, e_2, \dots be a basis in M . Put $M_0 = \{0\}$ and denote by M_k ($k = 1, 2, \dots$) the linear subspace of M spanned by the vectors e_1, \dots, e_k .

Let F be a finite-dimensional subspace of Z . For $j = 1, 2, \dots$ denote by $Y_{F,j}$ the set of all $y \in Y$ for which there exists $m \in M_j$, $m \neq 0$, such that $R(m)y \in F$ and $R(m')y \notin F$ for every $m' \in M_{j-1}$, $m' \neq 0$. By the assumption $\bigcup_{j=1}^{\infty} Y_{F,j} = Y$ so there exists $k = k(F)$ such that $Y_{F,k}$ is of the second category and $Y_{F,l}$ is of the first category for every $l < k$. Fix a finite-dimensional subspace $F \subset Z$ with the property that

$$k = k(F) = \min_{\substack{G \subset Z \\ \dim G < \infty}} k(G).$$

We have $Y_{F,k} = \bigcup_{s=1}^{\infty} Y_{F,k}^{(s)}$ where

$$Y_{F,k}^{(s)} = \left\{ y \in Y_{F,k}, \text{ there exists } m = e_k + \sum_{i=1}^{k-1} \alpha_i e_i \in M_k \right. \\ \left. \text{such that } \sum_{i=1}^{k-1} |\alpha_i| \leq s \text{ and } R(m)y \in F \right\}.$$

We prove that $Y_{F,k}^{(s)}$ is a closed set for every s . Let $y_j \in Y_{F,k}^{(s)}$ ($j = 1, 2, \dots$), $y_j \rightarrow y$. Then there exist elements $m_j \in M_k$, $m_j = e_k + \sum_{i=1}^{k-1} \alpha_{ji}e_i$ such that $\sum_{i=1}^{k-1} |\alpha_{ji}| \leq s$ and $R(m_j)y_j \in F$. Using the compactness argument it is possible to find a subsequence $\{y_{j_r}\}_{r=1}^\infty$ and a vector $m \in M_k$ such that $m_{j_r} \rightarrow m$ coordinate-wise and $R(m_{j_r}) \rightarrow R(m)$ in the norm topology. It is easy to show that

$$R(m)y = \lim_{r \rightarrow \infty} R(m_{j_r})y_{j_r} \in F;$$

hence $y \in Y_{F,k}^{(s)}$ and $Y_{F,k}^{(s)}$ is closed. Therefore there exists $w \in Y$, $r > 0$ and a positive integer s such that

$$\{y \in Y, \|y - w\| < r\} \subset Y_{F,k}^{(s)} \subset Y_{F,k}.$$

Let $a = e_k + \sum_{i=1}^{k-1} \alpha_i e_i$ be the element of M_k satisfying

$$(1) \quad R(a)w \in F.$$

Denote by $F' = F \vee \bigvee_{i=1}^k \{R(e_i)w\}$. Clearly $\dim F' \leq \dim F + k < \infty$. Put $V = Y_{F,k} - \bigcup_{l < k} Y_{F,l}$. It follows from the choice of the subspace F that V is of the second category. Let $v \in V$. Then $v \in Y_{F,k}$ and

$$(2) \quad R(b)v \in F$$

for some $b \in M_k$, $b = e_k + \sum_{i=1}^{k-1} \beta_i e_i$.

Further $w + \lambda v \in Y_{F,k}$ for some complex number $\lambda \neq 0$, i.e. there exists $c = e_k + \sum_{i=1}^{k-1} \gamma_i e_i \in M_k$ such that

$$(3) \quad R(c)(w + \lambda v) = R(c)w + \lambda R(c)v \in F.$$

This implies $R(c)v \in F'$ and together with (2) $R(c - b)v \in F'$ where $c - b = \sum_{i=1}^{k-1} (\gamma_i - \beta_i)e_i \in M_{k-1}$. Since $v \notin \bigcup_{l < k} Y_{F,l}$, we conclude $c - b = 0$, $c = b$.

By (2), (3) and (1) we have $R(c)v = R(b)v \in F$, $R(c)w \in F$ and $R(c - a)w \in F$, where $c - a \in M_{k-1}$. Since $w \notin \bigcup_{l < k} Y_{F,l}$ we conclude again that $c = a$, i.e. $R(a)v \in F$ for every $v \in V$. Thus $R(a)^{-1}F \supset V$ and $R(a)^{-1}F$ is a linear subspace of the second category in Y , therefore $R(a)^{-1}F = Y$, $R(a)Y \subset F$ and $R(a)$ is a finite dimensional operator.

REMARK. One is tempted to expect in Theorem 1 that there exists $m \in M$, $m \neq 0$, such that $R(m) = 0$. However, the following example shows that this is not true in general. Let $Y = Z$ be a separable

Hilbert space with an orthonormal basis $\{h_i\}_{i=1}^\infty$. Define operators $R(m)$, $m \in M$, by

$$\begin{aligned} R(e_1)h_1 &= h_1, & R(e_1)h_j &= 0 & (j \geq 2), \\ R(e_2)h_1 &= 0, & R(e_2)h_2 &= h_1, & R(e_2)h_j &= 0 & (j \geq 3), \\ R(e_i)h_j &= \delta_{ij}h_j & (i \geq 3; \delta_{ij} &\text{ means the Kronecker's symbol}). \end{aligned}$$

It is easy to show that the conditions of Theorem 1 are satisfied and $R(m) \neq 0$ ($m \neq 0$).

THEOREM 2. *Let X be a Banach space, $1 \leq n \leq \infty$. Let $T = \{T_i\}_{i=1}^n$ be a (finite or infinite) sequence of bounded operators on X . Then T is algebraic if and only if it is locally algebraic.*

Proof. Suppose T is locally algebraic. We prove that it is algebraic (the converse implication is trivial). Put $M = \mathcal{P}^{(n)}$, $Y = Z = X$. For $p \in \mathcal{P}^{(n)}$ put $R(p) = p(T)$. By Theorem 1 there exist a polynomial $p \in \mathcal{P}^{(n)}$, $p \neq 0$, such that $\dim p(T)X < \infty$. Hence $(q \circ p)(T) = 0$ where $q \in \mathcal{P}^{(1)}$ is the characteristic polynomial of the finite-dimensional operator $p(T)|_{p(T)X}$.

In [9], the following generalization of the Kaplansky's theorem was proved: Let $T \in B(X)$ be a non-algebraic operator. Then there exists a sequence x_1, x_2, \dots of elements of X such that $\sum_{i=1}^k p_i(T)x_i \neq 0$ for every $k \geq 0$ and for every polynomial $p_1, \dots, p_k \in \mathcal{P}^{(1)}$ not all of which are equal to 0.

This result can be extended to the case of more than one operator.

THEOREM 3. *Let X be a Banach space, $1 \leq n \leq \infty$. Let $T = \{T_i\}_{i=1}^\infty$ be a (finite or infinite) sequence of bounded operators on X which is not algebraic. Then there exist vectors $x_1, x_2, \dots \in X$ such that $\sum_{i=1}^k p_i(T)x_i \neq 0$ for every k and for every polynomial $p_1, \dots, p_k \in \mathcal{P}^{(n)}$ not all of which are equal to 0.*

Proof. Suppose on the contrary that for every sequence x_1, x_2, \dots of elements of X there exist k and polynomials $p_1, \dots, p_k \in \mathcal{P}^{(n)}$, $(p_1, \dots, p_k) \neq (0, \dots, 0)$ such that $\sum_{i=1}^k p_i(T)x_i = 0$.

Let M be the linear space of all sequences $\{p_i\}_{i=1}^\infty$ of polynomials $p_i \in \mathcal{P}^{(n)}$ only a finite number of which are non-zero. Put $Z = X$ and

$$Y = \{\{x_i\}_{i=1}^\infty, x_i \in X (i = 1, 2, \dots), \sup\{\|x_i\|, i = 1, 2, \dots\} < \infty\}.$$

Then Y with the norm $\|\{x_i\}_{i=1}^\infty\| = \sup\{\|x_i\|, i = 1, 2, \dots\}$ is a Banach space. For $p = \{p_i\}_{i=1}^\infty \in M$ and $y = \{x_i\}_{i=1}^\infty \in Y$ put $R(p)y = \sum_{i=1}^\infty p_i(T)x_i$ (in fact the sum is finite). By Theorem 1 there exist a finite-dimensional subspace $F \subset X$, a positive integer k and polynomials $p_1, \dots, p_k \in \mathcal{P}^{(n)}$, $(p_1, \dots, p_k) \neq (0, \dots, 0)$, such that

$$\sum_{i=1}^k p_i(T)x_i \in F \quad \text{for every } x_1, \dots, x_k \in X.$$

Choose $j \in \{1, \dots, k\}$ such that $p_j \neq 0$. Let $x \in X$ be arbitrary. If we put $x_j = x$, $x_i = 0$ ($i \neq j$) then we get $p_j(T)x \in F$ for every $x \in X$, i.e. $p_j(T)$ is a finite-dimensional operator. The rest is the same as in the proof of Theorem 2.

REMARK. Theorem 1 unifies some of the results of Kaplansky's type (cf. problem of Halmos [3]). On the other hand there are some results of this type which do not fit into this frame (see e.g. [10] where bounded analytic functions are used instead of polynomials or "approximative" results of Kaplansky's type [7], [11]). Another example will be the result for Banach PI-algebras which we prove in the following section.

Let A be a Banach algebra with the unit (we shall always assume that a Banach algebra has a unit element although this assumption is not essential). We say that A is PI if there exist a positive integer n and a non-zero polynomial $p \in \mathcal{P}^{(n)}$ such that $p(a_1, \dots, a_n) = 0$ for every $a_1, \dots, a_n \in A$. We say that A is locally PI if for every sequence $\{a_i\}_{i=1}^\infty$ of elements of A there exist n and a non-zero polynomial $p \in \mathcal{P}^{(n)}$ such that $p(a_1, \dots, a_n) = 0$ (both n and p depend on the sequence $\{a_i\}_{i=1}^\infty$).

THEOREM 4. *Let A be a Banach algebra with the unit. Then A is PI if and only if A is locally PI.*

Proof. The implication PI \Rightarrow locally PI is trivial. Suppose that A is locally PI. Denote by \tilde{A}

$$\tilde{A} = \{\{a_i\}_{i=1}^\infty, a_i \in A, i = 1, 2, \dots, \sup\{\|a_i\|, i = 1, 2, \dots\} < \infty\}.$$

Then \tilde{A} with the norm $\|\{a_i\}_{i=1}^\infty\| = \sup\{\|a_i\|, i = 1, 2, \dots\}$ is a Banach space. Further $\tilde{A} = \bigcup_{n=1}^\infty \tilde{A}_n$ where

$$\tilde{A}_n = \{\{a_i\}_{i=1}^\infty \in \tilde{A}, \text{ there exists } p \in \mathcal{P}^{(n)}, \deg p \leq n, n^{-1} \leq |p| \leq n, p(a_1, \dots, a_n) = 0\}$$

(we denote by $\deg p$ the degree of a polynomial p and $|p|$ denotes the sum of moduli of coefficients of p).

Since \tilde{A}_n is a closed subset for every n , Baire's theorem implies that there exist a positive integer n , $\tilde{y} \in \tilde{A}$ and $r > 0$ such that

$$\{\tilde{a} \in \tilde{A}, \|\tilde{a} - \tilde{y}\| < r\} \subset \tilde{A}_n.$$

Let $\tilde{z} = \{z_i\}_{i=1}^\infty \in \tilde{A}_n$. Then $p(z_1, \dots, z_n) = 0$ for some $p \in \mathcal{P}^{(n)}$, $p \neq 0$, $\deg p \leq n$, i.e. the set

$$C = \{z_{i_1}, \dots, z_{i_k}, 0 \leq k \leq n, i_1, \dots, i_k \in \{1, \dots, n\}\}$$

is linearly dependent and $\sum_{c \in C} \alpha_c c = 0$ where α_c denotes the coefficient of p standing at the term c . Therefore $\sum_{c \in C} \alpha_c (cz_{n+1} - z_{n+1}c) = 0$. Let $C = \{c_1, \dots, c_s\}$. Denote by

$$e_s(x_1, \dots, x_s) = \sum_{\sigma \in \mathcal{S}_s} (-1)^{\text{sign } \sigma} x_{\sigma(1)} \cdots x_{\sigma(s)}$$

the standard polynomial (the sum is taken over all permutations of the set $\{1, \dots, s\}$). Clearly,

$$e_s(c_1 z_{n+1} - z_{n+1} c_1, \dots, c_s z_{n+1} - z_{n+1} c_s) = 0,$$

i.e. there exists a non-zero polynomial $p_n \in \mathcal{P}^{(n+1)}$ such that $p_n(z_1, \dots, z_{n+1}) = 0$ for every sequence $\{z_i\}_{i=1}^\infty \in \tilde{A}_n$. Let $\tilde{a} = \{a_i\}_{i=1}^\infty \in \tilde{A}$ be arbitrary. Then $\tilde{y} + \lambda \tilde{a} \in \tilde{A}_n$ for all complex λ , $|\lambda| \|\tilde{a}\| < r$, i.e.

$$p_n(y_1 + \lambda a_1, \dots, y_{n+1} + \lambda a_{n+1}) = 0.$$

We can write

$$\begin{aligned} & p_n(y_1 + \lambda a_1, \dots, y_{n+1} + \lambda a_{n+1}) \\ &= p_n(y_1, \dots, y_{n+1}) + \lambda q^{(1)}(y_1, \dots, y_{n+1}, a_1, \dots, a_{n+1}) \\ & \quad + \dots + \lambda^{\deg p_n - 1} q^{(\deg p_n - 1)}(y_1, \dots, y_{n+1}, a_1, \dots, a_{n+1}) \\ & \quad + \lambda^{\deg p_n} p_n(a_1, \dots, a_{n+1}). \end{aligned}$$

Since this expression is equal to 0 for all λ such that $|\lambda| \|\tilde{a}\| < r$, we conclude that $p_n(a_1, \dots, a_{n+1}) = 0$ for every $(n + 1)$ -tuple a_1, \dots, a_{n+1} of elements of A . Thus A is a PI-algebra.

REMARK. In [2], S. Grabiner proved that a nil Banach algebra (i.e. consisting of nilpotent elements) is nilpotent (i.e. $A^n = 0$ for some n). The previous theorem is closely related to this result.

An algebra A is called algebraic if every element $a \in A$ is algebraic, i.e. $p(a) = 0$ for some non-zero polynomial $p \in \mathcal{P}^{(1)}$. An algebra

is called locally finite if every finite subset of A generates a finite-dimensional subalgebra.

Clearly, a locally finite algebra is algebraic.

As an easy corollary of the previous theorem we can obtain the following result of Dixon [1] that the converse implication is true for Banach algebras.

COROLLARY 5. *Let A be a Banach algebra with the unit. Then A is algebraic if and only if A is locally finite.*

Proof. If A is algebraic then A is locally PI and thus PI by Theorem 4. An algebraic PI-algebra is locally finite (see [4], X/12, Theorem 1).

REFERENCES

- [1] P. G. Dixon, *Locally finite Banach algebras*, J. London Math. Soc., **8** (1974), 325–328.
- [2] S. Grabiner, *The nilpotency of Banach nil algebras*, Proc. Amer. Math. Soc., **21** (1969), 510.
- [3] P. R. Halmos, *Capacity in Banach algebras*, Indiana Univ. Math. J., **20** (1971), 855–863.
- [4] N. Jacobson, *Structure of rings*, Amer. Math. Soc. Coll. Publ. 37, Providence, R.I. 1956.
- [5] I. Kaplansky, *Infinite Abelian Groups*, University of Michigan Press, Ann Arbor, 1954.
- [6] N. Ya. Krupnik, *Banach Algebras with Symbol and Singular Integral Operators*, Birkhäuser Verlag, Basel-Boston, 1987.
- [7] V. Müller, *On quasilgebraic operators in Banach spaces*, J. Operator Theory, **17** (1987), 291–300.
- [8] L. H. Rowen, *Polynomial Identities in Ring Theory*, Academic Press, New York, 1980.
- [9] A. M. Sinclair, *Automatic continuity of linear operators*, London Math. Soc. Lect. Note Series 21, Cambridge University Press, Cambridge, 1976.
- [10] B. Sz.-Nagy, C. Foias, *Local characterization of operators of class C_0* , J. Funct. Anal., **8** (1971), 76–81.
- [11] P. Vrbová, *On local spectral properties of operators in Banach spaces*, Czechoslovak Math. J., **25** (1973), 483–492.

Received March 23, 1988.

INSTITUTE OF MATHEMATICS,
 ČSAV, ŽITNÁ 25, 115 67
 PRAGUE 1, CZECHOSLOVAKIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1990 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 141, No. 2 December, 1990

| | |
|---|-----|
| Ulrich F. Albrecht , Locally A -projective abelian groups and generalizations | 209 |
| Michel Carpentier , Sommes exponentielles dont la géométrie est très belle: p -adic estimates | 229 |
| G. Deferrari, Angel Rafael Larotonda and Ignacio Zalduendo , Sheaves and functional calculus | 279 |
| Jane M. Hawkins , Properties of ergodic flows associated to product odometers | 287 |
| Anthony To-Ming Lau and Viktor Losert , Complementation of certain subspaces of $L_\infty(G)$ of a locally compact group | 295 |
| Shahn Majid , Matched pairs of Lie groups associated to solutions of the Yang-Baxter equations | 311 |
| Diego Mejia and C. David (Carl) Minda , Hyperbolic geometry in k -convex regions | 333 |
| Vladimír Müller , Kaplansky's theorem and Banach PI-algebras | 355 |
| Raimo Näkki , Conformal cluster sets and boundary cluster sets coincide | 363 |
| Tomasz Przebinda , The wave front set and the asymptotic support for p -adic groups | 383 |
| R. F. Thomas , Some fundamental properties of continuous functions and topological entropy | 391 |