

Pacific Journal of Mathematics

**UNITARY COBORDISM OF CLASSIFYING SPACES OF
QUATERNION GROUPS**

ABDESLAM MESNAOUI

UNITARY COBORDISM OF CLASSIFYING SPACES OF QUATERNION GROUPS

ABDESLAM MESNAOUI

The main purpose of this article is to prove that the complex cobordism ring of classifying spaces of quaternion groups Γ_k ($|\Gamma_k| = 2^k$) is a quotient of the graded ring $U^*(pt)[[X, Y, Z]]$ ($\dim X = \dim Y = 2, \dim Z = 4$) by a graded ideal generated by six homogeneous formal power series.

0. Introduction. Let Γ_k be the generalized quaternion group. Γ_k is generated by u, v , subject to the relations $u^t = v^2, uvu = v, t = 2^{k-2}$. In order to calculate $U^*(B\Gamma_k)$ we first consider the case $k = 3$, i.e. $\Gamma_3 = \Gamma$. We recall that $\Gamma = \{\pm 1, \pm i, \pm j, \pm k\}$ with the relations $i^2 = j^2 = k^2 = -1, ij = k, jk = i, ki = j$. We shall define $A \in \tilde{U}^2(B\Gamma), B \in \tilde{U}^2(B\Gamma), D \in \tilde{U}^4(B\Gamma)$ as Euler classes of complex vector bundles over $B\Gamma$ corresponding to unitary irreducible representations of Γ . Let Λ_* be the graded $U^*(pt)$ -algebra $U^*(pt)[[X, Y, Z]]$ with $\dim X = \dim Y = 2, \dim Z = 4, \Omega_* = U^*(pt)[[Z]] \subset \Lambda_*$ and $U^*(pt)[[D]] = \{P(D), P \in \Omega_*\}$. Then by using the Atiyah-Hirzebruch spectral sequence we obtain the following results where $T(Z) \in \Omega_4, J(Z) \in \Omega_0$ are well defined formal power series.

THEOREM 2.18. (a) *As graded $U^*(pt)$ -algebras we have:*

$$U^*(pt)[[D]] \simeq \Omega_*/(T(Z)).$$

(b) *As graded $U^*(pt)[[D]]$ -modules we have: $U^*(B\Gamma) \simeq U^*(pt)[[D]] \oplus U^*(pt)[[D]] \cdot A \oplus U^*(pt)[[D]] \cdot B$ and A, B have the same annihilator $(2 + J(D)) \cdot U^*(pt)[[D]]$.*

THEOREM 2.17. *The graded $U^*(pt)$ -algebra $U^*(B\Gamma)$ is isomorphic to Λ_*/I_* where I_* is a graded ideal generated by six homogeneous formal power series.*

The method used for Γ is extended to $\Gamma_k, k \geq 4$. As before we shall define $B_k \in \tilde{U}^2(B\Gamma_k), C_k \in \tilde{U}^2(B\Gamma_k), D_k \in \tilde{U}^4(B\Gamma_k)$ as Euler classes of complex vector bundles over $B\Gamma_k$ corresponding to unitary irreducible representations of Γ_k and elements $G'(Z) \in \Omega_2,$

$T_k(Z) \in \Omega_4$. If $B'_k = B_k + G'_k(D_k)$, $C'_k = C_k + G'_k(D_k)$ then we get:

THEOREM 3.14. (a) $U^*(pt)[[D_k]] \simeq \Omega_*/(T_k)$ as graded $U^*(pt)$ -algebras.

(b) As graded $U^*(pt)[[D_k]]$ -modules we have:

$$U^*(B\Gamma_k) \simeq U^*(pt)[[D_k]] \oplus U^*(pt)[[D_k]] \cdot B'_k \oplus U^*(pt)[[D_k]] \cdot C'_k$$

and B'_k, C'_k have the same annihilator $(2 + J(D_k)) \cdot U^*(pt)[[D_k]]$.

THEOREM 3.12. The graded $U^*(pt)$ -algebra $U^*(B\Gamma_k)$ is isomorphic to Λ_*/\tilde{I}_* where \tilde{I}_* is a graded ideal of Λ_* generated by six homogeneous formal power series.

In the appendix, part A, we give a new method of calculating $U^*(BZ_m)$. Let Λ'_* be the graded algebra $U^*(pt)[[Z]]$, $\dim Z = 2$.

THEOREM A.1. $U^*(BZ_m) \simeq \Lambda'_*/([m](Z))$ as graded $U^*(pt)$ -algebras.

In part B we show that:

THEOREM B.2.

$$U^{2i+2}(BSU(n)) \simeq U^{2i+2}(BU(n))/e(\Lambda^n \gamma(n)) \cdot U^{2i}(BU(n))$$

and $U^{2i+1}(BSU(n)) = 0$, $i \in \mathbb{Z}$.

In this theorem $e(\Lambda^n \gamma(n))$ is the Euler class of $\Lambda^n \gamma(n)$ where $\gamma(n)$ denotes the universal bundle over $BU(n)$.

In part C we calculate $H^*(B\Gamma_k)$, $k \geq 4$.

THEOREM C. If $k \geq 4$ then we have $H^*(B\Gamma_k) = \mathbb{Z}[x_k, y_k, z_k]$ with $\dim x_k = \dim y_k = 2$, $\dim z_k = 4$, subject to the relations:

$$2x_k = 2y_k = x_k y_k = 2^k z_k = 0, \quad x_k^2 = y_k^2 = 2^{k-1} z_k.$$

Theorem C is certainly known to workers in the field.

The layout is as follows:

- I Preliminaries and notations.
- II Calculation of $U^*(B\Gamma)$.
- III Calculation of $U^*(B\Gamma_k)$, $k \geq 4$.
- IV Appendix.

In the course of the computations we have determined the leading coefficients of some formal power series with the purpose of using them in a subsequent paper where the bordism groups $\tilde{U}_*(B\Gamma_k)$ are calculated.

We shall use the same notation for unitary irreducible representations of Γ_k and corresponding complex vector bundles over $B\Gamma_k$. The notation $\gamma(n)$ will be used for the universal complex vector bundle over $BU(n)$. The notation \mathbb{Z} will be for the ring of integers and \mathbb{C} for the complex number field.

The results of this paper have been obtained in 1983 under the supervision of Dr. L. Hodgkin, University of London. I thank him sincerely for having proposed the subject, for his advice and encouragement. I would like to express my deep thanks to the referee who made many useful suggestions; they helped to improve the exposition of this paper and the statement of some results, particularly Theorems 2.18 and 3.14.

I. Preliminaries and notations. 1. Let X be a CW-complex; we define a filtration on $U^n(X)$ by the subgroups

$$J^{p,q} = \text{Ker}(i^*: U^n(X) \rightarrow U^n(X_{p-1})),$$

X_p being the p -skeleton of X , $i: X_{p-1} \subset X$, $p+q=n$; $U^n(X)$ is a topological group, the subgroups $J^{p,q}$ being a fundamental system of neighbourhoods of 0; we denote this topology by T . If the U^* -Atiyah-Hirzebruch spectral sequence (denoted by U^* -AHSS) for X collapses then T is complete and Hausdorff (see [3]). The edge homomorphism $\mu: U^n(X) \rightarrow H^n(X)$ is defined by $\mu = 0$ if $n < 0$ and if $n \geq 0$ it is the projection $U^n(X) = J^{0,n} = J^{n,0} \rightarrow J^{n,0}/J^{n+1,-1} = E_\infty^{n,0} \subset E_2^{n,0} = H^n(X)$. By easy arguments involving spectral sequences we have the following basic result:

THEOREM 1.1. *Let X be a CW-complex such that:*

- (a) *The U^* -AHSS for X collapses.*
- (b) *For each $n \geq 0$ there are elements a_{in} generating the \mathbb{Z} -module $H^n(X)$.*

Then for each $n \geq 0$ there are elements $A_{in} \in U^n(X)$ such that:

- (a) $\mu(A_{in}) = a_{in}$.
- (b) *If E denotes the $U^*(pt)$ -submodule of $U^*(X)$ generated by the system (A_{in}) and if E_n is the n -component of E then $\overline{E}_n = U^n(X)$, \overline{E}_n being the closure of E_n for T .*

Moreover (b) is valid if we take any system (A'_{in}) , $A'_{in} \in U^n(X)$ such that $\mu(A'_{in}) = a_{in}$ for each (i, n) . \square

(See Theorem 2.5 for a proof of this result in a special case.)

2. Let X be a skeleton-finite CW-complex, which is the case we are interested in. There is a ring spectra map $f: MU \rightarrow H$ (see [1]); by naturality of AHSS the map $f^\#(X): U^*(X) \rightarrow H^*(X)$ induced by f is identical to the edge-homomorphism described above. Let ξ be a complex vector bundle over X of dimension n ; the Conner-Floyd characteristic classes of ξ will be denoted by $cf_i(\xi)$; the Euler class $e(\xi)$ of ξ for MU is $cf_n(\xi)$ and the Euler class $e_1(\xi)$ for H is the Chern class $c_n(\xi)$. As $f^\#(X)$ maps Euler classes on Euler classes we have $\mu(e(\xi)) = e_1(\xi)$ (see [7]).

3. Consider the formal power series ring $E_* = U^*(pt)[[c_1, c_2, \dots, c_r]]$ graded by taking $\dim c_1 = n_1 > 0, \dots, \dim c_r = n_r > 0$. Given $P(c_1, \dots, c_r) \in E_n$ with $P \neq 0$,

$$P = \sum a_u \cdot c_1^{u_1} \cdots c_r^{u_r}, \quad u = (u_1, \dots, u_r),$$

we define $\nu(P) = \{\inf(n_1 u_1 + \dots + n_r u_r), a_u \neq 0\}$ and $\nu(0) = +\infty$. Let J_p be $\{P \in E_n \mid \nu(P) \geq p\}$; we have $E_n = J_0 \supset J_1 \supset \dots$, and since $\bigcap_{p \geq 0} J_p = 0$, $E_n = \varprojlim E_n/J_p$, it follows that E_n is complete and Hausdorff for the topology defined by the filtration (J_p) .

Suppose that B is a CW complex such that the associated U^* -AHSS collapses; if $A_i \in U^{n_i}(B)$, $i = 1, 2, \dots, r$, then there is a unique continuous homomorphism $\psi: E_* \rightarrow U^*(B)$ such that $\psi(c_i) = A_i$, $i = 1, 2, \dots, r$.

Now in a different situation consider the case where B_1 is a CW-complex such that $U^*(B_1) = E_*$. There are two topologies on $U^*(B_1)$ defined respectively by the filtration (J_p) on E_* and by the filtration $(J_1^{p,q})$ deduced from the U^* -AHSS for B_1 . If B is a CW-complex such that the U^* -AHSS for B collapses, $(J^{p,q})$ the corresponding filtration on $U^*(B)$ (see §I) and g a continuous map: $B \rightarrow B_1$ then from $J_p \subset J_1^{p,q}$, $g^*(J_1^{p,q}) \subset J^{p,q}$ it follows that $g^*: E_n \rightarrow U^n(B)$ is continuous for the topologies defined by ν on E_n and $(J^{p,q})$ on $U^*(B)$. As a consequence if (P_m) is a sequence of polynomials such that $(P_m) \rightarrow P$ in E_n and if $g^*(c_i) = A_i$ then $P_m(A_1, \dots, A_r) \rightarrow g^*(P)$ in $U^*(B)$; so if $P = \sum a_u c_1^{u_1} \cdots c_r^{u_r} \in E_n$ we can write $g^*(P) = \sum a_u A_1^{u_1} \cdots A_r^{u_r}$.

In the sequel we shall also be concerned with $\Lambda_* = U^*(pt)[[X, Y, Z]]$, $\dim X = \dim Y = 2$, $\dim Z = 4$; Λ_* has the topology defined by ν .

The following assertions are clear:

- (a) In $\Lambda_{2n} : (R_p) \rightarrow 0$ iff $\nu(R_p) \rightarrow \infty$.
- (b) If $P(X, Y, Z) \in \Lambda_{2m+2n}$, $Q(X, Y, Z) \in \Lambda_{2n}$ and (R_p) a sequence in Λ_{2m} such that $R_p \rightarrow R$ and $\nu(P - R_p Q) \rightarrow \infty$ then $RQ = P$.
- (c) If $\nu(R_p) \rightarrow \infty$ then the sequence (M_p) defined by $M_p = R_0 + \dots + R_p$ converges to a unique limit denoted by $\sum_{p \geq 0} R_p$.

In Sections II and III we shall define three elements $A_k \in \tilde{U}^2(B\Gamma_k)$, $B_k \in \tilde{U}^2(B\Gamma_k)$, $D_k \in \tilde{U}^4(B\Gamma_k)$; as the U^* -AHSS for $B\Gamma_k$ collapses there is a unique continuous homomorphism φ of graded $U^*(pt)$ -algebras: $\Lambda_* \rightarrow U^*(B\Gamma_k)$ such that $\varphi(X) = A_k$, $\varphi(Y) = B_k$, $\varphi(Z) = D_k$.

The next well known result will be useful:

PROPOSITION 1.2. *Suppose X a CW-complex such that $H^*(X) = \mathbb{Z}[a]$. Then there is an element $A \in U^*(X)$ such that $\mu(A) = a$ and $U^*(X) = H^*(X) \hat{\otimes} U^*(pt) = U^*(pt)[[A]]$. Moreover for any $A' \in U^*(X)$ such that $\mu(A') = a$ we have $U^*(X) = U^*(pt)[[A']]$. \square*

II. Computation of $U^*(B\Gamma)$. We recall that the quaternion group Γ consists of $\{1, \pm i, \pm j, \pm k\}$ subject to the relations $ij = k$, $jk = i$, $ki = j$, $i^2 = k^2 = -1$. The irreducible unitary representations of Γ are $1: i \rightarrow 1, j \rightarrow 1$, $\xi_i: i \rightarrow 1, j \rightarrow -1$, $\xi_j: i \rightarrow -1, j \rightarrow 1$, $\xi_k: i \rightarrow -1, j \rightarrow -1$, $\eta: i \rightarrow \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, j \rightarrow \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$; the character table of Γ is:

(Conjugacy classes)

| | 1 | -1 | $\pm i$ | $\pm j$ | $\pm k$ |
|---------|---|----|---------|---------|---------|
| 1 | 1 | 1 | 1 | 1 | 1 |
| ξ_i | 1 | 1 | 1 | -1 | -1 |
| ξ_j | 1 | 1 | -1 | 1 | -1 |
| ξ_k | 1 | 1 | -1 | -1 | 1 |
| η | 2 | -2 | 0 | 0 | 0 |

We have the following relations in the representation ring $R(\Gamma)$:

$$\begin{aligned} \xi_i^2 = \xi_j^2 = \xi_k^2 = 1, \quad \xi_i \cdot \xi_j = \xi_k, \quad \xi_j \cdot \xi_k = \xi_i, \quad \xi_k \xi_i = \xi_j, \\ \eta \cdot \xi_i = \eta \cdot \xi_j = \eta, \quad \eta^2 = 1 + \xi_i + \xi_j + \xi_k \quad (\text{see [6], [2]}). \end{aligned}$$

We have $H^0(B\Gamma) = \mathbb{Z}$, $H^{4n}(B\Gamma) = \mathbb{Z}_8$, $n \geq 1$, $H^{4n+2}(B\Gamma) = \mathbb{Z}_2 \oplus \mathbb{Z}_2$, $n \geq 0$, $H^{2n+1}(B\Gamma) = 0$. Moreover if d is a generator of $H^4(B\Gamma)$ and if a, b are generators of $H^2(B\Gamma)$ then d^n is a generator of $H^{4n}(B\Gamma)$, $n \geq 1$, and ad^n, bd^n are generators of $H^{4n+2}(B\Gamma)$, $n \geq 0$ (see [5]).

Since $H^m(B\Gamma) = 0$, m odd we have:

PROPOSITION 2.1. *The U^* -AHSS for $B\Gamma$ collapses.* \square

There are four important complex vector-bundles $\xi_i, \xi_j, \xi_k: E\Gamma \times_{\Gamma} \mathbb{C} \rightarrow B\Gamma$ and $\eta: E\Gamma \times_{\Gamma} \mathbb{C}^2 \rightarrow B\Gamma$ where the actions of Γ on \mathbb{C} and \mathbb{C}^2 are induced by the representations ξ_i, ξ_j, ξ_k and η . We have a canonical inclusion $q: \mathbb{Z}_2 \subset \Gamma$ obtained by identifying $\{1, i^2\}$ with \mathbb{Z}_2 ; let ρ be the unitary representation of \mathbb{Z}_2 given by $\rho(1) = 1, \rho(i^2) = -1$; the restriction map: $R(\Gamma) \rightarrow R(\mathbb{Z}_2)$ sends ξ_i, ξ_j, ξ_k to 1 and η to 2ρ ; so:

PROPOSITION 2.2. *$(Bq)^*(\xi_h), h = i, j, k$, are trivial and $(Bq)^*(\eta) = 2\rho$.* \square

1. *Chern Classes of ξ_i, ξ_j, η .* The canonical isomorphism

$$\text{Hom}(\Gamma, U(1)) \rightarrow H^2(\Gamma)$$

is given by $\delta \rightarrow c_1(g(\delta))$ where g denotes the canonical map: $R(\Gamma) \rightarrow K^0(B\Gamma)$ and c_1 the first Chern class (Sec. [2]). Since $\text{Hom}(\Gamma, U(1)) = \{1, \xi_i, \xi_j, \xi_k\}$ and $H^2(B\Gamma) = \mathbb{Z}_2 \oplus \mathbb{Z}_2$ we have:

PROPOSITION 2.3. *$H^2(B\Gamma)$ is generated by $\{c_1(\xi_i), c_1(\xi_j)\}$.* \square

Now we consider the topological group $\text{Sp}(1)$ of quaternions of absolute value 1; $\text{Sp}(1)$ is homeomorphic to S^3 and $H^*(BS^3) = \mathbb{Z}[u]$, $\dim u = 4$, u being the first symplectic Pontrjagin class of the universal $\text{Sp}(1)$ -vector bundle θ . If we consider θ as a $U(2)$ -vector bundle, then $u = c_2(\theta)$ (see [12], page 179). Let $p: \Gamma \subset \text{Sp}(1) = S^3$ be the natural inclusion; then it is easily seen that $(Bp)^*(\theta) = \eta$, θ being regarded as a $U(2)$ -vector bundle.

PROPOSITION 2.4. *We have $c_1(\eta) = 0$ and $H^4(B\Gamma)$ is generated by $c_2(\eta)$.*

Proof. Since $\det \eta = 1$ we have $c_1(\eta) = 0$. From the transgression exact sequence of the fibration: $S^3/\Gamma \rightarrow B\Gamma \xrightarrow{Bp} BS^3$ we get the exact

sequence: $H^4(BS^3) \xrightarrow{(Bp)^*} H^4(B\Gamma) \rightarrow H^4(S^3/\Gamma) = 0$ and the result follows (see [11], page 519). \square

From 2.3, 2.4 we may take the Euler classes $e_1(\eta) = d$ as a generator of $H^4(B\Gamma)$ and $\{a = e_1(\xi_i), b = e_1(\xi_j)\}$ as a system of generators of $H^2(B\Gamma)$. Moreover $e_1(n \cdot \eta) = e_1(\eta)^n = d^n$ and $\{e_1(\xi_i + n \cdot \eta) = ad^n, e_1(\xi_j + n \cdot \eta) = bd^n\}$ are generators of $H^{4n}(B\Gamma)$, $n \geq 1$ and $H^{4n+2}(B\Gamma)$, $n \geq 0$, respectively.

2. *Computation of $U^*(B\Gamma)$.* Let A, B, D be the Euler classes for MU of ξ_i, ξ_j, η : $e(\xi_i) = A \in \tilde{U}^2(B\Gamma)$, $e(\xi_j) = B \in \tilde{U}^2(B\Gamma)$, $e(\eta) = D \in \tilde{U}^4(B\Gamma)$. We recall that $\Lambda_* = U^*(pt)[[X, Y, Z]]$ is graded by taking $\dim X = \dim Y = 2$, $\dim Z = 4$; there is a unique continuous homomorphism $\varphi: \Lambda_* \rightarrow U^*(B\Gamma)$ of graded $U^*(pt)$ -algebras such that $\varphi(X) = A$, $\varphi(Y) = B$, $\varphi(Z) = D$. In particular if $P(Z) = \alpha_0 + \alpha_1 Z + \dots + \alpha_i Z^i + \dots \in \Lambda_{2n}$ then $\varphi(P) = P(D) = \text{Lim}_{n \rightarrow \infty} (\alpha_0 + \dots + \alpha_n \cdot D^n)$ in $U^{2n}(B\Gamma)$. If $U^*(pt)[[D]] = \{R(D), R(Z) \in \Omega_*\}$, then $U^*(pt)[[D]]$ is a sub- $U^*(pt)$ -algebra of $U^*(B\Gamma)$.

THEOREM 2.5. *$U^*(B\Gamma)$ is concentrated in even dimensions and as a $U^*(pt)[[D]]$ -module $U^*(B\Gamma)$ is generated by 1, A, B .*

Proof. We have $U^{2n+1}(B\Gamma) = 0$ because $J^{p,q} = J^{p+1,q-1}$ if $p + q = 2n + 1$ and then $U^{2n+1}(B\Gamma) = J^{0,2n+1} = \bigcap_{p+q=2n+1} J^{p,q} = 0$ (see Section I).

Suppose $2n = 4m + 2 > 0$. If $x \in U^{4m+2}(B\Gamma) = J^{0,4m+2} = J^{4m+2,0}$ then $\mu(x) = \alpha_m ad^m + \beta_m bd^m = \mu(\alpha_m AD^m + \beta_m BD^m)$, $\alpha_m \in U^0(pt) = \mathbb{Z}$, $\beta_m \in U^0(pt) = \mathbb{Z}$. It follows that $\mu(x - (\alpha_m AD^m + \beta_m BD^m)) = 0$ and $x_1 = x - (\alpha_m AD^m + \beta_m BD^m) \in J^{4m+3,-1} = J^{4m+4,-2}$. Let s_1 be the quotient map: $J^{4m+4,-2} \rightarrow J^{4m+4,-2} / J^{4m+5,-3} = H^{4m+4}(B\Gamma, U^{-2}(pt)) = U^{-2}(pt) \otimes H^{4m+4}(B\Gamma)$. Then $s_1(x_1) = \gamma_{m+1} \otimes d^{m+1}$, $\gamma_{m+1} \in U^{-2}(pt)$. From the following commutative diagram where χ is induced by the $U^*(pt)$ -module-structure:

$$\begin{array}{ccc} U^{-2}(pt) \otimes U^{4m+4}(B\Gamma) = U^{-2}(pt) \otimes J^{4m+4,0} & \xrightarrow{\chi} & J^{4m+4,-2} \\ \downarrow 1 \otimes \mu & & \downarrow s_1 \\ U^{-2}(pt) \otimes H^{4m+4}(B\Gamma) & \xrightarrow{\sim} & H^{4m+4}(B\Gamma, U^{-2}(pt)) \end{array}$$

it follows that $s_1(x_1) = s_1(\gamma_{m+1} D^{m+1})$ and then $s_1(x_1 - \gamma_{m+1} D^{m+1}) = 0$; so $(x_1 - \gamma_{m+1} D^{m+1}) \in J^{4m+5,-3} = J^{4(m+1)+2,-4}$. We have $x_2 = x_1 - \gamma_{m+1} D^{m+1} = x - (A \cdot \alpha_m D^m + B \cdot \beta_m D^m + \gamma_{m+1} D^{m+1}) \in J^{4(m+1)+2,-4}$.

By using again the products χ we see that after a finite number of steps there are three polynomials in Z :

$$\begin{aligned} P_q(Z) &= \alpha_m Z^m + \alpha_{m+1} Z^{m+1} + \cdots + \alpha_{m+q-1} Z^{m+q-1}, \\ Q_q(Z) &= \beta_m Z^m + \beta_{m+1} Z^{m+1} + \cdots + \beta_{m+q-1} Z^{m+q-1}, \\ R_q(Z) &= \gamma_{m+1} Z^{m+1} + \cdots + \gamma_{m+q} Z^{m+q}, \quad \text{with} \\ \deg P_q &= m + (q - 1), \quad \deg Q_q = m + (q - 1), \\ \deg R_q &= m + q \quad \text{such that} \end{aligned}$$

$$(1) \ x - (A \cdot P_q(D) + BQ_q(D) + R_q(D)) \in J^{4(m+q)+2, -4q}.$$

Furthermore

$$\begin{aligned} P_{q+1}(Z) &= P_q(Z) + \alpha_{m+q} Z^{m+q}, \\ Q_{q+1}(Z) &= Q_q(Z) + \beta_{m+q} Z^{m+q}, \\ R_{q+1}(Z) &= R_q(Z) + \gamma_{m+q+1} Z^{m+q+1}. \end{aligned}$$

If

$$\begin{aligned} P(Z) &= \sum_{i=m}^{\infty} \alpha_i Z^i \in \Lambda_{4m} \\ Q(Z) &= \sum_{i=m}^{\infty} \beta_i Z^i \in \Lambda_{4m} \\ R(Z) &= \sum_{i=m+1}^{\infty} \gamma_i Z^i \in \Lambda_{4m+2} \end{aligned}$$

then by using (1) and Section I we have $x = AP(D) + BQ(D) + R(D)$.

The cases $2n = 4m + 2 < 0$ and $2n = 4m$ are similar. \square

The next two propositions will be used later on.

PROPOSITION 2.6. *If*

$$H(Z) = \sum_{i=0}^{\infty} \alpha_i Z^i \in \Lambda_{2n}$$

is such that $H(D) = 0$, then $\alpha_0 = 0$ and if α_p is the leading coefficient, we have $\alpha_p \in 8 \cdot U^(pt)$.*

Proof. Since $D \in \tilde{U}^*(B\Gamma)$ we have

$$\sum_{i=1}^{\infty} \alpha_i D^i = D \left(\sum_{i=1}^{\infty} \alpha_i D^{i-1} \right) \in \tilde{U}^*(B\Gamma);$$

then $\alpha_0 \cdot 1 \in \tilde{U}^*(B\Gamma) \cap U^*(pt) = \{0\}$ and $\alpha_0 \cdot 1 = 0$. If i denotes the inclusion $\{*\} \subset B\Gamma$ we have $i^*(\alpha_0 \cdot 1) = \alpha_0 = 0$. Then $H(Z) =$

$\alpha_p Z^p + \cdots + \alpha_m Z^m + \cdots$, $\alpha_p \neq 0$, $p \geq 1$. From $\alpha_q D^q \in J^{4q, 2n-4q} \subset J^{4p+4, 2n-(4p+4)}$, $q \geq p+1$, it follows that $t_q = \alpha_{p+1} D^{p+2} + \cdots + \alpha_q D^q \in J^{4p+4, 2n-(4p+4)}$, $q \geq p+1$. Since $J^{4p+4, 2n-(4p+4)}$ is closed for the topology T of $U^{2n}(B\Gamma)$ we have

$$\sum_{i=p+1}^{\infty} \alpha_i D^i \in J^{4p+4, 2n-(4p+4)} \subset J^{4p+1, 2n-(4p+1)}.$$

Let s be the quotient map

$$\begin{aligned} J^{4p, 2n-4p} &\rightarrow J^{4p, 2n-4p} / J^{4p+1, 2n-(4p+1)} \\ &= H^{4p}(B\Gamma, U^{2n-4p}(pt)) = H^{4p}(B\Gamma) \otimes U^{2n-4p}(pt) \\ &= \mathbb{Z}_8 \otimes U^{2n-4p}(pt) = U^{2n-4p}(pt) / 8 \cdot U^{2n-4p}(pt). \end{aligned}$$

Then:

$$0 = s(H(D)) = s(\alpha_p D^p) + s\left(\sum_{i=p+1}^{\infty} \alpha_i D^i\right) = s(\alpha_p D^p) = \alpha_p \otimes d^p;$$

since d^p is a generator of $H^{4p}(B\Gamma)$ we have $\alpha_p \in 8U^{2n-4}(pt)$. \square

Let F be the formal group law and $[2](Y) = F(Y, Y)$; if ρ is the nontrivial unitary irreducible representation for \mathbb{Z}_2 then we get (see [9]):

PROPOSITION 2.7. $U^*(B\mathbb{Z}_2) = U^*(pt)[[Y]]/([2](Y))$ and the image of Y by the quotient map: $U^*(pt)[[Y]] \rightarrow U^*(B\mathbb{Z}_2)$ is the Euler class $e(\rho)$. \square

We have adopted the following graduation in 2.7: if

$$F(X, Y) = X + Y + a_{11}XY + \sum_{i \geq 1, j \geq 1} a_{ij}X^iY^j,$$

then $|a_{ij}| = 2(1-i-j)$, $|X| = |Y| = 2$; so $F(X, Y) \in \Lambda_2$. We shall often make use of the coefficient a_{11} . We know that there is a unique formal power series $[-1](Y) \in U^*(pt)[[Y]] \subset \Lambda_2$ such that: $F(Y, [-1](Y)) = 0$.

PROPOSITION 2.8. *There is $P_0(Z) \in \Omega_2$, $P_0(Z) = b_1Z + \sum_{i \geq 1} b_i Z^i$ such that $cf_1(\eta) = P_0(D)$. The coefficients b_i , $i \geq 1$, are determined by the relation $\sum_{i \geq 1} b_i (Y \cdot [-1](Y))^i = Y + [-1]Y$; in particular $b_1 = -a_{11}$.*

Proof. We have seen that if θ is the universal $Sp(1)$ -bundle over $Sp(1) = BS^3$ considered as a $U(2)$ -vector bundle then $\eta = (Bp)^*(\theta)$,

$p: \Gamma \subset Sp(1)$. As $H^*(BS^3) = \mathbb{Z}[u]$, $u = c_2(\theta)$, we have $U^*(BS^3) = U^*(pt)[[V]]$, $V = e(\theta)$, the Euler class of θ for MU . Hence there is $P_0(Z) = \sum_{i \geq 1} b_i Z^i \in \Omega_2$ such that $P_0(V) = cf_1(\theta)$; it follows that

$$cf_1(\eta) = (Bp)^*(cf_1(\theta)) = (Bp)^* \left(\sum_{i \geq 1} b_i V^i \right) = \sum_{i \geq 1} b_i D^i = P_0(D).$$

The relation $\sum_{i \geq 1} b_i (Y \cdot [-1]Y)^i = Y + [-1](Y)$ is proved in the Appendix part B and gives $b_1 = -a_{11}$. \square

We recall that $A = cf_1(\xi_i) \in \tilde{U}^2(B\Gamma)$, $B = cf_1(\xi_j) \in \tilde{U}^2(B\Gamma)$, $D = cf_2(\eta) \in \tilde{U}^4(B\Gamma)$; let $C \in \tilde{U}^2(B\Gamma)$ be $cf_1(\xi_k)$.

PROPOSITION 2.9. (a) *There are $P(Z) \in \Omega_2$, $Q(Z) \in \Omega_4$, $P(Z) = -4a_{11}Z + \sum_{i \geq 2} \alpha_i Z^i$, $Q(Z) = 4Z + \sum_{i \geq 2} \beta_i Z^i$, $\beta_2 \notin 2U^*(pt)$, such that $cf_1(\eta^2) = P(D) = A + B + C$, $cf_2(\eta^2) = Q(D) = AB + BC + CA$.*

(b) $cf_3(\eta^2) = ABC = 0$,

(c) $A^3 = -AQ(D) + A^2P(D)$, $B^3 = -BQ(D) + B^2P(D)$.

Proof. (a) Let $g: B\Gamma \rightarrow BU(2)$ be a map classifying η ; then η^2 is classified by the composite: $B\Gamma \xrightarrow{\Delta} B\Gamma \times B\Gamma \xrightarrow{g \times g} BU(2) \times BU(2) \xrightarrow{m} BU(4)$, where m is a map classifying $\gamma(2) \otimes \gamma(2)$ and Δ the diagonal map. We have $U^*(BU(2) \times BU(2)) = U^*(pt)[[c_1, c_2, c'_1, c'_2]]$, c_1, c_2, c'_1, c'_2 being respectively the images of $cf_1(\gamma(2)) \otimes 1$, $cf_2(\gamma(2)) \otimes 1$, $1 \otimes cf_1(\gamma(2))$, $1 \otimes cf_2(\gamma(2))$ by the canonical map: $U^*(BU(2)) \otimes U^*(BU(2)) \xrightarrow{X} U^*(BU(2) \times BU(2))$. Since the following diagram commutes:

$$\begin{array}{ccc} U^*(BU(4)) \xrightarrow{m^*} U^*(BU(2) \times BU(2)) & \xrightarrow{(g \times g)^*} & U^*(B\Gamma \times B\Gamma) \xrightarrow{\Delta^*} U^*(B\Gamma) \\ & \uparrow X & \uparrow \nearrow \cup \\ U^*(BU(2)) \otimes U^*(BU(2)) & \xrightarrow{X} & U^*(B\Gamma) \otimes U^*(B\Gamma) \end{array}$$

we must substitute $cf_1(\eta)$ for c_1, c'_1 , $cf_2(\eta)$ for c_2, c'_2 in $m^*(cf_1(\gamma(4)))$, $m^*(cf_2(\gamma(4)))$, $m^*(cf_3(\gamma(4)))$ in order to calculate $cf_1(\eta^2)$, $cf_2(\eta^2)$, $cf_3(\eta^2)$ (see Sec. I).

We have $m^*(cf_1(\gamma(4))) = \sum a_{(u,v)} c_1^{u_1} c_2^{u_2} c'_1{}^{v_1} c'_2{}^{v_2}$, $u = (u_1, u_2)$, $v = (v_1, v_2)$, $u_1 \geq 0$, $u_2 \geq 0$, $v_1 \geq 0$, $v_2 \geq 0$. It is important to calculate $a_{(u,v)}$ when $u_1 = u_2 = 0$, or $v_1 = v_2 = 0$.

Suppose $u_1 = u_2 = 0$. We denote by 0 the pair $(0, 0)$. Then the coefficients $a_{(0,v)}$ are given by $i^* \circ m^*(cf_1(\gamma(4)))$, i being the natural inclusion:

$$\{*\} \times BU(2) \xrightarrow{i} BU(2) \times BU(2).$$

Since $i^* \circ m^*(\gamma(4)) = \gamma(2) + \gamma(2)$ we have $i^* \circ m^*(cf_1(\gamma(4))) = 2c'_1$. Similarly $a_{(u,0)} = 2c_1$. Hence

$$m^*(cf_1(\gamma(4))) = 2(c_1 + c'_1) + \sum_{\substack{\|u\| \geq 1 \\ \|v\| \geq 1}} a_{(u,v)} c_1^{u_1} c_2^{u_2} c'_1{}^{v_1} c'_2{}^{v_2}$$

where $\|u\| = u_1 + u_2$, $\|v\| = v_1 + v_2$.

We recall that $cf_1(\eta) = P_0(D)$, $P_0(Z) \in \Omega_2$, $\nu'(P_0) = 1$, $\nu' = \frac{1}{4}\nu$ (see Sec. I). Consider

$$\begin{aligned} P(Z) &= 2(P_0(Z) + P_0(Z)) + \sum_{\substack{\|u\| \geq 1 \\ \|v\| \geq 1}} a_{(u,v)} P_0^{u_1+v_1}(Z) Z^{u_2+v_2} \\ &= 4b_1 Z + \sum_{i \geq 2} \alpha_i Z^i, \end{aligned}$$

b_1 being the first coefficient $\neq 0$ of $P_0(Z)$ because $u_1 + v_1 + u_2 + v_2 \geq 2$ when $\|u\| \geq 1$, $\|v\| \geq 1$. Hence $cf_1(\eta^2) = P(D)$. We remark that $P(Z) \in \Omega_2$.

There are unique elements $b_{(u,v)} \in U^*(pt)$ such that $m^*(cf_2(\gamma(4))) = \sum b_{(u,v)} c_1^{u_1} c_2^{u_2} c'_1{}^{v_1} c'_2{}^{v_2}$. Then the coefficients $b_{(u,0)}$ and $b_{(0,v)}$ are given by $cf_2(\gamma(2) + \gamma(2)) = cf_1^2(\gamma(2)) + 2cf_2(\gamma(2))$. Hence

$$m^*(cf_2(\gamma(4))) = c_1^2 + c_1^2 + 2(c_2 + c_2^1) + \sum_{\|u\| \geq 1, \|v\| \geq 1} b_{u,v} c_1^{u_1} c_2^{u_2} c'_1{}^{v_1} c'_2{}^{v_2}.$$

Consider

$$\begin{aligned} Q(Z) &= 4Z + 2P_0^2(Z) + \sum_{\|u\| \geq 1, \|v\| \geq 1} b_{(u,v)} P_0^{u_1+v_1}(Z) Z^{u_2+v_2} \\ &= 4Z + \sum_{i \geq 2} \beta_i Z^i. \end{aligned}$$

Then $cf_2(\eta^2) = Q(D)$, $Q(Z) \in \Omega_4$.

Let q be the inclusion $\mathbb{Z}_2 \subset \Gamma$; since $(Bq)^*(\xi_h)$, $h = i, j, k$, are trivial by 2.2 we have $(Bq)^*(A) = (Bq)^*(B) = (Bq)^*(C) = 0$ and since $Q(D) = cf_2(\eta^2) = AB + BC + CA$ we have $(Bq)^*(Q(D)) = 0$. It follows by 2.7 that $(Bq)^*(D) = d^2$, d being the image of Y by the quotient map:

$$U^*(pt)[[Y]] \rightarrow U^*(pt)[[Y]]/([2](Y)).$$

Thus:

$$\begin{aligned} 4Y^2 + \sum_{i \geq 2} \beta_i \cdot Y^{2i} &= [2](Y) \cdot G(Y) \\ &= (2Y + a_{11}Y^2 + a_3Y^3 + \cdots)(\varepsilon_0Y + \varepsilon_1Y^2 + \varepsilon_2Y^3 + \cdots) \quad \text{and} \end{aligned}$$

$$\begin{aligned}\varepsilon_0 &= 2, & 0 &= 2\varepsilon_1 + a_{11}\varepsilon_0 = 2(\varepsilon_1 + a_{11}); & \text{so} \\ \varepsilon_1 &= -a_{11}, & \beta_2 &= 2\varepsilon_2 - a_{11}^2 + 2a_3;\end{aligned}$$

since $a_{11}^2 \notin 2U^*(pt)$ (because $U^*(pt) = [x_1, x_2, \dots]$, $a_{11} = -x_1$) it follows that $\beta_2 \notin 2U^*(pt)$. The relations $P(D) = A + B + C$, $Q(D) = AB + BC + CA$ are easy consequences of the relation $\eta^2 = 1 + \xi_i + \xi_j + \xi_k$.

(b) The above relation gives $cf_3(\eta^2) = ABC$; in order to show that $ABC = 0$ we consider the Boardman map $Bd: U^*(B\Gamma) \rightarrow K^*(B\Gamma) \hat{\otimes} \mathbb{Z}[a_1, a_2, \dots]$ (see [8], page 358). This map is a ring-homomorphism which is injective because $B\Gamma$ has a periodic cohomology; furthermore if τ is a line complex vector bundle over $B\Gamma$ we have:

$$Bd(e(\tau)) = (\tau - 1) + (\tau - 1)^2 \otimes a_1 + (\tau - 1)^3 \otimes a_2 + \dots;$$

as $(\xi_i - 1)(\xi_j - 1)(\xi_k - 1) = 0$ we get $Bd(ABC) = 0$ and $ABC = 0$.

(c) We have $Q(D) = A(B+C) + BC = A(P(D) - A) + BC$; as $ABC = 0$ we obtain $A^3 = -AQ(D) + A^2P(D)$; similarly $B^3 = -AQ(D) + A^2P(D)$. \square

PROPOSITION 2.10. *There is $S(Z) = -a_{11}Z + \sum_{i \geq 2} s_i \cdot Z^i \in \Omega_2$ such that $A^2 = AS(D)$, $B^2 = BS(D)$. Moreover:*

$$AB = (A + B)(P(D) - S(D)) - Q(D),$$

$P(Z)$, $Q(Z)$ being as in 2.9.

Proof. Consider the relation $\eta\xi_i = \eta$. If the vector bundle $\gamma(2) \otimes \gamma(1)$ over $B\mathbb{U}(2) \times B\mathbb{U}(1)$ is classified by $m_1: B\mathbb{U}(2) \times B\mathbb{U}(1) \rightarrow B\mathbb{U}(2)$ and if $g: B \rightarrow B\mathbb{U}(2)$, $h: B \rightarrow B\mathbb{U}(1)$ are classifying maps for η and ξ_i , then $\eta\xi_i$ is classified by:

$$B\Gamma \xrightarrow{\Delta} B\Gamma \times B\Gamma \xrightarrow{g \times h} B\mathbb{U}(2) \times B\mathbb{U}(1) \xrightarrow{m_1} B\mathbb{U}(2).$$

We have the following commutative diagram:

$$\begin{array}{ccc} U^*(B\mathbb{U}(2)) \xrightarrow{m_1^*} U^*(B\mathbb{U}(2)) \times U^*(B\mathbb{U}(1)) & \xrightarrow{(g \times h)^*} & U^*(B\Gamma \times B\Gamma) \xrightarrow{\Delta^*} U^*(B\Gamma) \\ & \uparrow x & \uparrow x \quad \nearrow \text{cup-product} \\ U^*(B\mathbb{U}(2)) \otimes U^*(B\mathbb{U}(1)) & \xrightarrow{g^* \times h^*} & U^*(B\Gamma) \otimes U^*(B\Gamma). \end{array}$$

Moreover $U^*(B\mathbb{U}(2) \times B\mathbb{U}(1)) = U^*(pt)[[c_1, c_2, c'_1]]$ where c_1, c_2, c'_1 are the images respectively of $cf_1\gamma(2) \otimes 1$, $cf_2\gamma(2) \otimes 1$, $1 \otimes cf_1\gamma(1)$ by the canonical map: $U^*(B\mathbb{U}(2)) \times U^*(B\mathbb{U}(1)) \xrightarrow{X} U^*(B\mathbb{U}(2) \times B\mathbb{U}(1))$. Then

$$m_1^*(cf_2(\gamma(2))) = \sum e_{(u,v)} c_1^{u_1} c_2^{u_2} c'_1{}^v, \quad u = (u_1, u_2).$$

If i and j are the natural inclusions: $BU(2) \times \{*\} \rightarrow BU(2) \times BU(1)$ and $\{*\} \times BU(1) \rightarrow BU(2) \times BU(1)$, then the coefficients $e_{(u,0)}$ and $e_{(0,v)}$ are given respectively by $i^* \circ m^*(cf_2(\gamma(2))) = cf_2(\gamma(2)) = c_2$ and $j^* \circ m^*(cf_2(\gamma(2))) = cf_2(\gamma(1) + \gamma(1)) = c_1'^2$. Hence

$$\begin{aligned} m_1^*(cf_2(\gamma(2))) &= c_2 + c_1'^2 + \sum_{\substack{\|u\| \geq 1 \\ v \geq 1}} e_{(u,v)} c_1^{u_1} c_2^{u_2} c_1'^v \\ &= c_2 + c_1'^2 + c_1' N_1(c_1, c_2) + c_1'^2 N_2(c_1, c_2) \\ &\quad + \cdots + c_1'^m N_m(c_1, c_2) + \cdots . \end{aligned}$$

To calculate $cf_2(\eta \cdot \xi_i)$ we substitute $cf_1(\eta)$, $cf_2(\eta)$, $cf_1(\xi_i)$, respectively for c_1 , c_2 , c_1' . We recall that $cf_1(\eta) = P_0(D)$, $\nu'(P_0) = 1$ ($\nu' = \frac{1}{4}\nu$; see Sec. I). We can substitute $P_0(Z)$ for c_1 and Z for c_2 in $N_m(c_1, c_2)$ to obtain $M_m(Z) \in \Omega_*$, $\nu'(M_m) \geq 1$, $m \geq 1$. We need to calculate the leading coefficient of $M_1(Z)$. To this purpose consider $T = BU(1) \times BU(1)$ and $r: T \rightarrow BU(2)$ a map classifying $\pi_1^*(\gamma(1)) + \pi_2^*(\gamma(1))$, π_1, π_2 being respectively the first and second projections $T \rightarrow BU(1)$; we have $U^*(T \times BU(1)) = U^*(pt)[[e_1, f_1, e_1']]$ with $(r \times 1)^*(c_1) = e_1 + f_1$, $(r \times 1)^*(c_2) = e_1 f_1$, $(r \times 1)^*(c_1') = e_1'$; it is easily seen that $(r \times 1)^*(m_1^* cf_2(\gamma(2))) = F(e_1, e_1') F(f_1, e_1')$ where F denotes the formal group law. It follows that $e_{((1,0),1)} = 1$, $e_{((0,1),1)} = 2a_{11}$ and $M_1(Z) = a_{11}Z + \sum_{i \geq 2} b_i' Z^i$, $\nu'(M_1) = 1$.

Now from the relation $A^3 = -AQ(D) + A^2P(D)$ we deduce that $A^n = AQ_n(D) + A^2P_n(D)$, $n \geq 3$, with $Q_n(Z) \in \Omega_{2n-2}$, $P_n(Z) \in \Omega_{2n-4}$, $Q_3(Z) = -Q(Z)$, $P_3(Z) = P(Z)$, $Q_{n+1}(Z) = -Q(Z)P_n(Z)$, $P_{n+1}(Z) = P(Z)P_n(Z) + Q_n(Z)$. Then $\nu'(P_{n+1}) \geq \inf(\nu'(P_n), \nu'(P_{n-1}))$ and $\nu'(P_{n+1}) \geq (n+1)/2$; so:

$$\lim_{n \rightarrow \infty} \nu'(P_n) = \lim_{n \rightarrow \infty} \nu'(Q_n) = +\infty.$$

Consider

$$\begin{aligned} M_n(X, Z) &= Z + X^2[1 + M_2(Z) + P(Z)M_3(Z) + \cdots + P_n(Z)M_n(Z)] \\ &\quad + X[M_1(Z) + Q_3(Z)M_3(Z) + \cdots + Q_n(Z)M_n(Z)] \in \Lambda_4. \end{aligned}$$

As

$$\lim_{n \rightarrow \infty} \nu(P_n M_n) = \lim_{n \rightarrow \infty} \nu(Q_n M_n) = +\infty$$

it follows that $\lim_{n \rightarrow \infty} M_n(X, Z)$ exists (see Sec. I) and may be written as: $Z + X^2[1 + H(Z)] + XH_1(Z)$ with $H(Z) \in \Omega_0$, $\nu'(H) \geq 1$. We remark that the leading coefficient of $H_1(Z)$ is that of $M_1(Z)$; so: $H_1(Z) = a_{11}Z + \sum_{i \geq 2} d_i Z^i \in \Omega_2$. Thus: $cf_2(\eta \xi_i) = D + A^2[1 + H(D)] + AH_1(D) = cf_2(\eta) = D$ and $A^2[1 + H(D)] =$

$-AH_1(D)$. Let $E(Z) \in \Omega_0$ be such that $E(Z)(1 + H(Z)) = 1$; hence $A^2 = AS(D)$ with $S(Z) = -H_1(Z)E(Z) = -a_{11}Z + \sum_{i \geq 2} s_i Z^i \in \Omega_2$. Similarly $B^2 = BS(D)$. Now

$$\begin{aligned} AB &= AB + BC + CA - C(A + B) \\ &= Q(D) - [P(D) - (A + B)] \cdot (A + B) \\ &= Q(D) - P(D) \cdot (A + B) + 2AB + (A + B)S(D) \\ &= 2AB + Q(D) + (A + B)(S(D) - P(D)). \end{aligned}$$

Then:

$$AB = (A + B)[P(D) - S(D)] - Q(D). \quad \square$$

LEMMA 2.11. *There is $T(Z) = 8Z + 2\lambda_2 Z^2 + \sum_{i \geq 3} \lambda_i Z^i \in \Omega_4$, $\lambda_2 \notin 2U^*(pt)$ and $T(D) = 0$.*

Proof. From $\eta^2 = 1 + \xi_i + \xi_j + \xi_k$ we get $\eta^3 = 4\eta$. Let $g_1: B\Gamma \rightarrow BU(4)$ and $g: B\Gamma \rightarrow BU(2)$ be classifying maps (respectively) for η^2 and η ; then η^3 is classified by: $B\Gamma \xrightarrow{\Delta} B\Gamma \times B\Gamma \xrightarrow{g_1 \times g} BU(4) \times BU(2) \xrightarrow{m_2} BU(8)$, m_2 being a map classifying $\gamma(4) \otimes \gamma(2)$. Then we get $m_2^*(cf_2(\gamma(8))) = \sum f_{(u,v)} c_1^{u_1} c_2^{u_2} c_3^{u_3} c_4^{u_4} c_1^{v_1} c_2^{v_2}$, with $u = (u_1, u_2, u_3, u_4)$, $v = (v_1, v_2)$. The coefficients $f_{(u,0)}$ and $f_{(0,v)}$ are given respectively by $cf_2(\gamma(4) + \gamma(4)) = c_1^2 + 2c_2$ and $cf_2(4\gamma(2)) = 6c_1'^2 + 4c_2'$. Thus

$$\begin{aligned} m_2^*(cf_2(\gamma(8))) &= c_1^2 + 2c_2 + 6c_1'^2 + 4c_2' \\ &\quad + \sum_{\substack{\|u\| \geq 1 \\ \|v\| \geq 1}} f_{(u,v)} c_1^{u_1} c_2^{u_2} c_3^{u_3} c_4^{u_4} c_1^{v_1} c_2^{v_2}. \end{aligned}$$

In order to calculate $cf_2(\eta^3)$ we must substitute $cf_1(\eta^2) = P(D)$, $cf_2(\eta^2) = Q(D)$, $cf_3(\eta^2) = 0$, $cf_4(\eta^2) = 0$, $cf_1(\eta) = P_0(D)$, $cf_2(\eta) = D$ respectively for $c_1, c_2, c_3, c_4, c_1', c_2'$. Consider

$$\begin{aligned} E(Z) &= P^2(Z) + 2Q(Z) + 6P_0^2(Z) + 4Z \\ &\quad + \sum_{\|u\| \geq 1, \|v\| \geq 1} f_{(u,v)} P^{u_1}(Z) Q^{u_2}(Z) P_0^{v_1}(Z) \cdot Z^{v_2}, \end{aligned}$$

$u = (u_1, u_2, 0, 0)$, $v = (v_1, v_2)$. Hence $E(D) = cf_2(\eta^3)$; but as the leading coefficients of $P(Z)$ and $Q(Z)$ belong to $4U^*(pt)$, $E(Z)$ has the form: $2Q(Z) + 6P_0^2(Z) + 4Z + 4\tau Z^2 + \sum_{i \geq 3} \tau_i Z^i$. So: $E(D) = 2Q(D) + 6P_0^2(D) + 4D + 4\tau D^2 + \sum_{i \geq 3} \tau_i D^i = cf_2(\eta^3) = cf_2(4\eta) = 6cf_1^2(\eta) + 4cf_2(\eta) = 6P_0^2(D) + 4D$. Hence if $T(Z) = 2Q(Z) + 4\tau Z^2 + \sum_{i \geq 3} \tau_i Z^i \in \Omega_4$, then $T(D) = 0$. As $Q(Z) = 4Z + \sum_{i \geq 2} \beta_i Z^i$, $\beta_2 \notin 2U^*(pt)$, we have: $T(D) = 8Z + 2\lambda_2 Z^2 + \sum_{i \geq 3} \lambda_i Z^i$, $\lambda_2 \notin 2U^*(pt)$. \square

THEOREM 2.12. *If $M(Z) \in \Omega_*$ is such that $M(D) = 0$, then $M(Z) \in \Omega_* T(Z)$.*

Proof. We may suppose $M(Z) \in \Omega_{2n}$, $n \in \mathbb{Z}$. If $M(Z) = \omega_0 + \sum_{i \geq 1} \omega_i Z^i$, then by 2.6 we have $\omega_0 = 0$ and the first coefficient $\omega_i \neq 0$, say ω_{P_0} , is such that $P_0 \geq 1$, $\omega_{P_0} \in 8U^*(pt)$. Thus $M(Z) = 8\omega'_{P_0} Z^{P_0} + \sum_{i > P_0} \omega_i Z^i$. Consider $M_1(Z) = M(Z) - \omega'_{P_0} \cdot Z^{P_0-1} \cdot T(Z) \in \Omega_{2n}$. We have $\nu(M_1(Z)) > \nu(M(Z))$ and $M_1(D) = 0$. Then $M_1(Z) = 8\omega'_{P_1} Z^{P_1} + \sum_{i > P_1} \theta_i \cdot Z^i$, $P_1 > P_0$. We form

$$M_2(Z) = M_1(Z) - \omega'_{P_1} Z^{P_1-1} T(Z)$$

and then $\nu(M_2) > \nu(M_1)$, $M_2(D) = 0$. After a finite number of steps we get $M_{r+1}(Z) = M(Z) - (\omega'_{P_0} Z^{P_0-1} + \dots + \omega'_{P_r} Z^{P_r-1}) T(Z)$ such that $P_r > P_{r-1} > \dots > P_1 > P_0$, $\nu(M_{r+1}) > \nu(M_r) > \dots > \nu(M_1) > \nu(M)$ and $M_{r+1}(D) = 0$. Since $\lim_{r \rightarrow \infty} \nu(M_r) = \infty$ it follows that $M(Z) = (\sum_{k \geq 0} \omega'_{P_k} \cdot Z^{P_k-1}) \cdot T(Z)$ (see Sec. I). \square

LEMMA 2.13. *There is $J(Z) = \mu_1 Z + \sum_{i \geq 2} \mu_i Z^i \in \Omega_0$, $\mu_1 \notin 2U^*(pt)$, such that $A[2 + J(D)] = B[2 + J(D)] = 0$.*

Proof. We have $[2](Y) = 2Y + a_{11} Y^2 + \sum_{i \geq 3} a_i Y^i$. As ξ_i^2 is trivial we have $[2](A) = 0$ and from $A^2 = AS(D)$ ($S(Z) \in \Omega_2$) we get $A^n = AS^{n-1}(D)$. Consider $H_n(X, Z) = X[2 + a_{11} S(Z) + \dots + a_n S^{n-1}(Z)]$. Since $\lim_{n \rightarrow \infty} \nu(S^n) = \infty$ it follows that $\lim_{n \rightarrow \infty} H_n(X, Z)$ exists and has the form $X[2 + J(Z)]$, with

$$J(Z) = a_{11} S(Z) + \sum_{n \geq 3} a_n S^{n-1}(Z) = -a_{11}^2 Z + \sum_{i \geq 2} \mu_i Z^i.$$

If $\mu_1 = -a_{11}^2$ we see that $\mu_1 \notin 2U^*(pt)$. Thus $A(2+J(D)) = [2](A) = 0$. Similarly $B(2+J(D)) = 0$. \square

LEMMA 2.14. *Suppose $XM(Z) + YN(Z) + E(Z) \in \Omega_*$ is such that $AM(D) + BN(D) + E(D) = 0$. Then the first coefficient $\neq 0$ of $M(Z)$ and the first coefficient $\neq 0$ of $N(Z)$ belong to $2U^*(pt)$.*

Proof. We may suppose $XM(Z) \in \Omega_{2n}$, $YN(Z) \in \Omega_{2n}$, $E(Z) \in \Omega_{2n}$, $n \in \mathbb{Z}$. We shall give a proof in the case: $0 \neq M(Z) = a_p Z^p + a_{p+1} Z^{p+1} + \dots$, $a_p \neq 0$, $0 \neq N(Z) = b_q Z^q + b_{q+1} Z^{q+1} + \dots$, $b_q \neq 0$ and $p \leq q$. We observe that if $s \geq p$ then $A(a_p D^p + \dots + a_{p+s} D^{p+s}) \in J^{4p+2, 2n-4p-2}$ and consequently $AM(D) \in J^{4p+2, 2n-4p-2}$ because the subgroups $J^{*,*}$ are closed in $U^*(B\Gamma)$. Similarly

$$A(a_{p+1} D^{p+1} + \dots + a_r D^r + \dots) \in J^{4p+6, 2n-4p-6}$$

and consequently

$$A(a_{p+1}D^{p+1} + \cdots + a_r D^r + \cdots) \in J^{4p+3, 2n-4p-3}.$$

There are similar remarks concerning $BN(D)$. Since by hypothesis $p \leq q$ we have $4p+2 \leq 4q+2$ and $J^{4p+2, 2n-4p-2} \supset J^{4q+2, 2n-4q-2}$. We shall denote by g the quotient map:

$$\begin{aligned} J^{4p+2, 2n-4p-2} &\rightarrow J^{4p+2, 2n-4p-2} / J^{4p+3, 2n-4p-3} \\ &= [U^h(pt)/2U^h(pt)] \oplus [U^h(pt)/2U^h(pt)], \end{aligned}$$

with $h = 2n - 4p - 2$. Then $g(AM(D)) = \bar{a}_p$, \bar{a}_p being the image of a_p by the quotient map

$$U^h(pt) \rightarrow U^h(pt)/2U^h(pt),$$

$U^h(pt)/2U^h(pt)$ being the first summand.

(a) Suppose $E(D) = 0$.

(i) $p = q$. We have $g(AM(D)) = \bar{a}_p$ and $g(BM(D)) = \bar{b}_p$ respectively in the first and second summand of the sum $[U^h(pt)/2U^h(pt)] \oplus [U^h(pt)/2U^h(pt)]$. Since $AM(D) + BM(D) = 0$ we have $\bar{a}_p = 0$, $\bar{b}_p = 0$ and thus $a_p \in 2U^*(pt)$, $b_p \in 2U^*(pt)$.

(ii) $p < q$. From $J^{4p+2, 2n-4p-2} \supset J^{4p+3, 2n-4p-3} \supset J^{4q+2, 2n-4q-2}$ it follows that $g(BN(D)) = 0$ and consequently $\bar{a}_p = 0$ which means that $a_p \in 2U^*(pt)$.

(b) Suppose $E(D) \neq 0$.

Take $E(Z) = d_0 + \sum_{i \geq 1} d_i Z^i$. As $E(D) = -(AM(D) + BN(D)) \in \tilde{U}^*(B\Gamma)$ we have $d_0 = 0$. Hence:

$$E(Z) = \sum_{i \geq r} d_i Z^i, \quad d_r \neq 0, \quad r \geq 1.$$

If $d_r = 8e_{r_1}$, we form

$$\begin{aligned} E_1(Z) &= E(Z) - e_{r_1} Z^{r-1} T(Z) \\ &= \sum_{i \geq r'} d'_i Z^i, \quad r' > r, \quad d'_{r'} \neq 0 \text{ or } \nu(E_1) > \nu(E). \end{aligned}$$

If $d'_{r'} = 8e_{r_2}$ we form $E_2(Z) = E_1(Z) - e_{r_2} Z^{r'-1} T(Z)$ and so on. But after a finite number of steps we have $E_{p_0}(Z) = \sum_{i \geq h} t_i Z^i$ and $t_h \notin 8U^*(pt)$ because, if not, we would have $E(Z) \in \Omega_* T(Z)$ and thus $E(D) = 0$ which contradicts the hypothesis (b): $E(D) \neq 0$ (see the proof of 2.12). Hence there is a formal power series $F(Z) \in \Omega_{2n}$ such that $F(D) = E(D)$ and $F(Z) = \sum_{i \geq h \geq 1} t_i Z^i$, $t_h \notin 8U^*(pt)$. This means that $E(D) \in J^{4h, 2n-4h}$ and $E(D) \notin J^{4h+1, 2n-4h-1}$.

(i) $p = q$, $4h < 4p + 2 = 4q + 2$.

Then: $J^{4h, 2n-4h} \supset J^{4h+1, 2n-4h-1} \supset J^{4p+2, 2n-4p-2}$. Since $E(D) = -(AM(D) + BN(D))$ we have $E(D) \in J^{4h+1, 2n-4h-1}$ which is impossible.

(ii) $p = q$, $4p + 2 = 4q + 2 < 4h$.

Then $J^{4p+2, 2n-4p-2} \supset J^{4p+3, 2n-4p-3} \supset J^{4h, 2n-4h}$ and $AM(D) + BN(D) = -E(D) \in J^{4p+3, 2n-4p-3}$. Consequently $\bar{a}_p = 0$, $\bar{b}_p = 0$ and thus $a_p \in 2U^*(pt)$, $b_p \in 2U^*(pt)$.

(iii) $p < q$, $4h < 4p + 2 < 4q + 2$.

Then $J^{4h, 2n-4h} \supset J^{4p+2, 2n-4p-2} \supset J^{4q+2, 2n-4q-2}$. From $E(D) = -(AM(D) + BN(D))$ it follows that

$$E(D) \in J^{4p+2, 2n-4p-2} \subset J^{4h+1, 2n-4h-1} (\subset J^{4h, 2h-4h})$$

which is impossible.

(iv) $p < q$, $4p + 2 < 4h < 4q + 2$ or $4p + 2 < 4q + 2 < 4h$.

We have either

$$J^{4p+2, 2n-4p-2} \supset J^{4p+3, 2n-4p-3} \supset J^{4h, 2n-4h} \supset J^{4q+2, 2n-4q-2}$$

or

$$J^{4p+2, 2n-4p-2} \supset J^{4p+3, 2n-4p-3} \supset J^{4q+2, 2n-4q-2} \supset J^{4h, 2n-4h}.$$

It follows in both cases that $\bar{a}_p = 0$ and $a_p \in 2U^*(pt)$. Hence we have proved that if $p \leq q$ we have $a_p \in 2U^*(pt)$ in both cases $E(D) = 0$, $E(D) \neq 0$. So $M(Z) = a_p Z^p + a_{p+1} Z^{p+1} + \dots$, $a_p = 2e_p \neq 0$. By 2.13 if $K(X, Z) = X(2 + J(Z))$ then $K(A, D) = 0$. We form $XM(Z) - e_p Z^p K(X, Z) = XM_1(Z)$, $M_1(Z) = e_{p_1} Z^{p_1} + \dots$, $p_1 > p$, and we get: $AM_1(D) + BN(D) + E(D) = 0$. If $p_1 < q$ we carry on the same process and after a finite number of steps there is $M_r(Z) \in \Lambda_{2n-2}$ such that $AM_r(D) + BN(D) + E(D) = 0$ and $q \leq p_r$, p_r being such that $M_r(Z) = \omega_{p_r} Z^{p_r} + \omega_{p_r+1} Z^{p_r+1} + \dots$, $\omega_{p_r} \neq 0$. Thus the argument used is the case $p \leq q$ (above) shows that $b_q \in 2U^*(pt)$. \square

Let I'_* the graded ideal of Λ_* generated by $K(X, Z) = X(2 + J(Z)) \in \Lambda_2$, $K(Y, Z) = Y \cdot (2 + J(Z)) \in \Lambda_2$ and $T(Z) \in \Omega_4$ (see 2.13, 2.12).

LEMMA 2.15. *Let $M(Z)$, $N(Z)$, $E(Z)$ be elements of Ω_* such that $AM(D) + BN(D) + E(D) = 0$. Then: $XM(Z) + YN(Z) + E(Z) \in K(X, Z)\Omega_* + K(Y, Z)\Omega_* + T(Z)\Omega_* \subset I'_*$ and $AM(D) = BN(D) = E(D) = 0$.*

Proof. Suppose $XM(Z) \in \Lambda_{2n}$, $YN(Z) \in \Lambda_{2n}$, $E(Z) \in \Lambda_{2n}$, $n \in \mathbb{Z}$. We shall give a proof in the case $M(Z) \neq 0$, $N(Z) \neq 0$, the other cases

being simpler. Take $P(X, Y, Z) = XM(Z) + YN(Z) + E(Z)$, $M(Z) = a_{p_0}Z^{p_0} + a_{p_0+1}Z^{p_0+1} + \dots$, $a_{p_0} \neq 0$, $N(Z) = b_{q_0}Z^{q_0} + b_{q_0+1}Z^{q_0+1} + \dots$, $b_{q_0} \neq 0$. By 2.14 we have $a_{p_0} = 2a'_{p_0}$, $b_{q_0} = 2b'_{q_0}$ and then: $P(X, Y, Z) - (a'_{p_0}Z^{p_0}K(X, Z) + b'_{q_0}Z^{q_0}K(Y, Z)) = X[M(Z) - a'_{p_0}Z^{p_0}(2 + J(Z))] + Y[N(Z) - b'_{q_0}Z^{q_0}(2 + J(Z))] + E(Z) = XM_1(Z) + YN_1(Z) + E(Z)$ with $\nu(M) < \nu(M_1)$, $\nu(N) < \nu(N_1)$. Moreover we have $AM_1(D) + BN_1(D) + E(D) = P(A, B, D) = 0$. The same process can be carried out for $XM_1(Z) + YN_1(Z) + E(Z)$ and after a finite number of operations we get $M_1(Z), M_2(Z), \dots, M_{r+1}(Z), N_1(Z), N_2(Z), \dots, N_{r+1}(Z)$,

$$P(X, Y, Z) - \left[\left(\sum_{i=0}^r a'_{p_i} Z^{p_i} \right) K(X, Z) + \left(\sum_{i=0}^r b'_{q_i} Z^{q_i} \right) K(Y, Z) \right] \\ = XM_{r+1}(Z) + YN_{r+1}(Z) + E(Z)$$

with $p_0 = \nu'(M) < p_1 = \nu'(M_1) < \dots < p_{r+1} = \nu'(M_{r+1})$, $q_0 = \nu'(N) < q_1 = \nu'(N_1) < \dots < q_{r+1} = \nu'(N_{r+1})$. Take

$$H_1(Z) = \sum_{i=0}^{\infty} a'_{p_i} Z^{p_i}, \quad H_2(Z) = \sum_{i=0}^{\infty} b'_{q_i} Z^{q_i}.$$

Since $\text{Lim}_{r \rightarrow \infty} \nu(M_r) = \text{Lim}_{r \rightarrow \infty} \nu(N_r) = +\infty$ we have $\text{Lim}_{r \rightarrow \infty} XM_r(Z) = \text{Lim}_{r \rightarrow \infty} YN_r(Z) = 0$ and there are $H_1(Z) \in \Omega_*$, $H_2(Z) \in \Omega_*$ such that: $P(X, Y, Z) - [H_1(Z)K(X, Z) + H_2(Z)K(Y, Z)] = E(Z)$. Since $P(A, B, D) = K(A, D) = K(B, D) = 0$ we have: $E(D) = 0$ and then by 2.12 there is $H_3(Z) \in \Omega_*$ such that $E(Z) = H_3(Z) \cdot T(Z)$. Finally we have $P(X, Y, Z) = H_1(Z)K(X, Z) + H_2(Z)K(Y, Z) + H_3(Z)T(Z) \in K(X, Z)\Omega_* + K(Y, Z)\Omega_* + T(Z)\Omega_* \subset I'_*$ and $XM(Z) = H_1(Z)K(X, Z)$, $YN(Z) = H_2(Z) \cdot K(Y, Z)$, $E(Z) = H_3(Z) \cdot T(Z)$. Consequently: $AM(D) = BN(D) = E(D) = 0$. \square

Consider $S(X, Z) = X^2 - XS(Z) \in \Lambda_4$, $S(Y, Z) = Y^2 - YS(Z) \in \Lambda_4$, $R(X, Y, Z) = XY - (X + Y)(P(Z) - S(Z)) + Q(Z) \in \Lambda_4$. By 2.10 we have: $S(A, D) = S(B, D) = R(A, B, D) = 0$. Let I''_* be the grade ideal of Λ_* generated by $S(X, Z), S(Y, Z), R(X, Y, Z)$.

LEMMA 2.16. *For any $P(X, Y, Z) \in \Lambda_*$ there are $M(Z), N(Z), E(Z)$, elements of Ω_* such that $P(X, Y, Z) - [XM(Z) + YN(Z) + E(Z)] \in I''_*$.*

Proof. From $X^2 - XS(Z) = S(X, Z)$ we see that there is $M_n(X, Z) \in \Lambda_*$ such that $X^n - XS^{n-1}(Z) = S(X, Z)M_n(X, Z)$, $n \geq 2$, with $M_2(X, Z) = 1$ and $M_{n+1}(X, Z) = S^{n-1}(Z) + XM_n(X, Z)$, $n \geq 2$. It is easily seen that $\text{Lim}_{n \rightarrow \infty} \nu(S^n) = \text{Lim}_{n \rightarrow \infty} \nu(M_n) = +\infty$. If $P(X, Y, Z) \in \Lambda_{2m}$ we

can write $P(X, Y, Z) = \sum_{i=0}^{\infty} X^i P_i(Y, Z)$ with $\dim P_i = 2(m - i)$. We have $X^i P_i(Y, Z) = X S^{i-1}(Z) P_i(Y, Z) + S(X, Z) M_i(X, Z) P_i(Y, Z)$, $i \geq 2$. From Section I and the fact that the multiplication by an element of Λ_* is continuous we see that there are $H(Y, Z)$, $H_1(X, Y, Z)$ such that: $P(X, Y, Z) = XH(Y, Z) + S(X, Z)H_1(X, Y, Z) + P_0(Y, Z)$. Similarly there are $F_0(Z)$, $F_1(Z)$, $F_2(Y, Z)$ such that $H(Y, Z) = YF_1(Z) + S(Y, Z)F_2(Y, Z) + F_0(Z)$ and $G_0(Z)$, $G_1(Z)$, $G_2(Y, Z)$ such that $P_0(Y, Z) = YG_1(Z) + S(Y, Z)G_2(Y, Z) + G_0(Z)$. Then a straightforward calculation shows that with $M(Z) = F_0(Z) + F_1(Z) \cdot (P(Z) - S(Z))$, $N(Z) = G_1(Z) + F_1(Z) \cdot (P(Z) - S(Z))$, $E(Z) = G_0(Z) - Q(Z) \cdot F_1(Z)$ we get $P(X, Y, Z) - [XM(Z) + YN(Z) + E(Z)] \in I_*''$. \square

Let I_* be $I_*' + I_*''$.

THEOREM 2.17. *The graded $U^*(pt)$ -algebra $U^*(B\Gamma)$ is isomorphic to Λ_*/I_* where I_* is a graded ideal generated by six homogeneous formal power series.*

Proof. Consider the map $\varphi: \Lambda_* \rightarrow U^*(B\Gamma)$ of graded $U^*(pt)$ -algebras such that $\varphi(X) = A$, $\varphi(Y) = B$, $\varphi(Z) = D$. By Theorem 2.5 φ is surjective and by Lemmas 2.15, 2.16 φ is injective. \square

REMARKS. (1) Consider the involution $h: \Lambda_* \rightarrow \Lambda_*$ such that $h(Y) = X$, $h(X) = Y$, $h(Z) = Z$. We have $h(I_*) = I_*$ and thus there is an isomorphism \bar{h} of graded $U^*(pt)$ -algebras: $U^*(B\Gamma) \rightarrow U^*(B\Gamma)$ such $\bar{h}(A) = B$, $\bar{h}(B) = A$, $\bar{h}(D) = D$. Consequently $\bar{h}^2 = \text{Id}$.

(2) If $q: \mathbb{Z}_2 \subset \Gamma$ denotes the canonical inclusion, then $(Bq)^*: U^*(B\Gamma) \rightarrow U^*(B\mathbb{Z}_2)$ is neither injective nor surjective.

An important and easy consequences of Theorem 2.12 and Lemma 2.15 is the following theorem which gives the structure of $U^*(pt)[[D]]$ -module of $U^*(B\Gamma)$.

THEOREM 2.18. (a) *As graded $U^*(pt)$ -algebras we have:*

$$U^*(pt)[[D]] \simeq \Omega_*/(T(Z)).$$

(b) *As graded $U^*(pt)[[D]]$ -modules we have: $U^*(B\Gamma) \simeq U^*(pt)[[D]] \oplus U^*(pt)[[D]]A \oplus U^*(pt)[[D]]B$ and: A and B have the same annihilator*

$$(2 + J(D))U^*(pt)[[D]].$$

\square

III. Computation of $U^*(B\Gamma_k)$, $k \geq 4$. The group Γ_k , $k \geq 4$, is generated by u , v , subject to the following relations $u^t = v^2$, $uvu = v$,

$t = 2^{k-2}$; $|\Gamma_k| = 2^k$. We have $H^0(B\Gamma_k) = \mathbb{Z}$, $H^{4p}(B\Gamma_k) = \mathbb{Z}_{2^k}$, $p > 0$, $H^{4p+2} = \mathbb{Z}_2 \oplus \mathbb{Z}_2$, $p \geq 0$, $H^{2p+1}(B\Gamma_k) = 0$, $p \geq 0$. Furthermore if $d_1, \{a_1, b_1\}$ are generators of respectively $H^4(B\Gamma_k)$ and $H^2(B\Gamma_k)$, then $d_1^p, \{a_1 d_1^p, b_1 d_1^p\}$ are generators of respectively $H^{4p}(B\Gamma_k)$ and $H^{4p+2}(B\Gamma_k)$, $p \geq 0$ (see [5]). The irreducible unitary representations of Γ_k are $1: u \rightarrow 1, v \rightarrow 1$, $\xi_1: u \rightarrow 1, v \rightarrow -1$, $\xi_2: u \rightarrow -1, v \rightarrow 1$, $\xi_3: u \rightarrow -1, v \rightarrow -1$,

$$\eta_r: u \rightarrow \begin{pmatrix} \omega^r & 0 \\ 0 & \omega^{-r} \end{pmatrix}, \quad v \rightarrow \begin{pmatrix} 0 & (-1) \\ 1 & 0 \end{pmatrix}, \quad r = 1, 2, \dots, 2^{k-2} - 1$$

and ω a primitive 2^{k-1} th root of unity (see [6]).

The relations between the irreducible unitary representations of Γ_k are as follows: $\xi_1^2 = \xi_2^2 = \xi_3^2 = 1$, $\xi_1 \cdot \xi_2 = \xi_3$, $\xi_2 \xi_3 = \xi_1$, $\xi_3 \cdot \xi_1 = \xi_2$; if we introduce $\eta_0 = 1 + \xi_1$, $\eta_{2^{k-2}} = \xi_2 + \xi_3$, then we can define $\eta_s, s \in \mathbb{Z}$, by the relations $\eta_{2^{k-2}+r} = \eta_{2^{k-2}-r}$, $\eta_r = \eta_{-r}$ and we have: $\eta_r \cdot \eta_s = \eta_{r+s} + \eta_{r-s}$, $r \in \mathbb{Z}$, $s \in \mathbb{Z}$ (see [10]). As in Section II we shall be working with $A_k = c f_1(\xi_1) \in \tilde{U}^2(B\Gamma_k)$, $B_k = c f_1(\xi_2) \in \tilde{U}^2(B\Gamma_k)$, $C_k = c f_1(\xi_3) \in \tilde{U}^2(B\Gamma_k)$, $D_k = c f_2(\eta_1) \in \tilde{U}^4(B\Gamma_k)$. We have as in 2.5 with $U^*(pt)[[D_k]] = \{R(D_k), R \in \Omega_*\}$:

THEOREM 3.1. $U^*(B\Gamma_k)$ is concentrated in even dimensions and as a module over $U^*(pt)[[D_k]]$, $U^*(B\Gamma_k)$ is generated by $1, B_k, C_k$. \square

The following proposition is proved in the same way as 2.8 and 2.6, $P_0(Z)$ being the formal power series of 2.8:

PROPOSITION 3.2. (a) We have $c f_1(\eta_1) = P_0(D_k)$.

(b) If $H(Z) = \sum_{i \geq 0} \alpha_i Z^i \in \Omega_{2n}$ is such that $H(D_k) = 0$, then $\alpha_0 = 0$ and the leading coefficient of $H(Z)$ belongs to $2^k U^*(pt)$. \square

LEMMA 3.3. For each $n \in \mathbb{Z}$ there is a polynomial $P_{2n+1}(X) \in \mathbb{Z}[X]$ such that $P_{2n+1}(0) = 0$, $P_{2n+1}(2) = 2$, $P_{2n+1}(\eta_1) = \eta_{2n+1}$.

Proof. Since $\eta_{-r} = \eta_r$, we may suppose $n \geq 0$. Then the assertion is evidently true if $n = 0$ with $P_1(X) = X$. Suppose that there are polynomials $P_{2i+1}(X) \in \mathbb{Z}[X]$, $0 \leq i \leq n-1$, such that $P_{2i+1}(\eta_1) = \eta_{2i+1}$, $P_{2i+1}(0) = 0$ and $P_{2i+1}(2) = 2$. Then $\eta_1^2 P_{2n-1}(\eta_1) = \eta_1^2 \eta_{2n-1} = (\eta_2 + \eta_0) \eta_{2n-1} = \eta_{2n+1} + 2\eta_{2n-1} + \eta_{2n-3}$. Hence if $P_{2n+1}(X) = (X^2 - 2)P_{2n-1}(X) - P_{2n-3}(X)$ we have $P_{2n+1}(X) \in \mathbb{Z}[X]$, $P_{2n+1}(0) = 0$, $P_{2n+1}(2) = 2$ and $P_{2n+1}(\eta_1) = \eta_{2n+1}$. \square

In the sequel we shall consider the sequence P_{2n+1} , $n \geq 0$, determined by $P_1(X) = X$, $P_3(X) = X^3 - 3X$ and the relation

$$(X^2 - 2)P_{2n-1}(X) - P_{2n-3}(X) = P_{2n+1}(X).$$

If $P(X) \in \mathbb{Z}[X]$ we shall denote by P' the derivatives of P .

PROPOSITION 3.4. *If ζ is a complex vector bundle over $B\Gamma_k$ such that $\zeta = P(\eta_1)$ where $P \in \mathbb{Z}[X]$, $P(0) = 0$, then there is a formal power series $P'(2)Z + \sum_{i \geq 2} \delta_i Z^i \in \Omega_4$ such that $cf_2(\zeta) = P'(2)D_k + \sum_{i \geq 2} \delta_i D_k^i$.*

Proof. For each $q \geq 1$ the complex bundle η_1^q is classified by the composite: $\Gamma_k \xrightarrow{\Delta} (B\Gamma_k)^q \xrightarrow{X^q g} (BU(2))^q \xrightarrow{m_q} BU(2^q)$ where Δ is the diagonal map, g a map classifying η_1 and m_q a map classifying $\otimes^q \gamma(2)$. We have $U^*(BU(2)^q) = U^*(pt)[[c_1^{(1)}, c_2^{(1)}, c_1^{(2)}, c_2^{(2)}, \dots, c_1^{(q)}, c_2^{(q)}]]$ where $c_k^{(i)}$, $k = 1$ or $k = 2$, is the image of $a_1 \otimes a_2 \cdots \otimes a_q$, $a_1 = a_2 = \cdots = a_{i-1} = 1$, $a_i = cf_k(\gamma(2))$ ($k = 1$ or $k = 2$), $a_{i+1} = \cdots = a_q = 1$, by the canonical product $\otimes^q U^*(BU(2)) \rightarrow U^*(BU(2^q))$. Then $m_q^*(cf_2\gamma(2^q)) = \sum a_u (c_1^{(1)})^{u_1^{(1)}} \cdot (c_2^{(1)})^{u_2^{(1)}} \cdots (c_1^{(q)})^{u_1^{(q)}} \cdot (c_2^{(q)})^{u_2^{(q)}}$. If we substitute Z for $c_2^{(i)}$ and $P_0(Z)$ for $c_1^{(i)}$, $i = 1, 2, \dots, q$, we have a formal power series $R_q(Z) \in \Omega_4$ such that $R_q(D_k) = cf_2(\eta_1^q)$. If $\{p_j\}$ denotes the base point of $BU(2)$ and k_i the inclusion:

$$\{p_1\} \times \{p_2\} \times \cdots \times \{p_{i-1}\} \times BU(2) \times \{p_{i+1}\} \times \cdots \times \{p_q\} \subset (BU(2))^q,$$

we see that $k_i^* \circ m_q^*(cf_2(\gamma(2^q))) = cf_2(2^{q-1}\gamma(2)) = 2^{q-1}cf_2(\gamma(2)) + 2^{q-2}(2^{q-1} - 1)cf_1^2(\gamma(2))$. Consequently $R_q(Z) = q2^{q-1}Z + \sum_{i \geq 2} \varepsilon_i Z^i$. Similarly there are formal powers series $H_1(Z) \in \Omega_2$, $H_s(Z) \in \Omega_{2s}$, $s \geq 3$, such that $H_1(D_k) = cf_1(\eta_1^q)$ and $H_s(D_k) = cf_s(\eta_1^q)$, $s \geq 3$; we have $\nu'(H_1) \geq 1$, $\nu'(H_s) \geq 2$, $s \geq 3$. (We recall that $\nu'(P(Z)) = \frac{1}{4}\nu P(Z)$.) It follows that if $\zeta = \sum_{i=1}^r m_i \eta_1^i$, $m_i \geq 0$, there is a formal power series $H(Z) \in \Omega_4$ such that $H(D_k) = cf_2(\zeta)$ and $H(Z) = (\sum_{i=1}^r i m_i 2^{i-1})Z + \sum_{i \geq 2} \varepsilon'_i Z^i$. Now suppose that ζ is a complex vector bundle such that $\zeta = \sum_{i=1}^r m_i \eta_1^i - \sum_{i=1}^r n_i \eta_1^i$, $m_i \geq 0$, $n_i \geq 0$. The above remarks show that

$$\begin{aligned} cf(\zeta) &= 1 + cf_1(\zeta) + cf_2(\zeta) + \cdots \\ &= [1 + M_1(D_k) + cf_2(\zeta_1) + M_2(D_k)] \\ &\quad \times [1 + M'_1(D_k) + cf_2(\zeta_2) + M'_2(D_k)]^{-1} \end{aligned}$$

with $\zeta_1 = \sum_{i=1}^r m_i \eta_1^i$, $\zeta_2 = \sum_{i=1}^r n_i \eta_1^i$, M_1, M_2, M'_1, M'_2 being elements of Ω_* such that $\nu'(M_1) \geq 1$, $\nu'(M'_1) \geq 1$, $\nu'(M_2) \geq 2$, $\nu'(M'_2) \geq 2$. It follows that $cf_2(\zeta) = M(D_k)$, with $M(Z) \in \Omega_4$ and $M(Z) =$

$\sum_{i=1}^r (im_i 2^{i-1} - in_i 2^{i-1})Z + \sum_{i \geq 2} \delta_i Z^i$. Then if $P(X) = \sum_{i=1}^r m_i X^i - \sum_{i=1}^r n_i X^i \in \mathbb{Z}[X]$ we see that $M(Z) = P'(2)Z + \sum_{i \geq 2} \delta_i Z^i$, $P'(X)$ being the derivative of $P(X)$. \square

LEMMA 3.5. *There is a formal power series*

$$Q_1(Z) = (1 + 2^2 n(n+1))Z + \sum_{i \geq 2} \beta'_i Z^i \in \Omega_4$$

such that $Q_1(D_k) = c f_2(\eta_{2n+1})$.

Proof. Since $\eta_{2n+1} = P_{2n+1}(\eta_1)$ with $P_{2n+1} \in \mathbb{Z}[X]$, $P_{2n+1}(0) = 0$, then by 3.4 it is enough to prove that $P'_{2n+1}(2) = 1 + 2^2 n(n+1)$. This assertion is true when $n = 0$ because $P_1(X) = X$. Suppose that $P'_{2i+1}(2) = 1 + 2^2 i(i+1)$, $0 \leq i \leq n-1$. We have $P_{2n+1} = (X^2 - 2)P_{2n-1} - P_{2n-3}$ and then $P'_{2n+1}(2) = 2^2 P_{2n-1}(2) + 2P'_{2n-1}(2) - P'_{2n-3}(2) = 2^3 + 2[1 + 2^2(n-1)n] - [1 + 2^2(n-2)(n-1)] = 1 + 2^2 n(n+1)$ ($P_{2n-1}(2) = 2$ by 3.3). Hence the lemma has been proved. \square

In Lemma 3.5 the coefficients β'_i depend on n ; however we have chosen not to complicate the notation.

PROPOSITION 3.6. *There is a formal power series*

$$T_k(Z) = 2^k Z + 2^{k-2} \lambda'_2 Z^2 + 2^{k-3} \lambda'_3 Z^3 \\ + \cdots + 2 \lambda'_{k-1} Z^{k-1} + \sum_{i \geq k} \lambda'_i Z^i \in \Omega_4,$$

with $\lambda'_2 \notin 2U^*(pt)$, such that $T_k(D_k) = 0$. Moreover if $R(Z) \in \Omega_*$ and $R(D_k) = 0$ then $R(Z) \in T_k(Z)\Omega_*$.

Proof. From 3.5 there is a formal power series

$$Q_1(Z) = [1 + 2^2(2^{k-3} - 2)(2^{k-3} - 1)]Z + \sum_{i \geq 2} \beta'_i Z^i \in \Omega_4$$

such that $Q_1(D_k) = c f_2(\eta_{2^{k-2-3}})$. We have $1 + 2^2(2^{k-3} - 2)(2^{k-3} - 1) = 9 + 2^{2k-4} - 3 \cdot 2^{k-1}$. Now

$$\eta_1^2 \eta_{2^{k-2-1}} = (\eta_2 + \eta_0) \eta_{2^{k-2-1}} \\ = \eta_{2^{k-2+1}} + \eta_{2^{k-2-3}} + 2\eta_{2^{k-2-1}} = 3\eta_{2^{k-2-1}} + \eta_{2^{k-2-3}}$$

and consequently if $P(X) = (X^2 - 3)P_{2^{k-2-1}}$, we have $P \in \mathbb{Z}[X]$, $P(0) = 0$ and $P(\eta_1) = \eta_{2^{k-2-3}}$. Then from 3.4 there is a formal power series $Q_2(Z) = P'(2) + \sum_{i \geq 2} \beta''_i Z^i \in \Omega_4$ such that $Q_2(D_k) = c f_2(\eta_{2^{k-2-3}})$. We

have $P'(2) = 2^2 P_{2^{k-2}-1}(2) + P'_{2^{k-2}-1}(2) = 2^3 + 1 + 2^2(2^{k-3} - 1)2^{k-3} = 9 + 2^{2k-4} - 2^{k-1}$. Hence

$$\begin{aligned} 0 &= Q_2(D_k) - Q_1(D_k) \\ &= [9 + 2^{2k-4} - 2^{k-1} - (9 + 2^{2k-4} - 3 \cdot 2^{k-1})]D_k \\ &\quad + \sum_{i \geq 2} (\beta''_i - \beta'_i)D_k^i \\ &= 2^k D_k + \sum_{i \geq 2} \mu'_i D_k^i, \quad \mu'_i = \beta''_i - \beta'_i. \end{aligned}$$

Then if $T_k(Z) = 2^k Z + \sum_{i \geq 2} \mu'_i Z^i \in \Omega_4$ then we have $0 = T_k(D_k)$. By 3.2 and a proof similar to that of 2.12, Section II, if $R(Z) \in \Omega_*$ is such that $R(D_k) = 0$ then $R(Z) \in T_k(Z)\Omega_*$. Now we want to show that $\mu'_2 = 2^{k-2}\lambda'_2$, $\lambda'_2 \notin 2U^*(pt)$, $\mu'_3 = 2^{k-3}\lambda'_3, \dots, \mu'_{k-1} = 2\lambda'_{k-1}$. Instead of $T_3(Z)$ we take the formal power series $T(Z)$ defined in Section II (see 2.11). We recall that $T(Z) = 2^3 Z + 2\lambda_2 Z^2 + \sum_{i \geq 3} \lambda_i Z^i$, $\lambda_2 \notin 2U^*(pt)$. Hence if $k = 3$ the assertion concerning the coefficients of $T_k(Z)$ is true. Suppose that

$$\begin{aligned} T_k(Z) &= 2^k Z + 2^{k-2}\lambda'_2 Z^2 + 2^{k-3}\lambda'_3 Z^3 \\ &\quad + \dots + 2\lambda'_{k-1} Z^{k-1} + \sum_{i \geq k} \lambda'_i Z^i, \quad \lambda'_2 \notin 2U^*(pt). \end{aligned}$$

Consider the inclusion

$$\begin{aligned} i_{k+1}: \Gamma_k &= \{(u^2)^m v^n, n = 0, 1, 0 \leq m \leq 2^{k-1} - 1\} \subset \Gamma_{k+1} \\ &= \{u^m v^n, n = 0, 1, 0 \leq m \leq 2^k - 1\}. \end{aligned}$$

It is easily seen that $(Bi_{k+1})^*(D_{k+1}) = D_k$. We have: $T_{k+1}(Z) = 2^{k+1}Z + \sum_{i \geq 2} \mu''_i Z^i$ and $T_{k+1}(D_{k+1}) = 0$. It follows that $T_{k+1}(D_k) = 0$ and consequently there is an element $\alpha'_0 + \alpha'_1 Z + \alpha'_2 Z^2 + \dots \in \Omega_0$ such that:

$$\begin{aligned} &2^{k+1}Z + \sum_{i \geq 2} \mu''_i Z^i \\ &= \left(2^k Z + 2^{k-2}\lambda'_2 Z^2 + \dots + 2\lambda'_{k-1} Z^{k-1} + \sum_{i \geq k} \lambda'_i Z^i \right) \left(\sum_{i \geq 0} \alpha'_i Z^i \right). \end{aligned}$$

Then $\alpha'_0 = 2$; $\mu''_2 = 2^k \alpha'_1 + 2^{k-2}\lambda'_2 \alpha'_0 = 2^{k-1}[2\alpha'_1 + \lambda'_2] = 2^{k-1}\lambda''_2$, $\lambda''_2 \notin 2U^*(pt)$; if $2 \leq i \leq k$ we have:

$$\mu''_i = 2^k \alpha'_{i-1} + 2^{k-2}\lambda'_2 \alpha'_{i-2} + 2^{k-3}\lambda'_3 \alpha'_{i-3} + \dots + 2^{k-i}\lambda'_i \alpha'_0 = 2^{(k+1)-i}\lambda''_i.$$

Hence the proposition has been proved. \square

Suppose $k \geq 4$; the inclusions $i_k: \Gamma_{k-1} \subset \Gamma_k$ and $j_k: \Gamma \subset \Gamma_k$ are given respectively by $\{(u^2)^m v^n, 0 \leq m \leq 2^{k-2} - 1, n = 0, 1\} \subset \{u^m v^n, 0 \leq m \leq 2^{k-1} - 1, n = 0, 1\}$ and $j_k = i_k \circ \cdots \circ i_4$; Γ_k is normal in Γ_{k+1} and $\Gamma_{k+1}/\Gamma_k = \{1, \bar{u}\} \simeq \mathbb{Z}_2$; if $q_k: \Gamma_k \rightarrow \Gamma_k$ is the conjugation by $u \in \Gamma_{k+1} - \Gamma_k$ then $q_k(u^2) = u^2$, $q_k(v) = v(u^2)^{-1}$. Let $f_k: B\Gamma_k \rightarrow B\Gamma_{k-1}$, $g_k: B\Gamma \rightarrow B\Gamma_k$, $h_k: B\Gamma_k \rightarrow B\Gamma_k$ be respectively Bi_k , Bj_k and Bq_k .

LEMMA 3.7. *Suppose $k \geq 4$.*

- (a) $f_k^*(A_k) = A_{k-1}$, $f_k^*(B_k) = 0$, $f_k^*(C_k) = A_{k-1}$, $f_k^*(D_k) = D_{k-1}$.
- (b) $g_k^*(A_k) = A$, $g_k^*(B_k) = 0$, $g_k^*(C_k) = A$, $g_k^*(D_k) = D$.
- (c) $h_k^*(A_k) = A_k$, $h_k^*(B_k) = C_k$, $h_k^*(C_k) = B_k$.

Proof. The proof is easy; for example $f_k^*(A_k) = A_{k-1}$ because $i_k^*: R(\Gamma_k) \rightarrow R(\Gamma_{k-1})$ maps ξ_1 to the similar representation: $u^2 \rightarrow 1$, $v \rightarrow -1$. ($R(\Gamma_k)$ and $R(\Gamma_{k-1})$ denote the representation rings). \square

The role played by A, B, C in Section II was symmetrical. Unfortunately this is not the case for A_k, B_k, C_k ($k \geq 4$) as we shall see in the forthcoming propositions. We recall that there are formal power series $S(Z) \in \Omega_2$, $J(Z) \in \Omega_0$ such that $A^2 = AS(D)$, $B^2 = BS(D)$, $C^2 = CS(D)$, $A(2 + J(D)) = B(2 + J(D)) = C(2 + J(D)) = 0$ (see 2.10, 2.13).

The formal power series $S(Z), J(Z)$ will play an important role in the calculations ahead.

PROPOSITION 3.8. *Suppose $k \geq 4$.*

- (a) $A_k B_k C_k = 0$.
- (b) $A_k(2 + J(D_k)) = 0$.
- (c) *There are $E_k \in \Omega_2$, $F_k \in \Omega_4$ such that $A_k = B_k + C_k - E_k(D_k)$, $B_k C_k = F_k(D_k)$.*

Proof. (a) The relation $A_k B_k C_k = 0$ is proved in exactly the same way as in 2.9(b).

(b) By 3.1 there are $H_0(Z) \in \Omega_2$, $H_1(Z) \in \Omega_2$, $H_2(Z) \in \Omega_4$ such that: $B_{k+1}^2 = B_{k+1}H_0(D_{k+1}) + C_{k+1}(D_{k+1}) + H_2(D_{k+1})$. By 3.7(c) we get $C_{k+1}^2 = C_{k+1}H_0(D_{k+1}) + B_{k+1}H_1(D_{k+1}) + H_2(D_{k+1})$ and $C_{k+1}^2 - B_{k+1}^2 = (C_{k+1} - B_{k+1})H_3(D_{k+1})$ with $H_3 = H_0 - H_1 \in \Omega_2$. By using 3.7(a) we see that: $A_k^2 = A_k \cdot H_3(D_k)$; as in 2.13 the relation $cf_1(\xi_1^2) = 0$ shows that there is $J_1(Z) \in \Omega_0$ depending on $H_3(Z)$ such that $A_k(2 + J_1(D_k)) = 0$ and by 3.7(b) we get $A(2 + J_1(D)) = 0$; so there is

$H_4(Z) \in \Omega_0$, $\nu'(H_4) \geq 1$ such that $2 + J_1(Z) = (2 + J(Z))(1 + H_4(Z))$ (see 2.15) and consequently $2 + J(Z) = (2 + J_1(Z))H_5(Z)$, $H_5(Z) \in \Omega_0$ being such that: $(1 + H_4(Z))(1 + H_5(Z)) = 1$. Hence $A_k(2 + J(D_k)) = 0$.

(c) By using the relations $\eta_r \cdot \eta_s = \eta_{r+s} + \eta_{r-s}$, $r \in \mathbb{Z}$, $s \in \mathbb{Z}$, $\eta_0 = 1 + \xi_1$, $\eta_{2^{k-2}} = \xi_2 + \xi_3$, then a straightforward calculation shows that there is a polynomial $R_m[X] \in \mathbb{Z}[X]$ such that $R_m(0) = 0$ and $\eta_{2^m} = R_m(\eta_1) + \eta_0$, $2 \leq m \leq k-2$; in fact $R_m(X)$ is determined by $R_2(X) = X^4 - 4X$, $R_m(X) = R_{m-1}^2(X) + 4R_{m-1}(X)$; so: $\xi_2 + \xi_3 = \eta_{2^{k-2}} = R_{k-2}(\eta_1) + \eta_0 = R_{k-2}(\eta_1) + 1 + \xi_1$. Then the proof of 3.4 shows that there are $E_k(Z) \in \Omega_2$, $F_k(Z) \in \Omega_4$ such that: $B_k + C_k = cf_1(R_{k-2}(\eta_1)) + A_k = E_k(D_k) + A_k$ and $B_k C_k = A_k E_k(D_k) + cf_2(R_{k-2}(\eta_1)) = A_k E_k(D_k) + F_k(D_k)$. As $0 = AE_k(D) + F_k(D)$ by 3.7(b) we see that $E_k(Z) \in (2 + J(Z))\Omega_*$ and consequently $B_k C_k = F_k(D)$ since $A_k(2 + J(D_k)) = 0$. Hence (c) is proved. \square

PROPOSITION 3.9. *Suppose $k \geq 4$.*

(a) *There is $M(Z) \in \Omega_2$ such that: $B_k(2 + J(D_k)) + M(D_k) = C_k(2 + J(D_k)) + M(D_k) = 0$ and $M(D_k) \neq 0$.*

(b) *There is $N(Z) \in \Omega_4$, such that: $B_k^2 = B_k S(D_k) + N(D_k)$, $C_k^2 = C_k S(D_k) + N(D_k)$ and $N(D_k) \neq 0$.*

(c) *There are $G_k(Z) \in \Omega_2$, $L_k(Z) \in \Omega_4$ the coefficients of which can be computed from those of $J(Z)$, $S(Z)$, $E_k(Z)$, $F_k(Z)$ and satisfying $G_k(D_k) = M(D_k)$, $L_k(D_k) = N(D_k)$.*

Proof. (a) As in 3.1 there are $H_1(Z) \in \Omega_2$, $K_0(Z) \in \Omega_2$, $K_1(Z) \in \Omega_4$ such that: $B_k^2 = B_k H_1(D_k) + A_k K_0(D_k) + K_1(D_k)$; hence: $AK_0(D) = 0$ which imply by 2.15 that $K_0(Z) \in (2 + J(Z))\Omega_*$; so: $B_k^2 = B_k H_1(D_k) + K_1(D_k)$ because $A_k(2 + J(D_k)) = 0$ by 3.8(b). We have $B_k^{n+1} = B_k H_n(D_k) + K_n(D_k)$ with $H_n(Z) \in \Omega_{2n}$, $K_n(Z) \in \Omega_{2n+2}$ satisfying: $H_n(Z) = H_1(Z)H_{n-1}(Z) + K_{n-1}(Z)$, $K_n(Z) = K_1(Z)H_{n-1}(Z)$, $n \geq 2$. It follows easily that $\text{Lim}_{n \rightarrow \infty} \nu(H_n) = \text{Lim}_{n \rightarrow \infty} \nu(K_n) = +\infty$; as $cf_1(\xi_2^2) = 0$ we have $2B_k + \sum_{n \geq 2} a_n B_k^n = 0$ with $a_n = \sum_{i+j=n} a_{ij}$, the a_{ij} , $i \geq 1$, $j \geq 1$, being the coefficients of the formal group law. A proof similar to that of 2.13 shows that there are $P_1(Z) \in \Omega_0$, $P_2(Z) \in \Omega_2$, $\nu'(P_1) \geq 1$, $\nu'(P_2) \geq 1$ such that $B_k(2 + P_1(D_k)) + P_2(D_k) = 0$; by 3.7(a) we have $C_k(2 + P_1(D_k)) + P_2(D_k) = 0$; hence $A(2 + P_1(D)) = 0$ and as a direct consequence of 2.15 there is $P_3(Z) \in \Omega_0$ such that $2 + J(Z) = (2 + P_1(Z))P_3(Z)$ and then: $B_k(2 + J(D_k)) + M(D_k) = C_k(2 + J(D_k)) + M(D_k) = 0$ with $M(Z) = P_2(Z)$. $P_3(Z) \in \Omega_2$. Suppose $M(D_k) = 0$; then $B_k(2 + J(D_k)) = C_k(2 + J(D_k)) = 0$; from

3.8(c) we have $A_k^2 = A_k(B_k + C_k) - A_k E_k(D_k)$ and consequently $A E_k(D) = 0$; so $E_k(Z) \in (2 + J(Z))\Omega_*$ and $A_k^2 = (B_k + C_k)^2$. Let $\theta: MU \rightarrow K$ being the canonical map between spectra; θ sends Euler classes to Euler classes; the relation $A_k^2 = (B_k + C_k)^2$ becomes by using θ : $1 + \xi_1 - \xi_2 - \xi_3 = 0$ in $K^0(B\Gamma_k)$ which is impossible since $1 + \xi_1 - \xi_2 - \xi_3 \neq 0$ in $R(\Gamma_k)$ (the canonical map from $R(\Gamma_k)$ to $K^0(B\Gamma_k)$ is injective). Hence $M(D_k) \neq 0$.

(b) We have seen in (a) that $B_k^2 = B_k H_1(D_k) + K_1(D_k)$; so: $C_k^2 = C_k H_1(D_k) + K_1(D_k)$ and: $A^2 = A H_1(D) + K_1(D) = A S(D)$; then $A[H_1(D) - S(D)] + K_1(D) = 0$ and there is $S_0(Z) \in \Omega_2$ such that $H_1(Z) = S(Z) + (2 + J(Z))S_0(Z)$; consequently: $B_k^2 = B_k S(D_k) - M(D_k)S_0(D_k) + K_1(D_k) = B_k S(D_k) + N(D_k)$ with: $N(Z) = K_1(Z) - M(Z)S_0(Z) \in \Omega_4$; by 3.7(c) $C_k^2 = C_k S(D_k) + N(D_k)$. If $N(D_k) = 0$ then as in 2.13 we would have $C_k(2 + J(D_k)) = 0$ and then $M(D_k) = 0$ which is false by (a). Hence: $N(D_k) \neq 0$.

(c) We need to show first that $T_k(Z) \notin 2\Omega_*$ ($T_3(Z) = T(Z)$ and $T_k(Z)$ are defined respectively in 2.11 and 3.6). Suppose $k = 3$; from $AB + BC + CA = Q(D)$ and $A(2 + J(D)) = B(2 + J(D)) = 0$ (see 2.9 and 2.13) we get $(2 + J(D))Q(D) = 0$; so:

$$\begin{aligned} (2 + J(Z))Q(Z) &= (2 + \mu_1 Z + \mu_2 Z^2 + \dots)(4Z + \beta_2 Z^2 + \beta_3 Z^3 + \dots) \\ &= 8Z + (2\beta_2 + 4\mu)Z^2 + (2\beta_3 + \mu_1\beta_2 + 4\mu_2)Z^3 \\ &\quad + \dots \in T(Z)\Omega_*; \end{aligned}$$

hence $T(Z) \notin 2\Omega_*$ since $\mu_1\beta_2 \notin 2U^*(pt)$ (see 2.9 and 2.13). Suppose that $T_i(Z) \notin 2\Omega_*$, $3 \leq i \leq k-1$, and $T_k(Z) \in 2\Omega_*$; as $A_k = B_k + C_k - E_k(D_k)$ (see 3.8(c)) we have $E_k(D_{k-1}) = 0$ and then $E_k(Z) \in T_{k-1}(Z)\Omega_*$; from $T_k(Z) \in T_{k-1}(Z)\Omega_*$, $T_k(Z) \in 2\Omega_*$ and $T_{k-1}(Z) \notin 2\Omega_*$ it follows easily that $2T_{k-1}(D_k) = 0$; consequently $2E_k(D_k) = 0$ and $2A_k = 2(B_k + C_k)$ which is impossible (it can be seen by using $\theta: MU \rightarrow K$ as in (a)). Hence $T_k(Z) \notin 2\Omega_*$, $k \geq 3$. Let $q: \Omega_* \rightarrow \Omega_*/2\Omega_* = (U^*(pt)/2U^*(pt))[[Z]]$ be the canonical projection and $\bar{R}(Z)$ the image of $R(Z)$ by q . Now it follows easily from 3.8(c) and (a) that: $2M(D_k) + E_k(D_k)(2 + J(D_k)) = 0$ and then $2M(Z) + E_k(Z)(2 + J(Z)) = T_k(Z) \cdot H(Z)$, $H(Z) \in \Omega_*$. Hence $\bar{E}_k(Z) \cdot \bar{J}(Z) = \bar{T}_k(Z) \cdot \bar{H}(Z)$; as $\bar{T}_k(Z) \neq 0$ the formal power series $\bar{H}(Z)$ is unique and its coefficients which belong to $U^*(pt)/2U^*(pt) = \mathbb{Z}_2[x_1, x_1, \dots]$ ($|x_i| = -2i$) are computable from those of \bar{E}_k , \bar{J} and \bar{T}_k ; if $\bar{H}(Z) = \sum \bar{d}_i Z^i$, $\bar{d}_i \neq 0$, then there is a unique element $e_i \in \mathbb{Z}[x_1, \dots, x_n, \dots] = U^*(pt)$ whose coefficients as a polynomial in x_1, \dots, x_n, \dots , are odd and such that $\bar{e}_i = \bar{d}_i$; it follows that

$E_k(Z)(2+J(Z))-T_k(Z)\cdot(\sum e_i Z^i) = -2G_k(Z)$ and $G_k(D_k) = M(D_k)$. The same method can be used to determine $L_k(Z)$ by considering the relation $2N(D_k) = E_k^2(D_k) - E_k(D_k)S(D_k) - 2F(D_k)$ which is an easy consequence of (b) and 3.8(c). \square

Let \tilde{I}'_* be the graded ideal of Λ_* generated by the homogeneous formal power series $G_k(X, Z) = X(2+J(Z))+G_k(Z) \in \Lambda_2$, $G_k(Y, Z) = Y(2+J(Z)) + G_k(Z) \in \Lambda_2$, $T_k(Z) \in \Lambda_4$ (see 3.6 and 3.9) and \tilde{I}''_* the graded ideal of Λ_* generated by the homogeneous formal power series $L_k(X, Z) = X^2 - XS(Z) - L_k(Z) \in \Lambda_4$, $L_k(Y, Z) = Y^2 - YS(Z) - L_k(Z) \in \Lambda_4$, $XY - F_k(Z) \in \Lambda_2$ (see 3.8(c) and 3.9). The proofs of the following lemmas are quite similar to those of 2.15, 2.16 and will be omitted.

LEMMA 3.10. *If $H_1(Z), H_2(Z), H_3(Z)$ are elements of Ω_* such that $B_k H_1(D_k) + C_k H_2(D_k) + H_3(D_k) = 0$ then $XH_1(Z) + YH_2(Z) + H_3(Z) \in G_k(X, Z)\Omega_* + G_k(Y, Z)\Omega_* + T_k(Z)\Omega_* \subset \tilde{I}'_*$.* \square

LEMMA 3.11. *For any $P(X, Y, Z) \in \Lambda_*$ there are $H_1(Z), H_2(Z), H_3(Z)$ elements of Ω_* such that $P(X, Y, Z) - [XH_1(Z) + YH_2(Z) + H_3(Z)] \in \tilde{I}''_*$.* \square

As a direct consequence of 3.10, 3.11 we get our main theorem where $\tilde{I}_* = \tilde{I}'_* + \tilde{I}''_*$ (see the proof of 2.17).

THEOREM 3.12. *The graded $U^*(pt)$ -algebra $U^*(B\Gamma_k)$ is isomorphic to Λ_*/\tilde{I}_* where \tilde{I}_* is a graded ideal of Λ_* generated by six homogeneous formal power series.* \square

REMARK. The homomorphism f_k^* induced by the inclusion $\Gamma_{k-1} \subset \Gamma_k$ (see 3.7) is such that $f_k^*(B_k) = 0$,

$$f_k^*(C_k) = B_{k-1} + C_{k-1} - E_{k-1}(D_{k-1})(E_{k-1}(D_{k-1}) \neq 0),$$

$f_k^*(D_k) = D_{k-1}$ if $k \geq 5$ (see 3.8). But $f_4^*(B_4) = 0$, $f_4^*(C_4) = P(D) - (B + C)$, $P(D) \neq 0$ (see 2.9, 2.6), $f_4^*(D_4) = D$.

Let $U^*(pt)[[D_k]]$ be $\{R(D_k), R(Z) \in \Omega_*\}$.

THEOREM 3.13. (a) $U^*(pt)[[D_k]] \simeq \Omega_*/(T_k)$ as graded $U^*(pt)$ -algebras.

(b) $U^*(B\Gamma_k)$ is generated by 1, A_k, B_k as a $U^*(pt)[[D_k]]$ -module. Moreover if $V_k = U^*(pt)[[D_k]]$ then:

$$V_k \cap V_k B_k = V_k \cap V_k C_k = V_k B_k \cap V_k C_k = G_k(D_k) \cdot V_k.$$

Proof. The assertion (a) is a consequence of 3.6; the first part of (b) is proven in 3.1 and the second part is a consequence of 3.10. \square

Now we are going to alter B_k, C_k in order to improve 3.13(b). From $B_k(2 + J(D_k)) + G_k(D_k) = 0$ it follows easily that $G_k(D) = 0$; so $AG_k(D) = 0$ and $G_k(Z) = (2 + J(Z))G'_k(Z), G'_k(Z) \in \Omega_2$; hence

$$(B_k + G'_k(D_k))(2 + J(D_k)) = (C_k + G'_k(D_k))(2 + J(D_k)) = 0.$$

Furthermore if $\mu: U^*(B\Gamma_k) \rightarrow H^*(B\Gamma_k)$ is the edge homomorphism (in connection with the U^* -AHSS for $B\Gamma_k$) then $\mu(B_k + G'_k(D_k)) = \mu(B_k), \mu(C_k + G'_k(D_k)) = \mu(C_k)$. This remark and Lemma 3.10 allow the following rearrangement of Theorem 3.13 with $B'_k = B_k + G'_k(D_k), C'_k = C_k + G'_k(D_k)$.

THEOREM 3.14. (a) $U^*(pt)[[D_k]] \simeq \Omega_*/(T_k)$ as graded $U^*(pt)$ -algebras.

(b) As graded $U^*(pt)[[D_k]]$ -modules we have:

$$U^*(B\Gamma_k) \simeq U^*(pt)[[D_k]] \oplus U^*(pt)[[D_k]] \cdot B'_k \oplus U^*(pt)[[D_k]] \cdot C'_k$$

and B'_k, C'_k have the same annihilator $(2 + J(D_k)) \cdot U^*(pt)[[D_k]]$. \square

Appendix.

(A) *Calculation of $U^*(B\mathbb{Z}_m)$ by a new method.* The method used in the case $G = \Gamma_k$ applies more simply in the case $G = \mathbb{Z}_m$. Let w be $\exp(2i/m)$ and ρ the irreducible unitary representation of \mathbb{Z}_m defined by $\rho(\bar{q}) = w^q, \bar{q} \in \mathbb{Z}_m$. Let η be the complex vector bundle over $B\mathbb{Z}_m$ corresponding to ρ and $D_1 = e(\eta) = cf_1(\eta) \in U^2(B\mathbb{Z}_m)$.

Let Λ'_* be $U^*(pt)[[Z]]$, graded by taking $\dim Z = 2$. There is a topology on $\Lambda'_{2n}, n \geq 0$, defined by the subgroups $J_r = \{P \in \Lambda'_{2n}, \nu(P) \geq r\}$, with $\nu(P) = 2s$ if $P(Z) = a_s Z^s + a_{s+1} Z^{s+1} + \dots, a_s \neq 0$; Λ'_{2n} is complete and Hausdorff. Furthermore, $U^{2n}(B\mathbb{Z}_m)$ is topologized by the subgroups $J^{r, 2n-r}$ induced by the U^* -AHSS for $B\mathbb{Z}_m$, taken as a system of neighbourhoods of 0. The group $U^{2n}(B\mathbb{Z}_m)$ is complete and Hausdorff because the U^* -AHSS for $B\mathbb{Z}_m$ collapses. Moreover there is a unique continuous homomorphism of graded $U^*(pt)$ -algebras $\varphi': \Lambda'_* \rightarrow U^*(B\mathbb{Z}_m)$ such that $\varphi'(Z) = D_1$ and φ' is surjective (see Sections I and II).

The complex vector bundle η^m is trivial ($\dim \eta^m = 1$) because $\rho^m = 1$. Hence $cf_1(\eta^m) = 0$. If m_0 denotes a map: $BU(1)^m \rightarrow BU(1)$ classifying $\otimes^m \gamma(1)$ ($\gamma(1)$ being a universal complex vector bundle over $BU(1)$) and if $c_1 = cf_1(\gamma(1))$ then:

$$m_0^*(c_1) = \sum a_{(u)} e_1^{u_1} e_2^{u_2} \dots e_m^{u_m}, \quad u = (u_1, \dots, u_m).$$

$u_1 \geq 0, \dots, u_m \geq 0$, e_i being the image of $a_1 \otimes a_2 \otimes \dots \otimes a_m$ with $a_1 = a_2 = \dots = a_{i-1} = 1$, $a_i = c_1$, $a_{i+1} = \dots = a_m = 1$, by the product: $\otimes^m U^*(BU(1)) \rightarrow UBU(1)^m$. The vector bundle η^m is classified by the composite:

$$\begin{array}{ccc} & m & \\ & \swarrow & \searrow \\ B\mathbb{Z}_m & \xrightarrow{d} (B\mathbb{Z}_m) \xrightarrow{\times g} BU(1)^m \xrightarrow{m_0} & BU(1), \end{array}$$

d being the diagonal map and g a map classifying η . It follows that if $T(Z) = \sum a_{(u)} Z^{u_1+u_2+\dots+u_m} \in \Lambda'_2$, we have $T(cf_1(\eta)) = T(e(\eta)) = T(D_1) = 0$. It is easily seen that $T(Z) = [m](Z)$.

THEOREM A.1. $U^*(B\mathbb{Z}_m) \simeq \Lambda'_*/([m](Z))$ as graded $U^*(pt)$ -algebras.

Proof. Let I_* be $([m](Z))$. The homomorphism $\varphi': \Lambda'_* \rightarrow U^*(B\mathbb{Z}_m)$ of graded $U^*(pt)$ -algebras, defined above, is surjective; moreover $\varphi'(I_*) = 0$. Hence φ' gives rise to a homomorphism of graded $U^*(pt)$ -algebras $\bar{\varphi}': \Lambda'_*/I_* \rightarrow U^*(B\mathbb{Z}_m)$. Let $P(Z)$ be any element of Λ'_{2n} ($n \geq 0$) such that $P(D_1) = 0$; if $P(Z) = a_0 + a_1 Z + a_2 Z^2 + \dots$, then $a_0 = 0$ because $a_0 = -(a_1 D_1 + a_2 D_1^2 + \dots) \in \dot{U}^*(B\mathbb{Z}_m) \cap U^*(pt) = 0$. It follows that $P(Z) = a_{p_0} Z^{p_0} + a_{p_0+1} Z^{p_0+1} + \dots$, with $p_0 \geq 1$, $a_{p_0} \neq 0$. We have

$$a_{p_0+1} D_1^{p_0+1} + \dots + a_{p_0+k} D_1^{p_0+k} \in J^{2(p_0+1), 2(n-p_0-1)};$$

since this group is closed in $U^{2n}(B\mathbb{Z}_m)$, it follows that

$$\sum_{i=1}^{\infty} a_{p_0+i} D_1^{p_0+i} \in J^{2(p_0+1), 2(n-p_0-1)} \subset J^{2p_0+1, 2(n-p_0)-1}.$$

Let s be the quotient map:

$$\begin{aligned} J^{2p_0, 2(n-p_0)} &\twoheadrightarrow J^{2p_0, 2(n-p_0)} / J^{2p_0+1, 2(n-p_0)-1} \\ &= H^{2p_0}(B\mathbb{Z}_m) \otimes U^{2(n-p_0)}(pt) = \mathbb{Z}_m \otimes U^{2(n-p_0)}(pt) \\ &= U^{2(n-p_0)}(pt) / mU^{2(n-p_0)}(pt) \end{aligned}$$

($H^{2p_0}(B\mathbb{Z}_m) = \mathbb{Z}_m$ because $p_0 \geq 1$). It follows from $s(P(D_1)) = 0$ that $a_{p_0} = m a'_{p_0}$. We form $P_1(Z) = P(Z) - a'_{p_0} Z^{p_0-1} T(Z)$; then $P_1(D_1) = 0$ and $\nu(P_1) > \nu(P)$. We repeat the same process, and there is an element $P_{r+1}(Z) \in \Lambda'_{2n}$, $r \geq 1$, such that

$$P_{r+1}(Z) = P(Z) - (a'_{p_0} Z^{p_0-1} + a'_{p_1} Z^{p_1-1} + \dots + a'_{p_r} Z^{p_r-1}) T(Z)$$

with the properties: $P_{r+1}(D_1) = 0$, $\nu(P_{r+1}) = p_{r+1} > p_r \dots > p_1 > p_0$. Hence $\lim_{r \rightarrow \infty} \nu(P_{r+1}) = +\infty$ and by Sec. I we have $P(Z) =$

$(\sum_{i=0}^{\infty} a'_p Z^{p_i-1})T(Z) \in I_{2n}$. It follows that $\bar{\varphi}'$ is injective and the theorem has been proved. \square

Note. P. S. Landweber has proved a similar result by using other methods (see [13]).

(B) *Calculation of $U^*(BSU(n))$. Particular case $n = 2$: $SU(2) = Sp(1)$. Consider the S^1 -bundle $U(n)/SU(n) = S^1 \rightarrow BSU(n) \xrightarrow{p} BU(n)$, $n \geq 2$, $p = Bi$ with $i: SU(n) \subset U(n)$; let ξ be the complex vector bundle $E = BSU(n) \times_{S^1} \mathbb{C} \xrightarrow{\pi} BU(n)$, where S^1 acts on \mathbb{C} by the multiplication in \mathbb{C} . If $E_0 = E - j(BU(n))$, j being the zero-section of ξ , then we have the Gysin exact sequence (see [4]):*

$$\begin{aligned} \cdots \rightarrow U^i(BU(n)) \xrightarrow{e(\xi)} U^{i+2}(BU(n)) \xrightarrow{\pi_0^*} U^{i+2}(E_0) \\ \rightarrow U^{i+1}(BU(n)) \rightarrow \cdots, \end{aligned}$$

where π_0 denotes $\pi|_{E_0}$. The map $g: BSU(n) \rightarrow E_0$ defined by $g(x) = [x, 1]$ is an embedding; take $E' = g(BSU(n))$, j' the inclusion: $E' \subset E_0$ and $h: E_0 \rightarrow E'$ the map defined by $h[x, z] = [xz/|z|, 1]$; then by using h and the homotopy $H: E_0 \times I \rightarrow E_0$ given by $H([x, z], t) = [x, tz + (1-t)z/|z|]$ we see that E' is a strong deformation retract of E_0 ; it is easily seen that $\pi' \circ h = \pi_0$ and $\pi' \circ g = p$ with $\pi' = \pi|_{E'}$, g being considered as a homeomorphism: $BSU(n) \xrightarrow{\sim} g(BSU(n))$. So: $\pi_0^* = h^* \circ g^{*-1} \circ p^*$ and since $h^* \circ g^{*-1}$ is an isomorphism the above exact sequence gives the following one:

$$\begin{aligned} \cdots \rightarrow U^i(BU(n)) \xrightarrow{e(\xi)} U^{i+2}(BU(n)) \xrightarrow{p^*} U^{i+2}(BSU(n)) \\ \rightarrow U^{i+1}(BU(n)) \rightarrow \cdots. \end{aligned}$$

Consider the canonical map of ring spectra $f: MU \rightarrow H$ (see [1]); $f^\#(-)$ maps Euler classes to Euler classes. Suppose $e(\xi) = 0$; then $f^\#(-)(e(\xi)) = 0$, which means that the Euler class of ξ for H is 0. From the Gysin exact sequence of ξ for H it follows easily that $H^2(BU(n)) \simeq H^2(BSU(n))$ which is impossible since $H^2(BU(n)) \neq 0$ and $H^2(BSU(n)) = 0$ (see [12], page 237). Hence $e(\xi) \neq 0$ and the map $\cdot - e(\xi)$ is injective. Consequently the sequence:

$$0 \rightarrow U^{2i}(BU(n)) \xrightarrow{e(\xi)} U^{2i+2}(BU(n)) \xrightarrow{p^*} U^{2i+2}(BSU(n)) \rightarrow 0$$

is exact and $U^{2i+1}(BSU(n)) = 0$, $i \geq 0$. So we have:

THEOREM B.1. *We have $U^{2i+1}(BSU(n)) = 0$, $i \geq 0$, and the map p^* induces an isomorphism:*

$$U^{2i+2}(BU(n))/e(\xi)U^{2i}(BU(n)) \simeq U^{2i+2}(BSU(n)), \quad i \in \mathbb{Z}.$$

Now let (g_{ij}) be a set of transition functions for a universal $U(n)$ -bundle: $EU(n) \rightarrow BU(n)$. If \bar{g}_{ij} denotes the image of g_{ij} by the quotient map $q: U(n) \rightarrow U(n)/SU(n) = S^1$ then (\bar{g}_{ij}) is a set of transition functions for ξ ; from $q(g_{ij}) = \det(g_{ij})$ and $\dim \xi = 1$, it follows that ξ is isomorphic to the complex vector bundle $\Lambda^n \gamma(n)$, $\gamma(n)$ being a universal vector bundle over $BU(n)$. Hence:

THEOREM B.2.

$$U^{2i+2}(BU(n))/e(\Lambda^n \gamma(n)) \cdot U^{2i}(BU(n)) \simeq U^{2i+2}(BSU(n)).$$

and $U^{2i+1}(BSU(n)) = 0$, $i \geq 0$. □

Particular Case $n = 2$; $Sp(1) = SU(2)$. By Section II we have $U^*(BSp(1)) = U^*(BSU(2)) = U^*(pt)[[V]]$, with $V = cf_2(\theta)$, θ being a universal $Sp(1)$ -vector bundle over $BS(1)$, regarded as a $U(2)$ -vector bundle. Then $cf_1(\theta) = P_0(V) = \sum_{i=1}^{\infty} b_i V^i \in U^2(BSU(2))$. If p denotes the projection: $BSU(2) \rightarrow BU(2)$, we have seen that the following sequence is exact: $0 \rightarrow U^{2i}(BU(2)) \xrightarrow{e(\Lambda^2 \gamma(2))} U^{2i+2}(BU(2)) \xrightarrow{p^*} U^{2i+2}(BSU(2)) \rightarrow 0$. We wish to calculate the coefficients b_i , $i \geq 1$. The $Sp(1)$ -vector bundle θ considered as a $SU(2)$ -vector-bundle is a universal $SU(2)$ -vector-bundle over $BSU(2)$ isomorphic to $p^*(\gamma(2))$ as a complex vector bundle. We have $U^*(BU(2)) = U^*(pt)[[c_1, c_2]]$. $c_1 = cf_1(\gamma(2))$, $c_2 = cf_2(\gamma(2))$ and consequently

$$\begin{aligned} p^*(c_1) &= \sum_{i \geq 1} b_i V^i = \sum_{i \geq 1} b_i (cf_2(\theta))^i = \sum_{i \geq 1} b_i p^*(c_2)^i \\ &= p^* \left(\sum_{i \geq 1} b_i c_2^i \right). \end{aligned}$$

It follows that: $c_1 - \sum_{i \geq 1} b_i c_2^i = e(\Lambda^2 \gamma(2)) \cdot H(c_1, c_2)$ with $H(c_1, c_2) \in U^0(BU(2))$.

Let $k: BU(1) \times BU(1) \rightarrow BU(2)$ be a map classifying $\gamma(1) \times \gamma(1)$. Hence $k^*(\Lambda^2 \gamma(2)) = \gamma(1) \otimes \gamma(1)$ and $k^*(e(\Lambda^2 \gamma(2))) = F(X, Y)$, the formal group law. Then $k^*(c_1 - \sum_{i \geq 1} b_i c_2^i) = F(X, Y)k^*(H(c_1, c_2))$; as $k^*(c_1) = X + Y$ and $k^*(c_2) = XY$ we get:

$$X + Y - \sum_{i \geq 1} b_i (XY)^i = F(X, Y)G(X, Y) \in U^*(pt)[[X, Y]].$$

If $i(X) = [-1](X)$ then we have:

$$X + i(X) = \sum_{i \geq 1} b_i (X \cdot i(X))^i.$$

This relation determines completely the coefficients b_i , $i \geq 1$; for example $b_1 = -a_{11}$, $b_2 = a_{11}a_{11}a_{21} - a_{22} \cdots$ the a_{ij} being the coefficients of the group law.

(C) *Ring Structure of $H^*(B\Gamma_k)$, $k \geq 3$.* M. Atiyah has determined the ring-structure of $H^*(B\Gamma_3)$ by using K -theory (see [2]); namely $H^*(B\Gamma_3) = \mathbb{Z}[x, y, z]$ subject to the relations $xy = 4z$, $2x = 2y = x^2 = y^2 = 8z = 0$, $\dim x = 2$, $\dim y = 2$, $\dim z = 4$. We want to give another proof of this result using complex cobordism and determine the ring structure of $H^*(B\Gamma_k)$, $k \geq 4$.

We have $H^2(B\Gamma) = \mathbb{Z}x \oplus \mathbb{Z}y$, $H^4(B\Gamma) = \mathbb{Z} \cdot z$ with $x = c_1(\xi_j)$, $y = c_1(\xi_k)$, $z = c_2(\eta)$ (see Section II). Moreover: $2x = 2y = 8z = 0$. We have

$$\begin{aligned} B^2 &= BS(D), & C^2 &= CS(D), \\ BC &= (B + C)[P(D) - S(D)] - Q(D) \end{aligned}$$

(A , B , C play a symmetrical role; see Section II). If μ is the edge homomorphism we have $x^2 = \mu(BS(D)) = 0$ ($\mu: J^{4,0} \rightarrow J^{4,0}/J^{5,-1} = H^4(B\Gamma_3)$; $BS(D) \in J^{6,-2} \subset J^{5,-1}$); similarly $y^2 = 0$; $xy = -\mu(Q(D)) = -4z_3 = -4z = 4z$ because $Q(D) = 4D + \sum_{i \geq 2} \beta_i Z^i$ (see 2.9).

Suppose $k \geq 4$. We have $H^2(B\Gamma_k) = \mathbb{Z}x_k \oplus \mathbb{Z}y_k$, $H^4(B\Gamma_k) = \mathbb{Z} \cdot z_k$ with $x_k = c_1(\xi_2)$, $y_k = c_1(\xi_3)$, $z_k = c_2(\eta_1)$ (see 2.3, 2.4). We have $2x_k = 2y_k = 2^k z_k = 0$. The proof of Proposition 3.8 shows that $x_k y_k = \mu(F_k(D_k))$, μ being the edge homomorphism, $F_k(D_k) = c f_2(R_{k-2}(\eta_1))$ with $R_{k-2}(X) \in \mathbb{Z}[X]$; $R_{k-2}(X)$ is determined inductively by $R_2(X) = X^4 - 4X^2$, $R_{m+1}(X) = R_m^2(X) + 4R_m(X)$, $m \geq 2$. By 3.4 we get $F_k(D_k) = R'_{k-2}(2) + \sum_{i \geq 2} \nu_i D_k^i$, $\nu_i \in U^*(pt)$, $R'_{k-2}(X)$ being the derivative of $R_{k-2}(X)$. An easy calculation shows that $R'_{k-2}(2) = 2^{2k-4}$. As $2k - 4 \geq k$ we get $x_k y_k = 2^{2k-4} z_k = 0$. As a consequence of the relations in $R(\Gamma_k)$ stated in the beginning of Section III we get: $\xi_2 \eta_1 = \eta_{2^{k-2}-1}$. Hence $x_k^2 + c_2(\eta_1) = c_2(\eta_{2^{k-2}-1})$ because $c_1(\eta_1) = 0$. By 3.5 $c f_2(\eta_{2^{k-2}-1}) = [1 + 2^{k-1}(2^{k-3} - 1)]D_k + \sum_{i \geq 2} \beta'_i D_k^i$ and consequently $c_2(\eta_{2^{k-2}-1}) = (1 - 2^{k-1})z_k$. Therefore: $x_k^2 = -2^{k-1}z_k = 2^{k-1}z_k$. Similarly: $y_k^2 = 2^{k-1}z_k$. Hence we have proved the following result:

THEOREM C. *If $k \geq 4$ we have $H^*(B\Gamma_k) = \mathbb{Z}[x_k, y_k, z_k]$, $\dim x_k = \dim y_k = 2$, $\dim z_k = 4$ subject to the relations: $2x_k = 2y_k = x_k y_k = 2^k z_k = 0$, $x_k^2 = y_k^2 = 2^{k-1}z_k$. \square*

REFERENCES

- [1] J. F. Adams, *Stable Homotopy and Generalized Homology*, University of Chicago Mathematics Lecture Notes, 1971.
- [2] M. F. Atiyah, *Characters and cohomology of finite groups*, I.H.E.S. Publ. Math., **9** (1961), 23–64.
- [3] N. A. Baas, *On the Stable Adams Spectral Sequence*, Aarhus Universitët Lecture Notes, 1969.
- [4] T. Bröcker and T. t. Dieck, *Kobordismen Theorie*, Lecture Notes in Math., Vol. 178, Springer Verlag, 1970.
- [5] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton University Press, 1956.
- [6] C. W. Curtis and I. Reiner, *Representation Theory of Finite Groups and Associative Algebras*, Wiley, New York, 1962.
- [7] T. t. Dieck, *Steenrod operationen in kobordismen teorien*, Math. Z., **107** (1968), 380–401.
- [8] ———, *Bordism of G-manifolds and integrality theorems*, Topology, **9** (1970), 345–358.
- [9] ———, *Actions of finite Abelian p-groups without stationary points*, Topology, **9** (1970), 359–366.
- [10] D. Pitt, *Free actions of generalized quaternion groups on spheres*, Proc. London Math. Soc., **26** (1973), 1–18.
- [11] E. H. Spanier, *Algebraic Topology*, McGraw-Hill, 1966.
- [12] R. E. Stong, *Notes on Cobordism Theory*, Mathematical Notes, Princeton University Press, 1968.
- [13] P. S. Landweber, *Coherence, flatness and cobordism of classifying spaces*, Proc. Adv. Study. Inst. Alg. Top. 256–269, Aarhus 1970.
- [14] D. C. Ravenel, *Complex Cobordism and Stable Homoty Groups of Spheres*, Academic Press, Inc., 1986.

Received October 5, 1986 and in revised form August 15, 1988.

UNIVERSITÉ MOHAMMED V
B.P. 1014
RABAT, MOROCCO

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Pacific Journal of Mathematics

Vol. 142, No. 1

January, 1990

| | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Marco Andreatta, Mauro Beltrametti and Andrew Sommese , Generic properties of the adjunction mapping for singular surfaces and applications | 1 |
| Chen-Lian Chuang and Pjek-Hwee Lee , On regular subdirect products of simple Artinian rings | 17 |
| Fernando Giménez and Vicente Miquel Molina , Volume estimates for real hypersurfaces of a Kaehler manifold with strictly positive holomorphic sectional and antiholomorphic Ricci curvatures | 23 |
| Richard J. Griego and Andrzej Korzeniowski , Asymptotics for certain Wiener integrals associated with higher order differential operators | 41 |
| Abdeslam Mesnaoui , Unitary bordism of classifying spaces of quaternion groups | 49 |
| Abdeslam Mesnaoui , Unitary cobordism of classifying spaces of quaternion groups | 69 |
| Jesper M. Møller , On equivariant function spaces | 103 |
| Bassam Nassrallah , A q -analogue of Appell's F_1 function, its integral representation and transformations | 121 |
| Peter A Ohring , Solvability of invariant differential operators on metabelian groups | 135 |
| Athanase Papadopoulos and R. C. Penner , Enumerating pseudo-Anosov foliations | 159 |
| Ti-Jun Xiao and Liang Jin , On complete second order linear differential equations in Banach spaces | 175 |
| Carl Widland and Robert F. Lax , Weierstrass points on Gorenstein curves | 197 |