

Pacific Journal of Mathematics

**ON COMPLETE SECOND ORDER LINEAR DIFFERENTIAL
EQUATIONS IN BANACH SPACES**

TI-JUN XIAO AND LIANG JIN

ON COMPLETE SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS IN BANACH SPACES

XIO TIJUN AND LIANG JIN

This paper is concerned with the complete second order equation $u''(t) + Bu'(t) + Au(t) = 0$ in a Banach space, where both A and B are densely defined closed linear operators. The main result is a theorem of Hill-Yosida-Phillips type for the Cauchy problem for the equation to be well posed.

1. Introduction and the main result. Consider the complete second order linear differential equation

$$(1.1) \quad u''(t) + Bu'(t) + Au(t) = 0 \quad (t \geq 0)$$

in a complete Banach space E , where A, B are densely defined closed linear operators. The equation has been extensively studied by semi-group methods during the last thirty years. A great amount of literature on it can be looked up in Fattorini's monograph [1] which was published in 1985. However, as stated in [1, Ch. VIII], the theory of (1.1) "can hardly be said in definitive form".

Let us begin with the restatements of some definitions in [1]:

DEFINITION 1. We say that an E -valued function $u(t)$ defined in $t \geq 0$ is a solution of (1.1) if $u(t)$ is twice continuously differentiable, $u(t) \in D(A)$, $u'(t) \in D(B)$, $Au(t)$ and $Bu'(t)$ are continuous and (1.1) is satisfied in $t \geq 0$.

DEFINITION 2. We say that the Cauchy problem for (1.1) is well posed if the following two assumptions are satisfied:

(a) There exist dense subspaces D_0, D_1 of E such that, for any $u_0 \in D_0, u_1 \in D_1$, there exists a solution $u(t)$ of (1.1) with $u(0) = u_0, u'(0) = u_1$.

(b) There exists a nondecreasing, nonnegative function $N(t)$ defined in $t \geq 0$ such that

$$(1.2) \quad \|u(t)\| \leq N(t)(\|u(0)\| + \|u'(0)\|) \quad (t \geq 0)$$

for any solution of (1.1).

Our definition of well posed Cauchy problem corresponds to that of uniformly well posed Cauchy problem in [1]. Thus, in quoting results from [1], “uniformly well posed” should be substituted by “well posed”. See also [2].

DEFINITION 3. Assume that the Cauchy problem for (1.1) is well posed. Define, for $t \geq 0$, $u \in D_0$, $v \in D_1$,

$$C(t)u = u(t), \quad S(t)v = v(t),$$

where $u(t)$ (resp. $v(t)$) is the solution of (1.1) with $u(0) = u$, $u'(0) = 0$ (resp. $v(0) = 0$, $v'(0) = v$). In view of (1.2), $C(t)$ (resp. $S(t)$) is a bounded operator in D_0 (resp. D_1). Since D_0 (resp. D_1) is dense in E we can extend $C(t)$ (resp. $S(t)$) to a bounded operator on E , which we denote by the same symbol. We call the operator-valued functions $C(t)$ and $S(t)$ the propagators of (1.1).

If $B = 0$, (1.1) becomes the incomplete equation

$$(1.3) \quad u''(t) + Au(t) = 0 \quad (t \geq 0).$$

According to [1, 8, 9], if the Cauchy problem for (1.3) is well posed then the solutions grow exponentially and a phase space exists; the well posedness is completely determined by the resolvent of A , that is

THEOREM A [1, 8, 9]. *The Cauchy problem for (1.3) is well posed if and only if there exist constants C , $\omega \geq 0$ such that for $\operatorname{Re} \lambda > \omega$, $(\lambda^2 I + A)^{-1} \in L(E)$ (the set of bounded linear operators on E) and*

$$\|[\lambda(\lambda^2 I + A)^{-1}]^{(n)}\| \leq Cn!(\operatorname{Re} \lambda - \omega)^{-n-1} \quad (n = 0, 1, 2, \dots).$$

However, for the complete equation (1.1), many problems are difficult to discuss if we use the same definition of well posed problem. We may encounter paradoxical situations entailing loss of exponential growth of solutions and nonexistence of phase spaces as has been illustrated by Fattorini [1] with a counterexample. For this, Fattorini has introduced the following

Assumption 3.1 [1, Ch. VIII]. (a) $S(t)u$ is continuously differentiable in $t \geq 0$ for all $u \in E$.

(b) $S(t)E \subseteq D(B)$ and $BS(t)$ is continuous in $t \geq 0$ for all $u \in E$. And he has shown that Assumption 3.1 guarantees exponential growth of solutions and existence of a state space.

DEFINITION 4. We say that the Cauchy problem for (1.1) is strongly well posed if it is well posed and Assumption 3.1 is satisfied.

When $B = 0$, strong well posedness is equivalent to well posedness.

The problem arises of giving necessary and sufficient conditions on A and B for strong well posedness of the Cauchy problem. We give a solution to this problem by proving

THEOREM 1. For equation (1.1) the following statements are equivalent:

- (i) The Cauchy problem for (1.1) is strongly well posed.
- (ii) There exists a complex number λ_0 such that $\Delta(\lambda_0) = \lambda_0^2 I + \lambda_0 B + A$ is closed and densely defined, and $\Delta(\lambda_0)(D(\Delta(\lambda_0))) = E$. The Cauchy problem for (1.1) is well posed and (1.1) has a solution for every initial value $(u_0, u_1) \in (D(A) \cap D(B))^2 = (D(A) \cap D(B)) \times (D(A) \cap D(B))$.
- (iii) $D(A) \cap D(B)$ is dense in E . There exist constants $C, \omega \geq 0$ such that for $\text{Re } \lambda > \omega, \Delta(\lambda)^{-1} = (\lambda^2 I + \lambda B + A)^{-1} \in L(E), \Delta(\lambda)^{-1} A$ is closable and

$$\begin{aligned} \|\lambda \Delta(\lambda)^{-1}\|^{(n)} &\leq Cn!(\text{Re } \lambda - \omega)^{-n-1} & (n = 0, 1, 2, \dots), \\ \|B\Delta(\lambda)^{-1}\|^{(n)} &\leq Cn!(\text{Re } \lambda - \omega)^{-n-1} & (n = 0, 1, 2, \dots), \\ \|\Delta(\lambda)^{-1}Bu\|^{(n)} &\leq Cn!(\text{Re } \lambda - \omega)^{-n-1}\|u\| \\ & & (u \in D(A) \cap D(B), n = 0, 1, 2, \dots). \end{aligned}$$

Moreover, if (iii) is satisfied, we have three kinds of explicit expressions for the propagators:

$$(1.4) \quad C(t) = \lim_{n \rightarrow \infty} \frac{(-1)^n}{n!} \left(\frac{n}{t}\right)^{n+1} \left[\frac{1}{\lambda} I - \frac{1}{\lambda} \overline{\Delta(\lambda)^{-1}A} \right]^{(n)} \Bigg|_{\lambda=n/t} \quad (t > 0),$$

$$(1.5) \quad S(t) = \lim_{n \rightarrow \infty} \frac{(-1)^n}{n!} \left(\frac{n}{t}\right)^{n+1} [\Delta(\lambda)^{-1}]^{(n)} \Bigg|_{\lambda=n/t} \quad (t > 0);$$

$$(1.6) \quad C(t)u = u - \lim_{\lambda \rightarrow \infty} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n!} e^{n\lambda t} \Delta(n\lambda)^{-1} Au \quad (u \in D(A), t \geq 0),$$

$$(1.7) \quad S(t)u = \lim_{\lambda \rightarrow \infty} \lambda \sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{(n-1)!} e^{n\lambda t} \Delta(n\lambda)^{-1} u \quad (u \in E, t \geq 0);$$

and for $\nu > \omega$,

$$(1.8) \quad C(t)u = u - \frac{1}{2\pi i} \int_{\nu-i\infty}^{\nu+i\infty} \frac{1}{\lambda} e^{\lambda t} \Delta(\lambda)^{-1} A u d\lambda \quad (u \in D(A), t \geq 0),$$

$$(1.9) \quad S(t)u = \frac{1}{2\pi i} \int_{\nu-i\infty}^{\nu+i\infty} e^{\lambda t} \Delta(\lambda)^{-1} u d\lambda \quad (u \in E, t \geq 0).$$

REMARK 1. Theorem 1 contains Theorem A.

REMARK 2. Although the implication (iii) \Rightarrow (i) could be possibly proved by direct application of the Hill-Yosida theorem, complications stemming from lack of commutativity of A and B make this approach impractical.

2. Proof of Theorem 1. First of all, we present three lemmas.

LEMMA 1. Let $f(t)$ be an E -valued continuous function defined in $t \geq 0$ such that $\int_0^\infty e^{-ct} f(t) dt$ exists for some positive c . Then, as $k \rightarrow \infty$,

$$(2.1) \quad M_k(t) = \left(\frac{k}{t}\right)^k \frac{1}{(k-1)!} \int_0^\infty e^{-ks/t} s^{k-1} f(s) ds \rightarrow f(t),$$

$$(2.2) \quad N_k(t) = \left(\frac{k}{t}\right)^{k+1} \frac{1}{k!} \int_0^\infty e^{-ks/t} s^k f(s) ds \rightarrow f(t),$$

uniformly on compact subsets of $t > 0$.

The proof of (2.2) is essentially the same as the one of [11, P. 285, Th. 5a] and we omit it. (2.1) follows immediately from $M_k(t) = N_{k-1}((k-1)t/k)$.

LEMMA 2. Let $f(t)$ be an E -valued continuous function with $\|f(t)\| \leq C e^{\omega t}$ in $t \geq 0$, where $C, \omega \geq 0$, then

$$\int_0^t f(s) ds = \text{Lim}_{\lambda \rightarrow \infty} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n!} e^{n\lambda t} \int_0^\infty e^{-n\lambda r} f(r) dr \quad (t \geq 0).$$

The proof is completely the same as the first part in the proof of (Phragmén's representation theorem, see [6]) and we also omit it.

LEMMA 3. Let $f(t)$ be an E -valued continuously differentiable function with $\|f(t)\| \leq Ce^{\omega t}$ in $t \geq 0$, where $C, \omega \geq 0$, then for $\bar{\omega} > \omega$

$$f(t) = \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} e^{\lambda t} \left[\int_0^\infty e^{-\lambda s} f(s) ds \right] d\lambda \quad (t > 0).$$

Proof. By [4, Th. 6.3.1],

$$f(t) = \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{\bar{\omega}-iT}^{\bar{\omega}+iT} e^{\lambda t} \left[\int_0^\infty e^{-\lambda s} f(s) ds \right] d\lambda \quad (t > 0).$$

It remains to show that the integral

$$\int_{\bar{\omega}+i\infty}^{\bar{\omega}+i\infty} e^{\lambda t} \left[\int_0^\infty e^{-\lambda s} f(s) ds \right] d\lambda$$

converges. We can prove this fact using arguments similar to those of [11, P. 68, Th. 7.5], noting that Riemann-Lebesgue theorem is applicable to vector-valued functions (see [3, P. 401]) and making use of the estimate $\|f(t)\| \leq Ce^{\omega t}$ for $t \geq 0$.

Proof of Theorem 1. (i) \Rightarrow (ii). By [1, §VIII.3].

(ii) \Rightarrow (i). Let $u(t)$ be a solution of (1.1) with $u(0), u'(0) \in D(A) \cap D(B)$, then $u(t) \in D(A)$ for $t \geq 0$, $Bu'(t)$ is continuous in $t \geq 0$ and therefore $Bu'(t)$ is integrable on any bounded interval of $t \geq 0$. Hence $u(t) = u(0) + \int_0^t u'(s) ds \in D(B)$ for $t \geq 0$. Set $v(t) = e^{-\lambda_0 t} u(t)$; then

$$\begin{aligned} u'(t) &= \lambda_0 e^{\lambda_0 t} v(t) + e^{\lambda_0 t} v'(t), \\ u''(t) &= \lambda_0^2 e^{\lambda_0 t} v(t) + 2\lambda_0 e^{\lambda_0 t} v'(t) + e^{\lambda_0 t} v''(t). \end{aligned}$$

Since $v(t) \in D(A) \cap D(B)$, we have

$$(2.3) \quad v''(t) + B_1 v'(t) + A_1 v(t) = 0,$$

where $B_1 = B + 2\lambda_0 I, A_1 = \Delta(\lambda_0) = \lambda_0^2 I + \lambda_0 B + A$. Obviously $u(0) = v(0), u'(0) = \lambda_0 v(0) + v'(0)$; hence for every initial value $(v(0), v'(0)) \in (D(A_1))^2 = (D(A) \cap D(B))^2$ the equation (2.3) has a solution. It is easily verified that if $v(t)$ is a solution of (2.3) then $u(t) = e^{\lambda_0 t} v(t)$ is a solution of (1.1). From these observations we deduce that the Cauchy problem for (2.3) is well-posed. Denoting the propagators of (2.3) by $C_1(t)$ and $S_1(t)$, clearly $S_1(t) = e^{-\lambda_0 t} S(t)$ for $t \geq 0$. Since $A_1(D(A_1)) = E$. Assumption 3.1 is then satisfied for equation (2.3) in view of [2, Th. 4.1(b)]. Thus Assumption 3.1 holds for equation (1.1) and this completes the proof.

(i) \Rightarrow (iii). According to [1], there exist constants $C, \omega \geq 0$ such that for $\operatorname{Re} \lambda > \omega, \Delta(\lambda)^{-1} \in L(E)$ and

$$\begin{aligned}\Delta(\lambda)^{-1}u &= \int_0^\infty e^{-\lambda t} S(t)u \, dt \quad (u \in E); \\ u - \int_0^t S(s)Au \, ds &= C(t)u \quad (t \geq 0, u \in D(A))\end{aligned}$$

and for $t \geq 0$.

$$\|S'(t)\| \leq Ce^{\omega t}, \quad \|BS(t)\| \leq Ce^{\omega t}, \quad \|C(t)\| \leq Ce^{\omega t}.$$

Consequently, for $u \in E$,

$$\begin{aligned}\lambda\Delta(\lambda)^{-1}u &= \int_0^\infty \lambda e^{-\lambda t} S(t)u \, dt = \int_0^\infty e^{-\lambda t} S'(t)u \, dt, \\ B\Delta(\lambda)^{-1}u &= \int_0^\infty e^{-\lambda t} BS(t)u \, dt.\end{aligned}$$

Hence

$$\begin{aligned}\|[\lambda\Delta(\lambda)^{-1}]^{(n)}\| &\leq \int_0^\infty t^n e^{-\operatorname{Re} \lambda t} Ce^{\omega t} \, dt \\ &= Cn!(\operatorname{Re} \lambda - \omega)^{-n-1} \quad (n = 0, 1, 2, \dots), \\ \|[B\Delta(\lambda)^{-1}]^{(n)}\| &\leq \int_0^\infty t^n e^{-\operatorname{Re} \lambda t} Ce^{\omega t} \, dt \\ &= Cn!(\operatorname{Re} \lambda - \omega)^{-n-1} \quad (n = 0, 1, 2, \dots).\end{aligned}$$

Also

$$\begin{aligned}\frac{1}{\lambda}\Delta(\lambda)^{-1}Au &= \frac{1}{\lambda} \int_0^\infty e^{-\lambda t} S(t)Au \, dt \\ &= \int_0^\infty e^{-\lambda t} \left[\int_0^t S(s)Au \, ds \right] dt \quad (u \in D(A)),\end{aligned}$$

Then for $u \in D(A), n = 0, 1, 2, \dots$,

$$\begin{aligned}\left\| \left[\frac{1}{\lambda}\Delta(\lambda)^{-1}Au \right]^{(n)} \right\| &\leq \int_0^\infty t^n e^{-\operatorname{Re} \lambda t} (1 + Ce^{\omega t}) \|u\| \, dt \\ &\leq (C + 1)n!(\operatorname{Re} \lambda - \omega)^{-n-1} \|u\|.\end{aligned}$$

But

$$\Delta(\lambda)^{-1}Bu = \frac{1}{\lambda}u - \lambda\Delta(\lambda)^{-1}u - \frac{1}{\lambda}\Delta(\lambda)^{-1}Au \quad (u \in D(A) \cap D(B)).$$

Thus, we obtain

$$\begin{aligned} \|\Delta(\lambda)^{-1}Bu\| \leq 2(C + 1)n!(\operatorname{Re} \lambda - \omega)^{-n-1}\|u\| \\ (u \in D(A) \cap D(B), n = 0, 1, 2, \dots). \end{aligned}$$

This ends the proof of the implication (i) \Rightarrow (iii).

(iii) \Rightarrow (i). We define a linear operator $G = \begin{pmatrix} 0 & -I \\ A & B \end{pmatrix}$ in the space $E \times E$ with domain $D(G) = D(A) \times (D(A) \cap D(B))$. It is easy to verify that for $\lambda > \omega$,

$$\begin{aligned} (\lambda I + G)^{-1} &= \begin{pmatrix} \frac{1}{\lambda}I - \Delta(\lambda)^{-1}A & \Delta(\lambda)^{-1} \\ -\Delta(\lambda)^{-1}A & \lambda\Delta(\lambda)^{-1} \end{pmatrix}, \\ D((\lambda I + G)^{-1}) &= D(A) \times E. \end{aligned}$$

By virtue of the equality $\frac{1}{\lambda}\Delta(\lambda)^{-1}Au = \frac{1}{\lambda}u - \lambda\Delta(\lambda)^{-1}u - \Delta(\lambda)^{-1}Bu$ for $u \in D(A) \cap D(B)$ and the fact that $\Delta(\lambda)^{-1}A$ is closable, we can extend $\Delta(\lambda)^{-1}A$ to a bounded operator on E and therefore can extend $(\lambda I + G)^{-1}$ to a bounded operator on $E \times E$, which is $\overline{(\lambda I + G)^{-1}}$. Accordingly, G is closable and $(\lambda I + \overline{G})^{-1} = \overline{(\lambda I + G)^{-1}}$. By [10, P. 73], for $t > 0, n > \omega t$,

$$\frac{d}{dt} \left[\left(\frac{n}{t}\right)^n \left(\frac{n}{t}I + \overline{G}\right)^{-n} \right] = - \left(\frac{n}{t}\right)^{n+1} \left(\frac{n}{t}I + \overline{G}\right)^{-n} \overline{G} \left(\frac{n}{t}I + \overline{G}\right)^{-1}.$$

Set, for $u \in D(A), v \in E$,

$$\begin{aligned} X_n(t; u, v) &= \left[\left(\frac{n}{t}\right)^n \left(\frac{n}{t}I + G\right)^{-n} \right] \begin{pmatrix} u \\ v \end{pmatrix}, \\ U_n(t; u, v) &= \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ \left[\frac{1}{\lambda}u - \frac{1}{\lambda}\Delta(\lambda)^{-1}Au \right]^{(n-1)} \right. \\ &\quad \left. + [\Delta(\lambda)^{-1}v]^{(n-1)} \right\} \Big|_{\lambda=n/t}, \\ V_n(t; u, v) &= \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \{ [-\Delta(\lambda)^{-1}Au]^{(n-1)} \\ &\quad + [\lambda\Delta(\lambda)^{-1}v]^{(n-1)} \} \Big|_{\lambda=n/t}. \end{aligned}$$

Then obviously

$$\frac{d}{dt}X_n(t; u, v) + GX_{n+1}\left(\frac{n+1}{n}t; u, v\right) = 0.$$

But

$$\begin{aligned}
X_n(t; u, v) &= \left\{ \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [(\lambda I + G)^{-1}]^{(n-1)} \Big|_{\lambda=n/t} \right\} \begin{pmatrix} u \\ v \end{pmatrix} \\
&= \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left(\begin{array}{l} [\frac{1}{\lambda}u - \frac{1}{\lambda}\Delta(\lambda)^{-1}Au]^{(n-1)} + [\Delta(\lambda)^{-1}v]^{(n-1)} \\ [-\Delta(\lambda)^{-1}Au]^{(n-1)} + [\lambda\Delta(\lambda)^{-1}v]^{(n-1)} \end{array} \right) \Big|_{\lambda=n/t} \\
&= \begin{pmatrix} U_n(t; u, v) \\ V_n(t; u, v) \end{pmatrix},
\end{aligned}$$

hence we have

$$(2.4) \quad \begin{cases} \frac{d}{dt}U_n(t; u, v) = V_{n+1}\left(\frac{n+1}{n}t; u, v\right), \\ \frac{d}{dt}V_n(t; u, v) = -BV_{n+1}\left(\frac{n+1}{n}t; u, v\right) - AU_{n+1}\left(\frac{n+1}{n}t; u, v\right). \end{cases}$$

Next, we shall discuss the convergence of (2.4) as $n \rightarrow \infty$. Define, for $t \geq 0$, $v \in D(A) \cap D(B)$,

$$\begin{aligned}
\tilde{S}(t)v &= tv - \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \frac{e^{\lambda t}}{\lambda^2} (\lambda\Delta(\lambda)^{-1})Bv \, d\lambda \\
&\quad - \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \frac{e^{\lambda t}}{\lambda^2} \Delta(\lambda)^{-1}Av \, d\lambda \\
&= \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} e^{\lambda t} \Delta(\lambda)^{-1}v \, d\lambda,
\end{aligned}$$

where $\bar{\omega} > \omega$. Clearly $\tilde{S}(0) = v = 0$, $\tilde{S}(t)v$ is continuous in $t \geq 0$ and for $t \geq 0$

$$\begin{aligned}
\int_0^t \tilde{S}(s)v \, ds &= \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \frac{e^{\lambda t}}{\lambda^2} \lambda\Delta(\lambda)^{-1}v \, d\lambda, \\
B \int_0^t \tilde{S}(s)v \, ds &= \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \frac{e^{\lambda t}}{\lambda} B\Delta(\lambda)^{-1}v \, d\lambda \\
&= \frac{1}{2}t^2Bv - \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \frac{e^{\lambda t}}{\lambda^2} B\Delta(\lambda)^{-1}Bv \, d\lambda \\
&\quad - \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \frac{e^{\lambda t}}{\lambda^3} B\Delta(\lambda)^{-1}Av \, d\lambda, \\
B \int_0^t (t-s)\tilde{S}(s)v \, ds &= \frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \frac{e^{\lambda t}}{\lambda^2} B\Delta(\lambda)^{-1}v \, d\lambda.
\end{aligned}$$

Making use of the Fubini theorem and the Cauchy formula, we obtain that for $\mu > \bar{\omega}$, $k = 0, 1, 2, \dots$,

$$\begin{aligned} & \int_0^\infty e^{-\mu t} (-t)^k \tilde{S}(t)v \, dt \\ &= \int_0^\infty e^{-\mu t} (-t)^k \left[\frac{1}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} e^{\lambda t} \Delta(\lambda)^{-1} v \, d\lambda \right] dt \\ &= \frac{(-1)^k}{2\pi i} \int_{\bar{\omega}-i\infty}^{\bar{\omega}+i\infty} \Delta(\lambda)^{-1} v \left[\int_0^\infty e^{(\lambda-\mu)t} t^k \, dt \right] d\lambda = [\Delta(\mu)^{-1} v]^{(k)}, \\ & \int_0^\infty e^{-\mu t} (-t)^k \left[\int_0^t \tilde{S}(s)v \, ds \right] dt = \left[\frac{1}{\mu} \Delta(\mu)^{-1} v \right]^{(k)}, \\ & \int_0^\infty e^{-\mu t} (-t)^k \left[B \int_0^t \tilde{S}(s)v \, ds \right] dt = \left[\frac{1}{\mu} B \Delta(\mu)^{-1} v \right]^{(k)}, \\ & \int_0^\infty e^{-\mu t} (-t)^k \left[B \int_0^t (t-s) \tilde{S}(s)v \, ds \right] dt = \left[\frac{1}{\mu^2} B \Delta(\mu)^{-1} v \right]^{(k)}. \end{aligned}$$

This and the obvious fact that $\tilde{S}(t)v$, $\int_0^t \tilde{S}(s)v \, ds$, $B \int_0^t \tilde{S}(s)v \, ds$ and $B \int_0^t (t-s) \tilde{S}(s)v \, ds$ are continuous in $t \geq 0$, in view of (2.1) in Lemma 1, show that as $n \rightarrow \infty$

$$(2.5) \quad \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [\Delta(\lambda)^{-1} v]^{(n-1)} \Big|_{\lambda=n/t} \rightarrow \tilde{S}(t)v \quad (t > 0),$$

$$(2.6) \quad \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left[\frac{1}{\lambda} \Delta(\lambda)^{-1} v \right]^{(n-1)} \Big|_{\lambda=n/t} \rightarrow \int_0^t \tilde{S}(s)v \, ds \quad (t > 0),$$

$$(2.7) \quad \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left[\frac{1}{\lambda} B \Delta(\lambda)^{-1} v \right]^{(n-1)} \Big|_{\lambda=n/t} \rightarrow B \int_0^t \tilde{S}(s)v \, ds \quad (t > 0),$$

$$(2.8) \quad \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left[\frac{1}{\lambda^2} B \Delta(\lambda)^{-1} v \right]^{(n-1)} \Big|_{\lambda=n/t} \rightarrow B \int_0^t (t-s) \tilde{S}(s)v \, ds \quad (t > 0),$$

uniformly on compact subsets of $t > 0$. It is easy to verify by the Leibniz formula that $\|[\Delta(\lambda)^{-1}]^{(n)}\|$, $\|[\frac{1}{\lambda} \Delta(\lambda)^{-1}]^{(n)}\|$ and $\|[\frac{1}{\lambda^2} B \Delta(\lambda)^{-1}]^{(n)}\|$ are all bounded by $Cn!(\text{Re } \lambda - \omega)^{-n-1}$ for $\text{Re } \lambda > \omega$, $n = 0, 1, 2, \dots$ (if the

constant is still denoted by C). Therefore for $t > 0$, $v \in D(A) \cap D(B)$,

$$\begin{aligned} \|\tilde{S}(t)v\| &= \left\| \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [\Delta(\lambda)^{-1}v]^{(n-1)} \Big|_{\lambda=n/t} \right\| \\ &\leq \operatorname{Lim}_{n \rightarrow \infty} \frac{C}{(n-1)!} \left(\frac{n}{t}\right)^n (n-1)! \left(\frac{n}{t} - \omega\right)^{-n} \|v\| \\ &= \operatorname{Lim}_{n \rightarrow \infty} C \left(1 - \frac{\omega t}{n}\right)^{-n} \|v\| = Ce^{\omega t} \|v\|, \\ \left\| \int_0^t \tilde{S}(s)v \, ds \right\| &\leq \operatorname{Lim}_{n \rightarrow \infty} C \left(1 - \frac{\omega t}{n}\right)^{-n} \|v\| = Ce^{\omega t} \|v\|, \\ \left\| B \int_0^t \tilde{S}(s)v \, ds \right\| &\leq Ce^{\omega t} \|v\|, \\ \left\| B \int_0^t (t-s)\tilde{S}(s)v \, ds \right\| &\leq Ce^{\omega t} \|v\|. \end{aligned}$$

Accordingly, $\tilde{S}(t)$, ($t \geq 0$) can be extended to all of E as a bounded operator which we denote by the same symbol; recalling that $\tilde{S}(0)v = 0$ and $\tilde{S}(t)v$ is continuous in $t \geq 0$ for $v \in D(A) \cap D(B)$, we can assert that $\tilde{S}(0) = 0$ and $\tilde{S}(t)$ is strongly continuous in $t \geq 0$. By virtue of the denseness of $D(A) \cap D(B)$ and the uniform boundedness of

$$\frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [\Delta(\lambda)^{-1}]^{(n-1)} \Big|_{\lambda=n/t}$$

on bounded subsets of $t > 0$, we deduce that (2.5) is valid for all $v \in E$. Similarly, (2.6), (2.7) and (2.8) also hold for all $v \in E$ (here, the closedness of B is used); moreover, $\int_0^t \tilde{S}(s)v \, ds$, $B \int_0^t \tilde{S}(s)v \, ds$ and $B \int_0^t (t-s)\tilde{S}(s)v \, ds$ are continuous in $t \geq 0$ for $v \in E$.

Based on the paragraph above, we shall define several operators. First, define, for $t > 0$, $u \in D(A)$,

$$\begin{aligned} (2.9) \quad \tilde{C}(t)u &= \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^n}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ \left[\frac{1}{\lambda} u \right]^{(n-1)} \right. \\ &\quad \left. - \left[\frac{1}{\lambda} \Delta(\lambda)^{-1} Au \right]^{(n-1)} \right\} \Big|_{\lambda=n/t} \\ &= u - \int_0^t \tilde{S}(s)Au \, ds. \end{aligned}$$

Since for $u \in D(A) \cap D(B)$, $\operatorname{Re} \lambda > \omega$, $n = 0, 1, 2, \dots$

$$(2.10) \quad \left\| \left[\frac{1}{\lambda} u - \frac{1}{\lambda} \Delta(\lambda)^{-1} A u \right]^{(n)} \right\| \\ = \| [\lambda \Delta(\lambda)^{-1} u + \Delta(\lambda)^{-1} B u]^{(n)} \| \\ \leq 2C n! (\operatorname{Re} \lambda - \omega)^{-n-1} \|u\|,$$

and $\Delta(\lambda)^{-1} A$ is closable, which implies that (2.10) is also valid for $u \in D(A)$, we have that for $u \in D(A)$, $t > 0$,

$$\|\tilde{C}(t)u\| \leq \operatorname{Lim}_{n \rightarrow \infty} 2C \frac{1}{(n-1)!} \left(\frac{n}{t}\right)^n (n-1)! \left(\frac{n}{t} - \omega\right)^{-n} \|u\| \\ = 2C e^{\omega t} \|u\|,$$

thus $\tilde{C}(t)$ ($t > 0$) can be extended to a bounded operator on E which we denote by the same symbol. Define $\tilde{C}(0) = I$. Then $\tilde{C}(t)$ is strongly continuous in $t \geq 0$.

Define, for $t > 0$, $v \in D(A) \cap D(B)$,

$$\begin{aligned} \tilde{K}(t)v &= \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [\lambda \Delta(\lambda)^{-1} v]^{(n-1)} \Big|_{\lambda=n/t} \\ &= \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ \left[\frac{1}{\lambda} v \right]^{(n-1)} - [\Delta(\lambda)^{-1} B v]^{(n-1)} \right. \\ &\quad \left. - \left[\frac{1}{\lambda} \Delta(\lambda)^{-1} A v \right]^{(n-1)} \right\} \Big|_{\lambda=n/t} \\ &= v - \tilde{S}(t) B v - \int_0^t \tilde{S}(s) A v \, ds, \\ \tilde{T}(t)v &= \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [B \Delta(\lambda)^{-1} v]^{(n-1)} \Big|_{\lambda=n/t} \\ &= \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ \left[\frac{1}{\lambda^2} B v \right]^{(n-1)} - \left[\frac{1}{\lambda} B \Delta(\lambda)^{-1} B v \right]^{(n-1)} \right. \\ &\quad \left. - \left[\frac{1}{\lambda^2} B \Delta(\lambda)^{-1} A v \right]^{(n-1)} \right\} \Big|_{\lambda=n/t} \\ &= -B \int_0^t \tilde{S}(s) B v \, ds - B \int_0^t (t-s) \tilde{S}(s) A v \, ds. \end{aligned}$$

Here the limits are uniform on compacts of $t > 0$. But

$$\begin{aligned} \|K(t)v\| &= \left\| \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [\lambda \Delta(\lambda)^{-1}v]^{(n-1)} \Big|_{\lambda=n/t} \right\| \\ &\leq \operatorname{Lim}_{n \rightarrow \infty} \frac{C}{(n-1)!} \left(\frac{n}{t}\right)^n (n-1)! \left(\frac{n}{t} - \omega\right)^{-n} \|v\| = Ce^{\omega t} \|v\|, \\ \|\tilde{T}(t)v\| &\leq \operatorname{Lim}_{n \rightarrow \infty} \frac{C}{(n-1)!} \left(\frac{n}{t}\right)^n (n-1)! \left(\frac{n}{t} - \omega\right)^{-n} \|v\| \\ &= Ce^{\omega t} \|v\|; \end{aligned}$$

therefore $\tilde{K}(t)$ and $\tilde{T}(t)$ ($t > 0$) can be extended to all of E as bounded operators which we denote by the same symbols. Define $\tilde{K}(0) = I$, $\tilde{T}(0) = 0$, then $\tilde{K}(t)$ and $\tilde{T}(t)$ are strongly continuous in $t \geq 0$. Arguing as in the treatment of (2.5), we have that for all $v \in E$,

$$(2.11) \quad \tilde{K}(t)v = \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [\lambda \Delta(\lambda)^{-1}v]^{(n-1)} \Big|_{\lambda=n/t},$$

$$(2.12) \quad \tilde{T}(t)v = \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [B\Delta(\lambda)^{-1}v] \Big|_{\lambda=n/t}.$$

The limits are uniform on compacts of $t > 0$.

Now, let us turn to (2.4). Define

$$\begin{aligned} (2.13) \quad u(t; u, v) &= \operatorname{Lim}_{n \rightarrow \infty} U_n(t; u, v) \\ &= \operatorname{Lim}_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ \left[\frac{1}{\lambda} u - \frac{1}{\lambda} \Delta(\lambda)^{-1} Au \right]^{(n-1)} \right. \\ &\quad \left. + [\Delta(\lambda)^{-1}v]^{(n-1)} \right\} \Big|_{\lambda=n/t} \\ &= \tilde{C}(t)u + \tilde{S}(t)v, \end{aligned}$$

$$(2.14) \quad v(t; u, v) = \operatorname{Lim}_{n \rightarrow \infty} V_n(t; u, v) = -\tilde{S}(t)Au + \tilde{K}(t)v,$$

where the limits are uniform on compacts of $t > 0$. By the closedness

of A, B , we obtain

$$\begin{aligned}
 AU_n(t; u, v) &= \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ \left[\frac{1}{\lambda} Au - \frac{1}{\lambda} A\Delta(\lambda)^{-1} Au \right]^{(n-1)} \right. \\
 &\quad \left. + [A\Delta(\lambda)^{-1} v]^{(n-1)} \right\} \Big|_{\lambda=n/t} \\
 &= \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ \left[\frac{1}{\lambda} Au - \frac{1}{\lambda} A\Delta(\lambda)^{-1} Au \right]^{(n-1)} \right. \\
 &\quad \left. + \left[\frac{1}{\lambda^2} Av - \frac{1}{\lambda} A\Delta(\lambda)^{-1} Bv \right. \right. \\
 &\quad \left. \left. - \frac{1}{\lambda^2} A\Delta(\lambda)^{-1} Av \right]^{(n-1)} \right\} \Big|_{\lambda=n/t}, \\
 BV_n(t; u, v) &= \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \{ [-B\Delta(\lambda)^{-1} Au]^{(n-1)} \\
 &\quad + [\lambda B\Delta(\lambda)^{-1} v]^{(n-1)} \} \Big|_{\lambda=n/t} \\
 &= \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n \left\{ [-B\Delta(\lambda)^{-1} Au]^{(n-1)} \right. \\
 &\quad \left. + \left[\frac{1}{\lambda} Bv - B\Delta(\lambda)^{-1} Bv \right. \right. \\
 &\quad \left. \left. - \frac{1}{\lambda} B\Delta(\lambda)^{-1} Av \right]^{(n-1)} \right\} \Big|_{\lambda=n/t}.
 \end{aligned}$$

Also

$$\frac{1}{\lambda} A\Delta(\lambda)^{-1} = \frac{1}{\lambda} I - B\Delta(\lambda)^{-1} - \lambda\Delta(\lambda)^{-1};$$

therefore we have

$$\begin{aligned}
 (2.15) \quad Au(t; u, v) &= \lim_{n \rightarrow \infty} AU_n(t; u, v) \\
 &= \tilde{T}(t)Au + \tilde{K}(t)Au - Bv + \tilde{T}(t)Bv + \tilde{K}(t)Bv \\
 &\quad + B \int_0^t \tilde{S}(s)Av ds + \tilde{S}(t)Av,
 \end{aligned}$$

$$\begin{aligned}
 (2.16) \quad Bv(t; u, v) &= \lim_{n \rightarrow \infty} BV_n(t; u, v) \\
 &= -\tilde{T}(t)Au + Bv - \tilde{T}(t)Bv - B \int_0^t \tilde{S}(s)Av ds,
 \end{aligned}$$

where the limits are uniform on compacts of $t > 0$. Accordingly, as $n \rightarrow \infty$, (2.4) becomes

$$(2.17) \quad \begin{cases} \frac{d}{dt}u(t; u, v) = v(t; u, v), \\ \frac{d}{dt}v(t; u, v) = -Bv(t; u, v) - Au(t; u, v), \end{cases}$$

in $t > 0$. Define $u(0; u, v) = u$, $v(0; u, v) = v$ for $u \in D(A)$, $v \in D(A) \cap D(B)$; then by (2.13), (2.14), (2.15) and (2.16), $u(t; u, v)$, $v(t; u, v)$, $Bv(t; u, v)$ and $Au(t; u, v)$ are continuous in $t \geq 0$, and therefore (2.17) holds in $t \geq 0$, i.e.

$$u''(t; u, v) + Bu'(t; u, v) + Au(t; u, v) = 0 \quad (t \geq 0).$$

We have then proved that the equation (1.1) has a solution for every initial value $(u(0), u'(0)) \in (D(A) \cap D(B))^2$. It remains to show continuous dependence on initial data, in view of the fact that the statement (ii) implies the statement (i) (as proved before). To this end we observe that, from (2.9) and combining with (2.11), (2.13), (2.14) and (2.17), we obtain

$$\begin{aligned} \tilde{C}(t)u &= \lim_{n \rightarrow \infty} \frac{(-1)^{n-1}}{(n-1)!} \left(\frac{n}{t}\right)^n [\Delta(\lambda)^{-1}(\lambda + B)u]^{(n-1)} \Big|_{\lambda=n/t} \\ &= \tilde{S}'(t)u + \tilde{S}(t)Bu \end{aligned}$$

for $u \in D(A) \cap D(B)$, $t > 0$. Hence,

$$(2.19) \quad \tilde{S}(t)u = \int_0^t [\tilde{C}(s)u - \tilde{S}(s)Bu] ds \quad (u \in D(A) \cap D(B), t \geq 0).$$

Let now $w(t)$ be an arbitrary solution of (1.1). Set, for $t \geq 0$, (n a natural number), $\varphi_n(t) = \int_0^{1/n} n(s+1)w'(t+s) ds$. It is clear that for $t \geq 0$, $\varphi_n(t) \in D(B)$, $\varphi_n(t) \rightarrow w'(t)$ and $B\varphi_n(t) \rightarrow Bw'(t)$ as $n \rightarrow \infty$. Moreover, integrating by parts, we obtain

$$\varphi_n(t) = n(s+1)w(t+s)|_0^{1/n} - \int_0^{1/n} nw(s+t) ds \in D(A) \quad (t \geq 0).$$

Thus (2.19) holds for $u \in w'(t)$ ($t \geq 0$). From this and (2.9), we deduce

$$\begin{aligned} &\frac{d}{ds}[\tilde{C}(t-s)w(s) + \tilde{S}(t-s)w'(s)] \\ &= -\tilde{C}'(t-s)w(s) + \tilde{C}(t-s)w'(s) - \tilde{S}'(t-s)w'(s) \\ &\quad + \tilde{S}(t-s)w''(s) \\ &= \tilde{S}(t-s)Aw(s) + \tilde{C}(t-s)w'(s) - \tilde{C}(t-s)w'(s) \\ &\quad + \tilde{S}(t-s)Bw'(s) + \tilde{S}(t-s)[-Bw'(s) - Aw(s)] \\ &= 0 \quad (0 \leq s \leq t). \end{aligned}$$

Consequently

$$w(t) = \tilde{C}(t)w(0) + \tilde{S}(t)w'(0) \quad (t \geq 0),$$

and this ends the proof of the implication (iii) \Rightarrow (i).

Finally, we show the explicit expressions (1.4) to (1.9). By virtue of the equivalence of (2.1) and (2.2) in Lemma 1, (1.4) (resp. (1.5)) results from (2.9) (resp. (2.5)). Recalling that for $\text{Re } \lambda > \omega$,

$$\begin{aligned} \Delta(\lambda)^{-1}u &= \int_0^\infty e^{-\lambda t} S(t)u \, dt, \\ \lambda \Delta(\lambda)^{-1}u &= \int_0^\infty e^{-\lambda t} S'(t)u \, dt \quad (u \in E), \\ \frac{1}{\lambda} \Delta(\lambda)^{-1}Au &= \int_0^\infty e^{-\lambda t} \left[\int_0^t S(s)Au \, ds \right] \quad (u \in D(A)), \end{aligned}$$

then (1.6) follows from (2.9) and Lemma 2, (1.7) from Lemma 2, (1.8) from (2.9) and Lemma 3, (1.9) from Lemma 3. Thus we have completed the proof of Theorem 1.

3. Applications of Theorem 1. F. Neubrander [5] has discussed the case $D(A) \supset D(B)$. He shows well posedness under the assumption that $-B$ is the generator of a strongly continuous semigroup and $R(\lambda, -B)A = AR(\lambda, -B)$ on $D(A)$ for $\text{Re } \lambda = \omega$ (ω a constant). As a consequence of Theorem 1, the following theorem generalizes the result.

THEOREM 2. *Suppose $D(A) \supset D(B)$ and there exists a complex number λ_0 such that $(\lambda_0 I + B)^{-1}A$ has bounded extension. Then the Cauchy problem for (1.1) is strongly well posed if and only if $-B$ is the generator of a strongly continuous semigroup.*

Proof. Necessity. According to [1, Ch. VIII, Corollary 3.5], for every initial value $(u_0, u_1) \in D(B) \times D(B)$ the Cauchy problem for (1.1) has a solution. This ends the proof by the well posedness and [5, Theorem 5].

Sufficiency. We assume $\lambda_0 = 0$ without the loss of generality. Let now $(T(t))$ be the semigroup generated by $-B$. It is well known that there exist $C, \omega \geq 0$ such that

$$\begin{aligned} (3.1) \quad (\lambda I + B)^{-1}u &= \int_0^\infty e^{-\lambda t} T(t)u \, dt \quad (u \in E, \text{Re } \lambda > \omega), \\ \|T(t)\| &\leq Ce^{\omega t} \quad (t \geq 0). \end{aligned}$$

Hence, we have

$$(3.2) \quad \begin{cases} \frac{1}{\lambda}A(\lambda I + B)^{-1}u = \frac{1}{\lambda}AB^{-1}B(\lambda I + B)^{-1}u \\ = \int_0^\infty e^{-\lambda t}AB^{-1}[I - T(t)]u dt \quad (u \in E, \operatorname{Re} \lambda > \omega), \\ \|\lambda AB^{-1}[I - T(t)]\| \leq Ce^{\omega t} \quad (t \geq 0), \end{cases}$$

(here and in the sequel, we denote by C a generic constant).

It is clear that for $\operatorname{Re} \lambda > \omega + C$,

$$(3.3) \quad \begin{aligned} \lambda \Delta(\lambda)^{-1} &= (\lambda I + B)^{-1} \left[I + \frac{1}{\lambda}A(\lambda I + B)^{-1} \right]^{-1} \\ &= (\lambda I + B)^{-1} \sum_{n=0}^{\infty} \left[-\frac{1}{\lambda}A(\lambda I + B)^{-1} \right]^n. \end{aligned}$$

In order to estimate $[\lambda \Delta(\lambda)^{-1}]^{(n)}$ for $n = 0, 1, 2, \dots$, we observe that, by (3.1),

$$[(\lambda I + B)^{-1}u]^{(m)} = \int_0^\infty (-t)^m e^{-\lambda t} T(t)u dt \quad (u \in E, m = 0, 1, 2, \dots),$$

and therefore

$$(3.4) \quad \begin{aligned} \|[(\lambda I + B)^{-1}]^{(m)}\| &\leq \int_0^\infty t^m e^{-\operatorname{Re} \lambda t} C e^{\omega t} dt \\ &= Cm! (\operatorname{Re} \lambda - \omega)^{-m-1} \quad (m = 0, 1, 2, \dots). \end{aligned}$$

Set $Q(t) = AB^{-1}[I - T(t)]$ for $t \geq 0$. Then by (3.2),

$$\left[\frac{1}{\lambda}A(\lambda I + B)^{-1} \right]^n u = \int_0^\infty e^{-\lambda t} [Q(t)]^{*n} u dt \quad (u \in E, n = 2, 3, \dots),$$

where $*n$ indicates the n th convolution power. Consequently, we obtain

$$\begin{aligned} \left\{ \left[\frac{1}{\lambda}A(\lambda I + B)^{-1} \right]^n u \right\}^{(m)} &= \int_0^\infty (-t)^m e^{-\lambda t} [Q(t)]^{*n} u dt \\ &\quad (u \in E, m = 1, 2, \dots, n = 2, 3, \dots). \end{aligned}$$

But for $u \in E, t \geq 0, n = 2, 3, \dots$,

$$\begin{aligned} [Q(t)]^{*n} u &= \int_0^t \int_0^{t_{n-2}} \cdots \int_0^{t_1} Q(t - t_{n-2}) Q(t_{n-2} - t_{n-3}) \\ &\quad \cdots Q(t_1 - t_0) Q(t_0) u dt_0 dt_1 \cdots dt_{n-2}, \end{aligned}$$

so that,

$$\begin{aligned} \|[Q(t)]^{*n}\| &\leq \int_0^t \int_0^{t_{n-2}} \cdots \int_0^{t_1} C^n e^{\omega t} dt_0 \cdots dt_{n-2} \\ &= C^n e^{\omega t} \frac{t^{n-1}}{(n-1)!}, \end{aligned}$$

hence for $m = 1, 2, \dots, n = 2, 3, \dots$,

$$\begin{aligned} \left\| \left\{ \left[\frac{1}{\lambda} A(\lambda I + B)^{-1} \right]^n \right\}^{(m)} \right\| &\leq \int_0^\infty t^m e^{-\operatorname{Re} \lambda t} C^n e^{\omega t} \frac{t^{n-1}}{(n-1)!} dt \\ &= \frac{C^n}{(n-1)!} (m+n-1)! (\operatorname{Re} \lambda - \omega)^{-m-n}. \end{aligned}$$

When $n = 1$, making use of (3.2) again, we obtain easily that for $m = 0, 1, 2, \dots$,

$$\begin{aligned} \left\| \left[\frac{1}{\lambda} A(\lambda I + B)^{-1} \right]^{(m)} \right\| &\leq \int_0^\infty t^m e^{-\operatorname{Re} \lambda t} C e^{\omega t} dt \\ &= Cm! (\operatorname{Re} \lambda - \omega)^{-m-1}. \end{aligned}$$

Accordingly, we have

$$\begin{aligned} (3.5) \quad &\left\| \left\{ \sum_{n=0}^\infty \left[-\frac{1}{\lambda} A(\lambda I + B)^{-1} \right]^n \right\}^{(m)} \right\| \\ &\leq \sum_{n=1}^\infty \frac{C^n}{(n-1)!} (m+n-1)! (\operatorname{Re} \lambda - \omega)^{-m-n} \\ &= \frac{C}{(\operatorname{Re} \lambda - \omega)^{m+1}} \sum_{n=0}^\infty \left[\left(\frac{C}{\operatorname{Re} \lambda - \omega} \right)^n (n+1)(n+2) \cdots (m+n) \right] \\ &= \frac{C}{(\operatorname{Re} \lambda - \omega)^{m+1}} \left(\frac{d^m}{dx^m} \sum_{n=0}^\infty x^{m+n} \right) \Big|_{x=C/\operatorname{Re} \lambda - \omega} \\ &= Cm! (\operatorname{Re} \lambda - \omega - C)^{-(m+1)} \quad (m = 1, 2, \dots), \end{aligned}$$

$$\begin{aligned} (3.6) \quad &\left\| \sum_{n=0}^\infty \left[-\frac{1}{\lambda} A(\lambda I + B)^{-1} \right]^n \right\| \\ &\leq \frac{1}{1 - C(\operatorname{Re} \lambda - \omega)^{-1}} = \frac{\operatorname{Re} \lambda - \omega}{\operatorname{Re} \lambda - \omega - C} \quad (m = 0). \end{aligned}$$

We now apply the Leibniz formula to (3.3), obtaining

$$\begin{aligned} [\lambda \Delta(\lambda)^{-1}]^{(n)} &= \sum_{k=0}^n C_n^k [(\lambda I + B)^{-1}]^{(k)} \\ &\quad \cdot \left\{ \sum_{m=0}^\infty \left[-\frac{1}{\lambda} A(\lambda I + B)^{-1} \right]^m \right\}^{(n-k)} \quad (n = 0, 1, 2, \dots). \end{aligned}$$

It is thus easy to see, using (3.4), (3.5) and (3.6), that

$$\begin{aligned}
& \left\| [\lambda \Delta(\lambda)^{-1}]^{(n)} \right\| \\
& \leq \sum_{k=0}^{n-1} C_n^k C k! (\operatorname{Re} \lambda - \omega)^{-k-1} C(n-k)! (\operatorname{Re} \lambda - \omega - C)^{-n+k-1} \\
& \quad + C n! (\operatorname{Re} \lambda - \omega)^{-n-1} (\operatorname{Re} \lambda - \omega) (\operatorname{Re} \lambda - \omega - C)^{-1} \\
& = C^2 n! (\operatorname{Re} \lambda - \omega - C)^{-n-1} (\operatorname{Re} \lambda - \omega)^{-1} \\
& \quad \times \left\{ \sum_{k=0}^{n-1} \left(\frac{\operatorname{Re} \lambda - \omega - C}{\operatorname{Re} \lambda - \omega} \right)^k + \sum_{k=n}^{\infty} \left(\frac{\operatorname{Re} \lambda - \omega - C}{\operatorname{Re} \lambda - \omega} \right)^k \right\} \\
& = C n! (\operatorname{Re} \lambda - \omega - C)^{-n-1} \quad (\operatorname{Re} \lambda > \omega + C, n = 0, 1, 2, \dots).
\end{aligned}$$

Similarly, we can obtain the other two estimations:

$$\begin{aligned}
\| [B \Delta(\lambda)^{-1}]^{(n)} \| & \leq C n! (\operatorname{Re} \lambda - \omega - C)^{-n-1} \\
& \quad (\operatorname{Re} \lambda > \omega + C, n = 0, 1, 2, \dots)
\end{aligned}$$

and from

$$\begin{aligned}
\Delta(\lambda)^{-1} B u & = \left(I + \frac{1}{\lambda} B (\lambda I + B)^{-1} \overline{B^{-1} A} \right)^{-1} \frac{1}{\lambda} (\lambda I + B)^{-1} B u \\
& \quad (u \in D(B), \operatorname{Re} \lambda > \omega + C),
\end{aligned}$$

we obtain

$$\begin{aligned}
\| [\Delta(\lambda)^{-1} B u]^{(n)} \| & \leq C n! (\operatorname{Re} \lambda - \omega - C)^{-n-1} \\
& \quad (u \in D(B), \operatorname{Re} \lambda > \omega + C, n = 0, 1, 2, \dots).
\end{aligned}$$

According to Theorem 1, the proof is now complete.

On the other hand, we discuss the case where $D(A) \subset D(B)$. We have shown in the proof of the implication (ii) \Rightarrow (i) in Section 2 that the general case can, in a certain way, be reduced to this one. The case where $B \in L(E)$ has been studied in [12], and the conclusion is that the Cauchy problem for (1.1) is well posed (or equivalently, in view of [2, Th. 4.1(a)], strongly well posed) if and only if $-A$ is the generator of a strongly continuous cosine function. As an example of the case when $D(B) \supset D(A)$ and B is unbounded, we now investigate

Then $(iF)^* = iF$, thus it follows from [7, P. 41, Th. 10.8] that F is the generator of a strongly continuous unitary group. Denote this group by

$$\begin{pmatrix} T_1(t) & T_2(t) \\ T_3(t) & T_4(t) \end{pmatrix} \quad \text{for } -\infty < t < +\infty.$$

Observing that

$$R(\lambda, F) = \begin{pmatrix} (\lambda I + B)\Delta_1(\lambda)^{-1} & B\Delta_1(\lambda)^{-1} \\ B\Delta_1(\lambda)^{-1} & \lambda\Delta_1(\lambda)^{-1} \end{pmatrix} \quad \text{for } \operatorname{Re} \lambda > 0,$$

where $\Delta_1(\lambda) = \lambda^2 I + \lambda B + A_2$, we obtain that for $\operatorname{Re} \lambda > 0$,

$$(3.8) \quad \begin{cases} \lambda\Delta_1(\lambda)^{-1}u = \int_0^\infty e^{-\lambda t} T_4(t)u dt & (u \in E) \\ & \|T_4(t)\| \leq 1 \quad (t \geq 0), \\ B\Delta_1(\lambda)^{-1}u = \int_0^\infty e^{-\lambda t} [T_1(t) - T_4(t)]u dt & (u \in E), \\ & \|T_1(t) - T_4(t)\| \leq 2 \quad (t \geq 0). \end{cases}$$

Therefore

$$(3.9) \quad \begin{cases} A_1\Delta_1(\lambda)^{-1}u = A_1B^{-1}B\Delta_1(\lambda)^{-1}u \\ & = \int_0^\infty e^{-\lambda t} A_1B^{-1}[T_1(t) - T_4(t)]u dt & (u \in E), \\ \|A_1B^{-1}[T_1(t) - T_4(t)]\| \leq 2\|A_1B^{-1}\| & (t \geq 0); \end{cases}$$

$$(3.10) \quad \begin{aligned} \overline{\Delta_1(\lambda)^{-1}A_1}u &= B\Delta_1(\lambda)^{-1}\overline{B^{-1}A_1}u \\ &= \int_0^\infty e^{-\lambda t} [T_1(t) - T_4(t)]\overline{B^{-1}A_1}u dt & (u \in E), \\ \|[T_1(t) - T_4(t)]\overline{B^{-1}A_1}\| &\leq 2\|\overline{B^{-1}A_1}\| & (t \geq 0). \end{aligned}$$

From (3.8), (3.9), (3.10) and the plain fact that

$$\begin{aligned} \lambda\Delta(\lambda)^{-1} &= \lambda\Delta_1(\lambda)^{-1}[I + A_1\Delta_1(\lambda)^{-1}]^{-1} & (\operatorname{Re} \lambda > 0), \\ B\Delta(\lambda)^{-1} &= B\Delta_1(\lambda)^{-1}[I + A_1\Delta_1(\lambda)^{-1}] & (\operatorname{Re} \lambda > 0), \\ \Delta(\lambda)^{-1}Bu &= [I + \overline{\Delta_1(\lambda)^{-1}A_1}]^{-1}B\Delta_1(\lambda)^{-1}u & (u \in E, \operatorname{Re} \lambda > 0), \end{aligned}$$

we can obtain the three estimations in (iii) of Theorem 1 using arguments similar to those in the proof of the sufficiency of Theorem 2. Then, applying Theorem 1, we conclude that (3.7) is strongly well posed and therefore the propagators grow exponentially.

Acknowledgment. We are grateful to our advisor Professor F. L. Huang for his generous guidance.

REFERENCES

- [1] H. O. Fattorini, *Second Order Linear Differential Equations in Banach Spaces*, (1985), Elsevier Science Publishers B. V., Amsterdam.
- [2] —, *Extension and behavior at infinity of solutions of certain linear operational differential equations*, Pacific J. Math., **33** (1970), 583–615.
- [3] E. Hewitt, *Real and Abstract Analysis*, (1965), Springer-Verlag, Berlin.
- [4] E. Hille and R. S. Phillips, *Functional analysis and semigroups*, (1957), Amer. Math. Soc. Colloquium Pubs. vol. 31, Providence.
- [5] F. Neubrander, *Well-posedness of higher order abstract Cauchy problems*, Trans. Amer. Math. Soc., **295** (1986), 257–290.
- [6] —, *On the relation between the semigroup and its infinitesimal generator*, Proc. Amer. Math. Soc., **100** (1987), 104–107.
- [7] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, (1983), Springer-Verlag, New York and Berlin.
- [8] G. Da Prato and E. Giusti, *Una caratterizzazione dei generatori di funzioni coseno astratte*. Boll. Un. Mat. Italiana (3), **22** (1967), 357–362.
- [9] M. Sova, *Cosine operator functions*, Rozprawy Mat., **49** (1966), 1–47.
- [10] —, *Problème de Cauchy pour équations hyperboliques opérationnelles a coefficients constants non-bornés*, Ann. Scuola Norm. Sup. Pisa, **22** (1968), 67–100.
- [11] D. V. Widder, *The Laplace transform*, (1946), Princeton University Press, Princeton.
- [12] T. J. Xiao, *On the well-posedness of the Cauchy problem for a kind of complete second order linear differential equation*, Sichuan University Natural Sci. Edition (Chinese) **25**, No. 4 (1988), 421–428.

Received December 28, 1987 and in revised form, March 13, 1989.

YUNNAN TEACHERS UNIVERSITY
KUNMING, PEOPLE'S REPUBLIC OF CHINA

AND

KUNMING ENGINEERING INSTITUTE
KUNMING, PEOPLE'S REPUBLIC OF CHINA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Pacific Journal of Mathematics

Vol. 142, No. 1

January, 1990

Marco Andreatta, Mauro Beltrametti and Andrew Sommese , Generic properties of the adjunction mapping for singular surfaces and applications	1
Chen-Lian Chuang and Pjek-Hwee Lee , On regular subdirect products of simple Artinian rings	17
Fernando Giménez and Vicente Miquel Molina , Volume estimates for real hypersurfaces of a Kaehler manifold with strictly positive holomorphic sectional and antiholomorphic Ricci curvatures	23
Richard J. Griego and Andrzej Korzeniowski , Asymptotics for certain Wiener integrals associated with higher order differential operators	41
Abdeslam Mesnaoui , Unitary bordism of classifying spaces of quaternion groups	49
Abdeslam Mesnaoui , Unitary cobordism of classifying spaces of quaternion groups	69
Jesper M. Møller , On equivariant function spaces	103
Bassam Nassrallah , A q -analogue of Appell's F_1 function, its integral representation and transformations	121
Peter A Ohring , Solvability of invariant differential operators on metabelian groups	135
Athanase Papadopoulos and R. C. Penner , Enumerating pseudo-Anosov foliations	159
Ti-Jun Xiao and Liang Jin , On complete second order linear differential equations in Banach spaces	175
Carl Widland and Robert F. Lax , Weierstrass points on Gorenstein curves	197