

Pacific Journal of Mathematics

**A GEOMETRIC BOUND FOR MAXIMAL FUNCTIONS
ASSOCIATED TO CONVEX BODIES**

DETLEF MÜLLER

A GEOMETRIC BOUND FOR MAXIMAL FUNCTIONS ASSOCIATED TO CONVEX BODIES

DETLEF MÜLLER

For a convex symmetric body B in \mathbb{R}^n let M_B denote the centered maximal operator

$$M_B f(x) = \sup_{t>0} \frac{1}{\text{Vol } B} \int |f(x + ty)| dy$$

for $f \in L^1_{\text{loc}}(\mathbb{R}^n)$. We associate with B two linear invariants $\sigma(B)$ and $Q(B)$, and show that for $p > 1$ the norm of the operator M_B on $L^p(\mathbb{R}^n)$ is bounded by a constant which may depend on $p, \sigma(B)$ and $Q(B)$, but not explicitly on the dimension n . In particular, if B_q denotes the unit ball in \mathbb{R}^n with respect to the l^q -norm, we can prove that M_{B_q} has a bound on $L^p(\mathbb{R}^n)$ which is independent of n , provided that $1 \leq q < \infty$.

The behaviour of maximal functions associated to convex bodies has been studied by various authors during recent years. When B is the Euclidean ball, i.e. $B = B_2$, Stein [9] has shown that M_B is bounded on $L^p(\mathbb{R}^n)$ uniformly in n for every $p > 1$, and Bourgain [2, 3, 4] and Carbery [6] have shown that the analogue of this holds for any convex body B , provided $p > 3/2$. Moreover, by a result of Stein and Strömberg [11] it is known that the L^p operator norm $\|M_B\|_{p,p}$ of M_B grows at most linearly in the dimension n for any $p > 1$.

Since the general estimates for convex bodies in [2] do not imply that $\|M_B\|_{p,p}$ has a bound independent of n , if $p \leq 3/2$, it is well possible that for $p \leq 3/2$ one can only hope for estimates of $\|M_B\|_{p,p}$ which depend on additional geometric invariants associated with the body B . In this article, we shall show that one can in fact prove an estimate of this kind:

We associate with B the following two linear invariants $\sigma(B)$ and $Q(B)$: There exists a regular linear transformation S of \mathbb{R}^n , which is unique modulo orthogonal transformations, and a unique constant $L(B)$ such that $\text{Vol}_n S(B) = 1$ and

$$\int_{S(B)} |\langle x, \xi \rangle|^2 dx = L(B)^2$$

for all unit vectors $\xi \in \mathbb{R}^n$. Let $1/\sigma(B)$ be the minimum of all $(n-1)$ -dimensional volumes of all sections of $S(B)$ by hyperplanes, and $Q(B)$

the maximum of the $(n - 1)$ -dimensional volumes of all orthogonal projections of $S(B)$ onto hyperplanes (we note that $\sigma(B) \approx L(B)$). Then, for $p > 1$, the operator norm $\|M_B\|_{p,p}$ can be estimated by a constant depending only on p , $\sigma(B)$ and $Q(B)$.

This criterion suffices for example to prove the uniform boundedness in n of the maximal function M_{B_q} , where B_q denotes the unit ball with respect to the l^q -norm on \mathbb{R}^n , $1 \leq q < \infty$. This extends a result of Bourgain [4] who proved it for $q \in 2\mathbb{N}$ by making use of an “extra” decay of the Fourier transform of $\chi_{B_{2k}}$, $\chi_{B_{2k}}$ denoting the characteristic function of B_{2k} . However, this extra decay depends on some “smoothness” of B_q for $q \in 2\mathbb{N}$, which can easily be destroyed by cutting off a small piece of B_{2k} along an affine hyperplane, whereas our result is invariant under such operations.

Moreover, since one can show that $Q(B_\infty) = \sqrt{n}$, this might indicate that the norm of the “cubic” maximal operator M_{B_∞} associated with the unit cube of L^p is possibly growing with the dimension, if $p \leq 3/2$, and our results give some hints how one might try to prove this.

I would like to express my gratitude to the Mathematical Sciences Research Institute in Berkeley for the warm hospitality during my stay there by which this paper was completed, and especially to E. M. Stein for hints concerning multipliers of Laplace-transform type.

2. The main theorem. Let B be a convex symmetric body in \mathbb{R}^n . Arguing as in [2], we see that there exist a linear transformation $S \in GL(\mathbb{R}^n)$ and a constant $L(B) > 0$ such that

$$(1) \quad \text{Vol}_n S(B) = 1 \quad \text{and} \quad \int_{S(B)} |\langle x, \xi \rangle|^2 dx = L(B)^2$$

for all unit vectors $\xi \in S^{n-1} = \{\xi \in \mathbb{R}^n : |\xi|^2 = \sum_j |\xi_j|^2 = 1\}$. It is easy to see that $L(B)$ is determined uniquely by (1), and that S is unique up to multiplication by an orthogonal transformation from the left.

For $\xi \in S^{n-1}$, we define similarly as in [2]

$$(2) \quad \varphi(u) := \varphi_\xi(u) := \text{Vol}_{n-1}(\{x \in S(B) : \langle x, \xi \rangle = u\}), \quad u \in \mathbb{R}.$$

Moreover, let π_ξ denote the orthogonal projection of \mathbb{R}^n onto the hyperplane perpendicular to ξ . Then the constants

$$(3) \quad \begin{aligned} 1/\sigma(B) &:= \max\{\varphi_\xi(0) : \xi \in S^{n-1}\}, \\ Q(B) &:= \max\{\text{Vol}_{n-1}(\pi_\xi(S(B))) : \xi \in S^{n-1}\} \end{aligned}$$

are obviously linear invariants for B , i.e. $\sigma(U(B)) = \sigma(B)$ and $Q(U(B)) = Q(B)$ for all $U \in GL(\mathbb{R}^n)$.

Since also $\|M_B\|_{p,p}$ is a linear invariant for B , we therefore may and shall assume in the sequel (except for §3) that $S(B) = B$. Then, by [2], Lemma 1, there exist two universal constants $0 < a, A < \infty$, such that

$$(4) \quad \varphi(u) \leq A\varphi(0)e^{-a\varphi(0)|u|}, \quad u \in \mathbb{R}.$$

Moreover, there is a universal constant $a_1 > 0$, such that with $L = L(B)$

$$(5) \quad a_1^{-1} \leq L \cdot \varphi_\xi(0) \leq a_1, \quad \xi \in S^{n-1}.$$

This implies in particular $\sigma(B) \approx L(B)$.

THEOREM 1. *Let $p > 1$. Then for all $f \in L^p(\mathbb{R}^n)$*

$$(6) \quad \|M_B f\|_p \leq C(p, \sigma(B), Q(B)) \|f\|_p,$$

where the constant $C = C(p, \sigma, Q)$ ¹ is independent of n and grows with σ and Q .

Note that, for $p > 3/2$, C can even be chosen to be independent of σ and Q by [3] or [6].

Let us fix some notation. We denote by m the multiplier

$$(7) \quad m(\xi) = \hat{\chi}_B(\xi) = \int_{\mathbb{R}^n} \chi_B(x) e^{-2\pi i \langle \xi, x \rangle} dx$$

associated to χ_B . If $w \in L^\infty(\mathbb{R}^n)$ is any multiplier, we define the corresponding multiplier operator T_w as

$$(8) \quad T_w(f) = \mathcal{F}^{-1}(w\hat{f}),$$

\mathcal{F}^{-1} denoting the inverse Fourier transform.

For $\rho \in \mathbb{R}$ with $\rho > 1/2$ let us define the ρ th fractional derivative $(\xi \cdot \nabla)^\rho m$ of m as in [6] by

$$(9) \quad \begin{aligned} (\xi \cdot \nabla)^\rho m(\xi) &= \left(\frac{d}{dr} \right)^\rho \Big|_{r=1} m(r\xi) \\ &= \int (-2\pi i \langle x, \xi \rangle)^\rho K(x) e^{-2\pi i \langle x, \xi \rangle} dx, \end{aligned}$$

where $K = \chi_B$. Then, by the results of [6], especially Theorem 2 and Proposition (ii), our Theorem 1 will be an immediate consequence of

¹Here and in the sequel constants will frequently be denoted by C , with the understanding that they may be different from statement to statement.

PROPOSITION 1. *Let $1/2 < \rho < 1$. Then for all $f \in L^p(\mathbb{R}^n)$*

$$\|T_{(\xi, \nabla)^\rho m} f\|_p \leq C_\rho(p, \sigma(B), Q(B)) \|f\|_p$$

if $1 < p < \infty$, where the constant C_ρ is again independent of n .

This proposition is closely related to the question raised in [6], whether it is possible to find a bound for $T_{(\xi, \nabla)_m}$ which is independent of n .

The proof of Proposition 1 will be based on analytic interpolation. We define a family of operators $T_\alpha = T_{m_\alpha}$, $\alpha \in \mathbb{C}$, by

$$(10) \quad m_\alpha(\xi) = (1 + |\xi|)^{1-\alpha} [I^{-\alpha} m(r\xi)]|_{r=1}, \quad \xi \neq 0.$$

Here, $I^{-\alpha}$ denotes the α th fractional Riesz derivative with base point 2, that is

$$(11) \quad I^{-\alpha} f(r) = \frac{-1}{\Gamma(-\alpha)} \int_r^2 (s-r)^{-\alpha-1} f(s) ds, \quad \operatorname{Re} \alpha < 0,$$

if $f \in C^\infty([0, 2])$.

It is well known that $I^{-\alpha}$ can be extended analytically to the whole complex plane, and that $I^{-k} = (d/dr)^k$ is the usual k th derivative for $k = 0, 1, \dots$. Note that $I^{-\alpha}$ and $(d/dr)^\alpha$ as defined in (9) do not agree. However, we shall show later that the difference of these two is unimportant for our problem. We also define $T_\alpha^\varepsilon = T_{m_\alpha^\varepsilon}$ by

$$(12) \quad m_\alpha^\varepsilon(\xi) = (1 + |\xi|)^{-\varepsilon} m_\alpha(\xi), \quad \varepsilon > 0.$$

The proof of Proposition 1 will essentially be contained in the Lemmas 2 and 4 to follow, which deal with the two endpoint cases for the interpolation. Lemmas 1 and 3 are more of a technical nature.

LEMMA 1. *Let $0 \leq \operatorname{Re} \alpha < 1$, $k \in \mathbb{N}$. Then for $u > 1$*

$$\left| \int_0^u \frac{s^{-\alpha}}{(1+s/u)^k} e^{-2\pi i s} ds - \frac{e^{\frac{\pi}{2}\alpha i}}{i} \Gamma(1-\alpha) \right| \leq C_k e^{(\pi/2)|\operatorname{Im} \alpha|} u^{-\operatorname{Re} \alpha}.$$

The proof of Lemma 1 is an easy consequence of Cauchy's integral theorem and follows by changing the path of integration from the interval $[0, u]$ to $-i[0, u]$, connecting those two paths by quarter circles of radii u and ε , $\varepsilon \rightarrow 0$. We shall omit the technical details.

LEMMA 2. Fix $N > 0$ and $0 < \varepsilon < 1/2$. Then

- (i) $\|m_\alpha\|_\infty \leq C_N(\sigma(B), Q(B))e^{2\pi|\operatorname{Im}\alpha|}$, $0 \leq \operatorname{Re}\alpha < N$,
- (ii) $\|m_\alpha^\varepsilon\|_\infty \leq C_N(\sigma(B), Q(B))e^{2\pi|\operatorname{Im}\alpha|}$, $-\varepsilon \leq \operatorname{Re}\alpha \leq N$.

Proof. Assume $\operatorname{Re}\alpha \geq -\varepsilon$, and let $k = [\operatorname{Re}\alpha]$ be the integer part of $\operatorname{Re}\alpha$. Then it follows easily by partial integration from (10) that

$$(13) \quad m_\alpha(\xi) = \sum_{j=0}^k \frac{(-1)^{j+1}}{\Gamma(j+1-\alpha)} (1+|\xi|)^{1-\alpha} \left(\frac{d}{dr}\right)^j m(r\xi)|_{r=2} \\ + \frac{(-1)^k (1+|\xi|)^{1-\alpha}}{\Gamma(k+1-\alpha)} \int_1^2 (s-1)^{k-\alpha} \left(\frac{d}{ds}\right)^{k+1} m(s\xi) ds.$$

By (1), with $\varphi = \varphi_{\xi/|\xi|}$, we have

$$(14) \quad m(\xi) = \int_{-\infty}^\infty e^{-2\pi i|\xi|u} \varphi(u) du,$$

hence

$$(15) \quad \left(\frac{d}{dr}\right)^j m(r\xi)|_{r=2} = (-2\pi i|\xi|)^j \int_{-\infty}^\infty e^{-4\pi i|\xi|u} u^j \varphi(u) du.$$

By partial integration this implies

$$(16) \quad \left(\frac{d}{dr}\right)^j m(r\xi)|_{r=2} = \frac{1}{2}(-2\pi i|\xi|)^{j-1} \int_{-\infty}^\infty e^{-4\pi i|\xi|u} (u^j \varphi)'(u) du.$$

(2) and (15) imply for $0 \leq j \leq N$

$$\left| \left(\frac{d}{dr}\right)^j m(r\xi)|_{r=2} \right| \leq C_N |\xi|^j \int_0^\infty u^j \varphi(0) e^{-a\varphi(0)u} du \\ \leq C_N \varphi(0)^{-j} |\xi|^j \leq C_N \sigma(B)^j |\xi|^j.$$

Moreover, since $(u^j \varphi)'(u) = ju^{j-1} \varphi(u) + u^j \varphi'(u)$, and since $\varphi'(u)$ has constant sign for $u \geq 0$ resp. $u \leq 0$, (16) and (4) yield

$$\left| \left(\frac{d}{dr}\right)^j m(r\xi)|_{r=2} \right| \leq C_N \varphi(0)^{-(j-1)} |\xi|^{j-1} \leq C_N \sigma(B)^{j-1} |\xi|^{j-1}.$$

Together, we obtain

$$\left| \left(\frac{d}{dr}\right)^j m(r\xi)|_{r=2} \right| \leq C_N (\sigma(B)) |\xi|^j / (1+|\xi|),$$

at least for $j \geq 1$. However, for $j = 0$, (15) and (16) easily imply

$$|m(\xi)| \leq C(1 + \varphi(0)) / (1 + |\xi|) \leq C \cdot Q(B) / (1 + |\xi|).$$

So, together we get

$$\left| \left(\frac{d}{dr} \right)^j m(r\xi)|_{r=2} \right| \leq C_N(\sigma(B), Q(B)) \frac{|\xi|^j}{1+|\xi|}, \quad 0 \leq j < N.$$

This implies, for $j = 0, \dots, k$,

$$(17) \quad \left| \frac{(1+|\xi|)^{\alpha-1}}{\Gamma(j+1-\alpha)} \left(\frac{d}{dr} \right)^j m(r\xi)|_{r=2} \right| \leq C_N(\sigma, Q) e^{\pi|\operatorname{Im} \alpha|},$$

where we made use of the well known asymptotics [8, p. 79]

$$(18) \quad |\Gamma(x+iy)| \sim e^{-(\pi/2)|y|} |y|^{(x-1/2)} \cdot \sqrt{2\pi} \quad \text{as } |y| \rightarrow \infty.$$

So, it remains to estimate the integral term in (13), which, up to the sign, is given by

$$J(\xi) = \frac{(1+|\xi|)^{1-\alpha}}{\Gamma(k+1-\alpha)} (-2\pi i |\xi|)^{k+1} \int_{-\infty}^{\infty} F(|\xi|u) u^{k+1} \varphi(u) du,$$

where

$$F(t) = \int_1^2 (s-1)^{k-\alpha} e^{-2\pi i t s} ds.$$

The estimate of $J(\xi)$ requires more technique, but is essentially based again on (4), so that the rest of the proof of the lemma could be skipped for a first reading. We set

$$G(u) = \int_0^u t^{k+1} F(t) dt, \quad u \in \mathbb{R}.$$

Then

$$\begin{aligned} & \int_{-\infty}^{\infty} F(|\xi|u) u^{k+1} \varphi(u) du \\ &= |\xi|^{-k-2} \int_{-\infty}^{\infty} F(u) u^{k+1} \varphi(u/|\xi|) du \\ &= -|\xi|^{-k-3} \int_{-\infty}^{\infty} G(u) \varphi'(u/|\xi|) du, \end{aligned}$$

and hence

$$(19) \quad |J(\xi)| \leq C_N |\xi|^{-2} (1+|\xi|)^{1-\operatorname{Re} \alpha} \times \left| \frac{1}{\Gamma(k+1-\alpha)} \int_{-\infty}^{\infty} G(u) \varphi'(u/|\xi|) du \right|.$$

Now

$$(20) \quad \int_0^u t^{k+1} e^{-2\pi i t s} dt = \left(\frac{i}{2\pi}\right)^{k+1} \left\{ (-1)^{k+1} (k+1)! s^{-(k+2)} (e^{-2\pi i u s} - 1) + \sum_{j=0}^k \binom{k+1}{j} (-1)^j j! (-2\pi i)^{k-j} \times s^{-(j+1)} u^{k+1-j} e^{-2\pi i u s} \right\}.$$

Let

$$(21) \quad G_j(u) = u^{k+1-j} \int_1^2 (s-1)^{k-\alpha} s^{-(j+1)} e^{-2\pi i u s} ds, \quad j = 0, \dots, k+1,$$

and

$$(21)' \quad G_{k+2}(u) = G_{k+2} = \int_1^2 (s-1)^{k-\alpha} s^{-(k+2)} ds = \int_0^1 \frac{s^{k-\alpha}}{(s+1)^{k+2}} ds,$$

and define for $j = 0, \dots, k+2$

$$(22) \quad J_j(\xi) = \frac{(1+|\xi|)^{1-\operatorname{Re}\alpha}}{|\xi|^2} \left| \frac{1}{\Gamma(k+1-\alpha)} \int_{-\infty}^{\infty} G_j(u) \phi'(u/|\xi|) du \right|.$$

By (20), G is a linear combination of the G_j , and so it remains only to show that all functions J_j have an estimate of the desired type.

For $j = 0, \dots, k+1$,

$$G_j(u) = u^{\alpha-j} e^{-2\pi i u} \int_0^u \frac{s^{k-\alpha}}{(1+s/u)^{j+1}} e^{-2\pi i s} ds,$$

so Lemma 1 implies for $|u| > 1$

$$(23) \quad G_j(u) = \pm i e^{(\pi/2)(\alpha-k)i} \Gamma(k+1-\alpha) u^{\alpha-j} e^{-2\pi i u} + O(e^{(\pi/2)|\operatorname{Im}\alpha|} |u|^{k-j}).$$

Moreover, if $|u| \leq 1$, then

$$G_j(u) = u^{\alpha-j} \frac{e^{-2\pi i u}}{k+1-\alpha} \left\{ \frac{u^{k+1-\alpha}}{2^{j+1}} e^{-2\pi i u} - \int_0^u s^{k+1-\alpha} \frac{d}{ds} \left[\frac{e^{-2\pi i s}}{(1+s/u)^{j+1}} \right] ds \right\},$$

which easily implies

$$(23)' \quad |G_j(u)| \leq C_N \frac{|u|^{k+1-j}}{|k+1-\alpha|}, \quad |u| \leq 1.$$

(23) and (23)' imply

$$(24) \quad |J_j(\xi)| \leq -C_N \frac{(1+|\xi|)^{1-\operatorname{Re} \alpha}}{|\xi|^2} \\ \times \left\{ \frac{1}{|\Gamma(k+2-\alpha)|} \int_0^1 u^{k+1-j} \varphi'(u/|\xi|) du + e^{(\pi/2)|\operatorname{Im} \alpha|} \right. \\ \left. \times \int_1^\infty \left[u^{\operatorname{Re} \alpha - j} + \frac{u^{k-j}}{|\Gamma(k+1-\alpha)|} \right] \varphi'(u/|\xi|) du \right\}.$$

However, if $j \leq k+1$, then

$$(25) \quad \left| \int_0^1 u^{k+1-j} \varphi'(u/|\xi|) du \right| \leq - \int_0^1 \varphi'(u/|\xi|) du \leq 2\varphi(0)|\xi|,$$

and similarly one shows by (4) that

$$\left| \int_1^\infty u^{\operatorname{Re} \alpha - j} \varphi'(u/|\xi|) du \right| = -|\xi|^{1+\operatorname{Re} \alpha - j} \int_{1/|\xi|}^\infty u^{\operatorname{Re} \alpha - j} \varphi'(u) du \\ \leq |\xi|^{1+\operatorname{Re} \alpha - j} \left\{ |\xi|^{j-\operatorname{Re} \alpha} \varphi(1/|\xi|) + |\operatorname{Re} \alpha - j| \varphi(0) \int_{1/|\xi|}^1 u^{\operatorname{Re} \alpha - j - 1} du \right. \\ \left. + |\operatorname{Re} \alpha - j| \int_1^\infty u^{\operatorname{Re} \alpha - j} \varphi(u) du \right\} \\ \leq C_N(\sigma, Q) |\xi|^{1+\operatorname{Re} \alpha - j} (1 + |\xi|^{j-\operatorname{Re} \alpha});$$

hence

$$(26) \quad \left| \int_1^\infty u^{\operatorname{Re} \alpha - j} \varphi'(u/|\xi|) du \right| \leq \begin{cases} C_N(\sigma, Q) |\xi|^{1+\operatorname{Re} \alpha}, & j \leq k, \\ C_N(\sigma, Q) |\xi|, & j = k+1. \end{cases}$$

Of course $|\int_1^\infty u^{k-j} \varphi'(u/|\xi|) du|$ is even dominated by (26). (24), (25) and (26) imply, for $|\xi| \geq 1$,

$$(27) \quad |J_j(\xi)| \leq C_N(\sigma, Q) e^{2\pi|\operatorname{Im} \alpha|} (1 + |\xi|^{-\operatorname{Re} \alpha}), \quad j = 0, \dots, k+1.$$

Moreover, since obviously $|G_{k+2}| \leq C_N/|k+1-\alpha|$, we have

$$(27)' \quad |J_{k+2}(\xi)| \leq C_N |\xi|^{-1-\operatorname{Re} \alpha} \frac{-1}{|\Gamma(k+2-\alpha)|} \int_0^\infty \varphi'(u/|\xi|) du \\ \leq C_N(\sigma) e^{\pi|\operatorname{Im} \alpha|} |\xi|^{-\operatorname{Re} \alpha}, \quad |\xi| \geq 1.$$

The last two estimates imply the desired uniform estimates of $m_\alpha(\xi)$ and $m_\alpha^\varepsilon(\xi)$ for $|\xi| \geq 1$.

There remains the case $|\xi| < 1$, which is easy: By partial integration

$$J(\xi) = \frac{(1 + |\xi|)^{1-\alpha}}{\Gamma(k + 2 - \alpha)} \left\{ \left(\frac{d}{ds}\right)^{k+1} m(s\xi)|_{s=2} - \int_1^2 (s - 1)^{k+1-\alpha} \left(\frac{d}{ds}\right)^{k+2} m(s\xi) ds \right\}$$

which, together with (15) and (2), implies

$$|J(\xi)| \leq C_N(\sigma, Q)e^{\pi|\operatorname{Im}\alpha|}|\xi|^{k+1};$$

this settles the case $|\xi| < 1$. □

LEMMA 3. *For each unit vector $\eta \in S^{n-1}$ define a distribution $\mu_\eta = \partial\chi_B/\partial\eta = (\eta \cdot \nabla)\chi_B$. Then μ_η is even a bounded measure, and*

$$\|\mu_\eta\|_{M(\mathbb{R}^n)} = 2 \operatorname{Vol}_{n-1}(\pi_\eta(B)).$$

Proof. Let $\varphi \in C_0^\infty(\mathbb{R}^n)$ with $\|\varphi\|_\infty = 1$. After rotating coordinates, we may assume that η is the n th coordinate vector. Writing $\mathbb{R}^n = \mathbb{R}^{n-1} \times \mathbb{R}$ with coordinates (x, u) , we then have

$$\langle \mu_\eta, \varphi \rangle = - \int_B \frac{\partial \varphi}{\partial \eta} = - \int_{\pi_\eta(B)} \int_{B_x} \frac{\partial \varphi}{\partial u}(x, u) du dx,$$

where B_x is the interval $B_x = \{u \in \mathbb{R} : (x, u) \in B\}$, with endpoints say $a(x) \leq b(x)$, unless $B_x = \emptyset$. So

$$|\langle \mu_\eta, \varphi \rangle| = \left| \int_{\pi_\eta(B)} [\varphi(b(x)) - \varphi(a(x))] dx \right| \leq 2 \operatorname{Vol}_{n-1}(\pi_\eta(B));$$

hence $\|\mu_\eta\|_M \leq 2 \operatorname{Vol}_{n-1}(\pi_\eta(B))$. Moreover, choosing φ to be linear on each section B_x such that $\varphi(b(x)) = 1$ and $\varphi(a(x)) = -1$ immediately also gives $\|\mu_\eta\|_M \geq 2 \operatorname{Vol}_{n-1}(\pi_\eta(B))$. □

LEMMA 4. *Let $0 < \varepsilon < 1/2$. Then*

$$(28) \quad \|T_{-\varepsilon+i\nu}^\varepsilon f\|_p \leq C_\varepsilon(p, \sigma(B), Q(B))e^{(\pi/2)|\nu|} \|f\|_p, \quad f \in L^p(\mathbb{R}^n),$$

for every $1 < p < \infty$.

Proof. Let $\alpha = -\varepsilon + i\nu$. Since

$$m_\alpha^\varepsilon(\xi) = -\frac{1}{\Gamma(-\alpha)} \int_1^2 (s - 1)^{-\alpha-1} (1 + |\xi|)^{1-\varepsilon-\alpha} m(s\xi) ds,$$

it clearly suffices to prove that the multiplier operator corresponding to $(1 + |\xi|)^{1-\varepsilon-\alpha}m(s\xi)$ satisfies (28) uniformly for $1 \leq s \leq 2$.

Consider the multiplier $M_\nu(\xi) = (1 + |\xi|)^{-i\nu}$. This multiplier is of Laplace-transform type in the sense of [8, Ch. II, §4], since one easily checks that

$$(1 + \lambda)^{-i\nu} = \lambda \int_0^\infty a(t)e^{-\lambda t} dt, \quad \lambda \geq 0, \quad \text{where}$$

$$a(t) = \frac{1}{\Gamma(1 + i\nu)} \left[t^{i\nu} e^{-t} + \int_0^t s^{i\nu} e^{-s} ds \right].$$

Since $\|a\|_\infty \leq Ce^{(\pi/2)|\nu|}$, the general theory of heat-diffusion semi-groups [8] implies for $1 < p < \infty$

$$(29) \quad \|T_{M_\nu} f\|_p \leq C_p e^{(\pi/2)|\nu|} \|f\|_p, \quad f \in L^p(\mathbb{R}^n),$$

where C_p is a constant depending only on p .

Since $(1 + |\xi|)^{1-\varepsilon-\alpha}m(s\xi) = (1 + |\xi|)^{-i\nu}(1 + |\xi|)m(s\xi)$, and since $\|T_{m(s\cdot)}\|_{p,p} = \|T_m\|_{p,p} \leq |B| = 1$ for all p , (29) reduces the proof of (28) finally to estimating the multiplier operator corresponding to

$$(30) \quad m_0(\xi) = -2\pi|\xi|m(\xi).$$

Define measures μ_j by $\mu_j = \partial\chi_B/\partial x_j$, $j = 1, \dots, n$. Since

$$m_0(\xi) = \sum_{j=1}^n \left(-i \frac{\xi_j}{|\xi|} \right) (-2\pi i \xi_j m(\xi)),$$

we have

$$(31) \quad T_{m_0} f = \sum_{j=1}^n R_j(\mu_j * f),$$

where R_j denotes the j th Riesz transform. By a result of Stein [10] (see also [7]), it is known that

$$(32) \quad \left\| \left(\sum_j |R_j f|^2 \right)^{1/2} \right\|_p \leq A_p \|f\|_p, \quad 1 < p < \infty,$$

where A_p is independent of n . Using a simple duality argument, (31) and (32) imply

$$(33) \quad \|T_{m_0} f\|_p \leq A_{p'} \left\| \left(\sum_j |\mu_k * f|^2 \right)^{1/2} \right\|_p, \quad 1 < p < \infty,$$

where $1/p + 1/p' = 1$. Let $g(f)^2(x) = \sum_j |\mu_j * f(x)|^2$. We want to estimate the L^p -operator norm of the sublinear operator g .

If $p = 2$, we obtain from (17)

$$(34) \quad \|g(f)\|_2 = \|T_{m_0}f\|_2 \leq \|m_0\|_\infty \|f\|_2 \leq C(\sigma, Q)\|f\|_2.$$

For $p = \infty$, we observe that

$$(35) \quad |g(f)(x)| = |(\nabla\chi_B) * f(x)| = \sup_{\eta \in S^{n-1}} |\mu_\eta * f(x)|,$$

where μ_η is defined as in Lemma 4. This in combination with Lemma 4 implies

$$(36) \quad \|g(f)\|_\infty \leq \sup_\eta \|\mu_\eta\|_M \|f\|_\infty = 2Q(B)\|f\|_\infty.$$

Interpolation between (34) and (36) yields

$$\|g(f)\|_p \leq C(p, \sigma, Q)\|f\|_p, \quad 2 \leq p \leq \infty;$$

hence, by (33), also

$$(37) \quad \|T_{m_0}f\|_p \leq C(p, \sigma, Q)\|f\|_p,$$

at least for $2 \leq p < \infty$, but by passing to the adjoint operator $T_{m_0}^*$, we get (37) also for $1 < p < 2$. This concludes the proof of Lemma 4. \square

Proof of Proposition 1. Let $\rho = 1 - \varepsilon \in]1/2, 1[$. From Lemma 2 (ii) and (13) it follows easily that the family $\{T_\alpha^\varepsilon\}$ in an admissible family (in the sense of [12, Ch. V]) on every strip $-\varepsilon \leq \text{Re } \alpha \leq N$, $N > 0$. Thus, choosing N sufficiently large and interpolating the estimates in Lemma 2 and Lemma 4 between $\text{Re } \alpha = -\varepsilon$ and $\text{Re } \alpha = N$, we obtain

$$(38) \quad \|T_{1-\varepsilon}^\varepsilon f\|_p \leq C_\varepsilon(p, \sigma(B), Q(B))\|f\|_p$$

for any $1 < p \leq 2$, hence, by duality, for any $1 < p < \infty$. But,

$$(39) \quad m_{1-\varepsilon}^\varepsilon(\xi) = [I^{-\rho} m(r\xi)]|_{r=1} \\ = -\frac{1}{\Gamma(\varepsilon)} m(\xi) + \frac{1}{\Gamma(\varepsilon)} \int_1^2 (s-1)^{-\rho} \frac{dm(s\xi)}{ds} ds.$$

Moreover, $(\xi \cdot \nabla)^\alpha m(\xi)$ is given by [5, p. 51]

$$(\xi \cdot \nabla)^\alpha m(\xi) = \frac{-1}{\Gamma(-\alpha)} \int_1^\infty (s-1)^{-\alpha-1} m(s\xi) ds \quad \text{if } -1 < \alpha < 0.$$

By partial integration, we see that

$$(\xi \cdot \nabla)^\alpha m(\xi) = \frac{1}{\Gamma(1-\alpha)} \int_1^\infty (s-1)^{-\alpha} \frac{dm(s\xi)}{ds} ds$$

for $0 < \alpha < 1$. A comparison with (39) shows that

$$\begin{aligned} (\xi \cdot \nabla)^\rho m(\xi) &= m_{1-\varepsilon}^\varepsilon(\xi) + \frac{1}{\Gamma(\varepsilon)} m(\xi) + \frac{1}{\Gamma(\varepsilon)} \int_2^\infty (s-1)^{-\rho} \frac{dm(s\xi)}{ds} ds \\ &= m_{1-\varepsilon}^\varepsilon(\xi) - \frac{1}{\Gamma(-\rho)} \int_2^\infty (s-1)^{-\rho-1} m(s\xi) ds. \end{aligned}$$

Since $\int_2^\infty (s-1)^{-\rho-1} ds < \infty$, this together with (38) implies

$$\|T_{(\xi \cdot \nabla)^\rho} f\|_p \leq C_\rho(p, \sigma, Q) \|f\|_p. \quad \square$$

3. Examples: The l^q -unit balls. In the sequel, let $1 \leq q \leq \infty$ be fixed, and let

$$B_q = B_q^n = \{x \in \mathbb{R}^n : |x|_q \leq 1\}$$

be the unit ball with respect to the l^q -norm $|x|_q = (\sum |x_j|^q)^{1/q}$ (resp. $|x|_\infty = \max |x_j|$, if $q = \infty$).

Let $\kappa(n) = \kappa_q(n)$ denote the volume of B_q^n . A straight-forward calculation, using induction on n , easily yields ($q < \infty$)

$$(40) \quad \kappa_q(n) = 2\Gamma\left(\frac{1}{q} + 1\right) \left[\frac{2}{q} \cdot \Gamma\left(\frac{1}{q}\right)\right]^{n-1} / \Gamma\left(\frac{n}{q} + 1\right).$$

Choose $m = m_q(n) > 0$ so, that the body $\tilde{B}_q = mB_q$ has volume 1. (40) implies $m \sim n^{1/q}$ up to a constant a_q (see [4]). Of course, if $q = \infty$, we have $\kappa_\infty(n) = 2^n$, and $m = 1/2$. Let us determine the constant L mentioned in (5):

Because of the symmetry properties of \tilde{B}_q , we have for any $\xi \in S^{n-1}$

$$\int_{\tilde{B}_q} \langle \xi, x \rangle^2 dx = \sum_j \int_{\tilde{B}_q} \xi_j^2 x_j^2 dx = \left(\sum_j \xi_j^2 \right) \int_{\tilde{B}_q} x_n^2 dx = \int_{\tilde{B}_q} x_n^2 dx,$$

and so we may choose $S(B_q)$ to be \tilde{B}_q , and obtain for $L = L(B_q)$

$$\begin{aligned} L^2 &= \int_{\tilde{B}_q} x_n^2 dx = 2 \int_0^m x_n^2 (m^q - |x_n|^q)^{(n-1)/q} \kappa_q(n-1) dx_n \\ &= 2m^{n+2} \kappa_q(n-1) \mathbf{B}\left(\frac{3}{q}, \frac{n-1}{q} + 1\right), \end{aligned}$$

where \mathbf{B} denotes the Beta-function.

Since $m^n \kappa_q(n) = 1$, this yields

$$L^2 = 2m^2 \frac{\kappa_q(n-1)}{\kappa_q(n)} \mathbf{B}\left(\frac{3}{q}, \frac{n-1}{q} + 1\right) \sim A_q^2$$

by Stirling's formula, and so by (5)

$$(41) \quad a_1^{-1} A_q \lesssim \sigma(B_q^n) \lesssim a_1 A_q,$$

at least for $q < \infty$. However, for $q = \infty$ clearly $L^2 = 1/2$, hence $\sigma(B_\infty^n) \approx (2\sqrt{3})^{-1}$.

In order to estimate $Q(\tilde{B}_q^n)$, we adapt an idea from [4]: Let $\tau : [0, \infty[\rightarrow [0, 1]$ be a smooth function satisfying the conditions ($q < \infty$)

$$(42) \quad \tau = 1 \quad \text{on } [0, m^q],$$

$$(42)' \quad \tau = 0 \quad \text{on } [m^q + 1, \infty[,$$

$$(42)'' \quad -2 \leq \tau' \leq 0,$$

and set $K(x) = \tau(\sum |x_j|^q)$, $x \in \mathbb{R}^n$. Note that by (42) $\chi_{\tilde{B}_q} \leq K$, and by (42)' $(m^q + 1)^{1/q} B_q \subset (1 + c/n)\tilde{B}_q = \tilde{\tilde{B}}_q$, hence $\|K\|_{L^1} \leq C$. Moreover, we have

$$(43) \quad \text{Vol}_{n-1}(\pi_\xi(\tilde{B}_q)) = \frac{1}{2} \left\| \frac{\partial K}{\partial \xi} \right\|_{L^1} \quad \text{for all } \xi \in S^{n-1}.$$

This is in fact true if $B = \tilde{B}_q$ is any convex body and K any function which is 1 on B , non-increasing with growing distance from B , and such that $\partial K/\partial \xi$ is integrable: We may assume without restriction that $\xi = e_n$. Then, adapting the notations from the proof of Lemma 4,

$$\begin{aligned} \int_{B_x} \left| \frac{\partial K}{\partial \xi}(x, t) \right| dt &= \int_{b(x)}^\infty \left| \frac{\partial K}{\partial t}(x, t) \right| dt + \int_{-\infty}^{a(x)} \left| \frac{\partial K}{\partial t}(x, t) \right| dt \\ &= K(x, b(x)) + K(x, a(x)) = 2; \end{aligned}$$

hence

$$\left\| \frac{\partial K}{\partial \xi} \right\|_{L^1} = 2 \text{Vol}_{n-1}(\pi_\xi(B)).$$

In order to estimate $\|\partial K/\partial \xi\|_{L^1}$, observe that

$$\partial K/\partial \xi = q\tau' \left(\sum |x_j|^q \right) \cdot \sum_j \xi_j \text{sgn}(x_j) |x_j|^{q-1},$$

and hence

$$\begin{aligned} \|\partial K/\partial \xi\|_{L^1} &\leq 2q \int_{\tilde{\tilde{B}}_q} \left| \sum_j \xi_j \text{sgn}(x_j) |x_j|^{q-1} \right| dx \\ &= \frac{2q}{2^n} \sum_{\varepsilon_j = \pm 1} \int_{\tilde{\tilde{B}}_q} \left| \sum_j \varepsilon_j \xi_j \text{sgn}(x_j) |x_j|^{q-1} \right| dx. \end{aligned}$$

However, Khintchine's inequality

$$2^{-n} \sum_{\varepsilon_i = \pm 1} \left| \sum_{j=1}^n \varepsilon_j \alpha_j \right| \leq C \left(\sum_j \alpha_j^2 \right)^{1/2}, \quad \alpha_j \in \mathbb{R},$$

implies

$$\begin{aligned} \|\partial K / \partial \xi\|_{L^1} &\leq Cq \int_{\tilde{B}_q} \left[\sum_j \xi_j^2 |x_j|^{2(q-1)} \right]^{1/2} dx \\ &\leq C'q \cdot \left[\int_{\tilde{B}_q} \sum_j \xi_j^2 |x_j|^{2(q-1)} dx \right]^{1/2} \end{aligned}$$

by Hölder's inequality, since $\text{Vol}_n(\tilde{B}_q) \leq C$. Because of the symmetry of \tilde{B}_q , this yields

$$\|\partial K / \partial \xi\|_{L^1} \leq C''q \cdot \left[\int_{\tilde{B}_q} |x_n|^{2(q-1)} dx \right]^{1/2},$$

and hence, because of (4), (5), (41) and (43),

$$(44) \quad Q(B_q^n) \leq Cq, \quad 1 \leq q < \infty,$$

independently of n . So Theorem 1 implies

COROLLARY 1. *Let $1 \leq q < \infty$. Then for all $f \in L^p(\mathbb{R}^n)$*

$$\|M_{B_q^n} f\|_p \leq C(p, q) \|f\|_p, \quad 1 < p \leq \infty,$$

independently of n .

What can be said about the case $q = \infty$?

In this case, an easy geometric consideration shows that for any $\xi \in S^{n-1}$ (see also [1], pp. 41, 45)

$$\text{Vol}_{n-1}(\pi_\xi(\tilde{B}_\infty)) = \sum_F \text{Vol}_{N-1}(F) \cdot \langle \xi, n(F) \rangle,$$

where summation is over all faces F of the cube \tilde{B}_∞ whose outward normal $n(F)$ satisfies $\langle \xi, n(F) \rangle \geq 0$. So, if we choose $\xi = n^{-1/2}(1, 1, \dots, 1)$, we get

$$\text{Vol}_{n-1}(\pi_\xi(\tilde{B}_\infty)) = \sum_j \xi_j = \sqrt{n}.$$

The same argument easily shows that $\text{Vol}_{n-1}(\pi_\eta(\tilde{B}_\infty)) \leq \sqrt{n}$ for any $\eta \in S^{n-1}$, and we get

$$(45) \quad Q(B_\infty^n) = \sqrt{n}.$$

So, our criterion gives a bound for $\|M_{B_\infty}\|_{p,p}$ which grows with n .

Let us conclude with a direct consequence of our results, which appears a bit surprising at the first glance (we do, however, not claim originality for this result). Let $\Sigma(\tilde{B}_q^n)$ denote the surface area of \tilde{B}_q^n .

COROLLARY 2. *If $1 \leq q < \infty$, then $c\sqrt{n} \leq \Sigma(\tilde{B}_q^n) \leq Cq\sqrt{n}$, whereas $\Sigma(\tilde{B}_\infty^n) = 2n$.*

Proof. By Cauchy's surface formula [1, p. 48]

$$\begin{aligned} \Sigma(\tilde{B}_q^n) &= \frac{1}{\kappa_2(n-1)} \int_{S^{n-1}} \text{Vol}_{n-1}(\pi_\xi(\tilde{B}_q^n)) \, d\xi \\ &\leq \frac{\Sigma(B_2^n)}{\kappa_2(n-1)} Q(\tilde{B}_q^n) = \frac{n\kappa_2(n)}{\kappa_2(n-1)} Q(\tilde{B}_q^n); \end{aligned}$$

hence, by (40), (44), for $q < \infty$

$$\Sigma(\tilde{B}_q^n) \leq C_q\sqrt{n}.$$

Moreover, it is well known [1, p. 104] that the Euclidean ball has minimal surface area among all convex bodies of given volume, and $\Sigma(\tilde{B}_2^n) = m^{n-1}\Sigma(B_2^n) = m^{n-1}n \cdot \kappa_2(n) = n/m \sim c \cdot \sqrt{n}$ by (40), where $m = m_2(n)$. So we also obtain

$$\Sigma(\tilde{B}_q^n) \geq \Sigma(\tilde{B}_2^n) \sim c \cdot \sqrt{n}. \quad \square$$

REFERENCES

- [1] T. Bonnesen and W. Fenchel, *Theorie der konvexen Körper*, Chelsea Publishing Company (1948).
- [2] J. Bourgain, *On high dimensional maximal functions associated to convex bodies*, Amer. J. Math., **108** (1986), 1467–1476.
- [3] —, *On the L^p -bounds for maximal functions associated to convex bodies in \mathbb{R}^n* , Israel J. Math., **54** (1986), 257–265.
- [4] —, *On dimension free maximal inequalities for convex symmetric bodies in \mathbb{R}^n* , Lecture Notes in Math. **1267**, Springer (1987), 168–176.

- [5] A. Carbery, *Radial Fourier multipliers and associated maximal functions*, Recent Progress in Fourier Analysis, North-Holland Math. Studies, **111** (1985), 49–56.
- [6] ———, *An almost-orthogonality principle with applications to maximal functions associated to convex bodies*, Bull. Amer. Math. Soc., **14** No 2 (1986), 269–273.
- [7] J. Duoandikoetxea and J. L. Rubio de Francia, *Estimations indépendantes de la dimension pour les transformées de Riesz*, C. R. Acad. Sc. Paris, t. 300, Série I, no 7 (1985), 193–196.
- [8] E. M. Stein, *Topics in harmonic analysis*, Annals of Math. Studies #63 (1970), Princeton Univ. Press.
- [9] ———, *The development of square functions in the work of A. Zygmund*, Bull. Amer. Math. Soc., **7** (1982), 359–376.
- [10] ———, *Three variations on the theme of maximal functions*, Recent Progress in Fourier Analysis, North-Holland Math. Studies **111** (1985), 49–56.
- [11] E. M. Stein and J. O. Strömberg, *Behavior of maximal functions in \mathbb{R}^n for large n* , Ark. f. Mat., **21** (1983), 259–269.
- [12] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton Univ. Press (1971).

Received April 29, 1988.

UNIVERSITÄT BIELEFELD
4800 BIELEFELD 1, FRG

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (6 Vols., 12 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 6 volumes per year. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1990 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 142, No. 2

February, 1990

Christopher J. Bishop , Bounded functions in the little Bloch space	209
Lutz Bungart , Piecewise smooth approximations to q -plurisubharmonic functions	227
Donald John Charles Bures and Hong Sheng Yin , Outer conjugacy of shifts on the hyperfinite II_1 -factor	245
A. D. Raza Choudary , On the resultant hypersurface	259
Luis A. Cordero and Robert Wolak , Examples of foliations with foliated geometric structures	265
Peter J. Holden , Extension theorems for functions of vanishing mean oscillation	277
Detlef Müller , A geometric bound for maximal functions associated to convex bodies	297
John R. Schulenberger , Time-harmonic solutions of some dissipative problems for Maxwell's equations in a three-dimensional half space	313
Mark Andrew Smith and Barry Turett , Normal structure in Bochner L^p -spaces	347
Jun-ichi Tanaka , Blaschke cocycles and generators	357
R. Z. Yeh , Hyperholomorphic functions and higher order partial differential equations in the plane	379