

Pacific Journal of Mathematics

**FIXED POINTS FOR ORIENTATION PRESERVING
HOMEOMORPHISMS OF THE PLANE WHICH INTERCHANGE
TWO POINTS**

MORTON BROWN

FIXED POINTS FOR ORIENTATION PRESERVING HOMEOMORPHISMS OF THE PLANE WHICH INTERCHANGE TWO POINTS

MORTON BROWN

Suppose h is an orientation preserving homeomorphism of the plane which interchanges two points p and q . If A is an arc from p to q , then h has a fixed point in one of the bounded complementary domains of $A \cup h(A)$.

1. Introduction. Brouwer's Lemma [2], one version of which is that each orientation preserving homeomorphism of the plane with a periodic point has a fixed point, has had much attention in the last few years. It has played a central role in some work of Fathi [7], Franks [8, 9], Pelikan and Slaminka [11], Slaminka [12] and the author [3, 4].

An interesting special case is when the periodic point has period two. Indeed, this case is at the heart of Fathi's argument in [7], and his proof of Brouwer's lemma requires a separate proof of this case. The purpose of this note is to show that this result follows from a particularly simple and elegant application of the notion of index of a homeomorphism along an arc. Furthermore, we get constructive information about the location of the fixed point. Our proof both simplifies and strengthens a result of Galliaro and Kottman [10].

In a final section we illustrate some techniques which can be used to locate fixed points more precisely.

2. The index. Let f, g be maps of the interval $[0, 1]$ into the plane such that $f(t)$ is distinct from $g(t)$ for each t in $[0, 1]$. Then $\text{index}(f, g)$ is defined to be the total winding number of the vector $g(t) - f(t)$ as t runs from 0 to 1. For example, in Figure 1 this vector makes a total of 1 and $1/2$ turns in the clockwise (i.e., negative) direction, so the index is $-(1 + 1/2)$. The reader who wishes a more precise definition of index and its properties should consult [5] and [6].

If f and f' are two maps of $[0, 1]$ into the plane such that $f(1) = f'(0)$ then we denote by $f * f'$ the map of $[0, 1]$ into the plane which is $f(2t)$ on $0 \leq t \leq 1/2$, and $f'(2t - 1)$ on $1/2 \leq t \leq 1$. Clearly, if $\text{index}(f, g)$ and $\text{index}(f', g')$ are defined then $\text{index}(f * f', g * g')$ is well defined and equal to $\text{index}(f, g) + \text{index}(f', g')$.




FIGURE 1

3. **LEMMA.** *Let h be an orientation preserving homeomorphism of the plane and let p, q be distinct points such that $h(p) = q$ and $h(q) = p$. Let f be a path from p to q whose image contains no fixed points of h . Then there exists an integer k such that*

$$\text{index}(f, hf) = \text{index}(hf, hhf) = 1/2 + k.$$

Proof. h interchanges p and q , so the vectors $hf(0) - f(0) = q - p$ and $hf(1) - f(1) = p - q$ point in opposite directions, i.e., $\text{index}(f, hf) = 1/2 + k$. Since h is orientation preserving, there is an isotopy g_s , $0 \leq s \leq 1$, connecting the identity to h . Then $\text{index}(g_s f, g_s hf)$ varies continuously from $\text{index}(f, hf)$ to $\text{index}(hf, hhf)$. On the other hand, for each s , the vectors $g_s hf(0) - g_s f(0) = g_s(q) - g_s(p)$ and $g_s hf(1) - g_s f(1) = g_s(p) - g_s(q)$ point in opposite directions, so, by continuity, $\text{index}(g_s f, g_s hf)$ is constant as s varies from 0 to 1. Hence

$$\text{index}(g_1 f, g_1 hf) = \text{index}(hf, hhf) = 1/2 + k.$$

4. **THEOREM.** *Let h, p, q, f be as in the Lemma. Then,*

$$\text{index}(f * hf, hf * hhf)$$

*is an odd integer, and h has a fixed point in a bounded complementary domain of the image of the loop $f * hf$.*

Proof. By the additivity of the index, $\text{index}(f * hf, hf * hhf) = \text{index}(f, hf) + \text{index}(hf, hhf) = 2(1/2 + k)$, which is an odd integer.

Since the image of $f * hf$ is locally connected, the set X consisting of the image of $f * hf$ and the union of its bounded complementary domains is a locally connected continuum ([13], p. 112–113). Since X does not separate the plane it is an absolute retract ([1]), and hence contractible. If h were fixed point free in each of the bounded complementary domains of the image of the loop $f * hf$, then the loop could be shrunk to a point within X , and $\text{index}(f * hf, hf * hhf)$ would be zero, a contradiction.

5. Examples. Let h, p, q, f be as in the Theorem.




FIGURE 2

In Figure 2 the curve f (more precisely the image of f) is a simple arc from p to q and intersects hf only at the endpoints which h interchanges. Then $\text{index}(f * hf, hf * hhf) = 1$, and there is a fixed point h inside the simple closed curve $f * hf$.




FIGURE 3

In Figure 3, f is again a simple arc and f intersects hf in one other point v . The index (f, hf) is seen by inspection to be $-1/2$ or $+1/2$ depending on whether $h(u) = v$ or $h(w) = v$, respectively. Hence, by the Lemma, $\text{index}(f * hf, hf * hhf) = -1$ or $+1$, respectively. Suppose $h(u) = v$. We wish to calculate the index of h “around” each of the domains, E, F ; that is, the index of positively oriented simple closed curves lying in and surrounding the fixed point sets of h in E, F respectively. Then

$$\begin{aligned} \text{index}(f * hf, hf * hhf) &= (\text{index of } h \text{ around } F) \\ &\quad - (\text{index of } h \text{ around } E) = 1. \end{aligned}$$

(Note that $f * hf$ goes around E in the negative direction.) It is not difficult to construct a homeomorphism g of the plane which equals h when restricted to $K = \text{image } f$, and such that g has index 1 around F , and 0 around E . I claim that this ensures that h has the same indicial values around E , F , respectively. The justification for the claim lies in the following Theorem.

THEOREM. *Let h, g be orientation preserving homeomorphisms of the plane and let K be an arc that K contains no fixed points of h , and $h = g$ on K . Let $X = K \cup h(K) = K \cup g(K)$. Then the maps*

$$\frac{x - h(x)}{\|x - h(x)\|} \quad \text{and} \quad \frac{x - g(x)}{\|x - g(x)\|}$$

are homotopic maps of X into the unit circle.

Proof. By a variation of Alexanders Isotopy Theorem ([3], page 38) h is isotopic to g relative to K . Let p_t denote the isotopy ($p_0 = h$, $p_1 = g$, and for each t , $p_t = h$ on K). Since p_t has no fixed points on K it has no fixed points on $p(K)$, so the required homotopy is $(x - p_t(x))/\|x - p_t(x)\|$.

A consequence of this result is that g and h have the same index around each complementary domain of $K \cup h(K)$.




FIGURE 4

In Figure 4, the calculation of the index (f, hf) depends again on the location of $f^{-1}(v)$. Let us suppose it is u , so that $\text{index}(f, hf) = 3/2$ and $\text{index}(f * hf, hf * hhf) = 3$. Notice that $f * hf$ winds twice positively around F and once positively around E , so that

$$(\text{index of } h \text{ around } E) + 2(\text{index of } h \text{ around } F) = 3.$$

With a bit more work than the previous case one can construct a homeomorphism g which equals h on $K = \text{image } f$ and which has index 1 around each of E and F . Thus, by the Theorem above, the same is true for h , and h has a fixed point in each of the domains E and F .

REFERENCES

- [1] Beverly Brechner and Morton Brown, *Mapping cylinder neighborhoods in the plane*, Proc. Amer. Math. Soc., **84** (1982), 433–436.
- [2] L. E. J. Brouwer, *Beweiss des ebenen Translationssatzes*, Math. Ann., **72** (1912), 37–54.
- [3] Morton Brown, *A new proof of Brouwer's lemma on translation arcs*, Houston J. Math., **10** (1984), 35–41.
- [4] —, *Homeomorphisms of two-dimensional manifolds*, Houston J. Math., **11** (1985), 455–469.
- [5] M. Brown and W. D. Neumann, *Proof of the Poincare Birkhoff fixed point theorem*, Michigan Math. J., **24** (1977), 21–31.
- [6] A. Dold, *Fixed point index and fixed point theorem for Euclidean neighborhood retracts*, Topology, **4** (1965), 1–8.
- [7] Albert Fathi, *An orbit closing proof of Brouwer's lemma on translation arcs*, L'Enseignement Mathematique, **33** (1987), 315–322.
- [8] John Franks, *Recurrence and fixed points of surface homeomorphisms*, to appear, Ergodic Theory Dynamical Systems.
- [9] —, *A generalization of the Poincare-Birkhoff theorem*, preprint.
- [10] E. Galliaro and C. Kottman, *Fixed points for orientation preserving homeomorphisms which interchange two points*, Pacific J. Math., **59** (1975), 27–32.
- [11] S. Pelikan and E. Slaminka, *A bound for the fixed point index of area preserving homeomorphisms of two manifolds*, Ergodic Theory and Dynamical Systems, **7** part 3 (1987), 463–479.
- [12] E. Slaminka, *A Brouwer translation theorem for free homeomorphisms*, Trans. Amer. Math. Soc., **306**, No. 1 (1988), 227–291.
- [13] G. T. Whyburn, *Analytic Topology*, American Math. Soc. Colloquium Pub., **28** (1942).

Received June 6, 1988 and, in revised form July 29, 1988.

UNIVERSITY OF MICHIGAN
ANN ARBOR, MI 48109

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Walter Bergweiler , <i>On the fix-points of composite functions</i>	1
Aldo Biancofiore, Maria Lucia Fania and Antonio Lanteri , <i>Polarized surfaces with hyperelliptic sections</i>	9
Ciprian Borcea , <i>Deforming varieties of k-planes of projective complete intersections</i>	25
Morton Brown , <i>Fixed points for orientation preserving homeomorphisms of the plane which interchange two points</i>	37
Hao Zhi Chuan , <i>Note on the inequality of the arithmetic and geometric means</i>	43
Paul Henry Edelman and Dennis E. White , <i>Codes, transforms and the spectrum of the symmetric group</i>	47
Pei-Kee Lin , <i>The isometries of $H^\infty(E)$</i>	69
James J. Moloney , <i>Residue class domains of the ring of convergent sequences and of $C^\infty([0, 1], \mathbf{R})$</i>	79
Zhong-Jin Ruan , <i>The structure of pure completely bounded and completely positive multilinear operators</i>	155
Wolfgang Ruess and William H. Summers , <i>Weakly almost periodic semigroups of operators</i>	175
Gideon Schwarz , <i>A pretender to the title "canonical Moebius strip"</i>	195
Ryszard Szwarc , <i>Banach algebras associated with spherical representations of the free group</i>	201