

# Pacific Journal of Mathematics

**AN ESTIMATE OF THE VOLUME OF A COMPACT SET IN  
TERMS OF ITS INTEGRAL OF MEAN CURVATURE**

YOUNG DO CHAI

# AN ESTIMATE OF THE VOLUME OF A COMPACT SET IN TERMS OF ITS INTEGRAL OF MEAN CURVATURE

Y. D. CHAI

A geometric inequality for a compact set in euclidean 3 space is obtained. The inequality involves volume and integral of mean curvature. Also some property of the compact set is studied. The method of outer parallel bodies is used in the proof.

**I. Introduction.** For a planar compact set  $K$  with area  $A$  and perimeter  $L$  the classical isoperimetric inequality states:

$$L^2 - 4\pi A \geq 0.$$

Equality in above inequality holds for regular disk. For proof of the inequality above see Guggenheimer [10].

The volume and surface area of the compact set  $W$  in euclidean 3 space,  $R^3$ , will be represented by the functionals  $V(W)$  and  $S(W)$ . The functional  $M(W)$  is the integral of the mean curvature and  $|M|(W)$  is the integral of absolute mean curvature [3].

Geometric inequality involving integral of absolute mean curvature  $|M|(W)$  and surface area  $S(W)$  in euclidean 3 space has been founded by Russia mathematician I. A. Danelich.

He found the following facts:

(1)  $|M|(W_2) \leq |M|(W_1)$  if  $W_1$  and  $W_2$  are compact sets and  $W_2$  is a convex set contained in  $W_1$  [4].

(2) If  $W$  is a compact set with bounded integral of absolute mean curvature, then

$$S(W) \leq 2/\pi^2 |M|^2 \quad [5].$$

The gist both of Danelich's result (2) and the classical geometric inequality  $36\pi[V(W)^2] \leq S(W)^3$ , [12], involving volume  $V(W)$  and surface area  $S(W)$  is contained in the following inequality:

$$(3) V(W) \leq \sqrt{2}/[3\pi^3\sqrt{\pi}] |M|^3(W).$$

In this paper we will derive a new upper bound for  $V(W)$  with compact set  $W$  having somewhat restricted conditions. It will be a sharper inequality than (3) in the previous paragraph and equality will hold for regular balls in  $R^3$ . It should be noted that Danelich's inequality

never decides when equality holds. We also improve Danelich's first result (1). In proving our results we will actually establish a bound for the volume of the outer parallel bodies of a given body.

**II. Geometric inequality for a compact set and its properties.** If the motion is determined by the position of the moving frame  $(Q, e_1, e_2, e_3)$ , the kinematic density has the form

$$dK = dP \wedge d\sigma \wedge d\pi$$

where  $dP$  is the volume element of  $R^3$  at the origin  $Q$  of the moving frame,  $d\pi$  is the area element of the unit sphere corresponding to the end point of  $e_3$ , and  $d\sigma$  is the element of rotation about  $e_3$ . For a compact set  $W$  parallel body of  $W$  at the distance  $r$  is defined to be set

$$W_r = \{x \in R^3 \mid |x - y| \leq r, y \in W\}.$$

Now we state *the kinematic fundamental formula* in  $R^3$  without proof which is the work of Blaschke.

**PROPOSITION 2.1.** *Let  $D_0$  and  $D_1$  be two domains of  $R^3$  bounded respectively by the surface  $\Sigma_0$  and  $\Sigma_1$  which we assume to be of class  $C^2$ . Let  $V_i, \chi_i$  be the volume and the euler characteristic of  $D_i$  and let  $F_i, M_i$  be the area and the integral of the mean curvature of  $\Sigma_i$  ( $i = 0, 1$ ) respectively. Suppose  $D_0$  is fixed and  $D_1$  is moving and let  $dK$  be the kinematic density for  $D_1$ . If  $\chi(D_0 \cap D_1)$  denotes the euler characteristic of the intersection  $D_0 \cap D_1$ , then*

$$\int_{D_0 \cap D_1 \neq \emptyset} \chi(D_0 \cap D_1) dk = 8\pi^2(V_0\chi_1 + V_1\chi_0) + 2\pi(F_0, M_1 + F_1M_0).$$

*Proof.* See [1].

Federer considered *sets with positive reach* in his article Curvature Measure [6]. *The reach of subset  $A$  of  $R^3$  is the largest number  $\varepsilon$  such that if point  $x$  is in  $R^3$  and the distance,  $\delta(x)$ , from  $x$  to  $A$  is smaller than  $\varepsilon$ , then  $A$  contains the unique point  $\zeta(x)$ , nearest to  $x$ .* It can be easily checked that compact set in  $R^3$  with  $C^2$  surface as its boundary has positive reach. Assuming that reach of  $A$ ,  $\text{reach}(A)$ , is positive, *the Steiner's formula* is established in the following form; for each topological ball  $A$  with  $C^2$ -boundary in  $R^3$  and for  $r, 0 \leq r < \text{reach}(A)$ , volume of the parallel body  $A_r$  of  $A$ ,  $A_r = \{x \in R^3 \mid |x - a| \leq r, a \in A\}$ , is given by a polynomial of degree 3 in  $r$ :

$$V(A_r) = V(A) + S(A) \cdot r + M(A) \cdot r^2 + (4/3)\pi \cdot r^3$$

where  $V$  is volume,  $M$  is integral of mean curvature.

In fact the above equality follows from Proposition 2.1 and the fact that  $A \cap D_r$  is contractible for all the positions of regular ball  $D_r$  of radius  $r$  which is shown in [6]. This suggests how we get geometric inequality involving volume and integral of mean curvature for special kind of compact sets by the usage of the estimate of volume of parallel bodies and the well known Minkowski inequality [11]. So we define a concept which generalizes the definition of compact convex sets in  $R^3$ .

**DEFINITION 2.2.** A topological 3-ball  $W$  in  $R^3$  with  $C^2$ -surface as its boundary is called an  $M(t)$ -compact set if each point in its boundary has at most one negative principal normal curvature with respect to the inward normal vector which is less than  $-1/t$  where  $t >$  diameter of  $W$ .

**REMARK 2.3.** (1) Every  $M(s)$ -compact set is an  $M(t)$ -compact set if  $s \geq t$ .

(2) Every compact convex set with  $C^2$ -surface is an  $M(\infty)$ -compact set.

(3) Every topological 3-ball with 2-convex surface, [14], is an  $M(\infty)$ -compact set.

(4) Every topological ball with next-to-convex surface, [9], is an  $M(\infty)$ -compact set.

**LEMMA 2.4.** *Let  $W$  be an  $M(t)$ -compact set and  $G$  be the set of all motions in  $R^3$ . Then for any  $g \in G$  and  $r < t - \text{diam}(W)$ ,*

$$\chi(W \cap gD_r) \geq 1 \quad \text{if } W \cap gD_r \text{ is nonempty}$$

where  $D_r$  is a solid regular ball with radius  $r$  and  $gD_r$  is the image of  $D_r$  by the motion  $g$ .

*Proof.* We will prove that  $\beta_i(W \cap gD_r) = 0$  for  $i \geq 1$  where  $\beta_i$  is the  $i$ th Betti number. For then  $\chi(W \cap gD_r) = \beta_0(W \cap gD_r) \geq 1$  if  $W \cap gD_r$  is nonempty. But we will show even stronger facts that  $H_i(W \cap gD_r) = 0$  for  $i \geq 2$ .

Consider the Mayer-Vietoris homology sequence with  $W$  and  $gD_r$  for any  $g \in G$  and  $r < t - \text{diam}(W)$  where  $W \cap gD_r$  is nonempty. Then we have the following exact homology sequence:

$$\begin{aligned} \rightarrow H_{i+1}(W) \oplus H_{i+1}(gD_r) &\rightarrow H_{i+1}(W \cup gD_r) \rightarrow H_i(W \cap gD_r) \\ \rightarrow H_i(W) \oplus H_i(gD_r) &\rightarrow \end{aligned}$$

Since  $H_{i+1}(W) = H_{i+1}(gD_r) = H_i(W) = H_i(gD_r) = 0$  for  $i \geq 2$ ,  $H_i(W \cap gD_r) \cong H_{i+1}(W \cup gD_r) = 0$  for  $i \geq 2$ .

Now suppose  $\chi(W \cap gD_r) < 1$  for some  $g \in G$  and  $r < t - \text{diam}(W)$ . Then  $r > 0$  and  $\beta_1(W \cap gD_r) > 0$  since  $\beta_i(W \cap gD_r) = 0$  for  $i \geq 2$ . So  $H_1(W \cap gD_r) \neq 0$ . Now we consider the following Mayer-Vietoris exact homology sequence with  $W$  and  $gD_r$ :

$$\begin{aligned} &\rightarrow H_2(W) \oplus H_2(gD_r) \rightarrow H_2(W \cup gD_r) \\ &\rightarrow H_1(W \cap gD_r) \rightarrow H_1(W) \oplus H_1(gD_r). \end{aligned}$$

Since

$$\begin{aligned} H_2(W) = H_2(gD_r) = H_1(W) = H_1(gD_r) = 0, \\ H_2(W \cup gD_r) \cong H_1(W \cap gD_r). \end{aligned}$$

Therefore

$$H_2(W \cup gD_r) \neq 0 \quad \text{and so} \quad H^2(W \cup gD_r) \neq 0.$$

By the Alexander duality theorem,

$$H^2(W \cup gD_r) \cong H_0^*(R^3 - W \cup gD_r).$$

So  $R^3 - (W \cup gD_r)$  has at least two open components. In fact  $R^3 - (W \cup gD_r)$  contains at least one bounded component whose boundary is the union of the subset of the boundary of  $gD_r$  and subset of the boundary of  $W$ . Let  $A$  be such a bounded component and  $g$  be the center of  $gD_r$ .

Consider the smallest sphere  $S_g$  with its center at  $g$  which circumscribes  $\bar{A} \cup gD_r$  where  $\bar{A}$  is the closure of  $A$ . Then the radius of  $S_g$  is greater than  $r$  and less than  $r + \text{diam}(W)$ . Let  $q$  be a point in  $S_g \cap (\bar{A} \cup gD_r)$ . Then  $q$  must be a point in the boundary of  $W$  since  $S_g$  never intersects the boundary of  $gD_r$ .

This means that all the principal normal curvatures at  $q$  with respect to inward normal vector are less than  $-1/[r - \text{diam}(W)]$ . But it is a contradiction because  $t$  is greater than  $r + \text{diam}(W)$  and  $W$  is an  $M(t)$ -compact set.

Now we use Proposition 2.1 and Lemma 2.4 to estimate the volume of  $W_r = \{x \in R^3 \mid |x - y| \leq r, y \in W\}$  in terms of the volume  $V(W)$  of  $W$ , area  $S(W)$  of the boundary of  $W$  and the integral of mean curvature  $M(W)$  defined by  $\int_{\text{boundary of } W} [(k_1 + k_2)/2] dw$ , where  $k_1$  and  $k_2$  are functions of principal normal curvatures on the boundary of  $W$ .

LEMMA 2.5. Let  $W$  be an  $M(t)$ -compact set in  $R^3$ .

Then  $V(W_r) \leq V(W) + S(W) \cdot r + M(W) \cdot r^2 + (4/3)\pi \cdot r^3$  for  $r, 0 \leq r < t - \text{diam}(W)$ .

*Proof.* Let  $W$  be fixed and  $D_r$  be moving by the motion  $g$ . So  $W \cap gD_r$  is the intersection of  $W$  and  $D_r$  at position  $g$ .

After putting  $\chi(W \cap gD_r) = 0$  if  $W \cap gD_r$  is empty if we apply Lemma 2.4, then we have for  $r, 0 \leq r < t - \text{diam}(W)$ ,

$$(1) \quad \int_{g \in G} \chi(W \cap gD_r) dk = \int_{W \cap gD_r \neq \emptyset} \chi(W \cap gD_r) dk \geq \int_{W \cap gD_r \neq \emptyset} dp \wedge d\sigma \wedge d\pi.$$

On the other hand,  $\chi(D_r) = \chi(W) = 1, V(D_r) = (4/3)\pi \cdot r^3$  and  $M(D_r) = 4\pi r$ . So by Proposition 2.1 we have

$$(2) \quad \int_{W \cap gD_r \neq \emptyset} \chi(W \cap gD_r) dk = 8\pi^2[V(W) + (4\pi/3)r^3 + S(W) \cdot r + M(W) \cdot r^2].$$

By (1) and (2), we have for  $r, 0 \leq r < t - \text{diam}(W)$

$$V(W_r) \leq V(W) + S(W) \cdot r + M(W) \cdot r^2 + (4\pi/3)r^3.$$

COROLLARY 2.6. If  $W$  is an  $M(\infty)$ -compact set, then

$$V(W_r) \leq V(W) + S(W) \cdot r + M(W) \cdot r^2 + (4\pi/3) \cdot r^3 \text{ for } r,$$

$0 \leq r < \infty$  and “=” holds if  $W$  is a compact convex subset in  $R^3$ .

*Proof.* Since  $W$  is a compact set  $\text{diam}(W)$  is finite. Now our corollary follows from Lemma 2.5. If  $W$  is a compact convex subset in  $R^3$ , (1) holds with equality since  $\chi(W \cap gD_r) = 1$  if  $W \cap gD_r$  is nonempty. So this fact and (2) prove the second statement.

One of our goals in this paper is to estimate the volume of  $M(\infty)$ -compact sets in terms of the integral of the mean curvature of the boundary of the set. For this purpose we may need the following inequality which is an estimate of the volume of the sum of sets from below.

**PROPOSITION 2.7.** *Let  $A$  and  $B$  be nonempty measurable sets in  $R^3$ . Then*

$$[V(A + B)]^{1/3} \geq [V(A)]^{1/3} + [V(B)]^{1/3}$$

where  $A + B = \{a + b \mid a \in A, b \in B\}$ .

*Proof.* See [11].

Note that in the proposition above if  $B$  is a regular ball centered at the origin of radius  $r$ , then  $A + B = A_r$ .

**THEOREM 2.8.** *Let  $W$  be an  $M(\infty)$ -compact set in  $R^3$ . Then*

$$V(W) \leq (1/48\pi^2)[M(W)]^3.$$

*Proof.* By Proposition 2.7 we have for  $r, 0 \leq r < \infty$

$$(3) \quad V(W_r) \geq V(W) + [3 \cdot (4\pi/3)^{1/3} \cdot (V(W))^{2/3}]r + [3(4\pi/3)^{2/3}(V(W))^{1/3}]r^2 + (4\pi/3)r^3.$$

On the other hand, if  $W$  is an  $M(\infty)$ -compact set, by Corollary 2.6, we have

$$(4) \quad V(W_r) \leq V(W) + S(W)r + M(W)r^2 + (4\pi/3)r^3 \quad \text{for } r, 0 \leq r < \infty.$$

From (3) and (4) we have

$$(5) \quad [3(4\pi/3)^{1/3}(V(W))^{2/3}]r + [3(4\pi/3)^{2/3}(V(W))^{1/3}]r^2 \leq S(W)r + M(W)r^2 \quad \text{for } r, 0 \leq r < \infty.$$

In (5) if we send  $r$  to  $\infty$ , then we have

$$(6) \quad [3(4\pi/3)^{2/3}(V(W))^{1/3}] \leq M(W).$$

Equivalently we have

$$V(W) \leq (1/48\pi^2)[M(W)]^3.$$

Inequality (6) is sharper than Danelich's result [4] for the  $M(\infty)$ -compact sets and equality holds when  $W$  is a regular ball.

**COROLLARY 2.9.** *If  $W$  is an  $M(\infty)$ -compact set and  $K$  is a convex subset of  $W$ , then*

$$M(K) \leq M(W).$$

*Proof.* It follows from the fact that  $V(K_r) \leq V(W_r)$  and Corollary 2.6 for  $r \rightarrow \infty$ .

## REFERENCES

- [1] W. Blaschke, *Vorlesungen über integralgeometrie*, 3rd ed. Berlin, 1955.
- [2] S. S. Chern, *Studies in global geometry and analysis*, M.A.A., **4** (1967), 16–56.
- [3] I. A. Danelich, *Surfaces of bounded absolute mean integral curvature and their plane sections*, *Sibirsk. Matem. Zh.*, **3** (1963), 519–538.
- [4] —, *An integral representation of the absolute mean integral curvature of a polyhedral surface and consequences of this representation*, *Sibirsk. Matem. Zh.*, **7** (1966), 954–959.
- [5] —, *An estimate for the area of a surface of bounded absolute mean integral curvature in terms of its absolute mean integral curvature and the sum of the boundary curve lengths*, *Sibirsk. Matem. Zh.*, **7** (1966), 1199–1203.
- [6] H. Federer, *Curvature measures*, *Trans. Amer. Math. Soc.*, **93** (1959), 418–491.
- [7] J. H. G. Fu, *Tubular neighborhoods in euclidean spaces*, *Duke Math. J.*, **52** (1985), 1025–1046.
- [8] M. J. Greenberg and J. R. Harper, *Algebraic Topology*, a first course, *Mathematics Lecture Note Series*, vol. 58, 1981.
- [9] M. Gromov, *Hyperbolic manifolds, groups and actions*, *Ann. of Math. Studies*, **97** (1981), 183–213.
- [10] H. W. Guggenheimer, *Differential Geometry*, Mc Graw-Hill, New York, 1963.
- [11] H. L. Montgomery, *The isoperimetric inequality and sum of sets*, preprint, (1979).
- [12] G. Polya and G. Szego, *Isoperimetric inequalities in mathematical physics*, *Ann. of Math.*, **27**, Princeton Univ. Press, Princeton, N.J., (1951).
- [13] L. A. Santalo, *Integral Geometry and Geometric Probability*, Addison-Wesley Publishing Co., 1976.
- [14] J. P. Sha, *p-convex Riemmanian manifolds*, *Invent. Math.*, **83** (1986), 437–447.

Received September 13, 1988. This work was partially supported by the Basic Science Research Institute Program, Ministry of Education, Korea, 1988-89.

SUNG KYUN KWAN UNIVERSITY  
300 CHUN CHUN DONG  
SUWON, 440-746 KOREA





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093

R. FINN  
Stanford University  
Stanford, CA 94305

HERMANN FLASCHKA  
University of Arizona  
Tucson, AZ 85721

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305

C. C. MOORE  
University of California  
Berkeley, CA 94720

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH  
(1906-1982)

B. H. NEUMANN

F. WOLF  
(1904-1989)

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1980 *Mathematics Subject Classification (1985 Revision)* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly. Regular subscription rate: \$190.00 a year (12 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1990 by Pacific Journal of Mathematics

<b>George E. Andrews and David M. Jackson</b> , An algebraically derived $q$ -analogue of a character sum associated with a class of semiregular permutations .....	207
<b>Fabio Bardelli and Andrea Del Centina</b> , The moduli space of genus four double covers of elliptic curves is rational .....	219
<b>Young Do Chai</b> , An estimate of the volume of a compact set in terms of its integral of mean curvature .....	229
<b>Salvador Comalada</b> , Elliptic curves with trivial conductor over quadratic fields .....	237
<b>Kahar El-Hussein</b> , Résolubilité semi-globale des opérateurs différentiels invariants sur les groupes de déplacements .....	259
<b>David M. Goldschmidt</b> , Classical link invariants and the Burau representation .....	277
<b>Liliana Janicka</b> , Radon-Nikodým problem for the variation of a vector measure .....	293
<b>Wacław Marzantowicz</b> , An almost classification of compact Lie groups with Borsuk-Ulam properties .....	299
<b>Akira Ohbuchi</b> , On the projective normality of some varieties of degree 5 ..	313
<b>Ken'ichi Ohshika</b> , Minimal measured laminations in geometric 3-manifolds .....	327
<b>Hal Leslie Smith</b> , A discrete Lyapunov function for a class of linear differential equations .....	345
<b>John Samuel Spielberg</b> , Diagonal states on $O_2$ .....	361
<b>Thomas Vogel</b> , A note on the sessile drop .....	383
<b>Gerold Wagner</b> , On means of distances on the surface of a sphere (lower bounds) .....	389