

Pacific Journal of Mathematics

DIAGONAL STATES ON O_2

JOHN SAMUEL SPIELBERG

DIAGONAL STATES ON O_2

JOHN S. SPIELBERG

In this paper we investigate certain states on the Cuntz algebra O_2 , and the von Neumann algebras obtained from their GNS representations.

Introduction. In this paper we investigate certain states on the Cuntz algebra O_2 , and the von Neumann algebras obtained from their GNS representations. The problem we begin with is that of finding different types of factor states on O_2 which extend the trace on Choi's algebra, viewed as a subalgebra of O_2 . The existence in general of such factor state extensions was established independently by Longo and Popa (see [1]).

The construction of specific examples, and classifying them as to type, has been done by several mathematicians. These examples arise by composing the expectation of O_2 onto the CAR algebra with a factor state on the CAR algebra. Work of Evans and Lance showed that by starting with the trace on the CAR algebra, a type $\text{III}_{1/2}$ factor state extension is obtained (see [1]). In [9], pure state extensions are constructed by a combinatorial argument. In [2], it is shown that uncountably many inequivalent pure state extensions are so obtained, and that they arise from certain pure states on the CAR algebra. Moreover, it is shown that if the Powers III_λ states on the CAR algebra are extended to O_2 , they result in factor states of type III_λ if $\lambda^{n+1} + \lambda^n = 1$, some n in \mathbb{Z}_+ , or III_1 , if $\log \lambda$ and $\log(\lambda + 1)$ are algebraically independent. In [11], a different collection of product states on the CAR algebra is shown to give rise to factor state extensions of type III_λ for all $0 \leq \lambda \leq 1$.

The techniques in [2] and [11] rely on the quasi-invariance, under the shift automorphism of [6], of an appropriate state or weight on the stabilized CAR algebra. In §1 of the present paper it is shown that arbitrary (infinite) Krieger factors can be obtained from factor state extensions on O_2 . The technique is the opposite of the above: namely, we use weights all of whose translates by powers of the shift are disjoint.

The remainder of the paper is a more detailed study of the states on O_2 which factor through product states on the diagonal of the CAR algebra. In §2 we use results of [11] to give a characterization of quasi-equivalence of product states on the CAR algebra somewhat different from that of [10] and [3]. We use these results in §3 to classify the corresponding states on O_2 up to quasi-equivalence. In §4 we apply these results to give a purely dynamical proof of the result of [2] on the existence of pure state extensions. In §5 we briefly indicate how the results of [11], computing invariants for the factors obtained from certain diagonal product states on O_2 , can be generalized to all such states.

Finally, we wish to thank Masamichi Takesaki for making the crucial suggestion which led to the results of §1.

Preliminaries. We recall some definitions and results from [11]. For $n \geq 0$, let $A_n = \bigotimes_{j=-n}^{\infty} M_2^{(j)}$, where $M_2^{(j)} = M_2(\mathbb{C})$, for all j . Let $i_n: A_n \rightarrow A_{n+1}$ be defined by $i_n(x) = e_{11}^{(-n-1)} \otimes x$, where for $b \in M_2$ we write $b^{(j)}$ for the same element viewed as an element of $M_2^{(j)}$. Let $A = \overline{\bigcup_n A_n}$. For $k \leq l$, and $b_k, b_{k+1}, \dots, b_l \in M_2(\mathbb{C})$, we write $\bigotimes_{j=k}^l b_j^{(j)}$ for the element

$$\left(\bigotimes_{j=-n}^{k-1} e_{11}^{(j)} \right) \otimes \left(\bigotimes_{j=k}^l b_j^{(j)} \right) \otimes \left(\bigotimes_{j=l+1}^{\infty} 1^{(j)} \right)$$

in A_n for suitably large n . Let $B_n = \bigotimes_{j=-n}^n M_2^{(j)}$, $B = \bigcup_n B_n$. Define $\alpha \in \text{Aut}(A)$ by letting $\alpha(\bigotimes_j x_j^{(j)}) = \bigotimes_j x_{j-1}^{(j)}$, for $x = \bigotimes_j x_j^{(j)}$ in B , and extending to A by linearity and continuity. Let $e_n = 1^{(-n)}$. Then as in [6], we have $A \times_{\alpha} \mathbb{Z} \cong \mathcal{K} \otimes O_2$, and $e_0(A \times_{\alpha} \mathbb{Z})e_0 \cong O_2$. Note also that A_0 is the CAR algebra, and $A \cong \mathcal{K} \otimes A_0$. We will use the notation

$$(i_0, j_0) \otimes \cdots \otimes (i_n, j_n)$$

for the element $\bigotimes_{k=0}^n e_{i_k, j_k}^{(k)}$ in A_0 . By a *weight sequence* we mean a sequence $\{t_j\}_{j \geq j_0}$ in $[0, 1]$. Given $\{t_j\}_{j \geq j_0}$, define $\Lambda_j \in M_2(\mathbb{C})$ for $j \in \mathbb{Z}$ by

$$\Lambda_j = \begin{cases} \text{diag}(t_j, 1 - t_j), & j \geq j_0, \\ 1, & j < j_0. \end{cases}$$

Define $f_n \in (A_n)_+^*$ by $f_n(\bigotimes_j x_j^{(j)}) = \prod_{j \geq -n} \text{Tr}(\Lambda_j x_j)$ for $\bigotimes_j x_j^{(j)}$ in $A_n \cap B$, where Tr is the non-normalized trace on $M_2(\mathbb{C})$. We define a weight f on A by $f(x) = \sup f_n(e_n x e_n)$, $x \in A_+$. (The fact that f is additive on A_+ follows from [11], Lemma 1.4.) Then f

is densely defined and lower semi-continuous. If $t_j \in (0, 1)$ for each j , then f is faithful. Let \tilde{f} be the canonical extension of f to a weight on $A \times_\alpha \mathbb{Z}$, and $\tilde{f}_0 = \tilde{f}|_{(e_0(A \times_\alpha \mathbb{Z})e_0)}$. Then the restriction of \tilde{f}_0 to Choi's algebra within O_2 is tracial, and hence is a multiple of the unique tracial state on Choi's algebra. The functionals on O_2 so constructed all factor through the conditional expectation $F_0: O_2 \rightarrow A_0$ of [6]. In fact, as noted in [2], these functionals factor further, through the conditional expectation E_0 of A_0 onto its diagonal ($= \overline{\text{span}}\{\bigotimes_j x_j^{(j)} \in B \cap A_0: x_j = e_{11} \text{ or } e_{22} \text{ for each } j\}$). It is shown in [2] that any state on O_2 which factors through the diagonal of A_0 extends the trace on Choi's algebra. We will make use of this fact. We remark that it also follows from our results mentioned above, as the pure states on the diagonal arise from weight sequences for which $t_j = 0$ or 1 for each j . We will refer to weights on a C^* -algebra as disjoint, quasi-equivalent, or unitarily equivalent, if their GNS representations are so related. Given a weight f on a C^* -algebra A , we will let π_f denote the representation of A given by the GNS construction applied to f , we will let H_f denote the Hilbert space of this representation, and $\eta_f: A \rightarrow H_f$ the canonical map of A onto a dense subspace of H_f . If X is a topological space we will let $C(X)$ denote the space of continuous complex-valued functions on X . If A is another topological space, we will let $C(X, A)$ denote the space of continuous functions from X into A . We will let $C_c(X, A)$ denote the space of functions in $C(X, A)$ having compact support. If T is a closed linear operator on a Hilbert space, we will let $\text{polar}(T)$ denote the partial isometry in the polar decomposition of T having the same kernel as T : $T = \text{polar}(T) \cdot |T|$.

1. Krieger factor states on O_2 .

LEMMA 1.1. *Let M be an infinite Krieger factor. Then there is a state f_0 on A_0 such that $\pi_{f_0}(A_0)''$ is isomorphic to M , and such that, letting $f = \text{Tr} \otimes f_0$ on A , we have that $f \circ \alpha^k$ is disjoint from f for all $k \neq 0$.*

Proof. Choose a sequence of integers $0 = n_0 < n_1 < n_2 < \dots$, with $\lim_{k \rightarrow \infty} (n_{k+1} - n_k) = \infty$. Let

$$\begin{aligned} I_1 &= \{n_{2k}: k = 0, 1, 2, \dots\}, \\ I_0 &= \{n_{2k-1}: k = 1, 2, 3, \dots\}, \\ I_2 &= \mathbb{N} \setminus (I_0 \cup I_1). \end{aligned}$$

For $j \in I_1 \cup I_2$ let μ_j be the measure on $\{1, 2\}$ given by

$$\mu_j = \begin{cases} \delta_1, & j \in I_1, \\ \delta_2, & j \in I_2. \end{cases}$$

By the diagonal of M_2 we mean $\text{span}\{e_{11}, e_{22}\}$. This is isomorphic to the space of continuous functions on the two-point space $\{1, 2\}$, where we may let e_{ii} represent the characteristic function of the clopen set $\{i\}$, for $i = 1, 2$. Let μ' be a measure on $\prod_{j \in I_0} \{1, 2\}^{(j)}$ such that $\pi_{\mu' \circ E_0}(\otimes_{j \in I_0} M_2^{(j)})''$ is isomorphic to M ([12]), where we let measures on a space X also denote functionals on $C(X)$. Let μ be the measure on $\prod_{j=0}^\infty \{1, 2\}^{(j)}$ given by $\mu = (\prod_{j \in I_1 \cup I_2} \mu_j) \times \mu'$. Let $f_0 = \mu \circ E_0$. Then $\pi_{f_0}(A_0)''$ is isomorphic to M , since the portions of μ over I_1 and I_2 yield type I factors, and M is an infinite factor. Let $f = \text{Tr} \otimes f_0$. Then B is contained in the definition domain of f . For $j \in I_1$ let $x_j = (\otimes_{p=-j}^{j-1} 1^{(p)}) \otimes e_{11}^{(j)}$. Let $y \in B$. Then for all large enough j , x_j commutes with y . We have then, for all large enough j ,

$$\begin{aligned} \|\pi_f(x_j) \eta_f(y) - \eta_f(y)\|^2 &= f(y^*(x_j - 1)^2 y) \\ &= f(y^* y) f((x_j - 1)^2 e_0) = 0. \end{aligned}$$

Since $\eta_f(B)$ is dense in H_f , it follows that $\pi_f(x_j)$ tends σ -strongly to the identity. Now let $k \neq 0$. Then $I_1 + k$ has only finitely many elements outside of I_2 . Thus for all large enough j ,

$$\|\pi_f \circ \alpha^k(x_j) \eta_f(y)\|^2 = \|\eta_f(y)\|^2 f(\alpha^k(x_j) e_0) = 0.$$

Thus $\pi_f \circ \alpha^k(x_j)$ tends σ -strongly to zero. It follows that π_f and $\pi_{f \circ \alpha^k}$ are disjoint [7, 5.2.4]. □

LEMMA 1.2. *Let A be a C^* -algebra, let $\alpha \in \text{Aut}(A)$, and let f be a lower semi-continuous weight on A such that f and $f \circ \alpha^k$ are disjoint for $k \neq 0$. Let \tilde{f} be the canonical extension of f to $A \times_\alpha \mathbb{Z}$. Then $\pi_{\tilde{f}}(A \times_\alpha \mathbb{Z})''$ is unitarily equivalent to $\pi_f(A)'' \otimes L(l^2(\mathbb{Z}))$, acting on $H_f \otimes l^2(\mathbb{Z})$.*

Proof. Let $H = H_f \otimes l^2(\mathbb{Z}) \cong l^2(\mathbb{Z}, H_f)$, let $\pi: A \rightarrow L(H)$ be given by $\pi(x) = \sum_{n \in \mathbb{Z}} \pi_f \circ \alpha^{-n}(x) \otimes \delta_n$, and let $U = 1 \otimes U_0$, where U_0 is the (rightward) shift on $l^2(\mathbb{Z})$. Then (π, U) is a covariant representation

of (A, α, \mathbb{Z}) . We claim first that for $x \in C_c(\mathbb{Z}, A) \subset A \times_\alpha \mathbb{Z}$, the map $\pi_{\tilde{f}}(x) \mapsto \sum_n \pi(x(n))U^n$ is unitarily implemented. To see this we will need to identify a suitable dense subspace of $H_{\tilde{f}}$. Let

$$L_f = \{x \in A : f(x^*x) < \infty\},$$

$$L_{\tilde{f}} = \left\{x \in A \times_\alpha \mathbb{Z} : \tilde{f}(x^*x) < \infty\right\}.$$

Let $E : A \times_\alpha \mathbb{Z} \rightarrow A$ be the canonical conditional expectation, so that $\tilde{f} = f \circ E$. If $\{\hat{\alpha}_t\}_{t \in [0, 2\pi)}$ is the dual action, then E is given by:

$$E(x) = \frac{1}{2\pi} \int_0^{2\pi} \hat{\alpha}_t(x) dt.$$

For $x \in A \times_\alpha \mathbb{Z}$, the elements

$$z_j = \frac{1}{2\pi} \int_0^{2\pi} \hat{\alpha}_t(x) e^{-ijt} dt$$

belong to $C_c(\mathbb{Z}, A)$, and $z_j(k) = \delta_{jk}x_j$, where the last equation defines the elements $\{x_j\}$ in A . The $\{x_j\}$ uniquely determine x . By Lemma 2.2, Chapter 1, of [8],

$$x = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \hat{\alpha}_t(x) k_n(t) dt,$$

where $\{k_n(t)\}$ is any summability kernel. Choosing $\{k_n\}$ to be Fejer's kernel,

$$k_n(t) = \sum_{j=-n}^n \left(1 - \frac{|j|}{n+1}\right) e^{ijt},$$

and letting

$$\sigma_n(x) = \frac{1}{2\pi} \int_0^{2\pi} \hat{\alpha}_t(x) k_n(t) dt,$$

a short computation yields $\sigma_n(x) \in C_c(\mathbb{Z}, A)$, and

$$\sigma_n(x)(j) = \begin{cases} \left(1 - \frac{|j|}{n+1}\right) x_j, & |j| \leq n, \\ 0, & |j| > n. \end{cases}$$

Another short computation yields

$$\begin{aligned} 0 &\leq E((x - \sigma_n(x))^*(x - \sigma_n(x))) \\ &= E(x^*x) - \sum_{j=-n}^n \left(1 - \left(\frac{|j|}{n+1}\right)^2\right) \alpha^{-j}(x_j^*x_j). \end{aligned}$$

Hence

$$\tilde{f}(x^*x) \geq \sum_{j=-n}^n \left(1 - \left(\frac{|j|}{n+1} \right)^2 \right) f \circ \alpha^{-j}(x_j^*x_j), \quad n = 0, 1, 2, \dots$$

It follows that $x \in L_{\tilde{f}} \Rightarrow x_j \in \alpha^{-j}(L_f)$ for all j . From the above and the lower semi-continuity of f , it follows that

$$\tilde{f}(x^*x) = \lim_{n \rightarrow \infty} \sum_{j=-n}^n \left(1 - \left(\frac{|j|}{n+1} \right)^2 \right) f \circ \alpha^{-j}(x_j^*x_j).$$

Hence letting

$$L_0 = \left\{ x \in C_c(\mathbb{Z}, A) : x(j) \in \alpha^j(L_f) \right\},$$

it follows that $\eta_{\tilde{f}}(L_0)$ is dense in $H_{\tilde{f}}$.

Now define $W_0: \eta_{\tilde{f}}(L_0) \rightarrow H$ by

$$\left(W_0 \eta_{\tilde{f}}(x) \right)(j) = \eta_f \left(\alpha^{-j}(x(j)) \right).$$

One easily verifies that W_0 is isometric, and that

$$W_0 \pi_{\tilde{f}}(x) \eta_{\tilde{f}}(y) = \left(\sum_n \pi(x(n)) U^n \right) W_0 \eta_{\tilde{f}}(y),$$

for $x \in C_c(\mathbb{Z}, A)$, $y \in L_0$. Thus W_0 extends by continuity to a unitary operator implementing the desired equivalence.

Now we note that by the disjointness hypothesized for the $\{\pi_f \circ \alpha^k : k \in \mathbb{Z}\}$, it follows (see [7, 5.3]) that

$$\pi(A)'' = \pi_f(A)'' \otimes l^\infty(\mathbb{Z}).$$

Hence

$$(\pi(A) \cup \{U\})'' = \pi_f(A)'' \otimes L(l^2(\mathbb{Z})). \quad \square$$

THEOREM 1.3. *For every infinite Krieger factor M , there is a state ρ on O_2 , with $\pi_\rho(O_2)''$ isomorphic to M , and such that ρ extends the (unique) tracial state on Choi's algebra.* □

Proof. Let M be an infinite Krieger factor. Choose a state f_0 on A_0 as in Lemma 1.1. Since M is infinite, $\pi_f(A)''$ is isomorphic to M , where f is as in the statement of Lemma 1.1. Since f is lower semi-continuous, and M infinite, Lemma 1.2 implies that $\pi_{\tilde{f}}(A \times_\alpha \mathbb{Z})''$ is isomorphic to M . Let $\rho = \tilde{f}_0 = \tilde{f}|_{O_2}$, as in the

section on preliminaries. Let P be the projection onto $\overline{\eta_{\tilde{f}}(O_2)}$, and let $Q = \pi_{\tilde{f}}(e_0)$. Then $\pi_{\rho}(O_2)''$ is unitarily equivalent to $PQ\pi_{\tilde{f}}(A \times_{\alpha} \mathbb{Z})''QP$. The latter is a factor and is infinite since O_2 is infinite. Therefore $\pi_{\rho}(O_2)''$ is isomorphic to M . As noted in the section on preliminaries, the restriction of ρ to Choi's algebra is tracial. Since f_0 is a state, so is ρ . Thus ρ extends the tracial state on Choi's algebra. \square

2. Equivalence of states on the CAR algebra. We will now consider unitary equivalence and quasi-equivalence of the states of O_2 obtained from different weight sequences. First we need to study the corresponding weights on A . We use the results of [11] to get a slightly different characterization than that of [10] and [3].

Let $\{t_j\}_{j \geq j_0}$ and $\{r_j\}_{j \geq j_1}$ be weight sequences in $(0, 1)$, and let f and g be the corresponding weights on A . In [11] it was shown that $\sum_{j=0}^{\infty} (t_j - r_j)^2 t_j^{-1} (1 - t_j)^{-1} < \infty$ implies the unitary equivalence of π_f and π_g . We wish to reformulate this convergence condition so as to obtain an equivalence relation.

DEFINITIONS. Let $\{t_j\}_{j \geq j_0}$ and $\{r_j\}_{j \geq j_1}$ be weight sequences in $(0, 1)$, and let f and g be the corresponding weights.

1. We write $\{t_j\} \mathcal{R} \{r_j\}$, or equivalently $f \mathcal{R} g$, if

$$\sum_{j=0}^{\infty} (t_j - r_j)^2 t_j^{-1} (1 - t_j)^{-1} < \infty,$$

2. We write $\{t_j\} \sim \{r_j\}$, or equivalently $f \sim g$, if there is a partition $\{I_1, I_2, I_3\}$ of \mathbb{N} (unrelated to the partition constructed in Lemma 1.1) such that

- (i) $\sum_{I_1} t_j < \infty$, and $\sum_{I_1} r_j < \infty$,
- (ii) $\sum_{I_2} (1 - t_j) < \infty$, and $\sum_{I_2} (1 - r_j) < \infty$,
- (iii) $\sum_{I_3} (t_j - r_j)^2 t_j^{-1} (1 - t_j)^{-1} < \infty$, and $\sum_{I_3} (t_j - r_j)^2 r_j^{-1} (1 - r_j)^{-1} < \infty$.

The fact that \sim is an equivalence relation on the weight sequences in $(0, 1)$ follows from Theorem 2.4 below. We remark here that \sim is clearly reflexive and symmetric.

LEMMA 2.1. *Let $\{t_j\}$ and $\{r_j\}$ be weight sequences in $(0, 1)$ with $\{t_j\} \mathcal{R} \{r_j\}$. Then $\{t_j\} \sim \{r_j\}$.*

Proof. Choose $p > 1$. Let

$$I_1 = \{j: t_j r_j^{-1} \geq p\}, \quad I_2 = \{j: (1 - t_j)(1 - r_j)^{-1} \geq p\},$$

and

$$I_3 = \mathbb{N} \setminus (I_1 \cup I_2).$$

(i):

$$\begin{aligned} \infty &> \sum_{I_1} (r_j - t_j)^2 t_j^{-1} (1 - t_j)^{-1} \\ &= \sum_{I_1} \left(1 - \frac{r_j}{t_j}\right)^2 t_j (1 - t_j)^{-1} \\ &\geq (1 - p^{-1})^2 \sum_{I_1} t_j (1 - t_j)^{-1} \\ &\geq (1 - p^{-1})^2 \sum_{I_1} t_j. \end{aligned}$$

Hence $\sum_{I_1} t_j < \infty$. But then $\sum_{I_1} r_j \leq p^{-1} \sum_{I_1} t_j < \infty$.

(ii): Similar to (i).

(iii):

$$\begin{aligned} &\sum_{I_3} (t_j - r_j)^2 r_j^{-1} (1 - r_j)^{-1} \\ &= \sum_{I_3} (t_j - r_j)^2 t_j^{-1} (1 - t_j)^{-1} (t_j r_j^{-1}) (1 - t_j) (1 - r_j)^{-1} \\ &< p^2 \sum_{I_3} (t_j - r_j)^2 t_j^{-1} (1 - t_j)^{-1} < \infty. \quad \square \end{aligned}$$

LEMMA 2.2. *Let $\{t_j\}$ and $\{r_j\}$ be weight sequences in $(0, 1)$ with $\{t_j\} \sim \{r_j\}$. Then there is a weight sequence $\{s_j\}$ in $(0, 1)$ such that $\{t_j\} \mathcal{R} \{s_j\}$ and $\{r_j\} \mathcal{R} \{s_j\}$.*

Proof. Let

$$s_j = \begin{cases} \min(r_j, t_j), & j \in I_1, \\ \max(r_j, t_j), & j \in I_2, \\ t_j, & j \in I_3. \end{cases}$$

Then

$$\begin{aligned} &\sum (s_j - t_j)^2 t_j^{-1} (1 - t_j)^{-1} \\ &= \sum_{I_1} (s_j - t_j)^2 t_j^{-1} (1 - t_j)^{-1} + \sum_{I_2} (s_j - t_j)^2 t_j^{-1} (1 - t_j)^{-1} \\ &\leq \sum_{I_1} t_j (1 - t_j)^{-1} + \sum_{I_2} t_j^{-1} (1 - t_j) < \infty, \quad \text{and} \end{aligned}$$

$$\begin{aligned} & \sum (s_j - r_j)^2 r_j^{-1} (1 - r_j)^{-1} \\ & \leq \sum_{I_1} r_j (1 - r_j)^{-1} + \sum_{I_2} r_j^{-1} (1 - r_j) \\ & \quad + \sum_{I_3} (t_j - r_j)^2 r_j^{-1} (1 - r_j)^{-1} < \infty. \end{aligned} \quad \square$$

THEOREM 2.3. *Let $\{t_j\}_{j \geq j_0}$ and $\{r_j\}_{j \geq j_1}$ be weight sequences in $(0, 1)$, and let f and g be the corresponding weights on A . Let $T: \eta_f(x) \in \eta_f(B) \mapsto \eta_g(x) \in H_g$. The following are equivalent:*

- (1) $\{t_j\} \mathcal{R} \{r_j\}$,
- (2) $\eta_g(B) \subset D(T^*)$ (and hence T is preclosed),
- (3) $\eta_g(B) \subset D(T^*)$, T^*T has dense range, and $\text{polar}(\overline{T})$ is a unitary operator (intertwining π_f and π_g).

Proof. (1) \Rightarrow (3): This implication is contained in the proof of Lemma 3.8 of [11].

(3) \Rightarrow (2): Immediate.

(2) \Rightarrow (1): The computation in the proof of Lemma 3.8(i) of [11] shows that if x is in B , then $\eta_g(x) \in D(T^*)$ implies that

$$\lim_{n \rightarrow \infty} \langle \eta_f(xh_n), \xi \rangle$$

exists for $\xi \in \eta_f(B)$, and defines a bounded linear function of ξ . (The elements $h_n \in B$ are given by

$$h_n = \bigotimes_{j=-n}^n (\Omega_j \Lambda_j^{-1})^{(j)},$$

where $\{\Omega_j\}$ are obtained from $\{r_j\}$ in the same way that $\{\Lambda_j\}$ are obtained from $\{t_j\}$.) It follows by the Banach-Steinhaus theorem that $\{\eta_f(xh_n): n \in \mathbb{N}\}$ is a bounded set. A computation similar to the first computation in the proof of Lemma 3.3 of [11] shows that if x is in B_k ,

$$\|\eta_f(xh_n)\|^2 = f(h_k x^* x h_k) \prod_{j=k+1}^n [1 + (t_j - r_j)^2 t_j^{-1} (1 - t_j)^{-1}].$$

Hence the infinite product $\prod [1 + (t_j - r_j)^2 t_j^{-1} (1 - t_j)^{-1}]$ converges. Equivalently, $\{t_j\} \mathcal{R} \{r_j\}$. □

THEOREM 2.4. *Let $\{t_j\}_{j \geq j_0}$ and $\{r_j\}_{j \geq j_1}$ be weight sequences in $(0, 1)$, and let f and g be the corresponding weights on A . Let $T: \eta_f(x) \in \eta_f(B) \mapsto \eta_g(x) \in H_g$. The following are equivalent:*

- (1) π_f and π_g are unitarily equivalent,
- (2) $\{t_j\} \sim \{r_j\}$,
- (3) T is preclosed,
- (4) T is preclosed, and $\text{polar}(\overline{T})$ is a unitary operator (intertwining π_f and π_g).

Proof. (2) \Rightarrow (4) : By Lemma 2.2 there is a weight sequence in $(0, 1)$ with corresponding weight γ such that $f \mathcal{R} \gamma$ and $g \mathcal{R} \gamma$. Let

$$\begin{aligned} S_f: \eta_f(x) \in \eta_f(B) &\mapsto \eta_\gamma(x), \quad \text{and} \\ S_g: \eta_g(x) \in \eta_g(B) &\mapsto \eta_\gamma(x). \end{aligned}$$

Note that $T = S_g^{-1} S_f$. By Theorem 2.3, $\eta_\gamma(B) \subset D(S_f^*) \cap D(S_g^*)$. Let $x, y \in B$. Then

$$\begin{aligned} \langle S_g^* \eta_\gamma(x), T \eta_f(y) \rangle &= \langle S_g^* \eta_\gamma(x), \eta_g(y) \rangle \\ &= \langle \eta_\gamma(x), \eta_\gamma(y) \rangle = \langle \eta_\gamma(x), S_f \eta_f(y) \rangle \\ &= \langle S_f^* \eta_\gamma(x), \eta_f(y) \rangle. \end{aligned}$$

This implies that $S_g^* \eta_\gamma(B) \subset D(T^*)$. By Theorem 2.3, again, $S_g^* \eta_\gamma(B) = R(S_g^* S_g)$ is dense in H_g . Thus T is preclosed. Since \sim is symmetric, T^{-1} is also preclosed. This implies that \overline{T} is one-to-one. Since \overline{T} has dense range, it follows that $\text{polar}(\overline{T})$ is unitary.

(4) \Rightarrow (3) : Immediate.

(3) \Rightarrow (1) : Since \overline{T} has dense range, $W = \text{polar}(\overline{T})$ is a co-isometry. From $T \pi_f(x)|_{\eta_f(B)} = \pi_g(x) T$, it follows routinely that $\pi_f|_{W^* W H_f}$ is unitarily equivalent to π_g . Since π_f and π_g are factor representations, π_f and π_g are quasi-equivalent. Since f and g are faithful and of product type, π_f and π_g are unitarily equivalent.

(1) \Rightarrow (2) : It is shown in [3] that π_f is unitarily equivalent to π_g if and only if

$$\sum \left(1 - (r_j t_j)^{1/2} - [(1 - r_j)(1 - t_j)]^{1/2} \right) < \infty.$$

(In [3] it is actually shown that this condition is equivalent to quasi-equivalence. Since we are working here with faithful product weights, this is the same as unitary equivalence.)

By rationalizing the numerator one obtains

$$\begin{aligned} & 1 - (ab)^{1/2} - [(1 - a)(1 - b)]^{1/2} \\ &= C^{-1} \left(a + b - 2ab - 2[ab(1 - a)(1 - b)]^{1/2} \right), \end{aligned}$$

where $1 \leq C \leq 2$ for $0 \leq a, b \leq 1$. Rationalizing again, one obtains

$$\begin{aligned} & 1 - (ab)^{1/2} - [(1 - a)(1 - b)]^{1/2} \\ &= C^{-1} (a - b)^2 \left((a - b)^2 + [(a - a^2)^{1/2} + (b - b^2)^{1/2}]^2 \right)^{-1}. \end{aligned}$$

Let

$$\begin{aligned} K_1 &= \{j: r_j - r_j^2 < t_j - t_j^2\}, \\ K_2 &= \{j: r_j - r_j^2 > t_j - t_j^2\}, \\ K_3 &= \{j: r_j = t_j\}, \\ K_4 &= \{j: r_j = 1 - t_j\}. \end{aligned}$$

We have

$$\begin{aligned} \infty &> \sum_{K_1} \frac{(r_j - t_j)^2}{(r_j - t_j)^2 + 4t_j(1 - t_j)} \\ &= \sum_{K_1} \frac{1}{1 + 4 \left((r_j - t_j)^2 t_j^{-1} (1 - t_j)^{-1} \right)^{-1}}. \end{aligned}$$

Equivalently,

$$\sum_{K_1} (r_j - t_j)^2 t_j^{-1} (1 - t_j)^{-1} < \infty.$$

Similarly,

$$\sum_{K_2} (r_j - t_j)^2 r_j^{-1} (1 - r_j)^{-1} < \infty.$$

Clearly,

$$\sum_{K_3} (r_j - t_j)^2 r_j^{-1} (1 - r_j)^{-1} < \infty.$$

Finally,

$$\begin{aligned} \infty &> \sum_{K_4} \frac{(1 - 2t_j)^2}{(1 - 2t_j)^2 + 2t_j(1 - t_j)} \\ &> \sum_{K_4} (1 - 2t_j)^2. \end{aligned}$$

Hence, $\sum_{K_4} (t_j - 1/2)^2 < \infty$, $\sum_{K_4} (r_j - 1/2)^2 < \infty$, and so

$$\sum_{K_4} (r_j - t_j)^2 r_j^{-1} (1 - r_j)^{-1} < \infty.$$

It is now clear from Lemma 2.1 that $\{t_j\} \sim \{r_j\}$. □

REMARKS. The relation \sim makes sense for arbitrary weight sequences, where we impose the following additional restriction on \sim in Definition 2: $t_j = 0$ or $r_j = 0 \Rightarrow j \notin I_3$. In this generality, the condition of [3] used in (1) \Rightarrow (2) above is equivalent to quasi-equivalence of the weights. It is easy to see that our proof of the equivalence of \sim and the condition of [3] holds for arbitrary weight sequences.

COROLLARY 2.5. *Let $\{t_j\}$ and $\{r_j\}$ be arbitrary weight sequences, and let f and g be the corresponding weights. The following are equivalent:*

- (1) π_f and π_g are quasi-equivalent,
- (2) $\{t_j\} \sim \{r_j\}$. □

3. Equivalence of states on O_2 .

DEFINITION. Let $\{t_j\}_{j \geq j_0}$ and $\{r_j\}_{j \geq j_1}$ be weight sequences. We say that $\{t_j\}$ and $\{r_j\}$ have *equivalent tails* if there is an integer n such that $\{t_j\}_{j \geq j_0} \sim \{r_{j-n}\}_{j \geq j_1+n}$.

LEMMA 3.1. *Let $\{r_j\}_{j \geq j_1}$ be a weight sequence. Let g and h be the weights corresponding to $\{r_j\}_{j \geq j_1}$ and $\{r_{j-n}\}_{j \geq j_1+n}$, respectively. Then $h = g \circ \alpha^n$.*

Proof. Let $l \geq |j_1| + |n|$. For $x \in (B_l)_+$, and $k > l + |n|$, a straightforward calculation shows that

$$h_k(e_k x e_k) = g_k(e_k \alpha^n(x) e_k).$$

Since h_k and g_k are bounded for each k , it follows that for $x \in (A_l)_+$, and $k > l + |n|$,

$$h_k(e_k x e_k) = g_k(e_k \alpha^n(x) e_k).$$

Now let $x \in A_+$. We have for $l \geq |j_1| + |n|$ and $k > l + |n|$:

$$\begin{aligned} h(e_l x e_l) &= \sup_k h_k(e_k e_l x e_l e_k) = \sup_k g_k(e_k \alpha^n(e_l x e_l) e_k) \\ &= \sup_k g_k(e_k e_{l-n} \alpha^n(x) e_{l-n} e_k) \\ &= g(e_{l-n} \alpha^n(x) e_{l-n}) \leq g \circ \alpha^n(x). \end{aligned}$$

It follows that $h(x) \leq g \circ \alpha^n(x)$. An analogous calculation shows that $g \circ \alpha^n(x) \leq h(x)$. \square

REMARKS. Let $J = C_c(\mathbb{Z}, B) \subset A \times_\alpha \mathbb{Z}$. Let f be a weight on A arising from a weight sequence. Since $\eta_f(B)$ is dense in H_f , and $\eta_f(B) \subset L_f$, it follows from the proof of Lemma 1.2 that $\eta_{\tilde{f}}(J)$ is dense in $H_{\tilde{f}}$. The map $W_f: H_{\tilde{f}} \rightarrow l^2(\mathbb{Z}, H_f)$ defined by

$$W_f \eta_{\tilde{f}}(x)(k) = \eta_f \circ \alpha^{-k}(x(k)), \quad x \in J,$$

extends to a unitary operator, and

$$W_f \pi_{\tilde{f}}(x) W_f^* = \sum_k \pi(x(k)) U^k$$

(see the proof of Lemma 1.2). We will let α also denote the automorphism of $A \times_\alpha \mathbb{Z}$ defined by $(\alpha(x))(k) = \alpha(x(k))$, for $x \in J$.

LEMMA 3.2. Let $\{t_j\}$ and $\{r_j\}$ be weight sequences in $(0, 1)$, and let f and g be the corresponding weights. Let $T: \eta_{\tilde{f}}(y) \in \eta_{\tilde{f}}(J) \mapsto \eta_{\tilde{g}}(\alpha^n(y))$. Suppose that $\{t_j\}$ and $\{r_j\}$ have equivalent tails. Then T is preclosed and $\text{polar}(\overline{T})$ is a unitary operator (intertwining $\pi_{\tilde{f}}$ and $\pi_{\tilde{g}} \circ \alpha^n$).

Proof. Let $T_0: \eta_f(b) \in \eta_f(B) \mapsto \eta_g(\alpha^n(b))$. By Lemma 3.1 and Theorem 2.4, T_0 is preclosed and $\text{polar}(\overline{T_0})$ is unitary. Let $\xi \in C_c(\mathbb{Z}, D(T_0^*))$ and let $y \in J$. Then

$$\begin{aligned} \langle W_g^* \xi, T \eta_{\tilde{f}}(y) \rangle &= \langle \xi, W_g \eta_{\tilde{g}}(\alpha^n(y)) \rangle \\ &= \sum_k \langle \xi(k), \eta_g \circ \alpha^{n-k}(y(k)) \rangle \\ &= \sum_k \langle \xi(k), T_0 \eta_f \circ \alpha^{-k}(y(k)) \rangle \\ &= \langle (T_0^* \otimes 1) \xi, W_f \eta_{\tilde{f}}(y) \rangle \\ &= \langle W_f^* (T_0^* \otimes 1) \xi, \eta_{\tilde{f}}(y) \rangle, \end{aligned}$$

where we let $T_0^* \otimes 1$ have domain

$$C_c(\mathbb{Z}, D(T_0^*)) \subset l^2(\mathbb{Z}, H_g) \cong H_g \otimes l^2(\mathbb{Z}).$$

Hence $W_g^* C_c(\mathbb{Z}, D(T_0^*)) \subset D(T^*)$, and

$$T^* W_g^* | C_c(\mathbb{Z}, D(T_0^*)) = W_f^*(T_0^* \otimes 1).$$

Since $\text{polar}(\overline{T_0})$ is unitary, we know that T_0^* has dense domain and dense range. Thus T is preclosed and \overline{T} is one-to-one. Since T already has dense range, it follows that $\text{polar}(\overline{T})$ is unitary. \square

THEOREM 3.3. *Let $\{t_j\}_{j \geq j_0}$ and $\{r_j\}_{j \geq j_1}$ be weight sequences in $(0, 1)$, let f and g be the corresponding weights on A , and let \tilde{f}_0 and \tilde{g}_0 be the restrictions of \tilde{f} and \tilde{g} , respectively, to $e_0(A \times_\alpha \mathbb{Z})e_0$.*

(1) *If $\{t_j\}$ and $\{r_j\}$ have equivalent tails, then \tilde{f}_0 and \tilde{g}_0 are quasi-equivalent.*

(2) *If $\{t_j\}$ and $\{r_j\}$ do not have equivalent tails, then \tilde{f}_0 and \tilde{g}_0 are disjoint.*

Proof. (1): Let $\{t_j\}_{j \geq j_0} \sim \{r_{j-n}\}_{j \geq j_1+n}$. By symmetry we may assume that $n \geq 0$. By altering $\{r_j\}$ up to \sim , we may assume that $j_1 = 0$. By Lemma 3.2 we have that $T: \eta_{\tilde{f}}(y) \in \eta_{\tilde{f}}(J) \mapsto \eta_{\tilde{g}}(\alpha^n(y))$ is preclosed, and that $\text{polar}(\overline{T})$ is unitary. Let $\overline{T} = V|\overline{T}|$ be the polar decomposition of \overline{T} , and let $D = e_0(A \times_\alpha \mathbb{Z})e_0$. Note that $\alpha(D) \subset D$, and that $H_{\tilde{f}_0} = \overline{\eta_{\tilde{f}}(D)}$, and $H_{\tilde{g}_0} = \overline{\eta_{\tilde{g}}(D)}$. It now follows that $V\pi_{\tilde{f}_0}(x) = \pi_{\tilde{g}_0}(\alpha^n(x))V|H_{\tilde{f}_0}$, $x \in D$. Let $V_0 = V|H_{\tilde{f}_0}: H_{\tilde{f}_0} \rightarrow H_{\tilde{g}_0}$. Then V_0 is an isometry. We then have

$$\begin{aligned} \pi_{\tilde{f}_0}(x) &= V_0^* \pi_{\tilde{g}_0}(\alpha^n(x)) V_0 \\ &= \left(\pi_{\tilde{g}_0}(S_1^*)^n V_0 \right)^* \pi_{\tilde{g}_0}(x) \left(\pi_{\tilde{g}_0}(S_1^*)^n V_0 \right). \end{aligned}$$

Note that

$$\begin{aligned} R(V_0) &= \overline{\eta_{\tilde{g}_0}(S_1^n D (S_1^*)^n)} \subset \overline{\eta_{\tilde{g}_0}(S_1^n D)} \\ &= R\left(\pi_{\tilde{g}_0}(S_1^n)\right). \end{aligned}$$

Hence $\pi_{\tilde{g}_0}(S_1^*)^n V_0$ is an isometry. It follows that $\pi_{\tilde{f}_0}$ is unitarily equivalent to the subrepresentation of $\pi_{\tilde{g}_0}$ obtained by restricting to

$$\text{range}\left(\pi_{\tilde{g}_0}\left(\left(S_1^*\right)^n V_0\right)\right) = \overline{\eta_{\tilde{g}_0}(D (S_1^*)^n)} = \overline{\pi_{\tilde{g}_0}(D) \eta_{\tilde{g}_0}\left(\left(S_1^*\right)^n\right)}.$$

We claim that the central support of this subrepresentation is 1. To see this we will show that

$$\overline{\pi_{\tilde{g}_0}(D) \pi_{\tilde{g}_0}(D)' \eta_{\tilde{g}_0} \left((S_1^*)^n \right)} = H_{\tilde{g}_0}.$$

In order to demonstrate this we will need certain elements of $\pi_{\tilde{g}_0}(D)'$. Let G_n be the set of words of length n in the elements S_1 and S_2 . Note that

$$(*) \quad \sum_{w \in G_n} ww^* = 1.$$

For $w, z \in G_n$ let

$$T_{w,z} : \eta_{\tilde{g}_0}(x) \in \eta_{\tilde{g}_0}(D) \mapsto \eta_{\tilde{g}_0}(xwz^*).$$

Let $x = S_{p_0} \cdots S_{p_k} S_{q_l}^* \cdots S_{q_0}^*$. Then $x^*x = (q_0, q_0) \otimes \cdots \otimes (q_l, q_l)$. Then

$$zw^*x^*xwz^* = \begin{cases} zz^* \otimes (q_{n+1}, q_{n+1}) \otimes \cdots \otimes (q_l, q_l), & \text{if } l \geq n \text{ and } w = S_{q_0} \cdots S_{q_n}, \\ zz^*, & \text{if } l < n \text{ and } w = S_{q_0} \cdots S_{q_l} S_{i_{l+1}} \cdots S_{i_n}, \\ 0, & \text{otherwise.} \end{cases}$$

Hence

$$\tilde{g}_0(zw^*x^*xwz^*) = \begin{cases} \frac{g_0(zz^*)}{g_0((q_0, q_0) \otimes \cdots \otimes (q_n, q_n))} \cdot g_0(x^*x), \\ \frac{g_0(zz^*)}{g_0((q_0, q_0) \otimes \cdots \otimes (q_l, q_l))} \cdot g_0(x^*x), \\ 0, \end{cases}$$

where the three lines in the bracket correspond to the three cases defined in the previous bracket. Note that for $l \leq n$,

$$g_0((q_0, q_0) \otimes \cdots \otimes (q_l, q_l)) \geq \prod_{m=0}^n \min(r_m, 1 - r_m) \stackrel{\text{def}}{=} C_n.$$

Thus

$$\tilde{g}_0(zw^*x^*xwz^*) \leq g_0(zz^*) (C_n)^{-1} \tilde{g}_0(x^*x).$$

Let y also be a product in the elements S_1, S_2, S_1^*, S_2^* . If $x \neq y$, a simple calculation shows that $\tilde{g}_0(y^*x) = 0$. (This relies heavily on the fact that g factors through the diagonal of A_0 .) It now follows easily that

$$\tilde{g}_0(zw^*x^*xwz^*) \leq g_0(zz^*) (C_n)^{-1} \tilde{g}_0(x^*x),$$

for x in the $*$ -algebra generated by S_1 and S_2 , and hence by continuity, for all x in D . This implies that $T_{w,z}$ extends to a bounded linear operator (also denoted $T_{w,z}$) in $\pi_{\tilde{g}_0}(D)'$.

By (*), we have that

$$\sum_{w \in G_n} \pi_{\tilde{g}_0}(w) T_{S_1^n, w} \eta_{\tilde{g}_0}((S_1^*)^n) = \eta_{\tilde{g}_0}(1).$$

Since $\eta_{\tilde{g}_0}(1)$ is cyclic for $\pi_{\tilde{g}_0}(D)$, the claim follows.

This implies that $\pi_{\tilde{f}_0}$ and $\pi_{\tilde{g}_0}$ are quasi-equivalent.

(2): Suppose that $\{t_j\}$ and $\{r_j\}$ do not have equivalent tails. Then $f \not\sim g \circ \alpha^n$ for all integers n . It follows from Theorem 2.3, and the fact that these are faithful factor weights on A , that f is disjoint from $g \circ \alpha^n$ for all n . We now need two lemmas.

LEMMA 3.4. *Let A be a C^* -algebra, let $\alpha \in \text{Aut}(A)$, let f, g be lower semi-continuous weights on $A \times_\alpha \mathbb{Z}$, and let \tilde{f}, \tilde{g} be the canonical extensions of f, g to weights on $A \times_\alpha \mathbb{Z}$. The following are equivalent:*

- (a) \tilde{f} is disjoint from \tilde{g} ,
- (b) f is disjoint from $g \circ \alpha^n$ for all integers n .

Proof. (a) \Rightarrow (b): If (b) is false then there is an integer n , and a non-zero operator T in $L(H_f, H_g)$, such that $T\pi_f(x) = \pi_g(\alpha^n(x))T$ for $x \in A$. We will use the notations and results of the proof of Lemma 1.2. Let $\tilde{T} = T \otimes U_0^{-n}$ on $H_f \otimes l^2(\mathbb{Z})$. Then for $x \in A$,

$$\begin{aligned} \tilde{T}\tilde{\pi}_f(x) &= \tilde{T} \left(\sum_{k \in \mathbb{Z}} \pi_f \circ \alpha^{-k}(x) \otimes \delta_k \right) \\ &= \sum_{k \in \mathbb{Z}} T\pi_f \circ \alpha^{-k}(x) \otimes U_0^{-n} \delta_k \\ &= \sum_{k \in \mathbb{Z}} \pi_g \circ \alpha^{n-k}(x) T \otimes \delta_{k-n} U_0^{-n} = \tilde{\pi}_g(x) \tilde{T}. \end{aligned}$$

It is clear that \tilde{T} commutes with $U = 1 \otimes U_0$. Therefore \tilde{T} intertwines $\pi_{\tilde{f}}$ and $\pi_{\tilde{g}}$.

(b) \Rightarrow (a): If (b) holds, then $\tilde{\pi}_f = \bigoplus_n \pi_f \circ \alpha^{-n}$ and $\tilde{\pi}_g = \bigoplus_n \pi_g \circ \alpha^{-n}$ are disjoint. Then there is a bounded sequence $\{x_j\}$ in A such that $\tilde{\pi}_f(x_j)$ tends strongly to 1, and $\tilde{\pi}_g(x_j)$ tends strongly to 0. Since

$(\tilde{\pi}_f, U)$ is unitarily equivalent to $\pi_{\tilde{f}}$ (by Lemma 1.2), and similarly for g , it follows that $\pi_{\tilde{f}}(x_j)$ tends strongly to 1, and $\pi_{\tilde{g}}(x_j)$ tends strongly to 0. Hence \tilde{f} and \tilde{g} are disjoint. \square

LEMMA 3.5. *Let C be a C^* -algebra, let $e \in C$ be a projection, and let $D = eCe$. Let π and ρ be representations of C , and let π_0 and ρ_0 be the restrictions of π and ρ to D , acting on eH_π and eH_ρ , respectively. If π and ρ are disjoint, then π_0 and ρ_0 are disjoint.*

Proof. There is a bounded sequence $\{x_j\}$ in C such that $\pi(x_j)$ tends strongly to 1, and $\rho(x_j)$ tends strongly to 0. Then $\{ex_j e\}$ is a bounded sequence in D , and $\pi_0(ex_j e)$ tends strongly to 1, and $\rho_0(ex_j e)$ tends strongly to 0. \square

End of proof of Theorem 3.3. By Lemma 3.4, \tilde{f} and \tilde{g} are disjoint. Then by Lemma 3.5, \tilde{f}_0 and \tilde{g}_0 are disjoint. \square

THEOREM 3.6. *Let $\{t_j\}_{j \geq 0}$ and $\{r_j\}_{j \geq 0}$ be weight sequences in $[0, 1]$. Let f and g be the corresponding weights on A , and let \tilde{f}_0 and \tilde{g}_0 be the restrictions of \tilde{f} and \tilde{g} , respectively, to $e_0(A \times_\alpha \mathbb{Z})e_0$.*

(1) *If $\{t_j\}$ and $\{r_j\}$ have equivalent tails, then \tilde{f}_0 and \tilde{g}_0 are quasi-equivalent.*

(2) *If $\{t_j\}$ and $\{r_j\}$ do not have equivalent tails, then \tilde{f}_0 and \tilde{g}_0 are disjoint.*

Proof. (1) By Corollary 2.5, we may modify $\{t_j\}$ and $\{r_j\}$ up to equivalence so as to obtain weight sequences in $(0, 1)$. In doing so we modify f and g up to quasi-equivalence. It is easily seen that \tilde{f} , \tilde{g} and \tilde{f}_0 , \tilde{g}_0 are all modified up to quasi-equivalence. Part (1) now follows from Theorem 3.3(1).

(2) By Corollary 2.5, f and $g \circ \alpha^n$ are disjoint for all n . Part (2) now follows from Lemmas 3.4 and 3.5. \square

4. Pure states on O_2 : a theorem of Archbold, Lazar, Tsui, and Wright. We now apply the results of the previous sections to give an alternative proof of a result of Archbold, Lazar, Tsui, and Wright ([2], Proposition 2.10). We will consider weight sequences $\{t_j\}_{j \geq j_0}$ with $t_j = 0$ or 1 for all j . It is clear that two such weight sequences are equivalent under \sim if and only if they are eventually equal, and have

equivalent tails if and only if they have equal tails. (We say that $\{t_j\}$ and $\{r_j\}$ have *equal tails* if $t_j = r_{j-n}$ for some n and all sufficiently large j .) Given such a weight sequence $\{t_j\}$, with corresponding weight f , we construct a pure state on A , denoted f' , as follows. Let

$$\Lambda'_j = \begin{cases} \Lambda_j, & j \geq j_0, \\ e_{11}, & j < j_0. \end{cases}$$

We construct a positive functional on $\bigcup A_n$ by means of $\{\Lambda'_j\}$ instead of $\{\Lambda_j\}$. Then the resulting functional is continuous. We let f' denote its extension to A , which is always a pure state. We remark that since

$$f'|_{A_0} = f|_{A_0} = f_0,$$

and

$$\tilde{f}'|_{e_0(A \times_\alpha \mathbb{Z})e_0} = f' \circ E|_{e_0(A \times_\alpha \mathbb{Z})e_0} = f_0 \circ F_0,$$

the functionals on O_2 obtained from f and f' are equal. We remark also that \sim is equivalent to quasi-equivalence for the primed functionals, as in §2.

LEMMA 4.1. *Let $\{t_j\}_{j \geq 0}$ and $\{r_j\}_{j \geq 0}$ be weight sequences with $t_j, r_j = 0$ or 1 for all j . Let f, g be the corresponding weights on A , and let \tilde{f}_0, \tilde{g}_0 be the resulting states on O_2 . Then*

- (1) \tilde{f}_0 is pure if and only if $\{t_j\}$ is not eventually repeating.
- (2) \tilde{f}_0 and \tilde{g}_0 are unitarily equivalent if $\{t_j\}$ and $\{r_j\}$ have equal tails, and are disjoint otherwise.

Proof. It is clear that $\{t_j\} \sim \{t_{j-n}\}$ for some non-zero n if and only if $\{t_j\}$ is eventually repeating. Suppose that $\{t_j\}$ is not eventually repeating. Then Corollary 2.5 implies that f' and $f' \circ \alpha^n$ are disjoint for $n \neq 0$. Now by Lemma 1.2 we conclude that \tilde{f}' is a pure state of $A \times_\alpha \mathbb{Z}$, and hence that $\tilde{f}_0 = \tilde{f}'|_{e_0(A \times_\alpha \mathbb{Z})e_0}$ is a pure state of O_2 . The proof of the other implication in (1) will follow from the proof of (2).

We know by Theorem 3.6 that \tilde{f}_0 and \tilde{g}_0 are quasi-equivalent or disjoint, depending on whether $\{t_j\}$ and $\{r_j\}$ have or do not have equal tails. From what we have already shown it follows that if $\{t_j\}$ and $\{r_j\}$ are not eventually repeating then \tilde{f}_0 and \tilde{g}_0 are pure, and hence quasi-equivalent if and only if unitarily equivalent. (We mention that what we have done so far already proves Theorem 4.2 below.)

We claim now that if $\{t_j\}$ is eventually repeating, then $\pi_{\tilde{f}'}(A \times_{\alpha} \mathbb{Z})'$ is non-trivial and abelian. Since $\pi_{\tilde{f}'}(e_0)$ has central cover 1, it follows that $\pi_{\tilde{f}'}(O_2)'$ is non-trivial and abelian. It then follows that repeating weight sequences can never yield pure states on O_2 , proving the reverse implication in (1). Moreover two repeating weight sequences with equal tails yield quasi-equivalent states on O_2 with multiplicity-free GNS representations, and hence are unitarily equivalent. Thus the lemma will follow from the claim.

So assume that $\{t_j\}$ eventually repeats with period n . Then the weight sequences $\{t_j\}$ and $\{t_{j-n}\}$ differ in only finitely many places. It follows that there is a unitary element W_0 in \tilde{A} so that

$$f' \circ \alpha^n = f' \circ \text{Ad}(W_0).$$

An easy calculation shows that the map

$$\eta_{f'}(x) \in \eta_{f'}(A) \mapsto \eta_{f'}(\alpha^{-n}(x)W_0)$$

defines a unitary operator W on $H_{f'}$, and that

$$\pi_{f'} \circ \alpha^{-n} = \text{Ad}(W) \circ \pi_{f'}.$$

Letting $\pi = \sum_{k \in \mathbb{Z}} \pi_{f'} \circ \alpha^{-k} \otimes \delta_k$ acting on $H_{f'} \otimes l^2(\mathbb{Z})$, we have that

$$\pi = \sum_{j=0}^{n-1} \sum_{k \in \mathbb{Z}} \text{Ad}(W^k) \circ \pi_{f'} \circ \alpha^{-j} \otimes \delta_{kn+j}.$$

Letting $\tilde{W} = W \otimes 1$, $U = 1 \otimes U_0$, and $M = \pi_{\tilde{f}'}(A \times_{\alpha} \mathbb{Z})''$, and using Lemma 1.2, it is clear that $\tilde{W}U^n \in M'$. We will finish the proof by showing that M' is generated as a von Neumann algebra by $\tilde{W}U^n$. By Lemma 1.2 we may work interchangeably with M or with $(\pi(A) \cup \{U\})''$.

Since n is the period of repetition in $\{t_j\}$, it is easily seen that

$$\pi(A)'' = \left\{ \sum_{k \in \mathbb{Z}} T_k \otimes \delta_k : T_k \in L(H_{f'}), T_{k+n} = WT_kW^* \right\}.$$

Let operators on $H_{f'} \otimes l^2(\mathbb{Z})$ have matrix decompositions along the basis $\{\delta_k\}$ of $l^2(\mathbb{Z})$. Then $T = (T_{ij})$ is in $\pi(A)''$ if and only if

$$T_{ij} = \begin{cases} 0, & i \neq j, \\ W^a T_{bb} W^{-a}, & i = j = an + b, \text{ with } 0 \leq b < n. \end{cases}$$

A routine calculation now shows that $S = (S_{ij})$ is in $\pi(A)'$ if and only if

$$S_{ij} = \begin{cases} 0, & i \not\equiv j \pmod{n}, \\ \lambda_{ij} W^{(i-j)/n}, & i \equiv j \pmod{n}, \end{cases}$$

where the λ_{ij} are constants. Additionally, S commutes with U if and only if

$$S_{ij} = \begin{cases} 0, & i \not\equiv j \pmod{n}, \\ \lambda(i-j) W^{(i-j)/n}, & i \equiv j \pmod{n}, \end{cases}$$

where $\lambda(p)$ is the (necessarily unique) value of $\{\lambda_{k, pn+k} : k \in \mathbb{Z}\}$. Thus $S \in M'$ if and only if there is a function $\lambda: \mathbb{Z} \rightarrow \mathbb{C}$ such that

$$S = \sum_p \lambda(p) (\widetilde{W} U^n)^p. \quad \square$$

THEOREM 4.2 ([2, Proposition 2.10]). *There are uncountably many inequivalent pure states of O_2 which extend the trace on Choi's algebra.* □

5. Invariants for diagonal product states on O_2 . We now indicate how the techniques of [11] can be extended to give information about the modular spectrum and period of the factors obtained from weight sequences.

DEFINITION. Let $\{t_j\}$ be a weight sequence. If there is a non-zero k for which $\{t_j\} \sim \{t_{j-k}\}$, we define the *period* of $\{t_j\}$ to be

$$\text{per} \{t_j\} = \inf \{k > 0 : \{t_j\} \sim \{t_{j-k}\}\}.$$

If there is no such non-zero k , we say that the period of $\{t_j\}$ is infinite.

THEOREM 5.1. *Let $\{t_j\}$ be a weight sequence such that $\sum t_j(1-t_j) = \infty$, and $\text{per}\{t_j\} < \infty$. Let f be the weight on A corresponding to $\{t_j\}$, and \tilde{f}_0 the state on O_2 constructed from f . Let $M_f = \pi_f(A)''$, and $M_0 = \pi_{\tilde{f}_0}(O_2)''$. Then M_0 is a type III factor, and*

$$\begin{aligned} S(M_0) &\supset S(M_f), \\ T(M_0) &\subset T(M_f). \end{aligned}$$

(S and T are modular spectrum and modular period, respectively. These invariants were defined in [5]. See also [11].)

- [9] A. Lazar, S.-K. Tsui, and S. Wright, *Pure state extensions of the trace on the Choi algebra*, Proc. Amer. Math. Soc., **102** No. 4 (1988), 957–964.
- [10] R. T. Powers and R. Størmer, *Free states of the canonical anticommutation relations*, Comm. Math. Phys., **16** (1970), 1–33.
- [11] J. S. Spielberg, *Type III factor states on O_2 which extend the trace on Choi's algebra*, J. Operator Theory, (to appear).
- [12] C. E. Sutherland, *Notes on orbit equivalence; Krieger's theorem*, Univ. of Oslo Lecture Notes, No. 23, 1976.

Received October 5, 1988 and in revised form September 11, 1989.

DALHOUSIE UNIVERSITY
HALIFAX, NOVA SCOTIA, CANADA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1980 *Mathematics Subject Classification* (1985 Revision) scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly. Regular subscription rate: \$190.00 a year (12 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1990 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 144, No. 2

June, 1990

George E. Andrews and David M. Jackson , An algebraically derived q -analogue of a character sum associated with a class of semiregular permutations	207
Fabio Bardelli and Andrea Del Centina , The moduli space of genus four double covers of elliptic curves is rational	219
Young Do Chai , An estimate of the volume of a compact set in terms of its integral of mean curvature	229
Salvador Comalada , Elliptic curves with trivial conductor over quadratic fields	237
Kahar El-Hussein , Résolubilité semi-globale des opérateurs différentiels invariants sur les groupes de déplacements	259
David M. Goldschmidt , Classical link invariants and the Burau representation	277
Liliana Janicka , Radon-Nikodým problem for the variation of a vector measure	293
Wacław Marzantowicz , An almost classification of compact Lie groups with Borsuk-Ulam properties	299
Akira Ohbuchi , On the projective normality of some varieties of degree 5 ..	313
Ken'ichi Ohshika , Minimal measured laminations in geometric 3-manifolds	327
Hal Leslie Smith , A discrete Lyapunov function for a class of linear differential equations	345
John Samuel Spielberg , Diagonal states on O_2	361
Thomas Vogel , A note on the sessile drop	383
Gerold Wagner , On means of distances on the surface of a sphere (lower bounds)	389