

Pacific Journal of Mathematics

**THE p -PARTS OF BRAUER CHARACTER DEGREES IN
 p -SOLVABLE GROUPS**

YOU-QIANG WANG

THE p -PARTS OF BRAUER CHARACTER DEGREES IN p -SOLVABLE GROUPS

YOU-QIANG WANG

Let G be a finite group. Fix a prime integer p and let e be the largest integer such that p^e divides the degree of some irreducible Brauer character of G with respect to the same prime p . The primary object of this paper is to obtain information about the structure of Sylow p -subgroups of a finite p -solvable group G in knowledge of e .

As applications, we obtain a bound for the derived length of the factor group of a solvable group G relative to its unique maximal normal p -subgroup in terms of the arithmetic structure of its Brauer character degrees and a bound for the derived length of the factor group of G relative to its Fitting subgroup in terms of the maximal integer e when p runs through the prime divisors of the order of G .

All groups considered are finite. Let G be a group and p be a prime. We denote by $\text{IBr}_p(G)$ the set of irreducible Brauer characters of G with respect to the prime p . For the same prime p , let $e_p(G)$ be the largest integer e such that p^e divides $\varphi(1)$ for some $\varphi \in \text{IBr}_p(G)$. Let P be a Sylow p -subgroup of G . Then the Sylow p -invariants of G are defined as follows:

- (1) $b_p(G)$, where $p^{b_p(G)}$ is the order of P ;
- (2) $c_p(G)$, the class of P , that is, the length of the (upper or) lower central series of P ;
- (3) $dl_p(G)$, the length of the derived series of P ;
- (4) $ex_p(G)$, where $p^{ex_p(G)}$ is the exponent of P , that is, the greatest order of any element of P .

For a p -solvable group G , we let $l_p(G)$ and $r_p(G)$ denote the p -length and p -rank (respectively) of G , i.e. $r_p(G)$ is the largest integer r such that p^r is the order of a p -chief factor of G .

We give a linear bound for $r_p(G/O_p(G))$ and a logarithmic bound for $l_p(G/O_p(G))$ in terms of $e_p(G)$. Then, using induction on $l_p(G/O_p(G))$, we obtain bounds for $c_p(G/O_p(G))$, $dl_p(G/O_p(G))$ and $ex_p(G/O_p(G))$ in terms of $e_p(G)$.

As one application, we bound the derived length of $G/F(G)$ for a solvable group G in terms of $f(G)$, where

$$f(G) = \max\{e_p(G) \mid p \mid |G|\}$$

and $F(G)$ is the Fitting subgroup of G .

1. p -rank and p -length. For p -solvable G , we bound in this section $r_p(G/O_p(G))$ and $l_p(G/O_p(G))$ in terms of $e_p(G)$.

LEMMA 1.1. *Let G have a nilpotent normal p -complement M and let $O_p(G) = 1$. Then*

- (1) $b_p(G) \leq 2e_p(G)$;
- (2) if $|G|$ is odd, then $b_p(G) \leq e_p(G)$;
- (3) if a Sylow p -subgroup of G is abelian, then $b_p(G) \leq e_p(G)$.

Proof. Let $\Phi(M)$ denote the Frattini subgroup of M . Since M is nilpotent, $M' \leq \Phi(M)$ and hence $M/\Phi(M)$ is abelian (see Gorenstein [3, Chapter 6, Theorem 1.6]). Notice that $\Phi(M) \triangleleft G$. Consider the factor group $G/\Phi(M)$. Let P be a Sylow p -subgroup of G . Then

$$G/\Phi(M) = MP/\Phi(M) = M/\Phi(M) \cdot P\Phi(M)/\Phi(M).$$

Write $G_1 = G/\Phi(M)$, $M_1 = M/\Phi(M)$ and $P_1 = P\Phi(M)/\Phi(M)$. Then M_1 is an abelian normal p -complement of G_1 , and $P_1 = P\Phi(M)/\Phi(M) \cong P/P \cap \Phi(M) \cong P$. By Huppert [5, Chapter 3, Satz 3.18], P acts faithfully on M_1 , and hence P_1 acts faithfully on M_1 . Then $O_p(G_1) = 1$ and P_1 acts faithfully on $\text{Irr}(M_1)$ which is an abelian p' -group.

By Corollary 2.4 of Passman [16], there exists $\theta \in \text{Irr}(M_1)$ such that $|I_{P_1}(\theta)| \leq |P_1|^{1/2}$. So $|P_1 : I_{P_1}(\theta)| \geq |P_1|^{1/2}$. By Clifford's Theorem, $p^{e_p(G_1)} \geq |P_1|^{1/2}$. Since $|P_1| = |P|$ and $e_p(G) \geq e_p(G_1)$, $2e_p(G) \geq b_p(G)$. This gives (1).

If $|G|$ is odd or P is abelian, then we can apply Lemma 2.2 and Corollary 2.4 of Passman [16] to conclude that there exists $\theta \in \text{Irr}(M_1)$ such that $I_{P_1}(\theta) = 1$. By Clifford's Theorem, $|P_1| \leq p^{e_p(G_1)}$, that is, $b_p(G) \leq e_p(G_1)$. So (2) and (3) hold. \square

LEMMA 1.2. *Let G have a normal p -complement M and let $O_p(G) = 1$, where p is an odd prime. Let $M = M_1 \times \cdots \times M_n$, where all M_i 's are isomorphic nonabelian simple groups. Then*

- (1) $b_p(G) \leq 2e_p(G)$;
- (2) if $e_p(G) = 1$, then $b_p(G) = 1$.

Proof. Let P be a Sylow p -subgroup of G . Notice that for any $x \in P$ and M_i , $xM_ix^{-1} \in \{M_1, \dots, M_n\}$. We write $N_i = \text{Stab}_P(M_i)$ and $C = \bigcap_{i=1}^n N_i$. Also, we let

$$C_i = C_C(M_i) = \{x \in C \mid xyx^{-1} = y \text{ for all } y \in M_i\}.$$

For each i , $C/C_i \leq \text{Out}(M_i)$. By Lemma 1.3 of Gluck and Wolf [2], C/C_i is a cyclic p -group. Let $\Omega_1(C/C_i) = \langle x \in C/C_i \mid x^p = 1 \rangle$, then $\Omega_1(C/C_i)$ is a cyclic group of order p .

On the other hand, since C/C_i acts faithfully on M_i and since $p \nmid |M_i|$, C/C_i acts faithfully on $\text{Irr}(M_i)$ (see Isaacs [9, Theorem 6.32]). Thus

$$\bigcap_{\theta \in \text{Irr}(M_i)} C_{C/C_i}(\theta) = 1.$$

So, there exists some $\theta_i \in \text{Irr}(M_i)$ such that

$$\Omega_1(C/C_i) \cap C_{C/C_i}(\theta_i) = 1.$$

This forces that $C_{C/C_i}(\theta_i) = 1$, that is, $C_C(\theta_i) = \check{C}_i$.

Let $\theta = \theta_1 \times \dots \times \theta_n$. Then $\theta \in \text{Irr}(M) = \text{IBr}_p(M)$ and $I_C(\theta) = \bigcap_{i=1}^n C_C(\theta_i) = \bigcap_{i=1}^n \check{C}_i$. Since P acts faithfully on M , $\bigcap_{i=1}^n C_i = 1$, and hence $I_C(\theta) = 1$. Since $C \triangleleft P$, $MC \triangleleft G$ and hence $e_p(MC) \leq e_p(G)$. Applying Clifford's Theorem to the group MC , we have $|C| = |C : I_C(\theta)| \leq p^{e_p(MC)}$. Hence

$$(A) \quad |C| \leq p^{e_p(G)}.$$

On the other hand, P/C is a permutation group on the set $\{M_1, \dots, M_n\}$ and p is an odd prime. So, by Corollary 1 of Gluck [1], we may assume without loss of generality that $\text{Stab}_{P/C}\{M_1, \dots, M_t\} = 1$, that is $\text{Stab}_P\{M_1, \dots, M_t\} = C$ for some suitable $t \in \{1, \dots, n\}$.

Choose $\theta_j \in \text{Irr}(M_j)$ with $\theta_j \neq 1$, $j = 1, \dots, t$. Let $\theta = \theta_1 \times \dots \times \theta_t \times 1 \times \dots \times 1$. Then $\theta \in \text{Irr}(M) = \text{IBr}_p(M)$ and $I_P(\theta) \leq \text{Stab}_P\{M_1, \dots, M_t\} = C$. Applying Clifford's Theorem to the group G , we get $|P : I_P(\theta)| \leq p^{e_p(G)}$. Hence

$$(B) \quad |P : C| \leq p^{e_p(G)}.$$

Combining (A) with (B), we obtain $|P| \leq p^{2e_p(G)}$. This gives (1).

Suppose that $e_p(G) = 1$. By (A) and (B), we know that $|C| \leq p$ and $|P : C| \leq p$.

In the following, we want to show that either $C = P$ or $C = 1$. Assume not. Then we have $|C| = p$ and $|P| = p^2$.

Since P acts faithfully on M , C acts faithfully on M . Since $|C| = p$, there exists some M_k such that $C_k = C_C(M_k) = 1$, that is, C acts faithfully on M_k . Thus C acts faithfully on $\text{Irr}(M_k)$. Hence there exists some $\theta_k \in \text{Irr}(M_k)$ such that $\theta_k \neq 1$ and $C_C(\theta_k) = 1$.

Since $\text{Stab}_p\{M_{t+1}, \dots, M_n\} = \text{Stab}_p\{M_1, \dots, M_t\}$, we may assume without loss of generality that $k = 1$. Choose $\theta_j \in \text{Irr}(M_j)$ with $\theta_j \neq 1$, $j = 2, \dots, t$. Let $\theta = \theta_1 \times \theta_2 \times \dots \times \theta_t \times 1 \times \dots \times 1$. Then $\theta \in \text{Irr}(M) = \text{IBr}_p(M)$ and $I_P(\theta) \leq \text{Stab}_P\{M_1, \dots, M_t\} = C$. So $I_P(\theta) = I_C(\theta) = \bigcap_{i=1}^t C_C(\theta_i) = 1$ (because of $C_C(\theta_1) = 1$). Thus there exists $\theta \in \text{Irr}(M) = \text{IBr}_p(M)$ such that $I_P(\theta) = 1$. By Clifford's Theorem, $|P| = |P : I_P(\theta)| \leq p$. This contradicts to $|P| = p^2$. So (2) holds. \square

THEOREM 1.3. *Let G be p -solvable. Then*

$$r_p(G/O_p(G)) \leq 2e_p(G).$$

Proof. By induction on $|G|$, we may assume without loss of generality that $O_p(G) = 1$.

If $p = 2$, then G is solvable, and we are done by Manz and Wolf [13, Theorem 2.3]. In the following, we assume that p is an odd prime.

Let M be a minimal normal subgroup of G and let $N/M = O_p(G/M)$. By the inductive hypothesis, we may assume that $N/M \neq 1$.

Since G is p -solvable and $O_p(G) = 1$, we have the following two cases:

Case 1. M is an elementary abelian q -group for some prime $q \neq p$;

Case 2. M is the direct product of isomorphic nonabelian simple p' -group.

Consider the group N . Notice that $M = O_{p'}(N)$ and $O_p(N) = 1$. Applying Lemma 1.1 (1) and Lemma 1.2 (1) to the group N , we get $b_p(N) \leq 2e_p(N)$. Since $N \triangleleft G$, $e_p(N) \leq e_p(G)$ by Clifford's Theorem. Hence $b_p(N) \leq 2e_p(G)$.

By the inductive hypothesis, the p -rank $r_p(G/N)$ of G/N does not exceed $2e_p(G/N) \leq 2e_p(G)$. Since $r_p(G) \leq \max\{r_p(G/N), b_p(N)\}$, $r_p(G) \leq 2e_p(G)$. \square

By using Lemma 1.1 (2) instead of Lemma 1.1 (1), the same proof yields the following improvement for groups of odd order.

THEOREM 1.4. *Let G be a group of odd order. Then*

$$r_p(G/O_p(G)) \leq e_p(G).$$

We note that Theorem 1.3 and Theorem 1.4 improve Theorem 2.1 of Manz [12], by the Fong-Swan Theorem.

THEOREM 1.5. *Let G be solvable and $e_p(G) < p$. Then*

$$r_p(G/O_p(G)) \leq e_p(G).$$

Proof. By induction on $|G|$, we may assume without loss of generality that $O_p(G) = 1$.

Let M be a minimal normal subgroup of G and let $N/M = O_p(G/M)$. By the inductive hypothesis, we may assume that $N/M \neq 1$. Since $O_p(G) = 1$, M is an elementary abelian q -group with $q \neq p$.

Consider the group N . Since $N \triangleleft G$ and $O_p(G) = 1$, $O_p(N) = 1$. Notice that N has a normal p -complement M and $e_p(N) \leq e_p(G) < p$.

Let P be a Sylow p -subgroup of N . Since $O_p(N) = 1$, P acts faithfully on M by conjugation. Hence P acts faithfully on $\text{Irr}(M)$. Since M is an elementary abelian q -group, $\text{Irr}(M)$ is an abelian q -group. Let $\Omega_1, \dots, \Omega_n$ be the P -orbits of $\text{Irr}(M)$ and $p^f = \max\{|\Omega_1|, \dots, |\Omega_n|\}$. We may assume without loss of generality that $|\Omega_1| = p^f$. Let $\theta_1 \in \Omega_1$. Applying Clifford's Theorem to the group N , we get $|\Omega_1| = |P : I_P(\theta_1)| \leq p^{e_p(N)}$. So $|\Omega_1| < p^p$. By Corollary 2.4 of Passman [16], there exists $\theta \in \text{Irr}(M)$ such that $I_P(\theta) = 1$. We apply Clifford's Theorem to conclude that $|P| = |P : I_P(\theta)| \leq p^{e_p(N)}$. So $b_p(N) \leq e_p(N)$, and hence $b_p(N) \leq e_p(G)$.

By the inductive hypothesis, the p -rank $r_p(G/N)$ of G/N does not exceed $e_p(G/N) \leq e_p(G)$. Since $r_p(G) \leq \max\{r_p(G/N), b_p(N)\}$, $r_p(G) \leq e_p(G)$. \square

Recall that the rank $r(G)$ of G is the maximum dimension of all chief-factors of G and $f(G) = \max\{e_p(G) \mid p \mid |G|\}$.

COROLLARY 1.6. *Let G be solvable. Then*

- (1) $r(G/F(G)) \leq 2f(G)$;
- (2) if $|G|$ is odd, then $r(G/F(G)) \leq f(G)$.

Proof. Let p be a prime number such that $p \mid |G|$. Then, by Theorem 1.3 and Theorem 1.4, $r_p(G/O_p(G)) \leq 2e_p(G) \leq 2f(G)$, and if $|G|$ is odd, $r_p(G/O_p(G)) \leq e_p(G) \leq f(G)$. Since $O_p(G) \leq F(G)$,

this yields that $r(G/F(G)) \leq 2f(G)$, and if $|G|$ is odd, $r(G/F(G)) \leq f(G)$. □

Combining Wolf [17, Theorem 2.3] with Theorem 1.3, we have

THEOREM 1.7. *Let G be p -solvable. Then*

- (1) $l_p(G/O_p(G)) \leq 1 + \log_p(2e_p(G))$ if p is not a Fermat prime; and
- (2) $l_p(G/O_p(G)) \leq 2 + \log_s(2e_p(G)/(p-1))$ where $s = (p^2 - p + 1)/p$.

2. Sylow p -invariants. In this section, we bound $c_p(G/O_p(G))$, $dl_p(G/O_p(G))$ and $ex_p(G/O_p(G))$ for a p -solvable group G in terms of $e_p(G)$. In particular, we show that if $e_p(G) = 1$, then a Sylow p -subgroup of $G/O_p(G)$ is elementary abelian. We also give bounds for $b_p(G/O_p(G))$.

LEMMA 2.1. *Let G have a normal p -complement and let $O_p(G) = 1$. Then*

- (1) $dl_p(G) \leq e_p(G)$;
- (2) $ex_p(G) \leq e_p(G)$;
- (3) $c_p(G) \leq p^{e_p(G)-1}$.

Proof. Let P be a Sylow p -subgroup of G and let H be the normal p -complement of G . Then P acts faithfully on $\text{Irr}(H)$.

Write $\Omega = \text{Irr}(H)$. Let $\Omega_1, \dots, \Omega_n$ be the P -orbits of Ω . Then P acts transitively on each Ω_i . Let $\varphi_i : P \rightarrow S(\Omega_i)$ be the homomorphism induced by the action, where $S(\Omega_i)$ is the permutation group on Ω_i . Then we can define a homomorphism from P into the direct product

$$S(\Omega_1) \times \cdots \times S(\Omega_n)$$

as follows:

$$\begin{aligned} \varphi : P &\rightarrow S(\Omega_1) \times \cdots \times S(\Omega_n), \\ \varphi(x) &= (\varphi_1(x), \dots, \varphi_n(x)), \quad x \in P. \end{aligned}$$

Since P acts faithfully on Ω , $\text{Ker } \varphi = 1$, and hence φ is an injection. By the definition of φ , we know that

$$P \cong \varphi(P) \leq \varphi_1(P) \times \cdots \times \varphi_n(P),$$

where each $\varphi_i(P) \leq S(\Omega_i)$ is a p -group.

Since $dl(\varphi_i(P)) \leq dl(P)$, $\max\{dl(\varphi_i(P)) : i = 1, \dots, n\} \leq dl(P)$. On the other hand, we have that

$$\begin{aligned} dl(P) &= dl(\varphi(P)) \\ &\leq dl(\varphi_1(P) \times \dots \times \varphi_n(P)) \\ &= \max\{dl(\varphi_i(P)) : i = 1, \dots, n\}. \end{aligned}$$

Hence, $dl_p(G) = \max\{dl(\varphi_i(P)) : i = 1, \dots, n\}$. Similarly,

$$ex_p(G) = \max\{ex_p(\varphi_i(P)) : i = 1, \dots, n\},$$

and

$$c_p(G) = \max\{c_p(\varphi_i(P)) : i = 1, \dots, n\}.$$

Let $p^f = \max\{|\Omega_i| : i = 1, \dots, n\}$. We may assume without loss of generality that $|\Omega_1| = p^f$. By Huppert [5, Chapter 3, Satz 15.3], $dl_p(G) \leq f$, $ex_p(G) \leq f$ and $c_p(G) \leq p^{f-1}$. Choose $\theta \in \Omega_1$. By Clifford's Theorem, $|\Omega_1| = |P : I_P(\theta)| \leq p^{e_p(G)}$. So $f \leq e_p(G)$. Thus the conclusions (1)–(3) hold. \square

THEOREM 2.2. *Let G be p -solvable. Then*

- (1) $dl_p(G/O_p(G)) \leq l_p(G/O_p(G))e_p(G)$;
- (2) $ex_p(G/O_p(G)) \leq l_p(G/O_p(G))e_p(G)$;
- (3) $c_p(G/O_p(G)) \leq l_p(G/O_p(G))p^{e_p(G)-1}$.

Proof. Since $e_p(G) = e_p(G/O_p(G))$, we may assume without loss of generality that $O_p(G) = 1$. We use induction on $l_p(G)$.

Write $E = O_{p',p}(G)$ and $M = O_p(G)$. Since $O_p(G) = 1$, $O_p(M) = 1$. Clearly, M has a normal p -complement E . Thus, by Lemma 2.1, we have that

$$dl_p(M) \leq e_p(M), \quad ex_p(M) \leq e_p(M), \quad \text{and} \quad c_p(M) \leq p^{e_p(M)-1}.$$

Since $M \triangleleft G$, $e_p(M) \leq e_p(G)$. Hence,

$$(A) \quad dl_p(M) \leq e_p(G), \quad ex_p(M) \leq e_p(G), \quad c_p(M) \leq p^{e_p(G)-1}.$$

Since $M = O_{p',p}(G)$, $O_p(G/M) = 1$ and $l_p(G/M) = l_p(G) - 1$. Then the induction yields that

$$\begin{aligned} dl_p(G/M) &\leq l_p(G/M)e_p(G/M), \\ ex_p(G/M) &\leq l_p(G/M)e_p(G/M), \quad \text{and} \\ c_p(G/M) &\leq l_p(G/M)p^{e_p(G/M)-1}. \end{aligned}$$

Hence, (B)

$$\begin{aligned} dl_p(G/M) &\leq (l_p(G) - 1)e_p(G), \\ ex_p(G/M) &\leq (l_p(G) - 1)e_p(G), \\ c_p(G/M) &\leq (l_p(G) - 1)p^{e_p(G)-1}. \end{aligned}$$

By (A) and (B), we have the conclusions. \square

Combining Theorem 1.7 with Theorem 2.2, we have

COROLLARY 2.3. *Let G be p -solvable. Then*

(1) *if p is not a Fermat prime, then*

$$\begin{aligned} dl_p(G/O_p(G)) &\leq (1 + \log_p(2e_p(G)))e_p(G), \\ ex_p(G/O_p(G)) &\leq (1 + \log_p(2e_p(G)))e_p(G), \\ c_p(G/O_p(G)) &\leq (1 + \log_p(2e_p(G)))p^{e_p(G)-1}; \end{aligned}$$

(2) *if p is a Fermat prime, then*

$$\begin{aligned} dl_p(G/O_p(G)) &\leq (2 + \log_s(2e_p(G)/(p-1)))e_p(G), \\ ex_p(G/O_p(G)) &\leq (2 + \log_s(2e_p(G)/(p-1)))e_p(G), \\ c_p(G/O_p(G)) &\leq (2 + \log_s(2e_p(G)/(p-1)))p^{e_p(G)-1}, \end{aligned}$$

where $s = (p^2 - p + 1)/p$.

In the rest of this section, we give some improvements on the bounds we just got for the cases $e_p(G) = 1, 2$ and $e_p(G) < p$.

COROLLARY 2.4. *Let G be solvable and $e_p(G) < p$. Then*

(1) *if p is not a Fermat prime, then*

$$\begin{aligned} dl_p(G/O_p(G)) &\leq e_p(G), \\ ex_p(G/O_p(G)) &\leq e_p(G), \\ c_p(G/O_p(G)) &\leq p^{e_p(G)-1}; \end{aligned}$$

(2) *if p is a Fermat prime, then*

$$\begin{aligned} dl_p(G/O_p(G)) &\leq 2e_p(G), \\ ex_p(G/O_p(G)) &\leq 2e_p(G), \\ c_p(G/O_p(G)) &\leq 2p^{e_p(G)-1}. \end{aligned}$$

Proof. By Theorem 1.5 and Wolf [17, Theorem 2.3], $l_p(G/O_p(G)) \leq 1 + \log_p(e_p(G))$, if p is not a Fermat prime; and $l_p(G/O_p(G)) \leq 2 + \log_s(e_p(G)/(p-1))$, if p is a Fermat prime, where $s = (p^2 - p + 1)/p$. Since $e_p(G) < p$, $l_p(G/O_p(G)) \leq 1$, if p is not a Fermat prime; and

$l_p(G/O_p(G)) \leq 2$, if p is a Fermat prime. Hence the conclusions follow from Theorem 2.2. \square

For a group G , Michler [14] and Okuyama [15] show that if $e_p(G) = 0$, then G has a normal Sylow p -subgroup. The assertion for p -solvable groups is elementary and well-known.

THEOREM 2.5. *Let G be p -solvable and $e_p(G) = 1$. Then*

$$r_p(G/O_p(G)) \leq 1.$$

Proof. The case $p = 2$ is done by Theorem 1.5. In the following, we assume that p is an odd prime. By induction on $|G|$, we may assume without loss of generality that $O_p(G) = 1$.

Let M be a minimal normal subgroup of G and $N/M = O_p(G/M)$. By the inductive hypothesis, we may assume that $N/M \neq 1$. Since G is p -solvable and $O_p(G) = 1$, we have the following two cases:

Case 1. M is an elementary abelian q -group for some prime $q \neq p$.

Case 2. M is the direct product of isomorphic nonabelian simple p' -groups.

Consider the group N . Since $N \triangleleft G$ and $O_p(G) = 1$, $O_p(N) = 1$. Then $1 \leq e_p(N) \leq e_p(G) = 1$. Thus $e_p(N) = 1$. Notice that N has a normal p -complement M . By Lemma 2.1, N/M is an abelian p -group. Applying Lemma 1.1 (3) and Lemma 1.2 (2) to the group N , we get $b_p(N) = 1$.

By the inductive hypothesis, the p -rank $r_p(G/N) \leq 1$. Since $r_p(G) \leq \max\{r_p(G/N), b_p(N)\}$, $r_p(G) \leq 1$. \square

Since $l_p(G) \leq r_p(G)$ (see Huppert [5, Chapter 6, Hauptsatz 6.6 (c)]), we get the following corollary by combining Theorem 2.2 with Theorem 2.5.

COROLLARY 2.6. *Let G be p -solvable and $e_p(G) = 1$. Then a Sylow p -subgroup of $G/O_p(G)$ is an elementary abelian p -group.*

For $e_p(G) = 2$, there is no general result similar to Corollary 2.6. Let $G = S_3 \text{ wr } Z_2$. Then $e_2(G) = 2$ and $O_2(G) = 1$. The Sylow 2-subgroup of G is $Z_2 \text{ wr } Z_2$, which is not abelian.

However, for solvable groups, we have the following corollary.

COROLLARY 2.7. *Let G be solvable and $e_p(G) = 2$ with $p \geq 5$. Then*

- (1) $dl_p(G/O_p(G)) \leq 2$;
- (2) $ex_p(G/O_p(G)) \leq 2$;
- (3) $c_p(G/O_p(G)) \leq p$.

Proof. For $p \geq 5$, by Theorem 1.5 and Wolf [17, Theorem 2.3], $l_p(G/O_p(G)) \leq 1$ if $e_p(G) = 2$. Hence the conclusions follow from Theorem 2.2. \square

In closing this section, we include the following remark, which tells us that logarithmic bounds for the Sylow p -invariants of $G/O_p(G)$ in terms of $e_p(G)$ are probably the best bounds we can expect.

REMARK 2.8. Fix a prime p . Let $G_0 \neq 1$ be a p' -group. We construct groups by iterated wreath products as follows: let $G_1 = G_0 \text{ wr } Z_p$ and $G_2 = G_1 \text{ wr } Z_p$. Following this way, we have $G_n = G_{n-1} \text{ wr } Z_p$ for any natural number n .

By Hall and Higman [4, Lemma 3.5.1], $dl_p(G_n) = ex_p(G_n) = n$. Since $O_p(G_1) = 1$ and $|G_1|_p = p$, $e_p(G_1) = 1$. In the following, we use an induction argument on n to show that

$$p^{n-1} \leq e_p(G_n) \leq (p^n - 1)/(p - 1).$$

Suppose that $p^{n-2} \leq e_p(G_{n-1}) \leq (p^{n-1} - 1)/(p - 1)$. By the definition of G_n , $G_n = (G_{n-1} \times \cdots \times G_{n-1}) \rtimes Z_p$. Let $H_n = G_{n-1} \times \cdots \times G_{n-1}$. Then $H_n \triangleleft G_n$ and $e_p(H_n) = pe_p(G_{n-1})$. Hence $p^{n-1} \leq e_p(H_n) \leq e_p(G_n)$. In particular, $p^{n-1} \leq e_p(G_n)$.

On the other hand, let $\varphi \in \text{IBr}_p(G_n)$ such that $\varphi(1) = p^{e_p(G_n)} m$. Choose $\theta \in \text{IBr}_p(H_n)$ such that $\varphi \in \text{IBr}_p(G_n | \theta)$. By Clifford's Theorem, $\varphi(1) = e\theta(1)$ with a positive integer e . Also, by Lemma 3.2 of Isaacs [8], φ is an irreducible constituent of θ^{G_n} . Thus $\varphi(1) \leq \theta^{G_n}(1) = |G_n : H_n| \theta(1) = p\theta(1)$. So $0 < e \leq p$. This yields that $e_p(G_n) \leq e_p(H_n) + 1$. Hence $e_p(G_n) \leq p(p^{n-1} - 1)/(p - 1) + 1 = (p^n - 1)/(p - 1)$.

Now we consider bounding the $b_p(G/O_p(G))$ for a p -solvable group G in terms of $e_p(G)$.

LEMMA 2.9. *Let G have a solvable normal p -complement H and let $O_p(G) = 1$. Then $b_p(G) \leq 2dl(H)e_p(G)$.*

Proof. We use an induction argument on $dl(H)$. Let P be a Sylow p -subgroup of G . Then P acts on H/H' by conjugation. Let $Q = C_p(H/H')$.

Since P/Q acts faithfully on H/H' , P/Q acts faithfully on $\text{Irr}(H/H')$ which is an abelian p' -group. By Corollary 2.4 of Passman [16], there exists $\theta \in \text{Irr}(H/H')$ such that $|C_{P/Q}(\theta)| \leq |P/Q|^{1/2}$. So $|P/Q : C_{P/Q}(\theta)| \geq |P/Q|^{1/2}$. Consider $\theta \in \text{Irr}(H)$ with $H' \leq \text{Ker } \theta$. By Clifford's Theorem, $|P : I_p(\theta)| \leq p^{e_p(G)}$. Since $|P : I_p(\theta)| = |P/Q : C_{P/Q}(\theta)|$, $|P/Q|^{1/2} \leq p^{e_p(G)}$. Hence $\log_p |P/Q| \leq 2e_p(G)$. If $Q = 1$, then we are done.

Next, we assume that $Q \neq 1$. We claim that Q acts faithfully on H' . Assume not. We may assume without loss of generality that Q acts trivially on H' . Since $Q = C_p(H/H')$, Q acts trivially on H/H' . Since $(|Q|, |H|) = 1$, Q acts trivially on H (see Huppert [5, Chapter 3, Hilfssatz 13.3 (b)]). But since P acts faithfully on H , we must have $Q = 1$. This contradicts to $Q \neq 1$.

Write $G_1 = H'Q$. Then G_1 has a normal p -complement H' and $O_p(G_1) = C_Q(H') = 1$. Furthermore, we claim that $G_1 \triangleleft G$. Since $H' \triangleleft G$ and $Q \triangleleft P$, we only need to show that $hQh^{-1} \subseteq H'Q$ for all $h \in H$. Let $q \in Q = C_p(H/H')$. Then $q^{-1}hqh^{-1} \in H'$. Hence $hqh^{-1} \in qH' \subseteq QH' = H'Q$. Thus $G_1 = H'Q \triangleleft G$. By Clifford's Theorem, $e_p(G_1) \leq e_p(G)$.

Since $dl(H') = dl(H) - 1 < dl(H)$, by induction, $\log_p |Q| \leq 2dl(H')e_p(G_1) = 2(dl(H) - 1)e_p(G_1)$. Hence,

$$\begin{aligned} b_p(G) &= \log_p |P| \\ &= \log_p (|Q| |P/Q|) \\ &= \log_p |Q| + \log_p |P/Q| \\ &\leq 2(dl(H) - 1)e_p(G_1) + 2e_p(G) \\ &\leq 2(dl(H) - 1)e_p(G) + 2e_p(G) \\ &= 2dl(H)e_p(G), \end{aligned}$$

which is the claim. □

The following Lemma is a corollary of Lemma 1.1.

LEMMA 2.10. *Let G be solvable, $O_p(G) = 1$ and P a Sylow p -subgroup of G . Let G have a normal p -complement. Then*

$$b_p(G) \leq 2e_p(F(G)P).$$

Proof. Since $O_p(G) = 1$, the Fitting subgroup $F(G)$ is a p' -group. By Huppert [5, Chapter 3, Satz 4.2], $C_G(F(G)) \leq F(G)$ and hence $C_P(F(G)) = 1$. Let $G_1 = F(G)P$. Then G_1 has a nilpotent normal p -complement $F(G)$ and $O_p(G_1) = C_P(F(G)) = 1$. By Lemma 1.1, $b_p(G_1) \leq 2e_p(G_1)$. Since $b_p(G_1) = b_p(G)$, $b_p(G) \leq 2e_p(F(G)P)$. \square

To handle p -solvable groups with arbitrary p -length, we introduce the following definition.

DEFINITION 2.11. For a prime p and a positive integer n , we define $\lambda_p(n)$ and $\beta_p(n)$ by

$$\lambda_p(n) = \sum_{i=1}^{\infty} [n/p^i]$$

and

$$\beta_p(n) = \sum_{i=0}^{\infty} [n/(p-1)p^i].$$

PROPOSITION 2.12. *If p is a prime and n is a positive integer, then*

$$\lambda_p(n) \leq n - 1 \quad \text{and} \quad \beta_p(n) \leq 2n - 1.$$

Proof. Since $\lambda_p(n) \leq (n-1)/(p-1)$, $\lambda_p(n) \leq n-1$. Since $p \geq 2$, $2(p-1)^2 - p \geq (p-1)^2 - (p-1)$, and hence $2n(p-1)^2 - np \geq (p-1)^2 - (p-1)$. So $(2n-1) \geq (np/(p-1)^2) - (1/(p-1))$. Since $\beta_p(n) \leq (np/(p-1)^2) - (1/(p-1))$, $\beta_p(n) \leq 2n-1$. \square

THEOREM 2.13. *Let G be p -solvable and $O_{p'}(G)$ be solvable. Suppose that $O_p(G) = 1$. Then*

- (1) $b_p(G) \leq 6dl(O_{p'}(G))e_p(O_{p',p}(G)) - 1$; and
- (2) $b_p(G) \leq 4dl(O_{p'}(G))e_p(O_{p',p}(G)) - 1$ unless p is a Fermat prime.

Proof. Write $E = O_{p'}(G)$ and $M = O_{p',p}(G)$. Since $O_p(G) = 1$, $O_p(M) = 1$. Clearly, M has a solvable normal p -complement E . Thus $b_p(M) \leq 2dl(O_{p'}(G))e_p(O_{p',p}(G))$ by Lemma 2.9.

Let $b_p(M) = m$, hence $|M/E| = p^m$. By Wolf [17, Corollary 2.1], we have that

- (1) $b_p(G) \leq m + \beta_p(m)$; and
- (2) $b_p(G) \leq m + \lambda_p(m)$ unless p is a Fermat prime.

Applying Proposition 2.12, we obtain

- (1) $b_p(G) \leq 3m - 1$; and
- (2) $b_p(G) \leq 2m - 1$ unless p is a Fermat prime.

Since $m = b_p(M)$, we get

- (1) $b_p(G) \leq 6dl(O_{p'}(G))e_p(O_{p',p}(G)) - 1$; and
- (2) $b_p(G) \leq 4dl(O_{p'}(G))e_p(O_{p',p}(G)) - 1$ unless p is a Fermat prime. □

Similarly, applying Lemma 2.10, we obtain the following Theorem.

THEOREM 2.14. *Let G be solvable, $O_p(G) = 1$ and P a Sylow p -subgroup of $O_{p',p}(G)$. Then*

- (1) $b_p(G) \leq 6e_p(F(G)P) - 1$; and
- (2) $b_p(G) \leq 4e_p(F(G)P) - 1$ unless p is a Fermat prime.

3. The derived length of solvable groups. Let $n = \prod_{i=1}^k p_i^{a_i}$ be the prime number decomposition of a natural number n ($a_i \neq 0$). We define

$$\omega(n) = \sum_{i=1}^k a_i.$$

For a group G , we let

$$\omega(G) = \max\{\omega(\chi(1)) \mid \chi \in \text{Irr}(G)\}$$

and

$$\omega_p(G) = \max\{\omega(\varphi(1)) \mid \varphi \in \text{IBr}_p(G)\}.$$

Recall that $f(G) = \max\{e_p(G) \mid p \mid |G|\}$.

For a solvable group G , we obtain a bound for the derived length of $G/O_p(G)$ in terms of $\omega_p(G)$ and a quadratic bound for the derived length of $G/F(G)$ in terms of $f(G)$.

LEMMA 3.1. *Let G be solvable with $O_p(G) = 1$ and $l_p(G) = 1$. Then $dl(G) \leq 5\omega_p(G)$.*

Proof. Since $O_p(G) = 1$ and $l_p(G) = 1$, $e_p(G) \geq 1$. Thus $\omega_p(G) \geq e_p(G) \geq 1$. If $\omega_p(G) = 1$, then $dl(G) \leq 4$ by Huppert [7, Theorem 1]. So, $dl(G) \leq 5\omega_p(G)$, and we are done in this case.

In the following, we assume that $\omega_p(G) \geq 2$. We have two cases to consider.

Case 1. $O_{p',p}(G) = G$.

By Lemma 2.1, $dl(G/O_{p'}(G)) \leq e_p(G)$, and hence $dl(G/O_{p'}(G)) \leq \omega_p(G)$. Since $O_{p'}(G)$ is a p' -group, $\omega_p(O_{p'}(G)) = \omega(O_{p'}(G))$.

If $\omega(O_{p'}(G)) \geq 2$, then, by Huppert [6, Theorem 3], $dl(O_{p'}(G)) \leq 2\omega(O_{p'}(G))$, and hence $dl(O_{p'}(G)) \leq 2\omega_p(O_{p'}(G))$. Since $O_{p'}(G) \triangleleft G$, $\omega_p(O_{p'}(G)) \leq \omega_p(G)$ by Clifford's Theorem. So $dl(O_{p'}(G)) \leq 2\omega_p(G)$. Thus

$$\begin{aligned} dl(G) &\leq dl(O_{p'}(G)) + dl(G/O_{p'}(G)) \\ &\leq 2\omega_p(G) + \omega_p(G) = 3\omega_p(G). \end{aligned}$$

If $\omega(O_{p'}(G)) \leq 1$, then, by Isaacs and Passman [10, Theorem 6.1], $dl(O_{p'}(G)) \leq 3$, and hence

$$dl(G) \leq dl(O_{p'}(G)) + dl(G/O_{p'}(G)) \leq 3 + \omega_p(G).$$

Since $\omega_p(G) \geq 2$, $dl(G) \leq 2\omega_p(G) + \omega_p(G) = 3\omega_p(G)$.

Case 2. $O_{p',p,p'}(G) = G$.

Write $M = O_{p',p}(G)$. By what we have just proved in the above, we have that

- (1) if $\omega_p(M) = 1$, then $dl(M) \leq 4$;
- (2) if $\omega_p(M) \geq 2$, then $dl(M) \leq 3\omega_p(M)$.

Since $M \triangleleft G$, $\omega_p(M) \leq \omega_p(G)$. Furthermore, since $\omega_p(G) \geq 2$, $3\omega_p(G) \geq 6$. Thus $dl(M) \leq 3\omega_p(G)$.

Since G/M is a p' -group, $\omega_p(G/M) = \omega(G/M)$. If $\omega(G/M) \leq 1$, $dl(G/M) \leq 3$ by Isaacs and Passman [10, Theorem 6.1]. If $\omega(G/M) \geq 2$, $dl(G/M) \leq 2\omega(G/M)$ by Huppert [6, Theorem 3], and hence $dl(G/M) \leq 2\omega_p(G/M)$. Since $\omega_p(G/M) \leq \omega_p(G)$ and $2\omega_p(G) \geq 4$, $dl(G/M) \leq 2\omega_p(G)$. Therefore,

$$\begin{aligned} dl(G) &\leq dl(M) + dl(G/M) \\ &\leq 3\omega_p(G) + 2\omega_p(G) = 5\omega_p(G). \end{aligned}$$

This completes the proof of the lemma. □

THEOREM 3.2. *Let G be solvable and $l_p(G/O_p(G)) \geq 1$. Then*

$$dl(G/O_p(G)) \leq 5l_p(G/O_p(G))\Omega_p(G).$$

Proof. We may assume without loss of generality that $O_p(G) = 1$. We use induction on $l_p(G)$. By Lemma 3.1, we can assume that $l_p(G) \geq 2$.

Write $M = O_{p',p}(G)$. Since $O_p(G) = 1$, $O_p(M) = 1$. Clearly, $l_p(M) = 1$. Thus $dl(M) \leq 5\omega_p(M)$ by Lemma 3.1. Since $M \triangleleft G$, $\omega_p(M) \leq \omega_p(G)$ by Clifford's Theorem. Hence $dl(M) \leq 5\omega_p(G)$.

Since $M = O_{p',p}(G)$, $O_p(G/M) = 1$ and $l_p(G/M) = l_p(G) - 1$. Notice that $1 \leq l_p(G/M) < l_p(G)$. Thus, by induction, $dl(G/M) \leq 5l_p(G/M)\omega_p(G/M)$. Since $l_p(G/M) = l_p(G) - 1$ and $\omega_p(G/M) \leq \omega_p(G)$, $dl(G/M) \leq 5(l_p(G) - 1)\omega_p(G)$. Hence,

$$\begin{aligned} dl(G) &\leq dl(M) + dl(G/M) \\ &\leq 5\omega_p(G) + 5(l_p(G) - 1)\omega_p(G) = 5l_p(G)\omega_p(G), \end{aligned}$$

and the assertion holds. \square

Combining Theorem 1.7 with Theorem 3.2, we get

COROLLARY 3.3. *Let G be solvable and $l_p(G/O_p(G)) \geq 1$. Then*

(1) *if p is not a Fermat prime, then*

$$dl(G/O_p(G)) \leq 5\omega_p(G)(1 + \log_p(2\omega_p(G)));$$

(2) *if p is a Fermat prime, then*

$$dl(G/O_p(G)) \leq 5\omega_p(G)[2 + \log_s(2\omega_p(G)/(p - 1))],$$

where $s = (p^2 - p + 1)/p$.

As usual, we denote by $F(G)$ the Fitting subgroup of G .

LEMMA 3.4. *Let G be solvable and $G/F(G) = F(G/F(G))$. Then*

$$dl(G/F(G)) \leq 2f(G)^2.$$

Proof. Let p be a prime number such that $p \mid |G|$. By Theorem 2.2, $dl_p(G/O_p(G)) \leq l_p(G/O_p(G))e_p(G)$. Combining $l_p(G/O_p(G)) \leq r_p(G/O_p(G))$ with Theorem 1.3, we have

$$dl_p(G/O_p(G)) \leq 2e_p(G)^2 \leq 2f(G)^2.$$

Since $dl_p(G/F(G)) = dl_p(G/O_p(G))$, $dl_p(G/F(G)) \leq 2f(G)^2$. Since $G/F(G) = F(G/F(G))$,

$$dl(G/F(G)) = \max\{dl_p(G/F(G)) \mid p \mid |G/F(G)|\}.$$

Thus $dl(G/F(G)) \leq 2f(G)^2$. \square

THEOREM 3.5. *Let G be solvable. Then*

$$dl(G/F(G)) \leq 2(f(G)^2 + f(G) + 1).$$

Proof. Let $F_2/F(G) = F(G/F(G))$. By Corollary 1.6, $r(G/F(G)) \leq 2f(G)$. We use Leisering and Manz [11, Lemma 2.3] to embed G/F_2 in the direct product of some $GL(2f(G), p)$, where p runs through the prime divisors of $|F_2/F(G)|$. Consequently, Theorem 2.5 of Leisering and Manz [11] yields that $dl(G/F_2) \leq 2f(G) + 2$.

Applying Lemma 3.4 to the group F_2 , we have $dl(F_2/F(F_2)) \leq 2f(F_2)^2$. Hence $dl(F_2/F(G)) \leq 2f(F_2)^2 \leq 2f(G)^2$. Finally,

$$\begin{aligned} dl(G/F(G)) &\leq dl(G/F_2) + dl(F_2/F(G)) \\ &\leq 2f(G) + 2 + 2f(G)^2 \\ &= 2(f(G)^2 + f(G) + 1). \end{aligned} \quad \square$$

Some remarks are appropriate for this theorem.

- (1) If $f(G) = 1$, then $dl(G/F(G)) \leq 2$.
- (2) If G has odd order, then $dl(G/F(G)) \leq f(G)^2 + f(G) + 2$.
- (3) Let $n(G)$ be the nilpotent length of G . Then $n(G) \leq 2(f(G) + 2)$.

Acknowledgment. This paper is a part of my dissertation at Ohio University. I would like to express my gratitude to my advisor Professor Thomas R. Wolf for his invaluable advice, patient guidance and encouragement.

I would like to thank the referee for the helpful suggestions for improving the exposition. These comments were greatly appreciated.

REFERENCES

- [1] D. Gluck, *Trivial set stabilizers in finite permutation groups*, Canad. J. Math., **35** (1983), 59–67.
- [2] D. Gluck and T. R. Wolf, *Brauer's height conjecture for p -solvable groups*, Trans. Amer. Math. Soc., **282** (1984), 137–152.
- [3] D. Gorenstein, *Finite Groups*, New York, 1968.
- [4] P. Hall and G. Higman, *On the p -length of p -solvable groups and reduction theorems for Burnside's problem*, Proc. London Math. Soc., **6** (1956), 1–40.
- [5] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin, 1967.
- [6] ———, *Inequalities for character degrees of solvable groups*, Arch. Math., **46** (1986), 387–392.
- [7] ———, *Solvable groups, all of whose irreducible representations in characteristic p have prime degrees*, J. Algebra, **104** (1986), 23–36.
- [8] I. M. Isaacs, *Lifting Brauer characters of p -solvable groups II*, J. Algebra, **51** (1978), 476–490.
- [9] ———, *Character Theory of Finite Groups*, Academic Press, New York, 1976.
- [10] I. M. Isaacs and D. Passman, *A characterization of groups in terms of the degrees of their characters II*, Pacific J. Math., **24** (1968), 467–510.

- [11] U. Leisering and O. Manz, *A note on character degrees of solvable groups*, Arch. Math., **48** (1987), 32–35.
- [12] O. Manz, *Degree problems: the p -rank in p -solvable groups*, Bull. London Math. Soc., **17** (1985), 545–548.
- [13] O. Manz and T. R. Wolf, *The q -parts of degrees of Brauer characters of solvable groups*, to appear in Illinois J. Math.
- [14] G. Michler, *A finite simple group of Lie-type has p -blocks with different defects, $p \neq 2$* , J. Algebra, **104** (1986), 220–230.
- [15] T. Okuyama, *On a problem of Wallace*, preprint.
- [16] D. Passman, *Groups with normal Hall- p' -subgroups*, Trans. Amer. Math. Soc., **123** (1966), 99–111.
- [17] T. R. Wolf, *Sylow- p -subgroups of p -solvable subgroups of $GL(n, p)$* , Arch. Math., **43** (1984), 1–10.

Received June 26, 1989 and in revised form November 3, 1989.

OHIO UNIVERSITY
ATHENS, OH 45701-2979

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906-1982)

B. H. NEUMANN

F. WOLF
(1904-1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1991 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 148, No. 2

April, 1991

Yuri A. Abramovich , Operators preserving disjointness on rearrangement invariant spaces	201
Andrew French Acker and Kirk Lancaster , Existence and geometry of a free boundary problem for the heat equation	207
So-Chin Chen , Real analytic regularity of the Szegő projection on circular domains	225
Chen-Lian Chuang , An independence property of central polynomials	237
Peter Larkin Duren and M. Schiffer , Robin functions and energy functionals of multiply connected domains	251
Johan Henricus Bernardus Kemperman , Sets of uniqueness and systems of inequalities having a unique solution	275
Ka-Lam Kueh , Fourier coefficients of nonholomorphic modular forms and sums of Kloosterman sums	303
Gerard J. Murphy , Ordered groups and crossed products of C^* -algebras ...	319
You-Qiang Wang , The p -parts of Brauer character degrees in p -solvable groups	351
Hidenobu Yoshida , Harmonic majorization of a subharmonic function on a cone or on a cylinder	369