

Pacific Journal of Mathematics

**CONJUGATES OF EQUIVARIANT HOLOMORPHIC MAPS OF
SYMMETRIC DOMAINS**

MIN HO LEE

CONJUGATES OF EQUIVARIANT HOLOMORPHIC MAPS OF SYMMETRIC DOMAINS

MIN HO LEE

In this paper we construct the conjugates of equivariant holomorphic maps of symmetric domains associated to morphisms of arithmetic varieties. We also prove that the conjugate of a Kuga fiber variety is another Kuga fiber variety.

0. Introduction. Let \mathbf{G} be a simply connected semisimple algebraic group over \mathbf{Q} that does not contain direct factors defined over \mathbf{Q} and compact over \mathbf{R} , and let K be a maximal compact subgroup of the semisimple Lie group $G = \mathbf{G}(\mathbf{R})$. We assume that the symmetric space $D = G/K$ has a complex structure. Let Γ be a torsion free arithmetic subgroup of \mathbf{G} and let $X = \Gamma \backslash D$ be the corresponding arithmetic variety. For each $\sigma \in \text{Aut}(X)$ it is known (cf. [5], [6], [7], [10]) that the conjugate X^σ of X is also an arithmetic variety.

Let \mathbf{G}' be another semisimple algebraic \mathbf{Q} -group, and consider the corresponding objects G', K', D', Γ' and X' as in the case of \mathbf{G} . Let $\rho: G \rightarrow G'$ be a homomorphism of Lie groups and $\tau: D \rightarrow D'$ a holomorphic map such that (ρ, τ) is an equivariant pair and $\rho(\Gamma) \subset \Gamma'$. Then τ induces the morphism $\phi: X \rightarrow X'$ of arithmetic varieties. Let D^σ and D'^σ be the universal covering spaces of X^σ and X'^σ respectively, and let $\tau^\sigma: D^\sigma \rightarrow D'^\sigma$ be the lifting of $\phi^\sigma: X^\sigma \rightarrow X'^\sigma$. Let G_0 and G'_0 be the connected components of the identity of $\text{Aut}(D^\sigma)$ and $\text{Aut}(D'^\sigma)$ respectively. If $\Gamma^\sigma \subset G_0$ and $\Gamma'^\sigma \subset G'_0$ are the fundamental groups of X^σ and X'^σ respectively, then we have the following result, Theorem 5.2 of this paper.

THEOREM. *There exist a finite covering G_1^σ of G_0^σ and a homomorphism $\rho_1^\sigma: G_1^\sigma \rightarrow G_1'^\sigma$ such that ρ_1^σ and τ^σ are equivariant and $\rho_1^\sigma(\Gamma^\sigma)$ is contained in Γ'^σ .*

As an application of this result we consider the conjugates of Kuga fiber varieties. Let $\mathbf{G}' = \text{Sp}(V, \beta)$ for some \mathbf{Q} -vector space V and a nondegenerate alternating bilinear form β , and assume that $X = \Gamma \backslash D$ is compact. Then from the equivariant pair (ρ, τ) we can construct a

Kuga fiber variety $\pi: Y \rightarrow X$ which is a fiber bundle such that X and Y are complex projective varieties, π is a morphism of varieties, and the fibers are polarized abelian varieties (see §6 for details). For each $\sigma \in \text{Aut}(\mathbb{C})$ we obtain the conjugate $\pi^\sigma: Y^\sigma \rightarrow X^\sigma$ of the Kuga fiber variety $\pi: Y \rightarrow X$. Then we have the following theorem, Theorem 6.3 of the text.

THEOREM. $\pi^\sigma: Y^\sigma \rightarrow X^\sigma$ is a Kuga fiber variety.

The above theorem is known when $\sigma: Y \rightarrow X$ is a family of abelian varieties associated to a PEL-type (cf. [14]) and it is also proved in [9] for Kuga fiber varieties constructed under certain assumptions. The theorem is also an immediate consequence of the main theorems in [5] and [10] in the case that ϕ arises from a homomorphism $\tilde{\rho}: \mathbf{G} \rightarrow \mathbf{G}'$ of algebraic groups defined over \mathbf{Q} and τ maps the CM-points of D to the CM-points of D' . In general, however, τ does not necessarily map the CM-points to CM-points (see [5, Proposition 1.11] for a necessary and sufficient condition for τ to map CM-points to CM-points).

This paper contains a part of the results in my Ph.D. thesis. I would like to thank my thesis advisor Professor Michio Kuga. I would also like to thank the referee for various helpful suggestions.

1. Prouniversal covering manifolds. In this section we shall review some of the results in [6] and [7]. Let \mathbf{G} be a simply connected semisimple algebraic group over \mathbf{Q} that does not contain direct factors defined over \mathbf{Q} and compact over \mathbf{R} , and let $G = \mathbf{G}(\mathbf{R})$ be the group of real elements of \mathbf{G} . If K is a maximal compact subgroup of G , we assume that the associated symmetric space $D = G/K$ has a G -invariant complex structure.

Let Γ be an arithmetic subgroup of \mathbf{G} that does not contain elements of finite order. Then the quotient space $X = \Gamma \backslash D$ has a natural structure of a complex manifold. Such complex manifolds are called arithmetic varieties. By a theorem of Baily and Borel ([3]) X has a structure of an algebraic variety over \mathbf{C} , and this structure is unique by a theorem of Borel ([4]).

Let $\{\Gamma_k | k = 1, 2, 3, \dots\}$ be an inductive system of subgroups of finite index of Γ such that each Γ_k is an arithmetic subgroup and

$$\Gamma_k \subset \Gamma_j \quad \text{for } j < k.$$

Then, for each k , the quotient space $X_k = \Gamma_k \backslash D$ is an arithmetic variety which is a finite unramified covering manifold of X , and the

collection $\{X_k | k \geq 1\}$ is a projective system of finite unramified covering manifolds of X . The projective limit

$$\widehat{D} = \varprojlim X_k$$

has a natural structure of a non-connected complex manifold, which does not depend on the representation of \widehat{D} as a projective limit. If

$$\hat{x} = \varprojlim x_k$$

is an element of \widehat{D} with $x_k \in X_k$ for each k , then the open neighborhoods of \hat{x} in \widehat{D} are determined as follows: If $\{U_k\}$ is a collection of open sets $U_k \subset X_k$ containing x_k for each k , then the open neighborhood of \hat{x} associated to $\{U_k\}$ is given by

$$N(\hat{x}, \{U_k\}) = \{\hat{y} = \varprojlim y_k \in \widehat{D} | y_k \in U_k \text{ for all } k\}.$$

Let $\pi: D \rightarrow X$ and $\pi_k: X_k \rightarrow X$ be the canonical projections. To construct a mapping $\hat{\mu}: D \rightarrow \widehat{D}$, we fix a point $x \in X$ and two elements $d_0 \in D$ and $\hat{d}_0 \in \widehat{D}$ such that

$$\pi(d_0) = \hat{\pi}(\hat{d}_0) = x.$$

Then for each k there is a unique map $\mu_k: D \rightarrow X_k$ such that $\pi_k \circ \mu_k = \pi$. We define the embedding $\hat{\mu}: D \rightarrow \widehat{D}$ by

$$\hat{\mu} = \varprojlim \mu_k.$$

PROPOSITION 1.1. (i) $\hat{\mu}(D)$ is a connected component of \widehat{D} . (ii) $\hat{\mu}(D)$ is dense in \widehat{D} .

Proof. (i) Let

$$\hat{d}_0 = \varprojlim d_{0k} \in \hat{\mu}(D) \subset \widehat{D}.$$

If

$$\hat{d}_1 = \varprojlim d_{1k} \in \widehat{D}$$

is in the connected component of \widehat{D} containing \hat{d}_0 , then there is a continuous map $\hat{c}: [0, 1] \rightarrow \widehat{D}$ such that $\hat{c}(0) = \hat{d}_0$ and $\hat{c}(1) = \hat{d}_1$. Then \hat{c} can be represented by

$$\varprojlim c_k,$$

where each c_k is a continuous map from $[0, 1]$ to X_k such that

$$\pi_{jk} \circ c_j = c_k \quad \text{for } j < k;$$

here the maps $\pi_{jk}: X_j \rightarrow X_k$ are the natural projections. If $c: [0, 1] \rightarrow D$ is the common lifting of the maps c_k , then $\hat{\mu}(c(0)) = \hat{d}_0$ and

$$\hat{\mu}(c(1)) = \hat{d}_1 \in \hat{\mu}(D).$$

Thus $\hat{\mu}(D)$ is a connected component of \hat{D} .

(ii) Let

$$\hat{x} = \varprojlim x_k \in \hat{D},$$

and let $N(\hat{x}, \{U_k\})$ be the neighborhood of \hat{x} associated to a collection $\{U_k\}$ of open sets with $x_k \in U_k \subset X_k$ for each k . If x_0 is an element of X with $\pi_k(x_k) = x_0$ for all k , we choose $y_k \in \pi_k^{-1}(x_0) \cap U_k$ for each k . If $y \in D$ is an element with $\mu_k(y) = y_k$ for all k , then $\hat{\mu}(y) \in N(\hat{x}, \{U_k\})$. Thus $\hat{\mu}(D)$ is dense in \hat{D} .

Let $\Gamma(X) \subset \text{Aut}(D)$ be the fundamental group of X and let

$$G_a(X) = \{g \in \text{Aut}(D) \mid [\Gamma(X): g\Gamma(X)g^{-1} \cap \Gamma(X)] < \infty \\ \text{and } [\Gamma(X): g^{-1}\Gamma(X)g \cap \Gamma(X)] < \infty\}.$$

The natural homomorphism $G \rightarrow \text{Aut}(D)$ induces the homomorphism

$$\alpha: \mathbf{G}(\mathbf{Q}) \rightarrow \text{Aut}(D).$$

Since \mathbf{G} has no factors defined over \mathbf{Q} and compact over \mathbf{R} , the kernel of α is the center $Z_{\mathbf{Q}}$ of $\mathbf{G}(\mathbf{Q})$ and therefore it is finite. The image of $\Gamma \subset \mathbf{G}(\mathbf{Q})$ under α coincides with $\Gamma(X)$. If $\hat{\pi}: \hat{D} \rightarrow X$ denotes the natural projection, we set

$$\hat{G}_a(X) = \text{Aut}(\hat{D}), \quad \hat{\Gamma}(X) = \{\hat{g} \in \hat{G}_a(X) \mid \hat{\pi} \circ \hat{g} = \hat{\pi}\}.$$

To define a homomorphism $\hat{\chi}: G_a(X) \rightarrow \hat{G}_a(X)$, we take an element $g \in G_a(X)$. For each k , g induces a map $\hat{\chi}_k(g): \hat{D} \rightarrow X_k$ (see [6, p. 158]). We set

$$\hat{\chi}(g) = \varprojlim \hat{\chi}_k(g) \in \text{Aut}(\hat{D}).$$

Then we have

$$\hat{\mu}(gd) = \hat{\chi}(g)\hat{\mu}(d) \quad \text{for } d \in D \text{ and } g \in G_a(X).$$

Since $\hat{\mu}(D)$ is dense in \hat{D} , we have

$$\hat{\chi}(g_1g_2) = \hat{\chi}(g_1)\hat{\chi}(g_2) \quad \text{for } g_1, g_2 \in G_a(X);$$

hence $\hat{\chi}: G_a(X) \rightarrow \hat{G}_a(X)$ is a homomorphism.

PROPOSITION 1.2. *Let \hat{d}_0 be an element of $\hat{\mu}(D)$. Then $\hat{\chi}$ is an isomorphism between $G_a(X)$ and the subgroup of $\hat{G}_a(X)$ consisting of all $\hat{g} \in \hat{G}_a(X)$ such that $\hat{d}_0 \hat{g} \in \hat{\mu}(D)$.*

Proof. See [6, Lemma 4].

The group $\hat{G}_a(X)$ is a complete locally compact topological group relative to the topology in which a basis of neighborhoods of the identity consists of the subgroups of finite index in $\hat{\Gamma}(X)$.

PROPOSITION 1.3. (i) $G_a(X)$ is dense in $\hat{G}_a(X)$.
 (ii) $\hat{\chi}$ induces an isomorphism between the double cosets $\Gamma(X) \backslash G_a(X) / \Gamma(X)$ and $\hat{\Gamma}(X) \backslash \hat{G}_a(X) / \hat{\Gamma}(X)$.

Proof. (i) See [6, Theorem 1].

(ii) This follows from [6, Lemma 3 and Lemma 3'].

Let $\sigma \in \text{Aut}(\mathbb{C})$. Then we consider the complex variety X^σ obtained from X by the base change. Let D^σ be the universal covering manifold of X^σ , and let

$$\Gamma^\sigma = \Gamma(X^\sigma) \subset \text{Aut}(D^\sigma)$$

be the fundamental group of X^σ . If the varieties X_k^σ are the conjugates of X_k , we set

$$\hat{D}^\sigma = \varprojlim X_k^\sigma, \quad \hat{G}_a(X^\sigma) = \text{Aut}(\hat{D}^\sigma).$$

Then $\hat{G}_a(X^\sigma)$ is a complete locally compact topological group in the topology of subgroups of finite index in

$$\hat{\Gamma}(X^\sigma) = \{ \hat{g}^\sigma \in \hat{G}_a(X^\sigma) \mid \hat{\pi}^\sigma \circ \hat{g}^\sigma = \hat{\pi}^\sigma \},$$

where $\hat{\Gamma}^\sigma: \hat{D}^\sigma \rightarrow X^\sigma$ is the natural projection. As in the case of X , we can construct the homomorphism $\hat{\chi}^\sigma: G_a(X^\sigma) \rightarrow \hat{G}_a(X^\sigma)$, where

$$G_a(X^\sigma) = \{ g^\sigma \in \text{Aut}(D^\sigma) \mid [\Gamma^\sigma: g^\sigma \Gamma^\sigma (g^\sigma)^{-1} \cap \Gamma^\sigma] < \infty \text{ and } [\Gamma^\sigma: (g^\sigma)^{-1} \Gamma^\sigma g^\sigma \cap \Gamma^\sigma] < \infty \}.$$

Let $V \subset G_a(X)$ be a subgroup with $[\Gamma(X): V \cap \Gamma(X)] < \infty$, and let \hat{V} be the closure of $\hat{\chi}(V)$ in $\hat{G}_a(X)$. If $\hat{g} \in \hat{V} \subset \hat{G}_a(X)$, then there are morphisms $g_k: Z_k \rightarrow X_k$ such that

$$\varprojlim Z_k = \hat{D} = \varprojlim X_k, \quad \hat{g} = \varprojlim g_k.$$

Applying σ to g_k , we obtain morphisms $g_k^\sigma: Z_k^\sigma \rightarrow X_k^\sigma$. Then

$$\hat{g}^\sigma = \varprojlim g_k^\sigma$$

is an element of $\widehat{G}(X^\sigma) = \text{Aut}(\widehat{D}^\sigma)$. We set $\widehat{V}^\sigma = \{\hat{g}^\sigma | \hat{g} \in \widehat{V}\}$ and define the subgroup V^σ of $G_a(X^\sigma)$ by

$$V^\sigma = (\hat{\chi}^\sigma)^{-1}(\widehat{V}^\sigma \cap \text{Im}(\hat{\chi}^\sigma)).$$

Now let G_a be a subgroup of finite index in $\alpha(\mathbf{G}(\mathbf{Q}))$ containing $\alpha(\Gamma) = \Gamma(X)$, and let

$$G_a^\sigma = (G_a)^\sigma \subset G_a(X^\sigma).$$

PROPOSITION 1.4. *G_a^σ is dense in the connected component of the identity of $\text{Aut}(D^\sigma)$ in the ordinary topology.*

Proof. G_a^σ is contained in the connected component of the identity of $\text{Aut}(D^\sigma)$ by [10, Lemma 3.7]. The density follows from [6, Theorem 5] (see also [6, Theorem A.7]).

THEOREM 1.5. *The group $\Gamma^\sigma = \Gamma(X^\sigma)$ is an arithmetic subgroup of the connected component of the identity of $\text{Aut}(D^\sigma)$.*

Proof. This follows from the main theorems in [6] and [7].

2. The homomorphism $\hat{\rho}$. Let \mathbf{G} , G , K , D , Γ and X be as in §1. We consider another semisimple algebraic \mathbf{Q} -group \mathbf{G}' and its arithmetic subgroup $\Gamma' \subset \mathbf{G}'(\mathbf{Q})$ that is torsion free. As in the case of \mathbf{G} , we associate G' , K' , D' and the arithmetic variety $X' = \Gamma' \backslash D'$ to \mathbf{G}' . Let $\rho: G \rightarrow G'$ be a homomorphism, $\phi: X \rightarrow X'$ a morphism of varieties, and $\tau: D \rightarrow D'$ a holomorphic lifting of ϕ such that ρ and τ are equivariant, i.e.,

$$\tau(gy) = \rho(g)\tau(y) \quad \text{for all } g \in G \text{ and } y \in D.$$

Let $\{\Gamma_k\}$, $\{X_k\}$ and \widehat{D} be as in §1, and let $\{\Gamma'_k\}$ be an inductive system of arithmetic subgroups of finite index of Γ' such that $\rho(\Gamma_k) \subset \Gamma'_k$ for each $k \geq 1$. The quotient spaces $X'_k = \Gamma'_k \backslash D'$ are arithmetic varieties and they form a projective system $\{X'_k\}$ of covering manifolds of X' . The holomorphic map $\tau: D \rightarrow D'$ induces a

morphism $\phi: X_k \rightarrow X'_k$ for each $k \geq 1$. We set

$$\widehat{D}' = \varprojlim X'_k, \quad \widehat{G}_a(X') = \text{Aut}(\widehat{D}'),$$

and define the holomorphic map $\hat{\tau}: \widehat{D} \rightarrow \widehat{D}'$ by

$$\hat{\tau} = \varprojlim \phi_k.$$

For each $\hat{g} \in \widehat{G}_a \subset \widehat{G}_a(X) = \text{Aut}(\widehat{D})$, there are elements

$$g_k \in \alpha^{-1}(G_a) \subset \mathbf{G}(\mathbf{Q})$$

(recall that $\text{Ker}(\alpha) = Z_{\mathbf{Q}}$ is finite) such that

$$\hat{g} = \varprojlim p_k(g_k),$$

where the morphisms $p_k(g_k)$ are determined as follows: Let

$$\begin{aligned} \Gamma_{g,k} &= \Gamma_k \cap g_k^{-1} \Gamma_k g_k, \\ X_{g,k} &= \Gamma_{g,k} \backslash D. \end{aligned}$$

Then, for each k , we define

$$p_k(g_k): X_{g,k} \rightarrow X_k$$

to be the morphism of arithmetic varieties induced by the left multiplication $g_k: D \rightarrow D$, $D = G/K$. We have

$$g_k = \gamma_{j,k} g_j \quad \text{with } \gamma_{j,k} \in \Gamma_j \text{ for } j \leq k.$$

Recall that $\{\Gamma'_k\}$ is an inductive system of arithmetic subgroups of finite index of Γ' such that $\rho(\Gamma_k) \subset \Gamma'_k$ for each k . We set

$$\begin{aligned} \Gamma'_{g,k} &= \Gamma'_k \cap \rho(g_k)^{-1} \Gamma'_k \rho(g_k), \\ X'_{g,k} &= \Gamma'_{g,k} \backslash D' \end{aligned}$$

for each k . Then we have $\rho(\Gamma_{g,k}) \subset \Gamma'_{g,k}$, and hence the left multiplication maps $\rho(g_k): D' \rightarrow D'$ induce the morphisms

$$p'_k(\rho g_k): X'_{g,k} \rightarrow X'_k.$$

Since we have

$$\rho(g_k) = \rho(\gamma_{j,k} g_j) = \rho(\gamma_{j,k}) \rho(g_j)$$

with

$$\rho(\gamma_{j,k}) \in \rho(\Gamma_j) \subset \Gamma'_j \quad \text{for } j \leq k,$$

the collection $\{\rho(g_k)\}$ of liftings $\rho(g_k)$ of $p'_k(\rho g_k)$ determines the projective limit

$$\varprojlim p'_k(\rho g_k) \in \widehat{G}_a(X') = \text{Aut}(\widehat{D}').$$

Thus we obtain a homomorphism $\hat{\rho}: \widehat{G}_a \rightarrow \widehat{G}'_a$ defined by

$$\hat{\rho}(\hat{g}) = \varprojlim p'_k(\rho g_k)$$

for all $\hat{g} = \varprojlim g_k \in \widehat{G}_a$.

PROPOSITION 2.1. *If $\hat{\tau}: \widehat{D} \rightarrow \widehat{D}'$ and $\hat{\rho}: \widehat{G}_a \rightarrow \widehat{G}'_a(X')$ are as described above, then*

$$\hat{\tau}(\hat{g}\hat{y}) = \hat{\rho}(\hat{g})\hat{\tau}(\hat{y}) \quad \text{for all } \hat{g} \in \widehat{G}_a \text{ and } \hat{y} \in \widehat{D}.$$

Proof. If $\mu_k: D \rightarrow X_k$, $\mu_{g,k}: D \rightarrow X_{g,k}$, $\mu'_k: D' \rightarrow X'_k$ and $\mu'_{g,k}: D' \rightarrow X'_{g,k}$ are the natural covering maps, then we have

$$\begin{aligned} \mu'_k \tau(g_k y) &= \phi_k(g_k \mu_{g,k}(y)), \\ \mu'_k \rho(g_k) \tau(y) &= (p'_k(\rho g_k)) \phi_{g,k}(\mu_{g,k}(y)) \end{aligned}$$

for all $y \in D$, where

$$\phi_{g,k}: X_{g,k} \rightarrow X'_{g,k}$$

is the morphism induced by $\tau: D \rightarrow D$ for each k with

$$\tau = \varprojlim \phi_{g,k}.$$

Since $\tau(g_k y) = \rho(g_k) \tau(y)$ for all $y \in D$, we have

$$\phi_k(g_k \mu_{g,k}(y)) = (p'_k(\rho g_k)) \phi_{g,k}(\mu_{g,k}(y))$$

for all $y \in D$. Since $\mu_{g,k}: D \rightarrow X_{g,k}$ is surjective and

$$\hat{\rho}(\hat{g}) = \varprojlim p'_k(\rho g_k),$$

it follows that

$$\hat{\tau}(\hat{g}\hat{y}) = \hat{\rho}(\hat{g})\hat{\tau}(\hat{y}) \quad \text{for all } \hat{y} = \varprojlim (\mu_{g,k}(y)) \in \widehat{D}.$$

3. Conjugates of $\hat{\rho}$. Let $\{X_k\}$ and $\{X'_k\}$ be the projective systems of finite unramified covering manifolds of X and X' , respectively, considered in §2. We fix an element $\sigma \in \text{Aut}(\mathbf{C})$. The varieties X^σ , X'^σ , X_k^σ and $X'_k{}^\sigma$ are arithmetic varieties, and the collections

$\{X_k^\sigma\}$ and $\{X'_k{}^\sigma\}$ are projective systems of finite unramified covering manifolds of X^σ and X'^σ respectively (see [5], [6], [7], [10]). Let $\phi_k^\sigma: X_k^\sigma \rightarrow X'_k{}^\sigma$ be the conjugate morphism of $\phi_k: X_k \rightarrow X'_k$ for each k . Let \widehat{D} , $\widehat{G}_a(X^\sigma)$ and $\widehat{\Gamma}(X^\sigma)$ be as in §1, and let

$$\widehat{D}'^\sigma = \varprojlim X'_k{}^\sigma, \quad \widehat{G}_a(X'^\sigma) = \text{Aut}(\widehat{D}'^\sigma),$$

$$\widehat{\Gamma}(X'^\sigma) = \{\hat{g}'^\sigma \in \widehat{G}_a(X'^\sigma) \mid \hat{\pi}'^\sigma \circ \hat{g}'^\sigma = \hat{\pi}'^\sigma\},$$

where $\hat{\pi}'^\sigma: \widehat{D}'^\sigma \rightarrow X'^\sigma$ is the natural projection. We define the map $\hat{\tau}^\sigma: \widehat{D}^\sigma \rightarrow \widehat{D}'^\sigma$ by

$$\hat{\tau}^\sigma = \varprojlim \phi_k^\sigma.$$

If $\hat{h} \in \widehat{G}_a^\sigma$, then there exist a projective system $\{Z_k\}$ of finite unramified covering manifolds of X^σ and morphisms $h_k: Z_k \rightarrow X_k^\sigma$ such that

$$\varprojlim Z_k = \widehat{D}^\sigma = \varprojlim X_k^\sigma, \quad \hat{h} = \varprojlim h_k.$$

The morphisms h_k induce the morphisms

$$h_k^{\sigma^{-1}}: Z_k^{\sigma^{-1}} \rightarrow X_k.$$

Let $\tilde{h}_k^{\sigma^{-1}}: D \rightarrow D$ be an element of $\alpha^{-1}(G_a(X)) \subset \mathbf{G}(\mathbf{Q})$ that is a lifting of $h_k^{\sigma^{-1}}$ for each k . We define the homomorphism $\hat{\rho}^\sigma: \widehat{G}_a^\sigma \rightarrow \widehat{G}_a(X'^\sigma)$ by

$$\hat{\rho}^\sigma(\hat{h}) = \varprojlim (p'_k(\rho(\hat{h}_k^{\sigma^{-1}})))^\sigma$$

where

$$p'_k(\rho(\tilde{h}_k^{\sigma^{-1}})): W'_k \rightarrow X'_k$$

is the morphism induced from $\rho(\tilde{h}_k^{\sigma^{-1}}): D' \rightarrow D'$ for each k .

PROPOSITION 3.1. $\hat{\tau}^\sigma(\hat{g}^\sigma \hat{y}^\sigma) = \hat{\rho}^\sigma(\hat{g}^\sigma) \hat{\tau}^\sigma(\hat{y}^\sigma)$ for all $\hat{g}^\sigma \in \widehat{G}_a^\sigma$ and $\hat{y}^\sigma \in \widehat{D}^\sigma$.

Proof. For each k the relation $\hat{\tau}(\hat{g}\hat{y}) = \hat{\rho}(\hat{g})\hat{\tau}(\hat{y})$ induces the following commutative diagram:

$$\begin{array}{ccc} X_{g,k} & \xrightarrow{\phi_{g,k}} & X'_{g,k} \\ p_k(g_k) \downarrow & & \downarrow p'_k(\rho g_k) \\ X_k & \xrightarrow{\phi_k} & X'_k \end{array}$$

Applying σ to this diagram, we obtain

$$\begin{array}{ccc} X_{g,k}^\sigma & \xrightarrow{\phi_{g,k}^\sigma} & X'_{g,k}^\sigma \\ (p_k(g_k))^\sigma \downarrow & & \downarrow (p'_k(\rho g_k))^\sigma \\ X_k^\sigma & \xrightarrow{\phi_k^\sigma} & X'_k{}^\sigma \end{array}$$

so that

$$\phi_k^\sigma((p_k(g_k))^\sigma x_k^\sigma) = (p'_k(\rho g_k))^\sigma \phi_{g,k}^\sigma(x_k^\sigma)$$

for all $x_k^\sigma \in X_{g,k}^\sigma$. By taking projective limits we obtain

$$\hat{\tau}^\sigma(\hat{g}^\sigma \hat{y}^\sigma) = \hat{\rho}^\sigma(\hat{g}^\sigma) \hat{\tau}^\sigma(\hat{y}^\sigma)$$

for all $\hat{y}^\sigma \in \hat{D}^\sigma$.

PROPOSITION 3.2. $\hat{\rho}(\hat{\Gamma}(X^\sigma))$ is contained in $\hat{\Gamma}(X'^\sigma)$.

Proof. If $\hat{h} \in \hat{\Gamma}(X^\sigma) \subset \hat{G}_a(X^\sigma)$, there are morphisms $h_k: Z_k \rightarrow X_k^\sigma$ such that

$$\varprojlim Z_k = \hat{D}^\sigma = \varprojlim X_k^\sigma, \quad \hat{h} = \varprojlim h_k \quad \text{and} \quad \pi_k^\sigma \circ h_k = \pi_k^0,$$

where $\pi_k^0: Z_k \rightarrow X^\sigma$ is the natural projection. Applying σ^{-1} , we obtain the morphisms $h_k^{\sigma^{-1}}: Z_k^{\sigma^{-1}} \rightarrow X_k$ such that

$$\pi_k \circ h_k^{\sigma^{-1}} = (\pi_k^0)^{\sigma^{-1}} \quad \text{for all } k,$$

where the maps $(\pi_k^0)^{\sigma^{-1}}: Z_k \rightarrow X$ are the natural projections. If the maps $\tilde{h}_k^{\sigma^{-1}}: D \rightarrow D$ with $\tilde{h}_k^{\sigma^{-1}} \in \mathbf{G}(\mathbf{Q})$ are as before, we have $\tilde{h}_k^{\sigma^{-1}}(\Gamma) \subset \Gamma$, and therefore

$$\rho(\tilde{h}_k^{\sigma^{-1}})(\Gamma') \subset \Gamma' \quad \text{for all } k.$$

Thus the morphisms $p'_k(\rho(\tilde{h}_k^{\sigma^{-1}})): W'_k \rightarrow X'_k$ satisfy

$$\pi_k'^0 \circ p'_k(\rho(\tilde{h}_k^{\sigma^{-1}})) = \pi_k' \quad \text{for all } k,$$

where the maps $\pi_k'^0: W'_k \rightarrow X'$ are the natural projections. Hence we have

$$(\pi_k'^0)^\sigma \circ (p'_k(\rho(\tilde{h}_k^{\sigma^{-1}})))^\sigma = (\pi_k')^\sigma$$

for all k . Thus it follows that $\hat{\rho}^\sigma(\hat{h}) \in \hat{\Gamma}(X'^\sigma)$.

4. The homomorphism ρ^σ . Let $X^\sigma, D^\sigma, \Gamma^\sigma, G_a^\sigma, G_a(X^\sigma), \widehat{D}^\sigma, \widehat{\Gamma}(X^\sigma), \widehat{G}_a(X^\sigma)$ be as in §1, and let $\widehat{D}'^\sigma, \widehat{\Gamma}_a(X'^\sigma), \widehat{G}(X'^\sigma)$ be as in §3. Let D'^σ be the universal covering space of X'^σ , and let

$$\Gamma'^\sigma = \Gamma(X'^\sigma) \subset \text{Aut}(D'^\sigma)$$

be the fundamental group of X'^σ . We set

$$G_a(X'^\sigma) = \{g'^\sigma \in \text{Aut}(D'^\sigma) \mid [\Gamma'^\sigma : g'^\sigma \Gamma'^\sigma (g'^\sigma)^{-1} \cap \Gamma'^\sigma] < \infty \\ \text{and } [\Gamma'^\sigma : (g'^\sigma)^{-1} \Gamma'^\sigma g'^\sigma \cap \Gamma'^\sigma] < \infty\}.$$

Since the map $\hat{\mu}: D \rightarrow \widehat{D}$ and the homomorphism $\hat{\chi}: G_a(X) \rightarrow \widehat{G}_a(X)$ described in §1 are injective, we shall identify each of $D^\sigma, D'^\sigma, G_a(X^\sigma), G_a(X'^\sigma)$ with its embedded image in $\widehat{D}^\sigma, \widehat{D}'^\sigma, \widehat{G}_a(X^\sigma), \widehat{G}_a(X'^\sigma)$ respectively. By Proposition 1.1(i) and the above identification, D^σ and D'^σ are connected components of \widehat{D}^σ and \widehat{D}'^σ respectively. We shall assume that D'^σ is the connected component of \widehat{D}'^σ containing the image of D^σ under the map τ^σ .

PROPOSITION 4.1. *If D'^σ is the connected component of \widehat{D}'^σ chosen as above, then $\hat{\rho}^\sigma(G_a^\sigma)$ is contained in $G_a(X'^\sigma)$.*

Proof. Let $d_0^\sigma \in D^\sigma \subset \widehat{D}^\sigma$. Then we have

$$\hat{\tau}^\sigma(d_0^\sigma) \in D'^\sigma \subset \widehat{D}'^\sigma.$$

By Proposition 1.2 we have

$$G_a^\sigma \subset G_a(X^\sigma) = \{\hat{g}^\sigma \in \widehat{G}_a(X^\sigma) \mid \hat{g}^\sigma d_0^\sigma \in D^\sigma\}, \\ G_a(X'^\sigma) = \{\hat{g}'^\sigma \in \widehat{G}_a(X'^\sigma) \mid \hat{g}'^\sigma \hat{\tau}^\sigma(d_0^\sigma) \in D'^\sigma\}.$$

If $g^\sigma \in G_a^\sigma$, then $g^\sigma d_0^\sigma \in D^\sigma$; hence from the continuity of $\hat{\tau}^\sigma$ it follows that

$$\hat{\tau}^\sigma(g^\sigma d_0^\sigma) \in \hat{\tau}^\sigma(D^\sigma) \subset D'^\sigma.$$

Thus we get

$$\hat{\rho}^\sigma(g^\sigma) \hat{\tau}^\sigma(d_0^\sigma) = \hat{\tau}^\sigma(g^\sigma d_0^\sigma) \in D'^\sigma.$$

Therefore it follows that

$$\hat{\rho}^\sigma(g^\sigma) \in G_a(X'^\sigma) \quad \text{for all } g^\sigma \in G_a^\sigma.$$

Now we define the homomorphism $\rho^\sigma: G_a^\sigma \rightarrow G_a(X'^\sigma)$ and $\tau^\sigma: D^\sigma \rightarrow D'^\sigma$ by

$$\rho^\sigma = \hat{\rho}^\sigma|_{G_a^\sigma}, \quad \tau^\sigma = \hat{\tau}^\sigma|_{D^\sigma}.$$

Then ρ^σ and τ^σ satisfy the relation

$$\tau^\sigma(g^\sigma d^\sigma) = \rho^\sigma(g^\sigma)\tau^\sigma(d^\sigma)$$

for all $g^\sigma \in G_a^\sigma$ and $d^\sigma \in D^\sigma$.

5. Conjugates of equivariant pairs. Let $\rho: G \rightarrow G'$ and $\tau: D \rightarrow D'$ be equivariant as before so that we have

$$\tau(gz) = \rho(g)\tau(z)$$

for all $g \in G$ and $z \in D$. Also as before let $\Gamma \subset G$ and $\Gamma' \subset G'$ be torsion free arithmetic subgroups with $\rho(\Gamma) \subset \Gamma'$, and let $\phi: X \rightarrow X'$ be the morphism induced by τ where $X = \Gamma \backslash D$ and $X' = \Gamma' \backslash D'$. If $\rho^\sigma: G_a^\sigma \rightarrow G_a'^\sigma$ and $\tau^\sigma: D^\sigma \rightarrow D'^\sigma$ are as in §4, we denote by $G_0^\sigma, G_0'^\sigma$ the connected components of the identity of $\text{Aut}(D^\sigma), \text{Aut}(D'^\sigma)$ respectively. By Proposition 1.4 G_a^σ is dense in G_0^σ ; hence by Proposition 3.1 and Proposition 4.1, we have

$$\tau^\sigma(hy) = \rho^\sigma(h)\tau^\sigma(y)$$

for all $h \in G_a^0$ and $y \in D^\sigma$. We set

$$H = \{(g, g') \in G_0^\sigma \times G_0'^\sigma \mid \tau^\sigma(gy) = g'\tau^\sigma(y) \text{ for all } y \in D^\sigma\}.$$

Since the set $\{(g, \rho^\sigma(g)) \mid g \in G_a^\sigma\}$ is contained in H and G_a^σ is dense in G_0^σ , the projection map $\text{pr}_1: H \rightarrow G_0^\sigma$ is surjective.

PROPOSITION 5.1. *H is a reductive Lie group.*

Proof. Let K be the kernel of the projection map $\text{pr}_1: H \rightarrow G_0^\sigma$. Then we have

$$\begin{aligned} K &= \{(1, g') \in G_0^\sigma \times G_0'^\sigma \mid \tau^\sigma(y) = g'\tau^\sigma(y) \text{ for all } y \in D^\sigma\} \\ &\cong \{g' \in G_0'^\sigma \mid \tau^\sigma(y) = g'\tau^\sigma(y) \text{ for all } y \in D^\sigma\} = \bigcap_{y \in D^\sigma} \text{Iso}(\tau^\sigma(y)), \end{aligned}$$

where $\text{Iso}(\tau^\sigma(y))$ is the isotropy subgroup of $\tau^\sigma(y)$ in $G_0'^\sigma$. Hence K is a compact Lie group, and therefore a reductive Lie group. Since there is an exact sequence

$$1 \rightarrow K \rightarrow H \xrightarrow{\text{pr}_1} G_0^\sigma \rightarrow 1$$

of Lie groups with K and G_0^σ reductive, it follows that H is reductive.

THEOREM 5.2. *There exist a finite covering G_1^σ of G_0^σ and a homomorphism $\rho_1^\sigma: G_1^\sigma \rightarrow G_0^\sigma$ of Lie groups such that ρ_1^σ and τ^σ are equivariant and $\rho_1^\sigma(\Gamma^\sigma)$ is contained in Γ^σ .*

Proof. Decompose the reductive Lie group H into an almost direct product of simple Lie groups and simple tori, and let G_1^σ be the product of the simple factors of H which map nontrivially to G_0^σ (see [2, III.2.2] for a similar argument). Then the kernel of the map from G_1^σ to G_0^σ is finite, and hence G_1^σ is a finite covering Lie group of G_0^σ . We define the action of G_1^σ on D^σ and the homomorphism $\rho_1^\sigma: G_1^\sigma \rightarrow G_0^\sigma$ by

$$(g, g') \cdot y = gy, \quad \rho_1^\sigma(g, g') = g'$$

for all $y \in D^\sigma$ and $(g, g') \in G_1^\sigma \subset H \subset G_0^\sigma \times G_0^{\prime\sigma}$. Then we have

$$\tau^\sigma((g, g') \cdot y) = \tau^\sigma(gy) = g'\tau^\sigma(y) = \rho_1^\sigma(g, g')\tau^\sigma(y)$$

for all $(g, g') \in G_1^\sigma$ and $y \in D^\sigma$. Hence we obtain a homomorphism of Lie groups ρ_1^σ from a finite covering G_1^σ of G_0^σ to $G_0^{\prime\sigma}$ such that

$$\tau^\sigma(h_1 y) = \rho_1^\sigma(h_1)\tau^\sigma(y)$$

for all $h_1 \in G_1^\sigma$ and $y \in D^\sigma$. Thus it follows that ρ_1^σ and τ^σ are equivariant. Now it remains to show that $\rho_1^\sigma(\Gamma^\sigma)$ is contained in Γ^σ . By Proposition 3.2, $\hat{\rho}(\widehat{\Gamma}(X^\sigma))$ is contained in $\widehat{\Gamma}(X^{\prime\sigma})$; hence $\hat{\rho}^\sigma: \widehat{G}_a^\sigma \rightarrow \widehat{G}_a(X^{\prime\sigma})$ induces the mapping

$$\hat{\rho}^\sigma: \widehat{\Gamma}(X^\sigma) \backslash \widehat{G}_a^\sigma / \widehat{\Gamma}(X^\sigma) \rightarrow \widehat{\Gamma}(X^{\prime\sigma}) \backslash \widehat{G}_a(X^{\prime\sigma}) / \widehat{\Gamma}(X^{\prime\sigma})$$

of double cosets. By Proposition 1.3(ii) $\hat{\rho}^\sigma$ induces the mapping

$$\hat{\rho}^\sigma: \Gamma(X^\sigma) \backslash G_a^\sigma / \Gamma(X^\sigma) \rightarrow \Gamma(X^{\prime\sigma}) \backslash G_a(X^{\prime\sigma}) / \Gamma(X^{\prime\sigma}).$$

Thus we have

$$\hat{\rho}(\Gamma(X^\sigma)) = \rho^\sigma(\Gamma^\sigma) \subset \Gamma(X^{\prime\sigma}) = \Gamma^{\prime\sigma}.$$

Since $\rho_1^\sigma(\Gamma^\sigma) = \rho^\sigma(\Gamma^\sigma)$, it follows that $\rho_1^\sigma(\Gamma^\sigma) \subset \Gamma^{\prime\sigma}$.

6. Conjugates of Kuga fiber varieties. First, we shall describe the construction of Kuga fiber varieties (see [1], [8], [11, Chapter 4] for details). Let V be a \mathbf{Q} -vector space of dimension $2n$, and let L be a lattice in V . Let β be a nondegenerate alternating bilinear form on V such that $\beta(L, L) \subset \mathbf{Z}$. Let

$$\mathrm{Sp}(V, \beta) = \{g \in \mathrm{GL}(V) \mid \beta(gx, gy) = \beta(x, y) \text{ for all } x, y \in V\}$$

be the symplectic group of the pair (V, β) , and let \mathcal{H} denote the Siegel space

$$\mathcal{H} = \{J \in \text{GL}(V_{\mathbf{R}}) \mid J^2 = -1, \beta(x, Jy) \text{ is a positive definite symmetric bilinear form in } x, y \in V_{\mathbf{R}}\}.$$

Then each element $J \in \mathcal{H}$ defines a complex structure on $V_{\mathbf{R}}$ and there is a unique complex analytic structure on $\mathcal{H} \times V_{\mathbf{R}}$ such that the projection $P: \mathcal{H} \times V_{\mathbf{R}} \rightarrow \mathcal{H}$ is a complex vector bundle over \mathcal{H} . For each J if we denote the complex vector space $(V_{\mathbf{R}}, J)$ by V_J , then the complex torus $A_J = V_J/L$ is an abelian variety with the polarization β . We set

$$A_{\mathcal{H}} = L \backslash \mathcal{H} \times V_{\mathbf{R}},$$

where the action of L on $\mathcal{H} \times V_{\mathbf{R}}$ is given by

$$l \cdot (J, v) = (J, v + l) \quad \text{for } j \in \mathcal{H} \text{ and } l \in L.$$

Then the vector bundle $P: \mathcal{H} \times V_{\mathbf{R}} \rightarrow \mathcal{H}$ induces the fiber bundle $\pi_{\mathcal{H}}: A_{\mathcal{H}} \rightarrow \mathcal{H}$ whose fibers are abelian varieties polarized by β . We set

$$\text{Sp}(L, \beta) = \{g \in \text{Sp}(V, \beta) \mid gL = L\},$$

and take a subgroup Γ_S of $\text{Sp}(L, \beta)$ of finite index that contains no elements of finite order. Then, as discussed in §1, the quotient $X = \Gamma_S \backslash \mathcal{H}$ is an arithmetic variety that can be considered as a Zariski open subset of a complex projective variety. Now the fiber bundle $\pi_{\mathcal{H}}: A_{\mathcal{H}} \rightarrow \mathcal{H}$ induces the standard family of abelian varieties $\pi_S: Y_S \rightarrow X_S$ over X_S .

Let \mathbf{G} , G and $D = G/K$ be as in §1. Let $\rho: G \rightarrow \text{Sp}(V, \beta)(\mathbf{R})$ be a symplectic representation of G , and let Γ be a torsion free arithmetic subgroup of \mathbf{G} with $\rho(\Gamma) \subset \Gamma_S$. We shall assume that the quotient $X = \Gamma \backslash D$ is compact. Let $\tau: D \rightarrow \mathcal{H}$ be a holomorphic map such that ρ and τ are equivariant, and let $\phi: X \rightarrow X_S$ be the morphism of varieties induced by τ . By pulling back the fiber bundle $\pi_S: Y_S \rightarrow X_S$ via the morphism $\phi: X \rightarrow X_S$, we obtain the fiber bundle over the arithmetic variety X whose fibers are abelian varieties polarized by β . It is known (cf. [8], [11, Theorem 8.6]) that the fiber space Y of the family of abelian varieties $\pi: y \rightarrow X$ has a structure of a complex projective variety. The fiber space Y (or the fiber bundle $\pi: Y \rightarrow X$ itself) is called a Kuga fiber variety.

To consider conjugates of Kuga fiber varieties, we shall first state some of the known results of G. Shimura about the families of abelian varieties associated to PEL-types (see e.g. [12], [15] for details).

THEOREM 6.1. *Let Ω be a normal admissible PEL-type (see [14] for its definition). Then there exists a family of abelian varieties $f: W \rightarrow U$ denoted by $\mathcal{F}(\Omega)$ with the following properties:*

(i) *For each $u \in U$, the fiber $f^{-1}(u)$ of $\mathcal{F}(\Omega)$ has a PEL-structure Q_u of type Ω .*

(ii) *For each PEL-structure Q of type Ω , there exists one and only one point $u \in U$ such that Q is isomorphic to Q_u .*

(iii) *U and W are Zariski open subset of projective varieties and f is a morphism of varieties.*

(iv) *There exists a number field k_Ω such that U, W and f are defined over k_Ω .*

(v) *For each $\sigma \in \text{Aut}(\mathbb{C})$ there exists a PEL-type Ω^σ and an isomorphism of $\mathcal{F}(\Omega)^\sigma$ to $\mathcal{F}(\Omega^\sigma)$ defined over k_{Ω^σ} .*

Proof. See [12, Theorem 5.3] and [14, Proposition 3.1]; see also [5, §8].

THEOREM 6.2. *Let Ω be as in Theorem 6.1, and let $f: W \rightarrow U$ be a family of abelian varieties associated to Ω . Let $f': W' \rightarrow U'$ be another family of abelian varieties such that each fiber $(f')^{-1}(u)$ for $u \in U'$ has a PEL-structure Q'_u . Suppose that at least one Q'_u is of type Ω and that the family $\{Q'_u | u \in U'\}$ of PEL-structures satisfies the property (ii) in Theorem 6.1. Then there is a biregular isomorphism between $f: W \rightarrow U$ and $f': W' \rightarrow U'$.*

Proof. See [12, Theorem 5.4 and Theorem 5.5].

THEOREM 6.3. *If $\pi: Y \rightarrow X$ is a Kuga fiber variety and $\sigma \in \text{Aut}(\mathbb{C})$, then the conjugate $\pi^\sigma: Y^\sigma \rightarrow X^\sigma$ is also a Kuga fiber variety.*

Proof. Let $G_S = \text{Sp}(V, \beta)(\mathbb{R})$. We shall apply Theorem 5.2 to the equivariant pair

$$\rho: G \rightarrow G_S, \quad \tau: D \rightarrow \mathcal{H}$$

and the morphism $\phi: X \rightarrow X_S$ of arithmetic varieties. For $\sigma \in \text{Aut}(\mathbb{C})$, we consider the conjugate morphism

$$\phi^\sigma: X^\sigma \rightarrow X_S^\sigma$$

and its lifting $\tau^\sigma : D^\sigma \rightarrow D_S^\sigma$. Since Γ_S and H can be obtained from an admissible and normal PEL-type (see [14, §4–§6]), X_S^σ is isomorphic to $\Gamma_S^{(\sigma)} \backslash \mathcal{H}'$ where $\mathcal{H}' = \mathcal{H}$ and $\Gamma_S^{(\sigma)}$ is an arithmetic subgroup of $(G_S)_0^\sigma$, the connected component of the identity of $\text{Aut}(D_S^\sigma)$ (see e.g. [14, (3.2)]). Let

$$\pi_S^{(\sigma)} : Y_S^{(\sigma)} \rightarrow X_S^\sigma$$

be the standard family of abelian varieties obtained from $\Gamma_S^{(\sigma)}$, \mathcal{H} and $\text{Sp}(V, \beta)$. By Theorem 5.2 there exist a Lie group G_1^σ and a homomorphism $\rho_1^\sigma : G_1^\sigma \rightarrow (G_S)_0^\sigma$ such that ρ_1^σ and τ^σ are equivariant and $\rho_1^\sigma(\Gamma^\sigma)$ is contained in $\Gamma_S^{(\sigma)}$. Hence we obtain a Kuga fiber variety $\pi^{(\sigma)} : Y^{(\sigma)} \rightarrow X^\sigma$ by pulling back the standard family

$$\pi_S^{(\sigma)} : Y_S^{(\sigma)} \rightarrow X_S^\sigma$$

via the morphism $\phi^\sigma : S^\sigma \rightarrow X_S^\sigma$:

$$\begin{array}{ccc} Y^{(\sigma)} & \xrightarrow{\phi_Y^{(\sigma)}} & Y_S^{(\sigma)} \\ \pi^{(\sigma)} \downarrow & & \downarrow \pi_S^{(\sigma)} \\ X^\sigma & \xrightarrow{\phi^\sigma} & X_S^\sigma \end{array}$$

On the other hand, by applying σ to the commutative diagram

$$\begin{array}{ccc} Y & \xrightarrow{\phi_Y} & Y_S \\ \pi \downarrow & & \downarrow \pi_S \\ X & \xrightarrow{\phi} & X_S \end{array}$$

we obtain the following commutative diagram:

$$\begin{array}{ccc} Y^\sigma & \xrightarrow{\phi_Y^\sigma} & Y_S^\sigma \\ \pi^\sigma \downarrow & & \downarrow \pi_S^\sigma \\ X^\sigma & \xrightarrow{\phi^\sigma} & X_S^\sigma \end{array}$$

If $\pi_S : Y_S \rightarrow X_S$ is a family of abelian varieties $\mathcal{F}(\Omega)$ associated to a PEL-type Ω , then by Theorem 6.1 the conjugate $\pi_S^\sigma : Y_S^\sigma \rightarrow X_S^\sigma$ is a family $\mathcal{F}(\Omega^\sigma)$ associated to the PEL-type Ω^σ . Note that the Lie group $(G_S)_0^\sigma$, the symmetric domain \mathcal{H} and the arithmetic subgroup

$\Gamma_S^{(\sigma)}$ coincide with the corresponding objects associated to the standard family $\pi_S^{(\sigma)}: Y_S^{(\sigma)} \rightarrow X_S^\sigma$. If the construction of the fiber variety associated to a PEL-type described in [12, (3.14)] and [13, (5.1)] is used, then Y_S^σ can be constructed in the same way as $Y_S^{(\sigma)}$. In particular, at least one fiber of $\pi_S^{(\sigma)}: Y_S^{(\sigma)} \rightarrow X_S^\sigma$ has a PEL-structure $(Q^{(\sigma)})_u$ ($u \in X_S^\sigma$) that is of type Ω^σ . The standard family $\pi_S^{(\sigma)}$ certainly satisfies the property (ii) in Theorem 6.1. To show that the conjugate family $\pi_S^\sigma: Y_S^\sigma \rightarrow X_S^\sigma$ satisfies the same property, let $(Q^\sigma)_u$ and $(Q^\sigma)_v$ be the PEL-structures of the fibers over $u, v \in X_S^\sigma$. Then u is equal to v by [13, Proposition 4.4] (see also [15, Theorem 4]); hence the conjugate family π_S^σ also satisfies the property (ii) in Theorem 6.1. Thus by Theorem 6.2 there is a biregular isomorphism between the families $\pi_S^\sigma: Y_S^\sigma \rightarrow X_S^\sigma$ and $\pi_S^{(\sigma)}: Y_S^{(\sigma)} \rightarrow X_S^\sigma$. Therefore there is an isomorphism between the diagram for $\pi^{(\sigma)}: Y^{(\sigma)} \rightarrow X^\sigma$ and the one for $\pi^\sigma: Y^\sigma \rightarrow X^\sigma$ given above; hence it follows that $\pi^\sigma: Y^\sigma \rightarrow X^\sigma$ is a Kuga fiber variety.

REFERENCES

- [1] S. Addington, *Equivariant holomorphic maps of symmetric domains*, Duke Math. J., **55** (1987), 65–88.
- [2] A. Ash, D. Mumford, M. Rapoport and Y. S. Tai, *Smooth Compactification of Locally Symmetric Varieties*, Math. Sci. Press, Brookline, 1975.
- [3] W. Baily and A. Borel, *Compactification of arithmetic quotients of bounded symmetric domains*, Ann. of Math., **84** (1966), 442–528.
- [4] A. Borel, *Some metric problems on arithmetic quotients of symmetric spaces and an extension theorem*, J. Differential Geom., **6** (1972), 543–560.
- [5] M. Borovoi, *Langlands' conjecture concerning conjugation of Shimura varieties*, Selecta Math. Soviet, **3** (1983/84), 3–39.
- [6] D. Kazhdan, *On arithmetic varieties*, in *Lie groups and their representations*, Halsted, New York, 1975.
- [7] —, *On arithmetic varieties II*, Israel J. Math., **44** (1983), 139–159.
- [8] M. Kuga, *Fiber Varieties Over a Symmetric Space Whose Fibers are Abelian Varieties I, II*, Lect. Notes, Univ. Chicago, 1963/64.
- [9] M. Kuga and S. Ihara, *Families of families of abelian varieties*, in *Algebraic number theory*, Japan Soc. for Prom. Sci., Tokyo, 1977.
- [10] J. Milne, *The action of an automorphism of \mathbb{C} on a Shimura variety and its special points*, in *Progr. Math.*, Vol. 35, Birkhäuser, Boston, 1983.
- [11] I. Satake, *Algebraic Structures of Symmetric Domains*, Princeton Univ. Press, 1980.
- [12] G. Shimura, *Moduli and fiber systems of abelian varieties*, Ann. of Math., **83** (1966), 294–338.
- [13] —, *On the field of definition for a field of automorphic functions II*, Ann. of Math., **81** (1965), 124–165.

- [14] G. Shimura, *On the field of definition for a field of automorphic functions III*, Ann. of Math., **83** (1966), 377–385.
- [15] —, *Moduli of abelian varieties and number theory*, in *Proc. Sympos. Pure Math.*, Vol. 9, Amer. Math. Soc., Providence, RI, 1966.

Received August 31, 1989 and in revised form March 15, 1990.

UNIVERSITY OF NORTHERN IOWA
CEDAR FALLS, IA 50614

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Takao Akahori and Harunori Ameku, On the Romanov kernel and Kuranishi's L^2 -estimate for $\bar{\partial}_b$ over a ball in the strongly pseudo convex boundary 1

Robert Emile Beaudoin, The proper forcing axiom and stationary set reflection 13

Hans Engler, A matrix Volterra integrodifferential equation occurring in polymer rheology 25

Zhong Ge, On a constrained variational problem and the spaces of horizontal paths 61

Yutaka Hemmi, Higher homotopy commutativity of H -spaces and the mod p torus theorem 95

G. D. Johnson, An intrinsic characterization of a class of minimal surfaces in constant curvature manifolds 113

Min Ho Lee, Conjugates of equivariant holomorphic maps of symmetric domains 127

Jacek Nikiel, H. Murat Tuncali and Edward D. Tymchatyn, On the rim-structure of continuous images of ordered compacta 145

Tara Lynn Smith, Generalized Clifford-Littlewood-Eckmann groups 157

Tara Lynn Smith, Generalized Clifford-Littlewood-Eckmann groups II: Linear representations and applications 185