

Pacific Journal of Mathematics

**ON THE RIM-STRUCTURE OF CONTINUOUS IMAGES OF
ORDERED COMPACTA**

JACEK NIKIEL, H. MURAT TUNCALI AND EDWARD D. TYMCHATYN

ON THE RIM-STRUCTURE OF CONTINUOUS IMAGES OF ORDERED COMPACTA

J. NIKIEL, H. M. TUNCALI, AND E. D. TYMCHATYN

Let X be a Hausdorff continuous image of an ordered continuum. Mardešić proved that X has a basis of open sets with metrizable boundaries. We use T-set approximations to obtain bases of open sets for X whose boundaries satisfy a variety of conditions. In particular, we prove that

$$\begin{aligned} \dim X &= \text{ind } X = \text{Ind } X \\ &= \max\{1, \sup\{\dim Y : Y \subset X \text{ is metrizable and closed}\}\}. \end{aligned}$$

1. Introduction. In this paper we study the rim-properties of images of ordered continua and, more generally, of compact ordered spaces. Mardešić proved in [M1] that a Hausdorff space which is a continuous image of a compact ordered space is rim-metrizable. In [N3], the first author proved that every hereditarily locally connected continuum is a continuous image of an ordered continuum. Then he used the approximation by T-sets of cyclic elements in images of ordered continua to prove that every hereditarily locally connected continuum is rim-countable. We use the techniques of [N3] to improve the result of Mardešić and to answer a question of Mardešić and Papić [MP] about dimension-theoretic properties of continuous images of ordered continua and ordered compacta. We improve a result of Simone [Si1] by proving that if X is a continuous image of an ordered continuum and X contains no nondegenerate metric continuum, then it is rim-finite. We also prove that if a rim-scattered space is a continuous image of an ordered compactum, then it is rim-countable.

All spaces in this paper are Hausdorff. A *continuum* is a compact connected (Hausdorff) space. An *ordered compactum* is a compact space which admits a linear ordering such that the order topology is the given topology. Ordered continua are locally connected; they are often called *arcs*.

A point p of a connected set X is a *separating point* of X if $X - \{p\}$ is not connected. We let $E(X)$ denote the set of all separating points of X .

Let X be a locally connected continuum. A connected subset Q of X is a *cyclic element* of X if Q is maximal with respect to containing

no separating points of itself. Each cyclic element of X is a locally connected continuum. The theory of cyclic elements is presented in [Wh1, Ch. 4] for the case of metric locally connected continua. We shall use some extensions of this theory to the non-metric setting as set out in [Wh2] and [C], see also [N4].

A collection \mathbf{A} of subsets of a compact space X is said to be a *null-family* in X if, for every open covering \mathbf{U} of X , the subcollection $\{B \in \mathbf{A} : B \text{ is not contained in any } V \in \mathbf{U}\}$ is finite.

Let A be a subset of a locally connected continuum X . We let $K(X - A)$ denote the set of all components of $X - A$. We will say that A is a *T-set* in X if A is closed and each component of $X - A$ has a two-point boundary.

Let Y be a cyclic element of a locally connected continuum X . We say that a sequence $\{A_1, A_2, \dots, A_n, \dots\}$ of T-subsets of Y *T-approximates* Y if

- (1) A_1 is metrizable,
- (2) $A_n \subset A_{n+1}$,
- (3) if $Z \in K(Y - A_n)$, then $E(\text{Cl}(Z)) \subset A_{n+1}$,
- (4) if $Z \in K(Y - A_n)$ and C is a nondegenerate cyclic element of $\text{Cl}(Z)$, then $C \cap A_{n+1}$ is a metrizable set which contains at least three points.

Note that the conditions of the above definition imply that $\text{Cl}(\bigcup_{n=1}^{\infty} A_n) = Y$ (see [N1, Lemma 3.4]).

In [N1], there are given several characterizations of continuous Hausdorff images of ordered continua. One of them is the following:

THEOREM 1 [N1, 1.1]. *Let X be a locally connected continuum. Then the following are equivalent:*

- (1) X is a continuous image of an ordered continuum,
- (2) if Y is a nondegenerate cyclic element of X , then there is a sequence $\{A_1, A_2, \dots\}$ of T-sets in Y which T-approximates Y .

Further properties of continuous images of arcs and ordered compacta can be found in survey articles [M3], [TrW] and [N4]; see also [N1].

Let \mathbf{P} be a property of sets. A space X is said to be *rim-P* if it has a basis of open sets whose boundaries have property \mathbf{P} . A set is said to be *scattered* if each of its non-empty closed subsets has an

isolated point. Recall that compact, metrizable, scattered spaces are countable. For definitions of dimensions \dim , Ind and ind , the reader is referred to [E].

For a compact space X , we define

$$\alpha(X) = \sup\{\dim Z : Z \text{ is a closed metrizable subset of } X\}.$$

We let $\alpha - 1 = \infty$ if $\alpha = \infty$.

We shall need the following lemmas.

LEMMA 1 [Tr2]. *Let X be a locally connected continuum and A a T-set in X . There exists an upper semi-continuous decomposition G_A of X into closed sets such that if X_A denotes the quotient space and $f : X \rightarrow X_A$ is the quotient map, then:*

- (1) $f|_A$ is one-to-one and $f(A)$ is a T-set in X_A ,
- (2) each $Z \in K(X_A - f(A))$ is homeomorphic to $]0, 1[$,
- (3) for each $Z \in K(X_A - f(A))$ there exists a unique $P_Z \in K(X - A)$ such that $f(P_Z) \subset \text{Cl}(Z)$, and each component of $X - A$ is a P_Z for some $Z \in K(X_A - f(A))$.

In the above lemma, $f(A)$ is a T-set in X_A , and we call f a T-map with respect to A . The space X_A is uniquely determined by X and A . If the set A is metrizable it follows, by local connectedness of X , that $K(X - A)$ is countable, [N1, 4.1].

LEMMA 2. *Let X be a locally connected continuum and, for every cyclic element Y of X , let \mathbf{B}_Y be a basis for Y . Then X has a basis \mathbf{B} such that, for each $U \in \mathbf{B}$, there exist a family \mathbf{A} of cyclic elements of X , non-negative integers m and n , nondegenerate cyclic elements Y_1, \dots, Y_m of X , sets $U_1 \in \mathbf{B}_{Y_1}, \dots, U_m \in \mathbf{B}_{Y_m}$, and separating points x_1, \dots, x_n of X such that*

$$U = \left(\bigcup \mathbf{A} \right) \cup U_1 \cup \dots \cup U_m \quad \text{and} \\ \text{Bd}(U) = \text{Bd}_{Y_1}(U_1) \cup \dots \cup \text{Bd}_{Y_m}(U_m) \cup \{x_1, \dots, x_n\}.$$

Proof. The lemma follows from the generalization, by Cornette [C, p. 225-6], of Whyburn's cyclic chain approximation theorem [Wh1, IV.7.1, p. 73] to the case of locally connected Hausdorff continua. \square

LEMMA 3. *Let γ be an infinite cardinal number and let \mathbf{P} be a hereditary property of compact sets that is preserved under unions of fewer than γ compact sets. Let X be a locally connected continuum,*

$\{A_i\}_{i=1}^{\infty}$ an increasing sequence of closed subsets of X , and $\{\mathbf{V}_i\}_{i=1}^{\infty}$ a sequence of collections of sets such that:

- (1) \mathbf{V}_i is a basis of open sets for A_i ,
- (2) $\text{Bd}(K)$ has property **P** for each $K \in K(X - A_i)$,
- (3) $V \in \mathbf{V}_i$ implies $\text{Bd}_{A_i}(V)$ has property **P**,
- (4) $V \in \mathbf{V}_i$ implies $\{K \in K(X - A_i) : \text{Bd}(K) \cap V \neq \emptyset \text{ and } \text{Bd}(K) \not\subset \text{Cl}(V)\}$ has cardinality less than γ ,
- (5) for each open cover \mathbf{W} of X there is an integer i such that $K(X - A_i)$ refines \mathbf{W} .

Then X admits a basis of open sets whose boundaries have property **P**.

Proof. Let $x \in X$ and let U be an open neighbourhood of x . Let W be an open neighbourhood of x such that $\text{Cl}(W) \subset U$.

Suppose that $x \notin \bigcup_{n=1}^{\infty} A_n$. For every n let $K_n \in K(X - A_n)$ be such that $x \in K_n$. Then $K_{n+1} \subset K_n$. By (5), there is an integer i such that K_i is contained either in U or in $X - \text{Cl}(W)$. Since $x \in \text{Cl}(W) \cap K_i$, it follows that $K_i \subset U$. Since X is locally connected, K_i is an open set. By (2), $\text{Bd}(K_i)$ has property **P**.

Now suppose that $x \in A_n$ for some integer n . By (5), we may take n to be such that no component of $X - A_n$ meets both $\text{Cl}(W)$ and $X - U$. Let $V \in \mathbf{V}_n$ be such that $x \in V \subset \text{Cl}(V) \subset W$. Let $V' = V \cup \bigcup \{K \in K(X - A_n) : \text{Bd}(K) \cap V \neq \emptyset\}$. Then $V' \subset U$. Since X is locally connected, V' is open and

$$\text{Bd}(V') \subset \text{Bd}_{A_n}(V)$$

$$\cup \bigcup \{\text{Bd}(K) : K \in K(X - A_n), \text{Bd}(K) \cap V \neq \emptyset \text{ and } \text{Bd}(K) \not\subset V\}.$$

By (3), (2) and (4), it follows that $\text{Bd}(V')$ has property **P**. \square

2. Main results. The proof of the following lemma uses some ideas from the proof of [N3, Theorem 4.1].

LEMMA 4. *Let Y be a continuum with no separating point which is a continuous image of an ordered continuum. Let $\alpha = \max\{1, \alpha(Y)\}$. Then Y has a basis \mathbf{V} of open sets whose boundaries are metrizable sets of $\dim \leq \alpha - 1$. Moreover, if Y admits a basis of open sets with scattered boundaries, then the boundaries of members of \mathbf{V} are countable.*

Proof. Let $\{A_1, A_2, \dots\}$ be a sequence of T-sets in Y which T-approximates Y . For each n , let $f_n : Y \rightarrow Y_{A_n} = Y_n$ be a T-map with

respect to A_n (see Lemma 1). We let $B_n^m = f_n(A_m) \subset Y_n$ provided $m \leq n$. Notice that Y_n has no separating point, each B_n^m is a T-set in Y_n provided $m \leq n$, $f_n|_{A_m} : A_m \rightarrow B_n^m$ is a homeomorphism, and every component of $Y_n - B_n^m$ is homeomorphic to $]0, 1[$. Since Y_n has no separating point, it follows that if P is a component of $Y_n - B_n^m$, $\text{Bd}(P) = \{a, b\}$, then $\text{Cl}(P)$ is a cyclic chain from a to b (in the case when $m = n - 1$, all cyclic elements of $\text{Cl}(P)$ are metrizable—see below).

First, we use an induction to show that, for $n = 1, 2, \dots$, Y_n has a basis \mathbf{B}_n such that $\text{Bd}_{Y_n}(V)$ is metrizable and $\dim(\text{Bd}_{Y_n}(V)) \leq \alpha - 1$ for each $V \in \mathbf{B}_n$.

Note that $Y_1 = B_1^1 \cup (Y_1 - B_1^1)$ is a metrizable space which is a union of the compact metrizable set B_1^1 (which is homeomorphic to A_1) and a countable family of copies of $]0, 1[$. By [E, 1.5.3, p. 42], $\dim Y_1 \leq \max\{1, \dim B_1^1\} \leq \alpha$. Hence, Y_1 has a basis \mathbf{B}_1 as required.

Suppose that the required basis \mathbf{B}_n for Y_n has been already defined. Let $y \in Y_{n+1}$ and let V be an open neighbourhood of y in Y_{n+1} . If $y \notin B_{n+1}^n$, then $y \in Q$ for some $Q \in K(Y_{n+1} - B_{n+1}^n)$. Let $\text{Bd}(Q) = \{a, b\}$. Then $\text{Cl}(Q)$ is a cyclic chain from a to b and $E(\text{Cl}(Q)) \subset B_{n+1}^n$. If Z is a nondegenerate cyclic element of $\text{Cl}(Q)$, then $B_Z = B_{n+1}^n \cap Z$ is a metrizable T-set in Z , $Z \cap (E(\text{Cl}(Q)) \cup \{a, b\})$ consists of exactly two points, and each component of $Z - B_Z$ is homeomorphic to $]0, 1[$. Hence, $K(Z - B_Z)$ is countable and Z is metrizable. Now, it is easy to find an open neighbourhood W of y in Y_{n+1} such that $W \subset V \cap Q$, $\text{Bd}_{Y_n}(W)$ is contained in two cyclic elements Z_1 and Z_2 of $\text{Cl}(Q)$ and for $i = 1, 2$

$$\begin{aligned} \dim(\text{Bd}_{Y_n}(W) \cap Z_i) &\leq \dim Z_i - 1 \leq \max\{1, \dim B_{Z_i}\} - 1 \\ &\leq \max\{1, \dim A_{n+1}\} - 1 \leq \alpha - 1 \end{aligned}$$

provided Z_i is nondegenerate (the case when Z_i is degenerate is trivial). Thus we have $\dim(\text{Bd}_{Y_n}(W)) \leq \alpha - 1$.

Now, suppose that $y \in B_{n+1}^n$. Let x denote the unique point of A_n such that $f_{n+1}(x) = y$. For every $P \in K(Y_{n+1} - B_{n+1}^n)$ let $Q_P \in K(Y - A_n)$ be a component such that $f_{n+1}(Q_P) \subset \text{Cl}(P)$ and let $R_P \in K(Y_n - B_n^n)$ be such that $f_n(Q_P) \subset \text{Cl}(R_P)$. Set $\text{Bd}_{Y_{n+1}}(P) = \{a_P, b_P\}$ and $\text{Bd}_{Y_n}(R_P) = \{a'_P, b'_P\}$, where $f_{n+1}^{-1}(a_n) \cap A_n = f_n^{-1}(a'_n) \cap A_n$, and let \leq denote the natural ordering on $\text{Cl}(R_P)$ from a'_P to b'_P . Choose $r_P \in R_P$ and let $I_P = \{r \in R_P : r < r_P\}$ and $J_P = \{r \in R_P : r_P < r\}$.

Let

$$\begin{aligned} V' &= f_n(f_{n+1}^{-1}(V) \cap A_n) \\ &\cup \bigcup \{R_P : P \in K(Y_{n+1} - B_{n+1}^n) \text{ and } \text{Cl}(P) \subset V\} \\ &\cup \bigcup \{I_P : P \in K(Y_{n+1} - B_{n+1}^n) \text{ and } a_P \in V\} \\ &\cup \bigcup \{J_P : P \in K(Y_{n+1} - B_{n+1}^n) \text{ and } b_P \in V\}. \end{aligned}$$

Since $\{\text{Cl}(R_P) : P \in K(Y_{n+1} - B_{n+1}^n)\}$ is a null-family, V' is an open subset of Y_n . Moreover, $f_n(x) \in V'$. By the inductive hypothesis, there is a connected open set W' in Y_n such that $f_n(x) \in W' \subset V'$, $\text{Bd}_{Y_n}(W')$ is metrizable and $\dim(\text{Bd}_{Y_n}(W')) \leq \alpha - 1$. Let

$$\begin{aligned} \mathbf{H}_1 &= \{P \in K(Y_{n+1} - B_{n+1}^n) : a'_P \in W' \text{ and } R_P \not\subset W'\}, \\ \mathbf{H}_2 &= \{P \in K(Y_{n+1} - B_{n+1}^n) : b'_P \in W' \text{ and } R_P \not\subset W'\} \end{aligned}$$

and

$$\mathbf{H}_3 = \{P \in K(Y_{n+1} - B_{n+1}^n) : R_P \subset W'\}.$$

Note that if $P \in \mathbf{H}_1 \cup \mathbf{H}_2$, then $R_P \cap \text{Bd}_{Y_n}(W')$ is a non-empty open subset of $\text{Bd}_{Y_n}(W')$. Since $\text{Bd}_{Y_n}(W')$ is compact and metrizable, $\mathbf{H}_1 \cup \mathbf{H}_2$ is countable. For every $P \in \mathbf{H}_1$ (resp. $P \in \mathbf{H}_2$), let W_P^1 (resp. W_P^2) be an open subset of $\text{Cl}(P)$ such that $a_P \in W_P^1 \subset V$ (resp. $b_P \in W_P^2 \subset V$), $\text{Bd}_{\text{Cl}(P)}(W_P^1)$ is metrizable and $\dim(\text{Bd}_{\text{Cl}(P)}(W_P^1)) \leq \alpha - 1$ (resp. $\text{Bd}_{\text{Cl}(P)}(W_P^2)$ is metrizable and $\dim(\text{Bd}_{\text{Cl}(P)}(W_P^2)) \leq \alpha - 1$). Note that $\text{Bd}_{\text{Cl}(P)}(W_P^i)$ may be assumed to be contained in one cyclic element Z of $\text{Cl}(P)$. By the fact that $K(Z - B_Z)$ is countable, it follows that Z is metrizable and $\dim Z \leq \alpha$. Let

$$W = f_{n+1}(f_n^{-1}(W') \cap A_n) \cup \bigcup_{P \in \mathbf{H}_1} W_P^1 \cup \bigcup_{P \in \mathbf{H}_2} W_P^2 \cup \bigcup \mathbf{H}_3.$$

Since $K(Y_{n+1} - B_{n+1}^n)$ is a null-family, W is open in Z . A straightforward argument shows that $y \in W \subset V$ (because if $P \in K(Y_{n+1} - B_{n+1}^n)$ is not contained in V , then $r_P \notin V'$ and so $R_P \not\subset W'$) and

$$\begin{aligned} \text{Bd}_{Y_{n+1}}(W) &= f_{n+1}(f_n^{-1}(\text{Bd}_{Y_n}(W') \cap A_n)) \\ &\cup \bigcup_{P \in \mathbf{H}_1} \text{Bd}_{\text{Cl}(P)}(W_P^1) \cup \bigcup_{P \in \mathbf{H}_2} \text{Bd}_{\text{Cl}(P)}(W_P^2). \end{aligned}$$

Thus $\text{Bd}_{Y_{n+1}}(W)$ is a union of countably many compact metrizable sets of $\dim \leq \alpha - 1$. It is well-known that each compact space which

can be covered by countably many closed and metrizable subsets is metrizable. Hence, $\text{Bd}_{Y_{n+1}}(W)$ is metrizable. By [E, 1.5.3, p. 42], $\dim(\text{Bd}_{Y_{n+1}}(W)) \leq \alpha - 1$. The inductive argument is complete.

Let \mathbf{P} be the following property of compact spaces: a space is metrizable of dimension $\leq \alpha - 1$. Let $\gamma = \aleph_1$ be the first uncountable cardinal number. Note that Y satisfies all the assumptions of Lemma 3. Indeed, the condition (2) of Lemma 3 follows immediately from the definition of a T-set. Let $V_n = \{A_n \cap f_n^{-1}(U) : U \in \mathbf{B}_n\}$ for $n = 1, 2, \dots$. Then V_n is a basis for A_n which satisfies the conditions (1) and (3). The condition (4) follows from [N1, 4.1], and the condition (5) is a consequence of [N1, 3.4]. By Lemma 3, Y has a basis \mathbf{V} of open sets with metrizable boundaries of dimension $\leq \alpha - 1$.

Suppose that Y is rim-scattered. Then Y_1 is metrizable and rim-scattered. Hence, Y_1 has a basis of open sets with countable boundaries. It is now easy to modify the above argument to show that each Y_n has a basis of open sets with countable boundaries. By Lemma 3, Y has a basis of open sets with countable boundaries. \square

Simone, [Si1] and [Si2], proved that if X is a continuum with degree of cellularity \aleph_0 , which is a continuous image of an ordered continuum and which contains no nondegenerate metric subcontinuum, then X has a basis of open sets with finite boundaries. Simone's theorem can be improved as follows:

THEOREM 2. *Let X be a continuum which is a continuous image of an arc and which contains no nondegenerate metric subcontinuum. Then X has a basis of open sets with finite boundaries.*

Proof. Let Y be a nondegenerate cyclic element of X . Since having a basis of open sets with finite boundaries is a cyclically extensible property (see Lemma 2), it suffices to prove that Y is rim-finite.

Let $\{A_1, A_2, \dots\}$ be a sequence of T-sets in Y which T-approximates Y and, for $n = 1, 2, \dots$, let $f_n: Y \rightarrow Y_n$ be a T-map with respect to A_n (see Lemma 1). Since A_1 is metrizable, and, hence, zero-dimensional, Y_1 has a basis of open sets with finite boundaries (see [N1, 4.3]). If U is an open set in Y_1 which has a finite boundary, then all but at most finitely many components of $Y_1 - A_1$ whose closures meet $U \cap A_1$ are contained in $\text{Cl}(U)$. An inductive argument similar to the one given in the proof of Lemma 4 shows that each Y_n is rim-finite. Taking \mathbf{P} to be the property of being a finite set and

$\gamma = \aleph_0$ in Lemma 3, it follows that Y has a basis of open sets with finite boundaries. \square

THEOREM 3. *If X is a nondegenerate continuous image of an ordered continuum, then*

$$\max\{1, \alpha(X)\} = \dim X = \text{Ind } X = \text{ind } X.$$

Proof. Let $\alpha = \max\{1, \alpha(X)\}$. Since X is a nondegenerate continuum, $\text{ind } X \geq 1$. By general facts (see [E, 3.1.4 on p. 209, 2.2.1 on p. 170, and 1.1.2 on p. 4]), it follows that $\dim X \geq \dim Z$, $\text{Ind } X \geq \text{Ind } Z$ and $\text{ind } X \geq \text{ind } Z$ for each closed subspace Z of X . Hence $\dim X$, $\text{Ind } X$, $\text{ind } X \geq \alpha$. For each normal space X , we have $\text{ind } X \leq \text{Ind } X$ [E, 1.6.3, p. 52] and $\dim X \leq \text{Ind } X$ [E, 3.1.28, p. 220]. Thus it suffices to show that $\text{Ind } X \leq \alpha$.

Let $x \in X$ and V be an open neighbourhood of x . By Lemmas 4 and 2, there exists an open set W such that $x \in W \subset V$, $\text{Bd}(W)$ is contained in the union of a finite collection $\{Z_1, \dots, Z_n\}$ of cyclic elements of X , $\text{Bd}(W) \cap Z_i$ is metrizable and $\dim(\text{Bd}(W) \cap Z_i) \leq \alpha - 1$ for $i = 1, \dots, n$. Hence, $\text{Bd}(W)$ is metrizable and $\text{Ind } \text{Bd}(W) = \dim \text{Bd}(W) \leq \alpha - 1$. By the sum theorem for separable metric spaces, [E, 1.5.3, p. 42], we have $\text{Ind } X \leq \alpha$. \square

REMARK. In Theorem 3, if $\alpha(X) = 0$, then X is rim-finite by Theorem 2.

THEOREM 4. *Let X be a continuum which is a continuous image of an arc. If X has a basis of open sets with scattered boundaries, then it has a basis of open sets with countable boundaries.*

Proof. By Lemma 4, each cyclic element of X is rim-countable. The theorem follows by Lemma 2. \square

The following theorem answers a question of Mardešić and Papić ([MP], see also [N4, Problem 4]):

THEOREM 5. *Let Z be a continuous image of a compact ordered space. Then*

- (1) $\dim Z = \text{Ind } Z = \text{ind } Z$. *If, moreover, $\dim Z > 0$ then $\dim Z = \max\{1, \alpha(Z)\}$.*
- (2) *If Z is rim-scattered, then it is rim-countable.*

Proof. For every compact space T , $\text{Ind } T = 0$ iff $\dim T = 0$ iff $\text{ind } T = 0$, [E, 3.1.30, p. 221]. Thus we may assume that Z is not zero-dimensional. Let $\alpha = \max\{1, \alpha(Z)\}$.

By [N2, Theorem 2], see also [M1, Lemma 8], there exists a space X such that X is a continuous image of an arc, $Z \subset X$, Z is a T-set in X , and each component of $X - Z$ is homeomorphic to $]0, 1[$. If Y is a closed metrizable subset of X , then Y is a union of $Z \cap Y$ and at most countably many closed sets which are homeomorphic to subsets of $]0, 1[$. Hence, $\dim Y \leq \max\{1, \dim(Y \cap Z)\}$. By Theorem 3, $\alpha = \dim X = \text{Ind } X = \text{ind } X$. Since Z is not zero-dimensional, $\alpha \leq \dim Z, \text{Ind } Z, \text{ind } Z$. However, $\dim Z \leq \dim X, \text{Ind } Z \leq \text{Ind } X$ and $\text{ind } Z \leq \text{ind } X$. This completes the proof of (1). A similar argument together with Theorem 4 show that (2) holds. \square

REMARKS. 1. In the case when $\alpha(Z) = 0$, the result (1) of Theorem 5 was obtained by Mardešić [M2, Corollary, p. 425].

2. The proofs of Lemma 4 and Theorems 3 and 5 show that if a space X is a continuous image of an ordered compactum, then it has a basis \mathbf{B} such that $\text{Bd}(U)$ is metrizable and $\dim \text{Bd}(U) \leq \dim X - 1$ for each $U \in \mathbf{B}$. This improves results of [M1].

3. Problems. Filippov gave in [F] an example of a locally connected continuum which admits a basis of open sets with metrizable zero-dimensional and perfect boundaries and which is not a continuous image of any ordered compactum.

In general, rim-scattered continua are not continuous images of ordered compacta. For example: the space $X = L \times S /_{\{0\} \times S}$, where L denotes the long interval and $S = \{\frac{1}{n} : n = 1, 2, \dots\} \cup \{0\}$, is a rim-countable continuum which is a continuous image of no ordered compactum. In fact, X contains a non-metric product of infinite compact spaces—see [Tr1]. However, the space X is not locally connected. In [Tu], it was proved that rim-scattered locally connected continua do not contain a non-metric product of nondegenerate continua. Hence we may ask the following question:

Question 1. Is every locally connected rim-scattered continuum a continuous image of an ordered continuum?

Filippov's example shows that rim-scattered locally connected continua are the largest possible class of spaces defined with the use of rim-properties that could be contained in the class of continuous images of ordered continua. Recall the following weaker question which is still open (see [N3] and [N4]).

Question 2. Is every locally connected, rim-countable continuum a continuous image of an ordered continuum?

Let us also pose the following problem:

Question 3. Is every locally connected and rim-scattered continuum a rim-countable space?

Recall that, by Theorem 4, Question 3 has a positive answer provided the space under consideration is a continuous image of an arc.

Added in proof. Recently the authors answered questions 1 and 2 in the negative in the paper: J. Nikiel, H. M. Tuncali, and E. D. Tymchatyn, *A locally connected rim-countable continuum which is the continuous image of no arc*, *Topology Appl.* (to appear). L. B. Treybig proved a result which implies Theorem 2 in *Proc. Amer. Math. Soc.* **74** (1979), 326–328.

REFERENCES

- [C] J. L. Cornette, “*Image of a Hausdorff arc*” is cyclically extensible and reducible, *Trans. Amer. Math. Soc.*, **199** (1974), 253–267.
- [E] R. Engelking, *Dimension theory*, North-Holland and PWN—Polish Scientific Publishers, 1978.
- [F] V. V. Filippov, *On perfectly normal compacta*, in Russian, *Dokl. Akad. Nauk SSSR*, **189** (1969), 736–739.
- [M1] S. Mardešić, *Images of ordered compacta are locally peripherally metric*, *Pacific J. Math.*, **23** (1967), 557–568.
- [M2] ———, *Continuous images of ordered compacta and a new dimension which neglects metric subcontinua*, *Trans. Amer. Math. Soc.*, **121** (1966), 424–433.
- [M3] ———, *On the Hahn-Mazurkiewicz theorem in non-metric spaces*, in *General Topology and its Relations to Modern Analysis and Algebra II*, Prague (1966), 248–255.
- [MP] S. Mardešić and P. Papić, *Some problems concerning mappings of ordered compacta*, in Serbo-Croatian, English summary, *Matematička Biblioteka*, **25** (1963), 11–22.
- [N1] J. Nikiel, *Images of arcs—a nonseparable version of the Hahn-Mazurkiewicz theorem*, *Fund. Math.*, **129** (1988), 91–120.
- [N2] ———, *Some problems on continuous images of compact ordered spaces*, *Questions Answers Gen. Topology*, **4** (1986/87), 117–128.
- [N3] ———, *The Hahn-Mazurkiewicz theorem for hereditarily locally connected continua*, *Topology Appl.*, **32** (1989), 307–323.
- [N4] ———, *On continuous images of arcs and compact orderable spaces*, *Topology Proceedings*, **14** (1989), 163–193.
- [Si1] J. N. Simone, *Metric components of continuous images of ordered compacta*, *Pacific J. Math.*, **69** (1977), 269–274.
- [Si2] ———, *Continuous images of ordered compacta and hereditarily locally connected continua*, *Colloq. Math.*, **40** (1978), 77–84.
- [Tr1] L. B. Treybig, *Concerning continuous images of compact ordered spaces*, *Proc. Amer. Math. Soc.*, **15** (1964), 866–871.

- [Tr2] L. B. Treybig, *Arcwise connectivity in continuous images of ordered compacta*, *Glasnik Mat.*, **21** (1986), 201–211.
- [TrW] L. B. Treybig and L. E. Ward, Jr., *The Hahn-Mazurkiewicz problem*, in *Topology and order structure*, part 1, Math. Centre Tracts, vol. 142. Amsterdam, 1981, pp. 95–105.
- [Tu] H. M. Tuncali, *Analogues of Treybig's product theorem*, *Proc. Amer. Math. Soc.*, **108** (1990), 855–858.
- [Wh1] G. T. Whyburn, *Analytic Topology*, Amer. Math. Soc., Providence, R. I., 1942.
- [Wh2] ———, *Cut points in general topological spaces*, *Proc. Nat. Acad. Sci. USA*, **61** (1968), 380–387.

Received January 26, 1990. This research was supported in part by NSERC grant number A5616 and by a grant from the University of Saskatchewan President's Fund.

UNIVERSITY OF SASKATCHEWAN
SASKATOON, SASKATCHEWAN
CANADA S7N 0W0

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Pacific Journal of Mathematics

Vol. 149, No. 1

May, 1991

Takao Akahori and Harunori Ameku , On the Romanov kernel and Kuranishi's L^2 -estimate for $\bar{\partial}_b$ over a ball in the strongly pseudo convex boundary	1
Robert Emile Beaudoin , The proper forcing axiom and stationary set reflection	13
Hans Engler , A matrix Volterra integrodifferential equation occurring in polymer rheology	25
Zhong Ge , On a constrained variational problem and the spaces of horizontal paths	61
Yutaka Hemmi , Higher homotopy commutativity of H -spaces and the mod p torus theorem	95
G. D. Johnson , An intrinsic characterization of a class of minimal surfaces in constant curvature manifolds	113
Min Ho Lee , Conjugates of equivariant holomorphic maps of symmetric domains	127
Jacek Nikiel, H. Murat Tuncali and Edward D. Tymchatyn , On the rim-structure of continuous images of ordered compacta	145
Tara Lynn Smith , Generalized Clifford-Littlewood-Eckmann groups	157
Tara Lynn Smith , Generalized Clifford-Littlewood-Eckmann groups II: Linear representations and applications	185