

Pacific Journal of Mathematics

**THE LARGEST DIGIT IN THE CONTINUED FRACTION
EXPANSION OF A RATIONAL NUMBER**

DOUGLAS AUSTIN HENSLEY

THE LARGEST DIGIT IN THE CONTINUED FRACTION EXPANSION OF A RATIONAL NUMBER

DOUGLAS HENSLEY

The finite continued fraction sequence of a reduced fraction a/b , with $0 \leq a < b$, is the sequence $d = (d(1), d(2), \dots, d(r))$ of positive integers such that $d(r) > 1$, and

$$a/b = 1/(d(1) + 1/(d(2) + \dots + 1/d(r))).$$

In the standard terminology of continued fractions, this is written as $[0; d(1), d(2), \dots, d(r)]$, which we abbreviate here to $[d(1), d(2), \dots, d(r)]$. Thus $[1, 4, 2] = 1/(1 + 1/(4 + 1/2)) = 9/11$. The empty sequence corresponds to $0/1$. For any other fraction, there will be $r \geq 1$ *digits* (also known as partial quotients) $d(j)$ in this expansion ($1 \leq j \leq r$). The largest of these we call $D(a/b)$ or $D(a, b)$. Thus $D(9/11) = D(9, 11) = 4$. The aim of this work is to elucidate the distribution of $D(a, b)$. Put informally, the main result is that $\text{Prob}[D(a, b) \leq \alpha \log b] \approx \exp(-12/\alpha\pi^2)$. More precisely, it is shown that for all $\varepsilon > 0$, and uniformly in $\alpha > \varepsilon$ as $x \rightarrow \infty$,

$$\#\{(a, b) : 0 \leq a < b \leq x, \text{gcd}(a, b) = 1, \text{ and } D(a, b) \leq \alpha \log x\} \\ \approx (3/\pi^2)x^2 \exp(-12/\alpha\pi^2).$$

The question of how often there are exactly M digits exceeding $\alpha \log b$ in the continued fraction expansion of a reduced fraction a/b with $0 \leq a < b \leq x$ is also touched on. Evidence points to the estimate

$$(3/\pi^2)x^2(M!)^{-1}(12/\alpha\pi^2)^M \exp(-12/\alpha\pi^2)$$

for the number of such fractions.

Previous work in a similar vein includes a result of Galambos [4, 5] concerning the distribution of the continued fraction partial quotients (digits) of a randomly chosen *real* number in the interval $(0, 1)$. Corresponding to any irrational ξ in $(0, 1)$ there is a unique sequence $d_\xi = d = (d(1), d(2), \dots)$ of positive integers such that

$$\xi = [d] = [d(1), d(2), \dots] = 1/(d(1) + 1/(d(2) + \dots)).$$

Galambos found that if X is a random variable uniformly distributed on $[0, 1]$ (in the statement of his result the random variable has the Gauss-Kuzmin distribution, but that was just a convenience), then

$$(1) \quad \lim_{r \rightarrow \infty} \text{Prob} \left(\max_{k \leq r} d_X(k) < \alpha r \right) = e^{-1/\alpha \log 2}.$$

There is also a literature concerning the distribution of pairs (a, b) for which, in the finite continued fraction expansion $d = d_{a/b}$ of a/b , all of the $d(j)$ are bounded by some fixed N . It is known [2, 3, 8] that for each $N \geq 2$ there exists a real number $H(N)$, $0 < H(N) < 1$, such that the number of pairs (a, b) for which $b \leq x$ is on the order of $x^{2H(N)}$, uniformly in N as $x \rightarrow \infty$. For each fixed N , there is also [9] a constant $C(N) > 0$ such that this pair count is $\approx C(N)x^{2H(N)}$, but it is not known how fast the convergence to this asymptotic behavior is, or whether it is uniform in N . There is no evident reason to suspect that it would not be uniform, but in any event numerical evidence suggests that x must be fairly large before the asymptotic trend takes hold. Recently, the author also showed [10] that

$$(2) \quad \lim_{n \rightarrow \infty} N(1 - H(N)) = 6/\pi^2.$$

As usual, $\Phi(x)$ denotes $\sum_{n \leq x} \phi(n) = \#\{(a, b) : 0 \leq a < b \leq x \text{ and } \gcd(a, b) = 1\}$, so that

$$(3) \quad \Phi(x) \approx (3/\pi^2)x^2 \quad \text{as } x \rightarrow \infty.$$

Now let

$$\Phi(x, \alpha) := \#\{(a, b) : 0 \leq a < b \leq x, \gcd(a, b) = 1, \text{ and } D(a, b) \leq \alpha \log x\}.$$

From the results mentioned above, it follows that there exists $C > 0$ such that for all sufficiently large x ,

$$(4) \quad (1/C)x^2 e^{-12/\alpha\pi^2} \leq \Phi(x, \alpha) \leq Cx^2 e^{-12/\alpha\pi^2}$$

whenever $\alpha \log x$ is an integer ≥ 2 . In view of the results just mentioned, our main result below fits in nicely:

THEOREM 1. *Uniformly in $\alpha \geq 4/\log \log x$ as $x \rightarrow \infty$,*

$$\Phi(x, \alpha) = (3/\pi^2)x^2 e^{-12/\alpha\pi^2} (1 + O((\alpha^{-2} + 1)e^{24/\pi^2\alpha} \log \log x / \log x))$$

as $x \rightarrow \infty$.

The result can also be put in a form which refers to the Diophantine approximation properties of a/b rather than to its continued fraction expansion. Let

$$\delta(a, b) := \min_{1 \leq k < b} \|ka/b\|,$$

where $\|u\|$ denotes the distance from u to the nearest integer. Let

$$(5) \quad F(x, \alpha) := \#\{(a, b) : 0 \leq a < b \leq x, \gcd(a, b) = 1, \\ \text{and } \delta(a, b) > 1/\alpha \log x\}.$$

Then for fixed $\alpha > 0$, as $x \rightarrow \infty$,

$$(6) \quad F(x, \alpha) \approx (3/\pi^2)x^2e^{-12/\alpha\pi^2}.$$

The basic idea of the proof is to count $\Phi(x, \alpha)$ by inclusion and exclusion, throwing out all fractions with at least one digit too large—once for each such digit—then restoring those with at least two—once for each such pair of digits—and so on. Term by term, these counts are asymptotic to the corresponding term in the identity

$$(7) \quad (3/\pi^2)x^2e^{-12/\alpha\pi^2} = (3/\pi^2)x^2 \sum_{j=0}^{\infty} (-12/\pi^2\alpha)^j/(j!).$$

REMARK. A more sophisticated version of inclusion and exclusion yields an asymptotic estimate of the number of fractions with exactly M digits $\geq \alpha \log x$, and denominator $\leq x$. Let $\mu_M: \mathbb{N} \rightarrow \mathbb{Z}$ satisfy

$$\sum_{k=0}^n \binom{n}{k} \mu_M(k) = \{1 \text{ if } n \leq M, 0 \text{ if not}\}.$$

This defines μ_M recursively, and it is not hard to see that $\mu_M(0) = 1$,

$$\mu_M(j) = 0 \quad \text{if } 1 \leq j \leq M, \quad \text{and} \\ \mu_M(j) = (-1)^{(j-M)} \binom{j-1}{M} \quad \text{for } j > M.$$

Following the proof given here for the case $M = 0$, but with μ_M in place of $(-1)^j$, leads to a main term of

$$(3x^2/\pi^2)(1/M!)(12/\pi^2\alpha)^M e^{-(12/\pi^2\alpha)}.$$

2. Inclusion and exclusion. Let $V_r := \{v : \{1, 2, \dots, r\} \rightarrow \mathbb{Z}^+\}$ be the set of all sequences of r positive integers, and let $V = \bigcup_{r=0}^{\infty} V_r$. For $v \in V_r$, let $\text{lex}(v) = r$, the lexicographic length of v .

Let $a_0(v) = 0$, $b_0(v) = 1$, $a_{-1}(v) = 1$, and $b_{-1}(v) = 0$. For $1 \leq i \leq r$ we define $a_i(v)$ and $b_i(v)$ recursively by the conditions

$$(8) \quad a_i(v) = d_i a_{i-1}(v) + a_{i-2}(v), \\ b_i(v) = d_i b_{i-1}(v) + b_{i-2}(v),$$

where $d_i = v(i)$ is the i th entry in the sequence v . Let

$$\langle v \rangle := b_{\text{lex}(v)}(v) = b_r(v),$$

say, let $[v] := a_r(v)/b_r(v)$, and $\{v\} := b_{r-1}(v)/b_r(v)$. By convention, if v is the empty sequence ($r = \text{lex}(v) = 0$) then $\langle v \rangle = 1$ and $[v] = \{v\} = 0$.

There is a two-to-one correspondence T between V and $\{(a, b) : 0 \leq a < b \text{ and } \gcd(a, b) = 1\}$. In one direction, we map $v \xrightarrow{T} (a_r, b_r)$ where $r = \text{lex}(v)$. In the other direction, $\text{cfx}(a, b)$ is defined as that one of the two v , mapped by T back to (a, b) , for which the last entry $v(r)$ is greater than one. The other, call it \tilde{v} , is obtained by replacing $v(r)$ with $v(r) - 1$, and appending 1 as $\tilde{v}(r + 1)$. Thus

$$(9) \quad \#\{v \in V : \langle v \rangle \leq x \text{ and } v(i) \leq \alpha \log x - 1 \text{ for } 1 \leq i \leq \text{lex}(v)\} \\ \leq 2\Phi(x, \alpha) \leq \#\{v \in V : \langle v \rangle \leq x \text{ and } v(i) \leq \alpha \log x \\ \text{for } 1 \leq i \leq \text{lex}(v)\}.$$

In (9) we don't get equality because there are some $v \in V$ such that $v(r) - 1 \leq \alpha \log x < v(r)$. Given two sequences $u, w \in V$, we write uw for their concatenation. That is, uw denotes the sequence v such that $v(j) = u(j)$ for $j \leq \text{lex}(u)$, $\text{lex}(v) = \text{lex}(u) + \text{lex}(w)$, and $v(j) = w(j - \text{lex}(u))$ for $\text{lex}(u) < j \leq \text{lex}(u) + \text{lex}(w)$. With this notation, a well-known identity reads

$$(10) \quad \langle uw \rangle = \langle u \rangle \langle w \rangle (1 + \{u\}[w]).$$

Now if $d_j = u(j) > N$, (where in the subsequent application, $N = \lceil \alpha \log x \rceil$), then $\{u\} = b_{j-1}(u)/(d_j b_{j-1}(u) + b_{j-2}(u)) < 1/N$, so that for $u \in V_j$ with $u(j) > N$, and $w \in V$,

$$(11) \quad \langle u \rangle \langle w \rangle \leq \langle uw \rangle \leq (1 + 1/N) \langle u \rangle \langle w \rangle.$$

This gives us a way to estimate, for $1 \leq l \leq L$ say, the number of constructions of the form

$$v = u_1 k_1 u_2 k_2 \cdots u_l k_l u_{l+1}, \quad \text{with } u_1, u_2, \dots, u_{l+1} \in V,$$

$k_1, k_2, \dots, k_l \in V(1)$ or \mathbb{Z}^+ (which we equate by a sleight of notation), with all $k_i > N$, and with $\langle v \rangle \leq x$. Note that since V includes the empty sequence, there need not be any genuine interposition between consecutive k_i . Note also that the same sequence, if more than l of the $v(i)$ are greater than N , can be expressed in the above form in more than one way.

Our inclusion and exclusion argument is based on counting representations of v of the above form. For every integer $l \geq 0$, and every $v \in V$, let $\sigma(v, l, N)$ denote the number of ways in which v can be written as $u_1 k_1 \cdots u_l k_l u_{l+1}$, with all $k_i > N$. Then

$$(12) \quad \sum_{l=0}^{\infty} (-1)^l \sigma(v, l, N) = \begin{cases} 1 & \text{if all } v(i) \leq N, \\ 0 & \text{if any } v(i) > N, \end{cases}$$

and $\sum_{l=0}^K (-1)^l \sigma(v, l, N)$ alternates about this, being $\geq \{1 \text{ resp. } 0\}$ for K even, and $\leq \{1 \text{ resp. } 0\}$ for K odd. Now let

$$\Phi^-(x, \alpha) := \frac{1}{2} \#\{v \in V : \langle v \rangle \leq x \text{ and } v(i) \leq \alpha \log x - 1 \\ \text{for } 1 \leq i \leq \text{lex}(v)\},$$

and

$$\Phi^+(x, \alpha) := \frac{1}{2} \#\{v \in V : \langle v \rangle \leq x \text{ and } v(i) \leq \alpha \log x \\ \text{for } 1 \leq i \leq \text{lex}(v)\}.$$

Then with $N = [\alpha \log x]$ or $[\alpha \log x] - 1$ respectively,

$$(13) \quad \Phi^\pm(x, \alpha) = \frac{1}{2} \sum_{v \in V} \chi(\langle v \rangle \leq x) \sum_{l=0}^{\infty} (-1)^l \sigma(v, l, N) \\ = \frac{1}{2} \sum_{l=0}^{\infty} (-1)^l \sum_{u_i \in V} \sum_{k_i > N} \sum_{u_2 \in V} \\ \cdots \sum_{k_i > N} \sum_{u_{l+1} \in V} \chi(\langle u_1 k_1 \cdots u_l k_l u_{l+1} \rangle \leq x).$$

Now let $W_l(x, N)$ denote the number of pairs $((u_1, u_2, \dots, u_{l+1}), (k_1, k_2, \dots, k_l))$ where the $u_i \in V$ and the $k_i > N$, and such that $\langle u_1 k_1 u_2 k_2 \cdots u_l k_l u_{l+1} \rangle \leq x$. Let $W'_l(x, N)$ denote the number of such pairs for which $\prod_{i=1}^l k_i \prod_{j=1}^{l+1} \langle u_j \rangle \leq x$. Then from (11), we see that

$$(14) \quad W'_l((1 + 1/N)^{-2l} x, N) \leq W_l(x, N) \leq W'_l(x, n).$$

But

$$(15) \quad W'_l(x, N) = 2^{l+1} \sum_{k_i > N} \sum_{b_1=1}^{\infty} \sum_{k_2 > N} \sum_{b_2=1}^{\infty} \\ \cdots \sum_{k_l > N} \sum_{b_{l+1}=1}^{\infty} \prod_{i=1}^{l+1} \phi(b_i) \chi \left(\prod_{i=1}^{l+1} b_i \prod_{j=1}^l k_j \leq x \right).$$

In view of this, it is natural to seek estimates for

$$A_m(y) := \sum_{b \in V_m} \prod_{i=1}^m \phi(b_i) \chi \left(\prod_{i=1}^m b_i \leq y \right),$$

and then to apply them with $y = (x/k_1 \cdots k_l)$ and $m = l+1$. Another way to write (15), using the definition above, is

$$(16) \quad 2^m A_m(y) = \# \left\{ (u_1, u_2, \dots, u_m) : u_i \in V \right. \\ \left. \text{for } 1 \leq i \leq m \text{ and } \prod_{i=1}^m \langle u_i \rangle \leq y \right\}.$$

Once we have suitable estimates for $A_m(y)$, we will use (13), together with (17) below:

$$(17) \quad 2^{l+1} \sum_{k_1=N+1}^{\infty} \cdots \sum_{k_l=N+1}^{\infty} A_{l+1}((1 + 1/N)^{-2l} x/k_1 k_2 \cdots k_l) \\ \leq W_l(x, N) \leq 2^{l+1} \sum_{k_1=N+1}^{\infty} \cdots \sum_{k_l=N+1}^{\infty} A_{l+1}(x/k_1 k_2 \cdots k_l).$$

(There are only finitely many nonzero terms in the sums of (17), as $A_m(y) = 0$ for $y < 1$.) But (13) can now be written as

$$(18) \quad \Phi^{\pm}(x, \alpha) = \frac{1}{2} \sum_{l=0}^{\infty} (-1)^l W_l(x, N^{\pm}),$$

where $N = [\alpha \log x]$ or $[\alpha \log x] - 1$ for Φ^+ or Φ^- respectively.

We are now in a position to sketch out the proof of Theorem 1. First we obtain an estimate of the form (with $\lambda = 6/\pi^2$)

$$(19) \quad A_m(y) \approx \frac{1}{2} y^2 \lambda^m (\log y)^{m-1} / (m-1)!,$$

by a study of the Dirichlet series

$$(20) \quad \int_1^{\infty} t^{-s} dA_k(t) = \left(\sum_{n=1}^{\infty} n^{-s} \phi(n) \right)^k = (\zeta(s-1)/\zeta(s))^k.$$

Next, we estimate the sums of (17), which from (19) are given approximately by

$$(21) \quad 2^{l+1} \sum_{\substack{k_i=N+1 \\ k_1 k_2 \cdots k_l \leq x}}^{\infty} \cdots \sum_{k_l=N+1}^{\infty} \frac{1}{2} x^2 \lambda^{l+1} (l!)^{-1} \prod_{i=1}^l k_i^{-2} (\log(x/k_1 \cdots k_l))^l,$$

as

$$(22) \quad \frac{1}{2}x^2(2\lambda)^{l+1}(l!)^{-1} \times \iint \cdots \int_R \left(\log x - \sum_{i=1}^l \log t_i \right)^l t_1^{-2} \cdots t_l^{-2} dt_1 \cdots dt_l,$$

where $R = \{(t_1, \dots, t_l) : t_1 \geq N, \dots, t_l \geq N \text{ and } t_1 t_2 \cdots t_l \leq x\}$. Calculus and simplifying estimates then reduce the integral expression above to about $\frac{1}{2}x^2(2\lambda)^{l+1}(l!)^{-1}\alpha^{-l}$. Finally, from (18) we expect to find that $2\Phi(x, \alpha)$ is given approximately by

$$(23) \quad x^2\lambda - \frac{1}{2}x^2 \sum_{l=1}^{\infty} (-2\lambda)^{l+1}\alpha^{-l}/l! = \lambda x^2 \sum_{l=0}^{\infty} (-2\lambda/\alpha)^l/l! = \lambda x^2 e^{-2\lambda/\alpha},$$

which is roughly what is claimed in Theorem 1.

In §3 we give details for the estimation of $A_m(y)$. In §4 we give details of the resulting estimates of $W_l(x, N)$, and tie it all together.

3. Bounds for $A_m(y)$. Recall that $A_m(y) = \sum_{b_1 b_2 \cdots b_m \leq y} \prod_1^m \phi(b_i)$.

LEMMA 1. *There is a positive absolute constant C such that for $1 \leq k \leq C\sqrt{\log y}$, $A_k(y) = (\frac{1}{2}\lambda^k \log^{k-1} y / (k-1)!)y^2(1 + O(k^2/\log y))$.*

Proof. First we note that if Lemma 1 holds for integer $y \geq 3$, then it holds for real $y \geq 3$ as well. Also, the case $k = 1$ is the well-known result $\sum_{n \leq y} \phi(n) = \frac{1}{2}\lambda y^2(1 + O(1/\log y))$. Now let

$$(24) \quad f(s, k) = \left(\sum_{n=1}^{\infty} n^{-s} \phi(n) \right)^k = (\zeta(s-1)/\zeta(s))^k = \sum_{n=1}^{\infty} a(k, n)n^{-s}, \text{ say.}$$

The series representations of $f(s, k)$ are absolutely convergent, uniformly in $\text{Re}(s) \geq c$ for each $c > 2$, and the zeta function representation provides the analytic continuation into the domain $\text{Re}(s) \geq 4/3$, apart from a single pole of order k at $s = 2$.

For the analysis ahead, it will be more convenient to first study $B_k(y) := \sum_{n=1}^y A_k(n)$, and to establish (for some fixed $C, 0 < C < 1$), the following lemma.

LEMMA 2. *Uniformly in*

$$k \leq C\sqrt{\log y}, \quad \text{as } y \rightarrow \infty,$$

$$B_k(y) = ((1/6)y^3\lambda^k(\log y)^{k-1}/(k-1)!(1 + O(k^2/\log y))).$$

Before proving Lemma 2, we show how Lemma 1 follows from this secondary lemma.

Since $A_k(n)$ is increasing in n , for any integer m , $0 < m \leq y$, we have

$$(25) \quad mA_k(y) \leq B_k(y+m) - B_k(y).$$

Now from Lemma 2,

$$(26) \quad B_k(y+m) - B_k(y) \\ = ((1/6)\lambda^k/(k-1)!)((y+m)^3 \log^k(y+m) - y^3 \log^k y) \\ + O(k^2 y^3 \lambda^k \log^{k-2} y / (k-1)!).$$

Taking $m = [ky/\log y]$, and bearing in mind that $k \leq C\sqrt{\log y}$, this gives

$$(27) \quad B_k(y+m) - B_k(y) \\ = \left(\frac{1}{2} m \lambda^k y^2 \log^{k-1} y / (k-1)! \right) (1 + O(k^2/\log y)).$$

Thus

$$(28) \quad A_k(y) \leq \left(\frac{1}{2} \lambda^k y^2 \log^{k-1} y / (k-1)! \right) (1 + O(k^2/\log y)).$$

A similar calculation, starting from $A_k(y) \geq B_k(y) - B_k(y-m)$, gives a reversed version of (28). Taken together, these constitute the conclusion of Lemma 1.

We now turn to the proof of Lemma 2. By Perron's formula, for $c > 2$ we have

$$(29) \quad B_k(y) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} (y^{s+1}/s(s+1))f(s, k) ds.$$

It is well known that $\zeta(s) = O(\theta^{-1}|s|^\theta)$, uniformly in $0 < \theta \leq 1/2$ and $\text{Re}(s) = 1 - \theta$. With $\theta = 1/2k$, it follows that for some fixed $C_1 > 1$, and uniformly in $k \geq 2$, $\text{Re}(s) = 2 - 1/2k$,

$$(30) \quad f(s, k) = O(C_1^k k^k |s|^{1/2}).$$

Although it is not essential to the proof, it will be convenient to have $C_1 = 4$. A little detail work, starting with the formula [11]

$$\zeta(s) = s/(s - 1) + s \sum_{n=1}^{\infty} \int_0^1 u(n + u)^{-s-1} du$$

valid for $\text{Re}(s) > 0$, is now in order. For $n \leq |s|$ in the sum, one uses integration by parts, and with the obvious bounds for the other terms, this gives, for $s = 1 - \theta$,

$$|\zeta(s)| \leq \theta^{-1} + \theta^{-1}(1 + |s|^\theta) + (1/2)|s|^\theta/(1 - \theta).$$

For $s > 1$, $|\zeta(s)| \leq \zeta(\sigma)$, so with $\theta = 1/2k$, the claim that C_1 can be 4 holds provided

$$4k + 2k|s|^\theta + \left(\frac{1}{2 - 2\theta}\right) |s|^\theta \leq 4k\zeta(2 - 1/2k)|s|^\theta.$$

But $4k\zeta(2 - 1/2k) > 4k\zeta(2) + 1$ since $\zeta'(\sigma) < -1/4$ for $1 < \sigma \leq 2$, so we just need

$$4 \leq (4\zeta(2) - 2)|s|^\theta.$$

The worst case is $k = 2$, $s = 3/4$, and even then $4 \leq 4.2619\dots$

Now let Γ be the linear path from $3 - i\infty$ to $3 + i\infty$, and let $\Gamma_{N,k}$ be the counterclockwise circuit of the rectangle with corners $3 - iN$, $3 + iN$, $(2 - 1/2k) + iN$, and $(2 - 1/2k) - iN$. Then

$$\begin{aligned} (31) \quad \lim_{N \rightarrow \infty} \left(\int_{\Gamma} - \int_{\Gamma_{N,k}} \right) (y^{s+1} f(s, k)/s(s + 1)) ds \\ = \int_{2-1/2k-i\infty}^{2-1/2k+i\infty} (y^{s+1} f(s, k)/s(s + 1)) ds = E_1(y, k), \text{ say.} \end{aligned}$$

In view of (30), $E_1(y, k) = O(C_1^k k^k y^{3-1/2k})$. For $k^2 \leq C_1^{-2} \log y$, a simple calculation now shows that

$$E_1(y, k) = O(y^3 \lambda^k (\log y)^{k-2} / k^2 (k - 1)!),$$

which is the error allowed for in Lemma 2.

REMARK. The argument fails here without some hypothesis on C_1 .

This brings us to the kernel of the matter: we must evaluate the integral over $\Gamma_{N,k}$ to within $O(y^3 \lambda^k (\log y)^{k-2} / k^2 (k - 1)!)$.

Let $\beta(s, k) = (s - 2)^k f(s, k)/s(s + 1)$. Then

$$\begin{aligned} (32) \quad \frac{1}{2\pi i} \int_{\Gamma_{N,k}} (y^{s+1} f(s, k)/s(s + 1)) ds \\ = y^3 \int_{\Gamma_{N,k}} ((s - 2)^{-k} \beta(s, k) y^{s-2} / 2\pi i) ds, \end{aligned}$$

and the latter integral is, by the residue theorem, equal to the $(s-2)^{k-1}$ coefficient, say $T_{k-1}(y)$, in the Taylor series expansion of $\beta(s, k)y^{s-2}$ about 2. To estimate this, we first note that for a complex analytic function ξ on a disk of radius r , if $|\xi| \leq K$ on the disk, then by the Plancherel formula, $|(d^j/ds^j)\xi(s)| \leq Kj!r^{-j}$ at the center.

Now $(s-2)\zeta(s-2)/\zeta(s-1) = \lambda(1+a(s-2) + O(s-2)^2)$, uniformly in $|s-2| \leq 1/2$, say. Thus for arbitrary j , $1 \leq j \leq k$, on the disk $|s-2| \leq j/2k$, we have

$$(33) \quad \beta(s, k) = O(\lambda^k \exp(O(j))),$$

so that from the observation above, if $D_j(k) = (d^j/ds^j)\beta(s, k)$ evaluated at $s = 2$, then for $j \leq k$,

$$(34) \quad D_j(k) = O((2k/j)^j j! \lambda^k \exp(O(j))).$$

(For $j = 0$, we have $D_j(k) = \lambda^k/6$, of course.) Now

$$T_{k-1}(y) = ((k-1)!)^{-1} (d^{k-1}/ds^{k-1})(y^{s-2}\beta(s, k)),$$

evaluated at $s = 2$. Expanding the iterated derivative of a product as in the binomial theorem, we get

$$(35) \quad (k-1)!T_{k-1}(y) = \sum_{j=0}^{k-1} \binom{k-1}{j} (\log y)^{k-1-j} D_j(k).$$

The main term here, corresponding to $j = 0$, is $(1/6)\lambda^k \log^{k-1} y$. For $j \geq 1$, we have, in the sum above,

$$(36) \quad \begin{aligned} \binom{k-1}{j} (\log y)^{k-1-j} D_j(k) &= O((k^j/j!)(\log y)^{k-1-j}(2k/j)^j j! \lambda^k \exp(O(j))) \\ &= (\log y)^{k-1} \cdot O_\varepsilon(k^{2j}(\log y)^{-j}(j^{-(1-\varepsilon)j})). \end{aligned}$$

Thus, for $k^2 \leq \log y$,

$$(37) \quad T_{k-1}(y) = (\lambda^k \log^{k-1} y / (k-1)!)(1 + O(k^2/\log y)),$$

which completes the proof of Lemma 2. With $C_1 = 4$, C in Lemma 1 becomes $1/4$. We need another estimate for the case of large k .

LEMMA 3. For $k \geq 1$ and $y \geq 1$,

$$A_k(y) \leq 4^{k+1} y^{2+3/2\pi^2}.$$

Proof. First note that this is trivial from the definition if $1 \leq y < 4$, or if $k = 1$. Now in (29), take $c = 2 + 3/2\pi^2$. From this, it follows that with $s = (2 + 3/2\pi^2) + i\tau$,

$$(38) \quad B_k(y) \leq (y^{c+1}/2\pi) \int_{-\infty}^{\infty} |f(s, k)|/|s(s + 1)| d\tau.$$

For $\sigma = \text{Re}(s) > 2$, by the product representation of the zeta function, and elementary properties of the linear fractional $(1 + zp^{-\sigma})/(1 + zp^{1-\sigma})$ on the circle $|z| = 1$, we have

$$|\zeta(s - 1)/\zeta(s)| \leq |\zeta(\sigma - 1)/\zeta(\sigma)|.$$

Thus from (38), $B_k(y) \leq (\frac{1}{2\pi})y^{c+1}(\zeta(c - 1)/\zeta(c))^k \cdot \pi/c$. Now taking $m = [y/4]$ in (26) gives

$$(39) \quad A_k(y) \leq (3/5)(7/2)^k m^{-1}((y + m)^{c+1} - y^{c+1}).$$

Since this expression is increasing in m/y ,

$$A_k(y) \leq (3/5)(7/2)^k 4((5/4)^{c+1} - 1)y^c,$$

which for $k \geq 2$ is $\leq 3 \cdot 4^k y^c < 4^{k+1} y^c$. This proves Lemma 3.

4. Estimation of $W_l(x, N)$. From (14), (15) and Lemma 1, we have

$$(40) \quad W_l(x, N) \leq \frac{x^2 \lambda^{l+1} 2^{l+1}}{2(l!)}$$

$$\sum_{\substack{k_1 k_2 \dots k_l \leq x \exp(-16(l+1)^2) \\ k_i > N \text{ for } 1 \leq i \leq l}} k_1^{-2} k_2^{-2} \dots k_l^{-2} \left(\log x - \sum_1^l \log k_i \right)^l$$

$$\cdot \left(1 + O \left(l^2 / \left(\log x - \sum_1^l \log k_i \right) \right) \right)$$

$$+ 2^{l+1} \sum_{\substack{x \exp(-16(l+1)^2) < k_1 k_2 \dots k_l \leq x \\ k_i > N \text{ for } 1 \leq i \leq l}} A_{l+1}(x/k_1 k_2 \dots k_l).$$

In the second term here, $u = \log x - \sum_1^l \log k_i < 16(l + 1)^2$, so that Lemma 1 is not applicable. Happily, for this term there is no need of

sharp estimates. We get a crude, but adequate, bound from

LEMMA 4. For $l \geq 1$, $N \geq 8$ and $x \geq (N + 1)^l e^{16l^2}$,

$$\begin{aligned}
 & 2^{l+1} \sum_{\substack{x \exp(-16(l+1)^2) < k_1 k_2 \cdots k_l \leq x \\ k_i > N \text{ for } 1 \leq i \leq l}} A_{l+1}(x/k_1 k_2 \cdots k_l) \\
 & \ll ((16)^l x \exp(16l^2(1 + 3/(2\pi^2))) (\log x)^l / l!).
 \end{aligned}$$

The application of the lemma will be to cases in which $N \leq (\log x)^2$ and $l \leq (\log x)^{1/3}$, so that the upper bound given in Lemma 4 comes to $O_\varepsilon(x^{1+\varepsilon})$, or what is good enough for our purposes, to $O(x^{3/2})$.

To prove Lemma 4, we first note that from Lemma 3,

$$(41) \quad A_{l+1}(y) \leq 4^{l+2} y^{2+\lambda/4}.$$

Thus

$$\begin{aligned}
 (42) \quad & 2^{l+1} \sum_{\substack{x \exp(-16(l+1)^2) < k_1 k_2 \cdots k_l \leq x \\ k_i > N \text{ for } 1 \leq i \leq l}} A_{l+1}(x/k_1 k_2 \cdots k_l) \\
 & \leq (16)^{l+1} x^2 \sum_{\text{(same range)}} k_1^{-2} \cdots k_l^{-2} \exp\left(\frac{\lambda}{4} \left(\log x - \sum_1^l \log k_i\right)\right) \\
 & \leq (16)^{l+1} x^{2+\lambda/4} \sum_{\text{all } k_i > N, \prod_1^l k_i > x \exp(-16(l+1)^2)} \prod_1^l k_i^{-(2+\lambda/4)}.
 \end{aligned}$$

The sum in the right side of (42) above is itself

$$\leq \iint \cdots \int_R \prod_{i=1}^l t_i^{-(2+\lambda/4)} dt_1 \cdots dt_l,$$

where $R = \{(t_1, t_2, \dots, t_l) : t_i \geq N \text{ for } 1 \leq i \leq l \text{ and } \prod_1^l t_i \geq x e^{-16l^2}\}$.

On setting $s_i = \log t_i$, $1 \leq i \leq l$, this integral becomes

$$\iint \cdots \int_{R'_l} \exp\left(-\beta \sum_{i=1}^l s_i\right) ds_l \cdots ds_1,$$

where $\beta = 1 + \lambda/4$, and where

$$R'_l = \left\{ (s_1, \dots, s_l) : s_i \geq \log N \text{ for } 1 \leq i \leq l \right. \\ \left. \text{and } \sum_1^l s_i \geq \log x - 16l^2 \right\}.$$

Seen as an iterated integral, the innermost integral is a function of s_1, s_2, \dots, s_{l-1} and is

$$\int_{\max(\log x - 16l^2 - s_1 - s_2 - \dots - s_{l-1}, \log N)}^{\infty} \exp(-\beta(s_1 + s_2 + \dots + s_{l-1})) e^{-\beta s_l} ds_l \\ \leq \min\{x^{-\beta} e^{16\beta l^2}, N^{-\beta} e^{-\beta(s_1 + s_2 + \dots + s_{l-1})}\}.$$

Thus the original multiple integral is

$$\leq x^{-\beta} e^{16\beta l^2} \text{Vol} \left(\left\{ (s_1, s_2, \dots, s_{l-1}) : \log x - 16l^2 - \log N \right. \right. \\ \left. \left. > \sum_1^{l-1} s_i \text{ and all } s_i < \log N \right\} \right) \\ + N^{-\beta} \iint \dots \int_{R'_{l-1}} e^{-\beta(s_1 + s_2 + \dots + s_{l-1})} ds_{l-1}, \dots ds_1,$$

where $R'_{l-1} := \{(s_1, s_2, \dots, s_{l-1}) : s_i \geq \log N \text{ and } \sum_1^{l-1} s_i \geq \log x - 16l^2 - \log N\}$. The first term above is just $x^{-\beta} \exp(16\beta l^2)(\log x)^l/l!$, while the second term is of the same form as the original integral. Hence, we proceed by induction. Let

$$F(l, z) := \iint \dots \int_{R(l, z)} \exp\left(-\beta \sum_{i=1}^l s_i\right) ds_l \dots ds_1,$$

where $R(l, z) := \{(s_1, \dots, s_l) : s_i \geq \log N \text{ for } 1 \leq i \leq l \text{ and } \sum_{i=1}^l s_i \geq z\}$. In this terminology, we have shown above that

$$(43) \quad F(l, z) \leq e^{-\beta z} (z + 16l^2)^l/l! + N^{-\beta} F(l-1, z - \log N).$$

Now $F(1, z) = \int_{\max(\log N, z)}^{\infty} e^{-\beta s} ds = \beta^{-1} \min(N^{-\beta}, e^{-\beta z})$, and in particular if $z > \log N$ then $F(1, z) = \beta^{-1} e^{-\beta z}$. Now from this and from (43), if $z > l \log N$ then

$$(44) \quad F(l, z) \leq e^{-\beta z} \sum_{j=2}^l (z + 16j^2)^j N^{\beta(j-L)} / j!.$$

Since $z > \log N > 2l$ under the assumption $N \geq 8$ in Lemma 4, the sum in (44) is dominated by the last term $(z + 16l^2)^l/l!$, so that

$$(45) \quad F(l, z) \ll e^{-\beta z}((z + 16l^2)^l/l!).$$

We apply (45) with $z = \log x = 16l^2$ to obtain, for $x > N^l e^{16l^2}$,

$$(46) \quad \iint \cdots \int_R \prod_{i=1}^l t_i^{-(2+\lambda/4)} dt_1 \cdots dt_l \ll x^{-\beta} \exp(16l^2\beta)((\log x)^l/l!).$$

In view of (42) and the following inequality, this proves Lemma 4.

For x sufficiently large, though, if $l \leq (\log x)^{1/3}$ and $N \leq (\log x)^2$, then

$$(47) \quad (16)^l x \exp(16\beta l^2)(N^\beta + (\log x)^l/l!) < x^{3/2}.$$

Thus for large x the second term in (40) is negligible, even in comparison to the potential error in the first term of (40). The main term of that, putting aside for now the contribution from the “ O ” in $(1 + O(l^2/(\log x - \sum_1^l \log k_i)))$, is

$$\sum_{\substack{k_1 k_2 \cdots k_l \leq x \exp(-16(l+1)^2) \\ k_i > N \text{ for } 1 \leq i \leq l}} k_1^{-2} k_2^{-2} \cdots k_l^{-2} \left(\log x - \sum_1^l \log(k_i) \right)^l.$$

But this is less than

$$\int_{\substack{t_1 t_2 \cdots t_l \leq x \exp(-16l^2) \\ t_i \geq N \text{ for } 1 \leq i \leq l}} \cdots \int (t_1^{-2} t_2^{-2} \cdots t_l^{-2})(\log x)^l dt_1 \cdots dt_l \leq (\log x/N)^l.$$

The error term just put aside is likewise

$$\ll l^2 \int \cdots \int (t_1^{-2} \cdots t_l^{-2})(\log x)^{l-1} dt_1 \cdots dt_l \ll (l^2/\log x)(\log x/N)^{l-2}$$

Thus for x sufficiently large, $l \leq (\log x)^{1/3}$ and $N \leq (\log x)^2$,

$$(48) \quad W_l(x, N) \leq (1 + O(l^2/\log x))(\log x/N)^l (2^l \lambda^{l+1}/l!) x^2 + O(x^{3/2}).$$

Next we obtain a similar lower bound for $W_l(x, N)$. From (17) and Lemma 1, we have

$$(49) \quad W_l(x, N) \geq (x^2 2^l \lambda^{l+1} / l!) \sum_{\substack{k_1 k_2 \dots k_l \leq x \exp(-16(l+1)^2)(1+1/N)^{-2l} \\ k_i > N \text{ for } 1 \leq i \leq l}} k_1^{-2} k_2^{-2} \dots k_l^{-2} \cdot \left(\log x - \sum_{i=1}^l \log((1 + 1/N)^2 k_i) \right)^l + \text{two error terms.}$$

Let $x' = x(1 + 1/N)^{-2l}$, and let $S_l := \{K = (k_1, k_2, \dots, k_l) : k_1 k_2 \dots k_l \leq x' \exp(-16(l + 1)^2) \text{ and } k_i > N \text{ for } 1 \leq i \leq l\}$. The first of the above-mentioned error terms stems from the factor $1 + O(k^2 / \log y)$ in Lemma 1. For $K \in S_l$, this factor, applied to each of the contributions to the sum in (49), is $1 + O((l + 1)^2 / \log x)$ so that the whole sum is also perturbed by only a factor of $(1 + O((l + 1)^2 / \log x))$ due to that source of error. The other term in (49) is the contribution to $\sum \sum \dots \sum A_{l+1}(x' / k_1 k_2 \dots k_l)$ due to $K = (k_1, k_2, \dots, k_l)$ for which $k_i > N$, $1 \leq i \leq l$, but $\prod_{i=1}^l k_i > x' \exp(-16(l + 1)^2)$.

For x sufficiently large, if $N \leq \log^2 x$ and $l \leq \log^{1/3} x$, then the hypotheses of Lemma 4 are satisfied, so that this error term is $O((16)^l x' e^{16l^2 \beta} ((\log x')^l / l!))$ and thus $O(x^{3/2})$ as before. Hence, for such x , N and l ,

$$(50) \quad W_l(x, N) \geq (2^l (x')^2 \lambda^{l+1} / l!) (1 + O((l + 1)^2 / \log x)) P + O(x^{3/2}),$$

where

$$P = \sum_{\substack{k_1 k_2 \dots k_l \leq x' \exp(-16(l+1)^2) \\ k_i > N \text{ for } 1 \leq i \leq l}} k_1^{-2} k_2^{-2} \dots k_l^{-2} \left(\log(x') - \sum_{i=1}^l \log k_i \right)^l.$$

Now we need a lower bound for P . Clearly,

$$P \geq \int_R t_1^{-2} t_2^{-2} \dots t_l^{-2} \left(\log(x') - \sum_{i=1}^l \log t_i \right)^l dt_l \dots dt_1,$$

where $R = \{(t_1, t_2, \dots, t_l) : t_1 t_2 \dots t_l \leq x' e^{-16(l+1)^2} \text{ and } t_i \geq N + 1 \text{ for } 1 \leq i \leq l\}$. After a change of variables ($u_i = \log t_i - \log(N + 1)$),

$1 \leq i \leq l$) this integral becomes

$$(N + 1)^{-l} \int_U e^{-(u_1+u_2+\dots+u_l)} \cdot \left(\log(x') - l \log(N + 1) - \sum_{i=1}^l u_i \right)^l du_l \cdots du_1,$$

where $U = \{(u_1, u_2, \dots, u_l) : \sum_{i=1}^l u_i \leq \log(x') - 16(l + 1)^2 - l \log(N + 1) \text{ and } u_i \geq 0 \text{ for } 1 \leq i \leq l\}$. This, though, is just

$$(N + 1)^{-l} \int_0^L (u^{l-1}/(l - 1)!) e^{-u} (M - u)^l du,$$

where $L = \log(x') = l \log(N + 1) - 16(l + 1)^2$ and $M = \log(x') - l \log(N + 1)$. Thus

$$(51) \quad P \geq ((N + 1)^{-l}/(l - 1)!) \left\{ \int_0^\infty - \int_L^\infty (u^{l-1} e^{-u} (M - u)^l du \right\}.$$

The $-\int_L^\infty$ contribution here is quite small. In fact, for large x , for $N \leq \log^2 x$ and for $l \leq (\log x)^{1/3}$,

$$\int_L^\infty u^{l-1} (M - u)^l e^{-u} du \leq \int_L^\infty u^{2l-1} e^{-u} du \leq 2^{2l-1} e^{-L},$$

this last because $(1 + 1/u)^{2l-1} e^{-1} < \frac{1}{2}$ throughout the interval of integration. But in view of the constraints on l and N , $2^{2l-1} e^{-L} \leq x^{-3/4}$ for large x . The main term in our lower bound for P is

$$\begin{aligned} & ((N + 1)^{-l}/(l - 1)!) \int_0^\infty u^{l-1} e^{-u} (M - u)^l du \\ &= ((N + 1)^{-l}/(l - 1)!) \sum_{j=0}^l (-1)^j \binom{l}{j} M^{l-j} \int_0^\infty u^{l-1+j} e^{-u} du \\ &= (N + 1)^{-l} M^l \sum_{j=0}^l (-M)^{-j} l(l - 1 + j)! / ((j!)(l - j)!). \end{aligned}$$

In view of the constraints on l and N , we have $M \geq (1 - \varepsilon) \log x$ (ε may be taken as small as we please by choosing a large enough lower bound for x). Thus the last sum above is dominated by its first term, and it simplifies to $(1 + O(l^2/\log x))$. Thus

$$(52) \quad P = (1 + O(l^2/\log x))(N + 1)^{-l} M^l.$$

From (50), we now conclude that

$$(53) \quad W_l(x, N) \geq (2^l(x')^2\lambda^{l+1}/l!)(1 + O((l + 1)^2/\log x))(N + 1)^{-l} \cdot (\log(x') - l \log(N + 1))^l + O(x^{3/2}).$$

Since $x' = (1 + 1/N)^{-2l}x$, this boils down to

$$(54) \quad W_l(x, N) \geq (2^l x^2 \lambda^{l+1} (\log x)^l / N^l (l!)) (1 + 1/N)^{-3l} \times (1 + O(l^2 \log \log x / \log x)) + O(x^{3/2}),$$

for $1 \leq l \leq (\log x)^{1/3}$, $N \leq (\log x)^2$ and x sufficiently large. Together with (48), and under the same constraints, this gives

$$(55) \quad W_l(x, N) = \frac{x^2 2^l \lambda^{l+1}}{(l!)} \left(\frac{\log x}{N} \right)^l \times \exp \left(O \left(\frac{l^2 \log \log x}{\log x} \right) + O \left(\frac{l}{N} \right) \right) + O(x^{3/2}).$$

Now from (13),

$$\Phi^\pm(x, \alpha) = \frac{1}{2} \sum_{l=0}^\infty (-1)^l W_l(x, N^\pm),$$

with $N^+ = [\alpha \log x]$ for Φ^+ and $N^- = [\alpha \log x] - 1$ for Φ^- . From (12), if we truncate this sum we get lower and upper bounds: if A is odd and $B = A + 1$, then

$$(56) \quad \frac{1}{2} \sum_{l=0}^A (-1)^l W_l(x, N^-) < \Phi(x, \alpha) < \frac{1}{2} \sum_{l=0}^B (-1)^l W_l(x, N^+).$$

If we choose $B = [(\log x)^{1/3}]$, then for $\alpha > 4/\log \log x$ and $l \leq B$, the $l^2 \log \log x / \log x$ contribution to the error factor in (55) dominates that from l/N , and both are small, so that (55) boils down to

$$(57) \quad W_l(x, N^\pm) = \frac{x^2 2^l \lambda^{l+1}}{(l!)} \left(\frac{\log x}{N^\pm} \right)^l \left(1 + O \left(\frac{l^2 \log \log x}{\log x} \right) \right) + O(x^{3/2})$$

for such l , N , and x . Thus in (56), the main terms are

$$\frac{1}{2} \sum_{l=0}^A x^2 2^l \lambda^{l+1} (-\log x / N^-)^l / l! \quad \text{and} \\ \frac{1}{2} \sum_{l=0}^B x^2 2^l \lambda^{l+1} (-\log x / N^+)^l / l!,$$

and the error factor in (57) perturbs these by at most

$$O\left(x^2 \sum_{l=0}^{\infty} 2^l \lambda^{l+1} (\log x / N^-)^l (l^2 \log \log x / \log x) / l!\right).$$

But this is $O((x^2 \log \log x / \log x)(z^2 + 1)e^z)$, where $z = 2\lambda \log x / N^-$. Now for $\alpha \geq 4 / \log \log x$, $N^- \geq (4 \log x / \log \log x) - 2$ so that $z \leq \frac{1}{2} \lambda \log \log x$, and

$$(x^2 \log \log x / \log x)(z^2 + 1)e^z \ll x^2 (\log \log x (1 + \alpha^{-2}) / \log x) e^{2\lambda/\alpha}.$$

This brings us to the main terms in (56). They are

$$\frac{1}{2} \lambda x^2 \sum_{l=0}^A \left(\frac{2\lambda \log x}{N^-}\right)^l / l!, \quad \text{and} \quad \frac{1}{2} \lambda x^2 \sum_{l=0}^B \left(\frac{2\lambda \log x}{N^+}\right)^l / l!$$

respectively. If we replace A and B with ∞ in these sums, the resulting change is $O((2\lambda/\alpha)^B / B!)$, and with $B = [\log x^{1/3}]$, that is $\ll 1 / \log x$. Thus the main terms above are

$$\frac{1}{2} \lambda x^2 (\exp(-2\lambda \log x / N^\pm) + O(1 / \log x)).$$

Replacing $(\log x / N^\pm)$ with $1/\alpha$ here introduces an error factor of $\exp(O(1/\alpha^2 \log x))$, so that the main terms boil down to

$$\frac{1}{2} \lambda x^2 e^{-2\lambda/\alpha} (1 + O((1 + \alpha^{-2}) / \log x)).$$

That is,

$$\begin{aligned} (58) \Phi(x, \alpha) &= \frac{1}{2} \lambda x^2 e^{-2\lambda/\alpha} (1 + O(1/\alpha^2 \log x) + O(1 / \log x)) \\ &\quad + O(x^2 \log \log x (1 + \alpha^{-2}) / \log x) e^{2\lambda/\alpha} + O(x^{3/2}) \\ &= \frac{1}{2} \lambda x^2 e^{-2\lambda/\alpha} (1 + O(e^{4\lambda/\alpha} \log \log x (1 + \alpha^{-2}) (\log x)^{-1})), \end{aligned}$$

for all sufficiently large x and all α , $4 / \log \log x \leq \alpha \leq (\log x)^2$. The condition $N \leq (\log x)^2$, which roughly coincides with $\alpha \leq \log x$, has been necessary in the workings of the main argument. But for $\alpha \geq \log x$, the claim made by Theorem 1 reduces to an assertion that $\Phi(x, \alpha) = \frac{1}{2} \lambda x^2 (1 + O(\log \log x / \log x))$. Now $\Phi(x, \alpha)$ is a nondecreasing function of α . But the upper bound part of this follows from Lemma 1, while the required lower bound follows from what we have proved above, on taking $\alpha = \log x$. Thus the theorem, while of no interest in this case, happens nonetheless to hold.

REFERENCES

- [1] R. Bumby, *Hausdorff dimension of sets arising in number theory*, Lecture Notes in Mathematics, **1135** (1-8) New York Number Theory Seminar, Springer NY 1985.
- [2] T. W. Cusick, *Continuants with bounded digits*, *Mathematika*, **24** (1977), 166–172.
- [3] —, *Continuants with bounded digits II*, *Mathematika*, **25** (1978), 107–109.
- [4] J. Galambos, *The distribution of the largest coefficient in continued fraction expansions*, *Quart. J. Math. Oxford Ser. (2)*, **23** (1972), 147–151.
- [5] —, *The largest coefficient in continued fractions and related problems, Diophantine approximation and its applications* (Proc. Conf. Washington D. C. 1972) 101–109, Academic Press, NY 1973.
- [6] I. J. Good, *The fractional dimension theory of continued fractions*, *Proc. Cambridge Phil. Soc.*, **37** (1941), 199–228.
- [7] D. Hensley, *A truncated Gauss-Kuzmin law*, *Trans. Amer. Math. Soc.*, **306** (1) 1988.
- [8] —, *The distribution of badly approximable numbers and continuants with bounded digits*, *Proc. Int. Conference on Number Theory, Quebec 1987* (371–385).
- [9] —, *The distribution of badly approximable numbers and continuants with bounded digits, II*, *J. Number Theory*, **34** (1990), 293–334.
- [10] —, *Continued fraction Cantor sets, Hausdorff dimension, and functional analysis*, *J. Number Theory*, to appear.
- [11] Aleksandr Ivic, *The Riemann Zeta Function*, Chapter 1, Wiley, NY 1985.

Received March 7, 1990 and in revised form October 25, 1990.

TEXAS A & M UNIVERSITY
COLLEGE STATION, TX 77843-3368

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN

(Managing Editor)

University of California

Los Angeles, CA 90024-1555-05

HERBERT CLEMENS

University of Utah

Salt Lake City, UT 84112

THOMAS ENRIGHT

University of California, San Diego

La Jolla, CA 92093

NICHOLAS ERCOLANI

University of Arizona

Tucson, AZ 85721

R. FINN

Stanford University

Stanford, CA 94305

VAUGHAN F. R. JONES

University of California

Berkeley, CA 94720

STEVEN KERCKHOFF

Stanford University

Stanford, CA 94305

C. C. MOORE

University of California

Berkeley, CA 94720

MARTIN SCHARLEMANN

University of California

Santa Barbara, CA 93106

HAROLD STARK

University of California, San Diego

La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

(1906-1982)

B. H. NEUMANN

F. WOLF

(1904-1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1991 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 151, No. 2 December, 1991

Michael G. Eastwood and A. M. Pilato , On the density of twistor elementary states	201
Brian E. Forrest , Arens regularity and discrete groups	217
Yu Li Fu , On Lipschitz stability for F.D.E	229
Douglas Austin Hensley , The largest digit in the continued fraction expansion of a rational number	237
Uwe Kaiser , Link homotopy in \mathbb{R}^3 and S^3	257
Ronald Leslie Lipsman , The Penney-Fujiwara Plancherel formula for abelian symmetric spaces and completely solvable homogeneous spaces	265
Florin G. Radulescu , Singularity of the radial subalgebra of $\mathcal{L}(F_N)$ and the Pukánszky invariant	297
Albert Jeu-Liang Sheu , The structure of twisted $SU(3)$ groups	307
Morwen Thistlethwaite , On the algebraic part of an alternating link	317
Thomas (Toma) V. Tonev , Multi-tuple hulls	335
Arno van den Essen , A note on Meisters and Olech's proof of the global asymptotic stability Jacobian conjecture	351
Hendrik J. van Maldeghem , A characterization of the finite Moufang hexagons by generalized homologies	357
Bun Wong , A note on homotopy complex surfaces with negative tangent bundles	369
Chung-Tao Yang , Any Blaschke manifold of the homotopy type of CP^n has the right volume	379