

Pacific Journal of Mathematics

LINK HOMOTOPY IN \mathbb{R}^3 AND S^3

UWE KAISER

LINK HOMOTOPY IN \mathbb{R}^3 AND S^3

UWE KAISER

We give the general homotopy classification of 2-component link maps in \mathbb{R}^3 and study 3-component link maps in S^3 .

Introduction. For any sequence of integer numbers $p_1 \geq p_2 \geq \dots \geq p_r \geq 0$ by an r -link map is meant a collection of continuous maps

$$f = \coprod_{1 \leq j \leq r} f_j: \coprod_{1 \leq j \leq r} S^{p_j} \rightarrow \mathbb{R}^3 \text{ or } S^3$$

with mutually disjoint images. A link homotopy is a homotopy through link maps.

In [M] J. Milnor studied the case $p_1 = \dots = p_r = 1$ and classified links up to homotopy for $r = 2$ and $r = 3$. The classification in case $r > 3$ has recently been given by N. Habegger and S. Lin. Note that for $p_1 \leq 1$ the classifications in \mathbb{R}^3 and S^3 coincide. Moreover in this case all involved 0-spheres can be omitted by transversality.

We write (p, q) and (p, q, r) instead of (p_1, p_2) and (p_1, p_2, p_3) . Let $E(p, q)$, resp. $L(p, q, r)$, denote the set of link homotopy classes of link maps $S^p \amalg S^1 \rightarrow \mathbb{R}^3$, resp. $S^p \amalg S^q \amalg S^r \rightarrow S^3$.

The starting point is the following easy consequence of the sphere theorem (compare [Ko1]).

PROPOSITION. *If $q > 0$, and $p > 1$, then every link map $f: S^p \amalg S^q \rightarrow S^3$ is link homotopic to a trivial link map.*

Furthermore link maps $S^p \amalg S^0 \rightarrow S^3$ are easily seen to be classified by the homotopy group $\pi_p S^2$.

It is a remarkable fact that link homotopy in \mathbb{R}^3 contains a considerable amount of additional information. This is solely caused by the hole at $\infty \in S^3$ (compare [K1, K2]). On the other hand the strength of the sphere theorem implies that expectable phenomena are fully present, at least for $r = 2$.

There are two obvious constructions briefly described as follows: for $q < 3$ take the standard embedding $S^1 \subset S^3$ and map S^p into the complement which contains an embedded $S^{3-q-1} \vee S^2$ as deformation

retract. This defines

$$e_* : [S^p, S^{3-q-1} \vee S^2] \rightarrow E(p, q),$$

[,] is the set of unbased homotopy classes. In the general situation we map one of the spheres onto the origin of \mathbb{R}^3 and wrap the second sphere into $S^2 \subset \mathbb{R}^3$. This defines

$$pt_* : \pi_p S^2 \vee \pi_q S^2 \rightarrow E(p, q).$$

Here, for two based sets M, N , i.e. sets with distinguished elements m_0, n_0 , let $M \vee N$ denote $\{(m, n) \in M \times N \mid m = m_0 \text{ or } n = n_0\}$. If M, N are topological spaces, then $M \vee N$ is the usual wedge.

THEOREM 1. *The following assignments are 1-1 and onto:*

$$\begin{aligned} e_* : [S^p, S^{3-q-1} \vee S^2] &\rightarrow E(p, q), & \text{if } q \leq 1, \\ pt_* : \pi_p S^2 \vee \pi_q S^2 &\rightarrow E(p, q), & \text{if } q > 1. \end{aligned}$$

Note that the nontrivial elements of $[S^p, S^1 \vee S^2]$ are in 1-1 correspondence with sequences $(a_k)_{k \in \mathbb{N}}$, such that $a_1 \neq 0$, $a_k \in \pi_p S^2$ for $k \in \mathbb{N}$, almost all a_k trivial.

The techniques we develop to handle 2-link maps in \mathbb{R}^3 can easily be applied to 3-link maps in S^3 . Define pt_* into $L(p, q, r)$ as above by mapping two spheres constantly. Let $j_* : E(p, q) \rightarrow L(p, q, 1)$ be defined by mapping the q -sphere onto $\infty \in S^3$ and identify $S^3 \setminus \infty \approx \mathbb{R}^3$. Define e_* into $L(p, 1, 1)$ by taking the unlinked disjoint union L of two unknotted circles and then mapping S^p into an embedded $S^2 \vee S^1 \vee S^1$, which is a deformation retract of $S^3 \setminus L$.

THEOREM 2. *The following assignments are 1-1 and onto:*

$$\begin{aligned} \text{(a)} \quad pt_* : \pi_p S^2 \vee \pi_q S^2 \vee \pi_r S^2 &\rightarrow L(p, q, r), & \text{if } r > 1, \\ j_* : E(p, 1) \vee E(q, 1) &\rightarrow L(p, q, 1), & \text{if } q > 1. \end{aligned}$$

Moreover, the map

$$\text{(b)} \quad j_* \vee e_* : E(1, 1) \vee [S^p, S^2 \vee S^1 \vee S^1] \rightarrow L(p, 1, 1)$$

is onto for $p > 1$.

In a future paper we will study r -link maps in \mathbb{R}^3 and S^3 for $r \geq 3$. For instance, if $p_r > 1$, the sphere theorem implies a funny general “periodicity” as follows: The natural map

$$\bigvee_{1 \leq i < j \leq r} L(p_1, \dots, \hat{p}_i, \dots, \hat{p}_j, \dots, p_r, 0) \rightarrow L(p_1, \dots, p_r)$$

is onto. Here $\hat{}$ means “omit the corresponding sphere.”

I would like to thank N. Habegger and U. Koschorke for many helpful hints and suggestions. This work was made possible by support from the scientific committee of NATO via DAAD.

NOTATION. \simeq means homotopic or homotopically equivalent, \approx diffeomorphic. For each manifold M let $\text{int}(M)$ denote the interior and ∂M denote the boundary. 1 is the identity map and $[]$ is a homotopy or link homotopy class.

Proof of Theorem 1. The result is obvious for $q = 0$ and is known for $(p, q) = (1, 1)$. Assume $q > 1$, so that also $p > 1$. Recall the definition of a belt projection of a 2-component link map $g: S^p \amalg S^q \rightarrow S^3$. Just take a path $\gamma: I \rightarrow S^3$, such that $\gamma(0) \in g(S^p)$, $\gamma(1) \in g(S^q)$, $\gamma(0, 1) \cap g(S^p \amalg S^q) = \emptyset$, and define the belt projection of g to be the oriented stereographic projection from $\gamma(\frac{1}{2})$. This is well-defined up to link homotopy (compare [Ko2] or [K1]). So, if $f: S^p \amalg S^q \rightarrow \mathbb{R}^3$ maps each sphere into the unbounded component of the second sphere, then f is belt projection of a link map in S^3 , thus trivial by the proposition. So we assume that f maps S^p into a bounded component of $\mathbb{R}^3 \setminus f(S^q)$, which is a component of the complement of $f(S^q)$ in S^3 , thus aspherical [P]. Contract the map of S^p into a constant map on some point and deform the q -sphere into a surrounding 2-sphere. This proves $[f] \in pt_*(\pi_q S^2)$. It is proved in [K1] that pt_* injects.

As expected the only interesting case involves a circle S^1 . A link map $f: S^p \amalg S^1 \rightarrow \mathbb{R}^3$ is called *proper*, if f is differentiable and embeds the circle. We may replace link homotopy of link maps by link homotopy of proper link maps. Let $f: S^p \amalg S^1 \rightarrow \mathbb{R}^3$ be proper, $K := f(S^1) \subset \mathbb{R}^3$.

To prove that e_* maps onto we have to unknot K by a link homotopy. Let T be a tubular neighborhood of K , such that $T \cap f(S^p) = \emptyset$. Choose an arc σ in $X := S^3 \setminus \text{int } T$, which joins ∞ to a point on ∂T . Now deform X along this path to get a manifold $X' \subset \mathbb{R}^3 \setminus \text{int } T$ diffeomorphic to X . Let S_∞ be a small sphere around ∞ . We have the obvious embedding (see Figure 1) $e: X \vee S^2 \approx X' \vee S_\infty \rightarrow \mathbb{R}^3$ (\approx means diffeomorphic outside the basepoints), such that $\mathbb{R}^3 \setminus K \simeq e(X \vee S^2) =: Y$. Thus we may assume that f maps S^p into Y . Let $p: \tilde{X} \rightarrow X$ be the universal cover. The universal cover \tilde{Y} of Y can be described as follows (Figure 2): $p^{-1}(*) = \{*_j\}_{j \in \mathbb{Z}}$ is a countable set in \tilde{X} . To each point $*_j$ we attach a separate 2-sphere S_j . Note that \tilde{X} is contractible. Let $r_t: \tilde{X} \rightarrow \tilde{X}$, $0 \leq t \leq 1$, be




FIGURE 1




FIGURE 2

a contraction, $r_0 = 1$, $r_1(\tilde{X}) = *_1$. For each $*_j \in p^{-1}(*)$ there is the path $\sigma_j: I \ni t \rightarrow r_t(*_j) \in \tilde{X}$. Define $\tilde{r}_1: \tilde{Y} \rightarrow \bigvee_{j \in \mathbb{Z}} (S^2)_j$ as follows: $\tilde{r}_1|_{\tilde{X}} = r_1$, r_1 maps S_j onto $(S^2)_j$ by a degree 1 map.

Similarly let $i: \bigvee_{j \in \mathbb{Z}} (S^2)_j \rightarrow \tilde{Y}$ be the map which takes the upper hemispheres with degree 1 onto S_j . The restriction of i on the lower hemispheres maps the geodesic lines from the equator of S_j to the common basepoint onto the path σ_j . By homotopy extension it follows that $i \circ \tilde{r}_1 \simeq 1$. Lift f_1 to $\tilde{f}_1: S^p \rightarrow \tilde{Y}$. Since S^p is compact, $\tilde{f}_1(S^p) \cap p^{-1}(*) = \{*_j\}_{j \in J}$, $J \subset \mathbb{Z}$ finite, and $\tilde{r}_1 \circ \tilde{f}_1$ maps into $\bigvee_{j \in J} (S^2)_j$. Thus $(i \circ \tilde{r}_1) \circ \tilde{f}_1$ maps into $\bigcup_{j \in J} (S_j \cup \sigma_j(I))$. The projection of the homotopy $1 \circ \tilde{f}_1 \simeq i \circ \tilde{r}_1 \circ \tilde{f}_1$ is a homotopy of f_1 in $\mathbb{R}^3 \setminus K$ to a map into the union of S_∞ and a finite collection of loops $p(\sigma_j(t))$ based in $* \in S_\infty \cap X'$. Now we can unknot K . This proves that e_* maps onto.

To prove injectivity of e_* we have to take advantage once more of the structure of knot complements. Recall that a knot $K \subset S^3$ comes naturally equipped with a Seifert map, i.e. a differentiable map $h = h(K): X \rightarrow S^1$, which restricts to the meridional projection $\partial X \rightarrow S^1$ associated to a special framing. h is well defined up to homotopy [Z]. Recall that $h^{-1}(y)$ is a Seifert-surface of K for some regular value $y \in S^1$.

DEFINITION. A *based knot* is a pair (K, τ) , such that $K \subset \mathbb{R}^3$ is an oriented differentiable knot and τ , the *basing*, is an arc in $X = S^3 \setminus \text{int } T$ for some tubular neighborhood $T \subset \mathbb{R}^3$; τ joins $\infty \in S^3$ to some point on ∂T . \square

To each based knot we associate an unbased map $g = g(K, \tau): Y := \mathbb{R}^3 \setminus \text{int}(T) \rightarrow S^1 \vee S^2$ as follows: Use τ to construct $X' \vee S_\infty \approx X \vee S^2 \simeq Y$ as above. We can assume that $h(K)$ maps a closed tubular neighborhood N of τ onto $(-1) \in S^1$. Define $g(x) = h(x)$ for $x \in Y \setminus \text{int}(N)$. Let $B_\infty \subset S^3$ denote the ball bounding S_∞ . The cell $N' = N \setminus \text{int}(B_\infty)$ can be collapsed onto $(\partial N') \setminus (N' \cap \partial X)$. Similarly we have the retraction $B_\infty \setminus \infty \rightarrow S_\infty$. This defines $g': \text{int}(N) \setminus \infty \rightarrow \partial X' \vee S_\infty$. We compose g' and $h \vee d$, where $d: S_\infty \rightarrow S^2$ is a diffeomorphism, to get $g: \text{int}(N) \setminus \infty \rightarrow S^1 \vee S^2$. It is easy to check that the unbased homotopy class of $g(K, \tau)$ does not depend on the choice of $h(K)$. Note that we may move τ in $S^3 \setminus K$ fixing $\tau(0)$ and restricting $\tau(1)$ to ∂X without changing $[g(K, \tau)] \in [Y, S^1 \vee S^2]$. Thus in case of an unknot $K = U$ the homotopy class of $g(K, \tau) \circ f_1$ does not depend on the choice of τ . This follows from the fact that any two arcs can be deformed into each other in $S^3 \setminus K$ by a move as above and a homotopy fixing endpoints.

It is convenient to introduce the following

DEFINITION. A *based homotopy* of based knots (K_0, τ_0) and (K_1, τ_1) is a pair (F, τ) consisting of:

- (i) $F : S^1 \times I \rightarrow \mathbb{R}^3$ is a homotopy, which restricts to K_0 , resp. K_1 , on $S^1 \times 0$, resp. $S^1 \times 1$.
- (ii) $\tau : I \times I \rightarrow S^3$ is an isotopy of arcs and restricts to τ_0 , resp. τ_1 , on $I \times 0$, resp. $I \times 1$. Furthermore $\tau(0, t) = \infty$ for all $t \in I$ and $\tau(1, t)$ is a point on a meridional curve over some regular point of $F|_{S^1 \times t}$. \square

LEMMA 1. Let $\bar{F} : (S^p \amalg S^1) \times I \rightarrow \mathbb{R}^3$ be a link homotopy between proper link maps and $(\bar{F}|_{S^1}, \tau)$ be a based homotopy of knots. Then $g(K_0, \tau_0) \circ (\bar{F}|_{S^p \times 0})$ and $g(K_1, \tau_1) \circ (\bar{F}|_{S^p \times 1})$ are homotopic maps.

Proof. The crucial point is already in [M]. The homomorphisms $H_1(S^3 \setminus K_0) \rightarrow \mathbb{Z}$ and $H_1(S^3 \setminus K_1) \rightarrow \mathbb{Z}$ corresponding to Seifert-maps for K_0 and K_1 extend to a map $H_1(S^3 \times I \setminus \bar{F}(S^1 \times I))$ onto \mathbb{Z} .¹ This can be proved by elementary obstruction theory and Poincaré duality. The resulting map $S^3 \times I \setminus \bar{F}(S^1 \times I) \rightarrow S^1$ and the basing τ can be used to construct $\mathbb{R}^3 \times I \setminus \bar{F}(S^1 \times I) \rightarrow S^1 \vee S^2$. Composition with the trace of $\bar{F}|_{S^p \times I}$ yields the desired homotopy. \square

LEMMA 2. Let $f = f_1 \amalg f_2 : S^p \amalg S^1 \rightarrow \mathbb{R}^3$ be proper, $K = f(S^1)$. Then $g(K, \tau) \circ f_1 \simeq g(K, \sigma) \circ f_1$ for any two basings σ, τ .

Proof. We know already that f can be homotoped into f' , such that $f'(S^1)$ is the unknot U . A corresponding differentiable generic link homotopy can be split up into link homotopies which either restrict to isotopy on S^1 or involve a single crossing change of a knot. Since isotopies are ambient we get induced deformations of the basings σ, τ . If a crossing change is involved we may first move a given basing (at the corresponding stage of the homotopy) away from the singularity. This is possible because of transversality. Thus the link homotopy from f to f' induces based knot homotopies from (K, σ) to (U, σ') and (K, τ) to (U, τ') . By Lemma 1 we know $g(K, \tau) \circ f_1 \simeq g(U, \tau') \circ f'_1$ and $g(K, \sigma) \circ f_1 \simeq g(U, \sigma') \circ f'_1$. Now the assertion follows by a previous remark. \square

¹ This observation is due to N. Habegger.

Lemmas 1 and 2 and the fact that the arguments in the proof of Lemma 2 can be applied to arbitrary link homotopies show that the assignment

$$\begin{aligned}\lambda: E(p, 1) &\rightarrow [S^p, S^1 \vee S^2], \\ \lambda[f] &= [g(K, \tau) \circ f_1], \quad K = f(S^1)\end{aligned}$$

is well defined, i.e. independent of all involved choices (f is assumed proper!).

From the construction above follows immediately

LEMMA 3. *The composition*

$$[S^p, S^1 \vee S^2] \xrightarrow{e_*} E(p, 1) \xrightarrow{\lambda} [S^p, S^1 \vee S^2]$$

is given by the identity map. □

This proves the rest of Theorem 1. □

Proof of Theorem 2. If $r > 1$, thus $p, q, r > 1$, we consider a path σ in S^3 which meets the image of each component sphere. We assume $\sigma(0) \in f(S^p)$, $\sigma(t_0) \in f(S^q)$ and $\sigma[0, t_0] \cap f(S^r) = \emptyset$. Then, $f(S^p) \cup \sigma[0, t_0] \cup f(S^q) \subset S^3$ is a path connected subset of S^3 . By [Pa] each component of the complement of this set is aspherical, so $f|_{S^r}$ can be homotoped into a constant. Thus $[f]$ is in the image of $j_*: E(p, q) \rightarrow L(p, q, r)$. But $pt_*: \pi_p S^2 \vee \pi_q S^2 \rightarrow E(p, q)$ is 1-1 and onto by Theorem 1. If we take into consideration all possibilities, clearly we have that $pt_*: \pi_p S^2 \vee \pi_q S^2 \vee \pi_r S^2 \rightarrow L(p, q, r)$ is onto. The map, which restricts each component to a map into the complement of the images of the basepoints of the other two components, is a two-sided inverse of pt_* .

Now assume $r = 1$ and $p, q > 1$. As above, a path σ which starts in $f(S^1)$ and meets each component sphere, has empty intersection with one of the remaining spheres for $t \leq t_0$. So we may assume that $f|_{S^q}$ maps into a component of $S^3 \setminus (f(S^1) \cup \sigma[0, t_0] \cup f(S^p))$, which is aspherical by [Pa]. This proves that j_* is onto. Again, a two-sided inverse is obvious.

The proof of (b) is very similar to the proof of Theorem 1. If the link of the two circles does not split, then $[f]$ is in the image of j_* . Note that the complement of an unsplit link is aspherical by [Pa], 27. Thus, we may assume that there is a 2-sphere S embedded in S^3 , which separates two knots K_1, K_2 . Choose a basepoint $x \in S$ and arcs σ_1, σ_2 , which join points in tubular neighborhoods of the knots to




FIGURE 3

$*$ $\in S$ and meet S only in their endpoints. Let X_1 , resp. X_2 , denote $S^3 \setminus K_1$, resp. $S^3 \setminus K_2$. Clearly, $S^3 \setminus (K_1 \cup K_2) \simeq X'_1 \vee X'_2 \vee S$, $X'_i \approx X_i$ for $i = 1, 2$. The covering space argument of Theorem 1 carries over first to deform $f|S^p$ and then unknot K_1 and K_2 . Note that $X'_1 \vee X'_2$ is homotopically equivalent to the complement of $K_1 \cup \sigma'_1 \cup \sigma'_2 \cup K_2$, when σ'_1, σ'_2 are canonical extensions of σ_1, σ_2 inside the tubular neighborhoods. This shows $[f] \in \text{Im}(e_*)$ and completes the proof. \square

REFERENCES

- [H] N. Habegger, Private communication (1988).
- [K1] U. Kaiser, *Verschlingungsabbildungen im Euklidischen Raum*, doctoral thesis, Siegen (1989).
- [K2] —, *Link homotopy of Euclidean spaces*, in preparation.
- [Ko1] U. Koschorke, *Link maps and the geometry of their invariants*, *Manuscripta Math.*, **61** (1988), 383–415.
- [Ko2] —, *Desuspending the a -invariant of link maps*, preprint (1987).
- [M] J. Milnor, *Link groups*, *Ann. of Math.*, **59** (1954), 177–195.
- [P] C. Papakyriakopoulos, *On Dehn's lemma on the asphericity of knots*, *Ann. of Math.*, **66** (1957), 1–26.
- [Z] E. C. Zeeman, *Twisting spun knots*, *Trans. Amer. Math. Soc.*, **115** (1965), 471–495.

Received November 1, 1989 and in revised form July 1, 1990.

UNIVERSITÄT SIEGEN
 FB6, HOLDERLINSTR. 3
 D-5900 SIEGEN, W. GERMANY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN

(Managing Editor)

University of California

Los Angeles, CA 90024-1555-05

HERBERT CLEMENS

University of Utah

Salt Lake City, UT 84112

THOMAS ENRIGHT

University of California, San Diego

La Jolla, CA 92093

NICHOLAS ERCOLANI

University of Arizona

Tucson, AZ 85721

R. FINN

Stanford University

Stanford, CA 94305

VAUGHAN F. R. JONES

University of California

Berkeley, CA 94720

STEVEN KERCKHOFF

Stanford University

Stanford, CA 94305

C. C. MOORE

University of California

Berkeley, CA 94720

MARTIN SCHARLEMANN

University of California

Santa Barbara, CA 93106

HAROLD STARK

University of California, San Diego

La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

(1906-1982)

B. H. NEUMANN

F. WOLF

(1904-1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1991 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 151, No. 2 December, 1991

Michael G. Eastwood and A. M. Pilato , On the density of twistor elementary states	201
Brian E. Forrest , Arens regularity and discrete groups	217
Yu Li Fu , On Lipschitz stability for F.D.E	229
Douglas Austin Hensley , The largest digit in the continued fraction expansion of a rational number	237
Uwe Kaiser , Link homotopy in \mathbb{R}^3 and S^3	257
Ronald Leslie Lipsman , The Penney-Fujiwara Plancherel formula for abelian symmetric spaces and completely solvable homogeneous spaces	265
Florin G. Radulescu , Singularity of the radial subalgebra of $\mathcal{L}(F_N)$ and the Pukánszky invariant	297
Albert Jeu-Liang Sheu , The structure of twisted $SU(3)$ groups	307
Morwen Thistlethwaite , On the algebraic part of an alternating link	317
Thomas (Toma) V. Tonev , Multi-tuple hulls	335
Arno van den Essen , A note on Meisters and Olech's proof of the global asymptotic stability Jacobian conjecture	351
Hendrik J. van Maldeghem , A characterization of the finite Moufang hexagons by generalized homologies	357
Bun Wong , A note on homotopy complex surfaces with negative tangent bundles	369
Chung-Tao Yang , Any Blaschke manifold of the homotopy type of CP^n has the right volume	379