

Pacific Journal of Mathematics

**NOTE ON THE DISCRIMINANT OF CERTAIN CYCLOTOMIC
PERIOD POLYNOMIALS**

JOHN DAVID BRILLHART

NOTE ON THE DISCRIMINANT OF CERTAIN CYCLOTOMIC PERIOD POLYNOMIALS

JOHN BRILLHART

In this note we determine the discriminant of the period polynomial for certain binomial periods in the n th cyclotomic field, where $n = pq$ and p, q are distinct, odd primes.

1. Introduction. The binomial cyclotomic periods η_j for an odd prime p are defined as

$$(1) \quad \eta_j = \theta_j = \zeta^j + \zeta^{-j}, \quad 1 \leq j \leq \frac{p-1}{2},$$

where $\zeta = \zeta_p$ denotes a primitive p th root of unity, which in the standard Gaussian notation is the case where $e = (p-1)/2$ and $f = 2$. The associated period polynomial is defined to be

$$(2) \quad \Psi_p(x) = \prod_{j=1}^{\frac{p-1}{2}} (x - \theta_j) = x^e + a_1 x^{e-1} + \cdots + a_e,$$

where

$$a_k = (-1)^{\lfloor \frac{k}{2} \rfloor} \binom{e - \lfloor \frac{k+1}{2} \rfloor}{\lfloor \frac{k}{2} \rfloor}, \quad 1 \leq k \leq e.$$

(Here we use θ_j when η_j is real. The square bracket is the greatest integer function.)

The formula for the discriminant of this polynomial is

$$(3) \quad D(\Psi_p(x)) = p^{\frac{e-3}{2}}.$$

It is also useful to introduce the polynomial

$$(4) \quad \Psi_p^{(m)}(x) = \prod_{j=1}^{\frac{p-1}{2}} (x - \theta_j^m), \quad m \geq 0.$$

(See [4] for references to the above.)

In [4], the Lehmers investigated the period polynomial for another species of binomial period, the case where $n = pq$, p and q being distinct, odd primes. As in the prime case, $\zeta = \zeta_n$ denotes a primitive n th root of unity and $f = 2$. Here $e = \phi(n)/2 = (p-1)(q-1)/2$.

To define the periods in this case, first solve the congruence $x^2 \equiv 1 \pmod{n}$ for the two values of $x = a \not\equiv \pm 1 \pmod{n}$. (Sylvester [7] had investigated the case in which $a \equiv -1 \pmod{n}$.) In what follows, we will only treat the case where $a \equiv -1 \pmod{p}$ and $a \equiv 1 \pmod{q}$ as in [4]. (The other case is the same with the values of p and q interchanged.) Then the binomial periods $\eta_j(a)$ are defined by

$$(5) \quad \eta_j(a) = \zeta^j + \zeta^{aj}, \quad 1 \leq j \leq \frac{n-1}{2}, \quad (j, n) = 1,$$

and the associated period polynomial is

$$(6) \quad \Psi_n(a, x) = \prod_{\substack{j=1 \\ (j, n)=1}}^{\frac{n-1}{2}} (x - \eta_j(a)).$$

Also established is the useful formula

$$(7) \quad \Psi_n(a, x) = \Psi_p^{(q)}(x^q) / \Psi_p(x).$$

(Observe that the polynomial division on the right is exact.) It is also shown in [4] that $D(\Psi_n(a, x))$ is divisible by the product $p^{(p-3)(q-1)/2} q^{(p-1)(q-2)/2}$.

This note had its origin in the observation that when the quotient

$$D(\Psi_n(a, x)) / p^{\frac{(p-3)(q-1)}{2}} q^{\frac{(p-1)(q-2)}{2}}$$

was computed for small values of p and q , it was always found to be a power of some integer. In what follows, we will prove that this is true in general by showing that the quotient is actually $(U_q^{(p)})^{2q-4}$, where the integer $U_q^{(p)}$ is a particular function of the θ_i 's, a function mentioned by E. Lucas in [6, p. 317]. Unfortunately, Lucas never fully discussed these and related functions before his tragic and untimely death in 1891 or explained their uses in developing the properties of a certain species of elliptic-like functions he claimed he had discovered and their connection with Fermat's Last Theorem. (See Bell [1].)

The multiplicative properties of the resultant were used in this note, instead of the more usual direct arguments based on the discriminant defined in terms of the roots, because this "exterior" approach gives the results almost automatically when the discriminant is computed in terms of the resultant using (7). (The form of (12) was originally found in this way.)

2. The discriminant formula. In the following derivation we will use some of the familiar properties of the discriminant and the resultant. (See [7, pp. 279-289], [2, p. 51].)

Let $f(x) = a_0x^r + \cdots + a_r$ and $g(x) = b_0x^s + \cdots + b_s$, $r, s \geq 1$. Then it is not difficult to prove that

$$(8) \quad D(f(g(x))) = (-1)^{r\binom{s}{2}} a_0^{s-1} b_0^{r(rs-s-1)} [D(f)]^s R(f(g(x)), g'(x)),$$

from which we get the special case

$$(9) \quad D(f(x^s)) = (-1)^{r\binom{s}{2}} (a_0 a_r)^{s-1} s^{rs} [D(f)]^s, \quad s \geq 1.$$

We will also need the following

LEMMA. Let $h(x) = c_0x^k + \cdots + c_k$, $k \geq 1$, and suppose that $f(x) = g(x)h(x)$ has no multiple zeros. Then

$$(10) \quad D\left(\frac{f}{g}\right) = b_0^2 \frac{D(f)D(g)}{R^2(f', g)}.$$

Proof. Since f has no multiple zero, we have that $0 \neq D(f) = D(g)D(h)R^2(g, h)$, so

$$(11) \quad D(h) = \frac{D(f)}{D(g)R^2(g, h)}.$$

Also, $f'(x) = g(x)h'(x) + g'(x)h(x)$, so, if α_i are the (distinct) zeros of $g(x)$, then $f'(\alpha_i) = g'(\alpha_i)h(\alpha_i)$. Since $g'(\alpha_i) \neq 0$,

$$\begin{aligned} R(g, h) &= b_0^k \prod_{i=1}^n h(\alpha_i) = b_0^k \prod_{i=1}^s \frac{f'(\alpha_i)}{g'(\alpha_i)} = \frac{b_0^k \prod_{i=1}^s f'(\alpha_i)}{(-1)^{\frac{s(s-1)}{2}} b_0^{2-s} D(g)} \\ &= \frac{(-1)^{\frac{s(s-1)}{2}} b_0^{s+k-2} \prod_{i=1}^s f'(\alpha_i)}{D(g)} = \frac{(-1)^{\frac{s(s-1)}{2}} R(g, f')}{b_0 D(g)}. \end{aligned}$$

(Note $\deg f' = s + k - 1$.) Substituting this result into (11) gives (10). \square

THEOREM 1.

$$(12) \quad D(\Psi_n(a, x)) = (-1)^{\frac{(p-1)(q-1)}{4}} p^{\frac{(p-3)(q-1)}{2}} q^{\frac{(p-1)(q-2)}{2}} (U_q(p))^{2q-4},$$

where

$$U_q^{(p)} = \prod_{1 \leq i < j \leq \frac{p-1}{2}} \left(\frac{\theta_j^q - \theta_i^q}{\theta_j - \theta_i} \right)$$

is a rational integer.

Proof. Since $\Psi_n(a, x) = \Psi_p^{(q)}(x^q)/\Psi_p(x)$, then (10) gives

$$D(\Psi_n(a, x)) = \frac{D(\Psi_p(x))D(\Psi_p^{(q)}(x^q))}{[R(\Psi_p(x), (\Psi_p^{(q)}(x^q))')]^2}.$$

In the numerator, $D(\Psi_p(x)) = p^{(p-3)/2}$ by (3), and using (9) with $s = q$ and $r = (p-1)/2$,

$$D(\Psi_p^{(q)}(x^q)) = (-1)^{\frac{p-1}{2} \frac{q-1}{2}} q^{q(\frac{p-1}{2})} [D(\Psi_p^{(q)}(x))]^q.$$

In the denominator, the second argument in the resultant will have the factor qx^{q-1} because of the chain rule. Thus, we can use the multiplicativity of the resultant to obtain:

$$\begin{aligned} (13) \quad R(\Psi_p(x), (\Psi_p^{(q)}(x^q))') &= R(\Psi_p(x), qx^{q-1}) R\left(\Psi_p(x), \frac{(\Psi_p^{(q)}(x^q))'}{qx^{q-1}}\right) \\ &= R(\Psi_p(x), qx^{q-1}) R(\Psi_p^{(q)}(x), (\Psi_p^{(q)}(x))'). \end{aligned}$$

Now, using the properties in [2, p. 51],

$$R(\Psi_p(x), qx^{q-1}) = q^{\frac{p-1}{2}} [\Psi_p(0)]^{q-1} = q^{\frac{p-1}{2}}.$$

Thus, (13) becomes

$$R(\Psi_p(x), (\Psi_p^{(q)}(x^q))') = q^{\frac{p-1}{2}} D(\Psi_p^{(q)}(x)).$$

Substituting these results in (11) gives:

$$(14) \quad D(\Psi_n(a, x)) = (-1)^{\frac{(p-1)(q-1)}{4}} p^{\frac{p-3}{2}} q^{\frac{(p-1)(q-2)}{2}} [D(\Psi_p^{(q)}(x))]^{q-2}.$$

But

$$\begin{aligned} D(\Psi_p^{(q)}(x)) &= \prod_{1 \leq i < j \leq \frac{p-1}{2}} (\theta_j^q - \theta_i^q)^2 \\ &= \prod_{1 \leq i < j \leq \frac{p-1}{2}} (\theta_j - \theta_i)^2 \prod_{1 \leq i < j \leq \frac{p-1}{2}} \left(\frac{\theta_j^q - \theta_i^q}{\theta_j - \theta_i} \right)^2 \\ &= D(\Psi_p(x)) (U_q^{(p)})^2 = p^{\frac{p-3}{2}} (U_q^{(p)})^2. \end{aligned}$$

Substituting this result into (14) gives the theorem. Note that $U_q^{(p)}$ is symmetric in the θ 's, so it is a rational integer. \square

In [5] the Lehmers published the special case when $p = 5$, i.e. when $n = 5q$. They observed in this case that the polynomial $P_n(x) = \Psi_n(a, x)$ has the Fibonacci numbers F_n as their coefficients. They also established that $5^{n-1}n^{2n-4}$ divides $D(P_n(x))$. In [3] we proved that the discriminant in this case has the simple formula

$$D(P_n(x)) = 5^{n-1}n^{2n-4}F_n^{2n-4}.$$

REFERENCES

- [1] E. T. Bell, *Analogies between the U_n , V_n of Lucas and elliptic functions*, Bull. Amer. Math. Soc., **29** (1923), 401-406.
- [2] J. Brillhart, *On the Euler and Bernoulli polynomials*, J. Reine Angew. Math., **234** (1969), 45-64.
- [3] ———, *Letter to the Editor*, Fibonacci Quart., **21** (1983), 259.
- [4] D. H. and Emma Lehmer, *Cyclotomy with short periods*, Math. Comp., **41** (1983), 743-758.
- [5] ———, *Properties of polynomials having Fibonacci numbers for coefficients*, Fibonacci Quart., **21** (1983), 62-64.
- [6] E. Lucas, *Théorie des fonctions numériques simplement périodiques*, Amer. J. Math., **1** (1878), 184-240, 289-321.
- [7] J. J. Sylvester, *On certain ternary cubic equations*, Collected Papers, vol. 3, Cambridge, 1904; Amer. J. Math., **2** (1879), 357-381.
- [8] J. V. Uspensky, *Theory of Equations*, McGraw-Hill, New York, 1948.

Received May 9, 1990 and in revised form February 10, 1991.

UNIVERSITY OF ARIZONA
TUCSON, AZ 85721

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721

R. FINN
Stanford University
Stanford, CA 94305

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Pacific Journal of Mathematics

Vol. 152, No. 1

January, 1992

B. V. Rajarama Bhat , On a characterization of velocity maps in the space of observables	1
John David Brillhart , Note on the discriminant of certain cyclotomic period polynomials	15
Steven R. Costenoble and Stefan Waner , The equivariant Thom isomorphism theorem	21
Karl Heinz Dovermann and Dong Youp Suh , Smith equivalence for finite abelian groups	41
Klaus-Jochen Engel , On singular perturbations of second order Cauchy problems	79
Angel Ferrandez and Pascual Lucas , On surfaces in the 3-dimensional Lorentz-Minkowski space	93
Brian Hartley , A general Brauer-Fowler theorem and centralizers in locally finite groups	101
Joel Hass , Intersections of least area surfaces	119
Edward Kissin , Indices of unbounded derivations of C^* -algebras	125
Erhard Luft and Denis Karmen Sjerve , On regular coverings of 3-manifolds by homology 3-spheres	151
Daniel Pecker , Sur la projection de variétés algébriques réelles	165
Claus Scheiderer , Some remarks on orderings under finite field extensions	175
Fernando Serrano , Elliptic surfaces with an ample divisor of genus two	187