

Pacific Journal of Mathematics

**THE BOUNDARY OF A SIMPLY CONNECTED DOMAIN AT AN
INNER TANGENT POINT**

JOHN MARAFINO

THE BOUNDARY OF A SIMPLY CONNECTED DOMAIN AT AN INNER TANGENT POINT

JOHN MARAFINO

Let T^* be the set of accessible boundary points at which the inner tangent to ∂D exists. That is, if $a \in T^*$ and $w(a)$ represents its complex coordinate, then there exists a unique $\nu(a)$, $0 \leq \nu(a) < 2\pi$, such that for each $\varepsilon > 0$ ($\varepsilon < \frac{\pi}{2}$) there exists a $\delta > 0$ such that

$$\Delta = \left\{ w(a) + \rho e^{i\nu} : 0 < \rho < \delta, |\nu - \nu(a)| < \frac{\pi}{2} - \varepsilon \right\} \subset D$$

and $w \rightarrow a$ as $\rho \rightarrow 0$, $w \in \Delta$.

Let $\gamma(a, r)$ represent the unique component of $D \cap \{|w - w(a)| = r\}$ that intersects the inner normal $\{w(a) + \rho e^{i\nu(a)} : \rho > 0\}$, $L(a, r)$ denote the length of $\gamma(a, r)$ and set $A(a, r) = \int_0^r L(a, r') dr'$. Finally let AD^* be those points of T^* at which a non-zero angular derivative exists.

Our main result is a *purely geometrical* proof of a theorem that describes the boundary of D near $a \in T^*$. As a consequence we have

(1) a geometric description of the boundary of D near almost every $a \in AD^*$ that is a generalization of the geometric behavior of a smooth curve,

(2) an answer on T^* and hence on AD^* of the two open questions and conjectures made by McMillan in [3, p. 739] concerning the length and area ratios

$$\frac{L(a, r)}{2\pi r} \quad \text{and} \quad \frac{A(a, r)}{\pi r^2} \quad \text{as } r \rightarrow 0.$$

1. Introduction.

1.1. Many of the definitions introduced in §§1.1 to 1.3 can be found in McMillan's papers.

Let D be a simply connected plane domain, not the whole plane, and define on D the relative metric d_D , the relative distance between two points of D being defined as the infimum of the Euclidean diameters of curves that lie in D and join these two points. Let (D^*, d_{D^*}) be the completion of the metric space (D, d_D) . Now $D^* = D \cup A^*$ where D is an isometric copy of D in D^* and A^* is the set of accessible boundary points of D . Any limits involving accessible boundary points are taken in d_{D^*} .

Let T^* be the set of accessible boundary points of D at which the inner tangent to ∂D exists. That is, if $a \in T^*$ and $w(a)$ represents its complex coordinate, then there exists a unique $\nu(a)$, $0 \leq \nu(a) < 2\pi$, such that for each $\varepsilon > 0$ ($\varepsilon < \pi/2$) there exists a $\delta > 0$ such that

$$\Delta = \{w(a) + \rho e^{i\nu} : 0 < \rho < \delta, |\nu - \nu(a)| < \pi/2 - \varepsilon\} \subset D$$

and $w \rightarrow a$ as $w \rightarrow w(a)$, $w \in \Delta$.

With $\arg(w - w(a))$ defined and continuous in D , we let R^* be the set of accessible boundary points of D such that

$$\liminf_{\substack{w \rightarrow a \\ w \in D}} \arg(w - w(a)) = -\infty \quad \text{and}$$

$$\limsup_{\substack{w \rightarrow a \\ w \in D}} \arg(w - w(a)) = +\infty.$$

Points of R^* are often called twist points. Using a one-to-one conformal mapping f of the unit disk onto D , one can establish using a result of Koebe a one-to-one correspondence between A^* and a dense subset of measure 2π of the unit circle. We shall say that a set $B^* \subset A^*$ is a D -conformal null set provided that it corresponds to a set of measure zero on $\{|z| = 1\}$ under this correspondence. This definition is independent of the map f . Let $z = g(w)$ be the inverse of f that maps D one-to-one and conformally onto the unit disk. Then for each $a \in A^*$, the limit

$$\lim_{\substack{w \rightarrow a \\ w \in D}} g(w) = g(a)$$

exists. We say that $g(w)$ has a finite non-zero angular derivative at $a \in A^*$ provided there exists a finite non-zero complex number $g'(a)$ such that for each Stolz angle \check{A} at a contained in D ,

$$\lim_{\substack{w \rightarrow a \\ w \in \check{A}}} g'(w) = g'(a).$$

Let AD^* be those points in A^* at which g has a finite non-zero angular derivative. For each $a \in AD^*$, it follows that g' has a finite non-zero asymptotic value along some curve ending at a . Since f' is normal [2, p. 50] we have by [7, p. 267] that f' has a finite non-zero asymptotic value along some path ending at $g(a)$ which in turn

implies that f is conformal at $g(a)$. Consequently, $a \in T^*$ and $AD^* \subset T^*$. The first part of McMillan's twist point theorem [2, p. 44] nicely ties together all the concepts introduced in this section. It states that $A^* = T^* \cup R^* \cup N^*$, where N^* is a D -conformal null set and that $T^* \setminus AD^*$ is a D conformal null set.

1.2. Let B be a subset of the plane and ε be an arbitrary positive number. Define

$$\lambda(\varepsilon) = \inf \sum_k d_k,$$

where the infimum is taken over all countable coverings of B by disks Δ_k of diameter $d_k < \varepsilon$. Clearly, $0 \leq \lambda(\varepsilon) \leq +\infty$ and $\lambda(\varepsilon)$ increases as ε decreases so that

$$\Lambda^*(B) = \lim_{\varepsilon \rightarrow 0} \lambda(\varepsilon) \quad (0 \leq \Lambda^*(B) \leq +\infty)$$

exists. Λ^* is a metric outer measure whose σ -field of measurable sets include the Borel sets [9, p. 64]. Moreover, Λ^* is outer regular relative to the class of G_δ sets. The restriction of Λ^* to this σ -field is denoted by Λ and is called either the linear measure or the one dimensional Hausdorff measure. The second part of McMillan's twist point theorem [2, p. 44] states that a subset of T^* is D -conformal null if and only if the set of complex coordinates of its points has linear measure zero. Since $T^* \setminus AD^*$ is a D conformal null set we have by this result that

$$(1) \quad \Lambda(T^* \setminus AD^*) = 0.$$

In the remainder of this paper we will restrict our attention to T^* . A proposition $P(a)$ will be said to hold for almost every $a \in T^*$ if $\{w(a) : a \in T^* \text{ and } P(a) \text{ is false}\}$ has linear measure zero. By (1) any proposition holding for almost every $a \in T^*$ will also hold for almost every $a \in AD^*$. Consequently, Theorem 1, Corollaries 1 and 2, and Theorem 2 in §1.3 can be restated using AD^* in place of T^* .

1.3. The main result of this paper is a geometric proof of a theorem describing the behavior of ∂D in a neighborhood of almost every $a \in T^*$. For each $a \in T^*$ and $r > 0$, let $\alpha(a, r)$ be the measure of the largest angle such that the sector

$$\{w(a) + \rho e^{i\nu} : 0 < \rho < r, |\nu - \nu(a)| < \alpha(a, r)\} \subset D.$$

If no such angle exists, set $\alpha(a, r) = 0$. Note that for each $a \in T^*$ there exists an r_a such that $\alpha(a, r) > 0$ for $r < r_a$. For each $a \in T^*$ and $r < r_a$, let $\gamma(a, r)$ be the unique component of $D \cap \{|w - w(a)| = r\}$ that intersects the inner normal $\{w(a) + \rho e^{i\nu(a)} : \rho > 0\}$. We denote the length of $\gamma(a, r)$ by $L(a, r)$ and set $A(a, r) = \int_0^r L(a, r') dr'$. Measurability of the integrand is shown in [3, p. 730].

For each $r < r_a$, we parameterize $\gamma(a, r)$ by $w_r(t) = w(a) + re^{it}$, $t_0(r) < t < t_1(r)$. For $S \subset (0, r_a)$, let $E_L^*(S) = \{\zeta \in A^* : w(\zeta) = w(a) + re^{it_0(r)}, r \in S, \text{ and } w(a) + re^{it} \rightarrow \zeta \text{ as } t \rightarrow t_0(r)^+\}$. That is, $E_L^*(S)$ is the set of accessible boundary points determined by $w_r(t)$ as $t \rightarrow t_0(r)^+$, $r \in S$. Similarly, $E_R^*(S)$ is the set of accessible boundary points determined by $w_r(t)$ as $t \rightarrow t_1(r)^-$, $r \in S$. Finally, set $E^*(S) = E_L^*(S) \cup E_R^*(S)$. In what follows $h(t)$ is a positive real valued function defined on $(0, +\infty)$ with the property that $\lim_{t \rightarrow 0} h(t) = 0$. An interesting case is when $h(t) = kt$ for large positive k .

THEOREM 1. *For almost every $a_0 \in T^*$, there exists a set $S \subset (0, r_{a_0})$, closed relative to $(0, r_{a_0})$, such that*

(i) $E^*(S) \subset T^*$ and

$$\lim_{r \rightarrow 0} \frac{m(S \cap (0, r))}{r} = 1,$$

where m denotes Lebesgue measure.

(ii)

$$\lim_{\substack{a \rightarrow a_0 \\ a \in E_R^*(S)}} \arg(w(a) - w(a_0)) = \nu(a_0) + \pi/2$$

and

$$\lim_{\substack{a \rightarrow a_0 \\ a \in E_L^*(S)}} \arg(w(a) - w(a_0)) = \nu(a_0) - \pi/2.$$

(iii)

$$\lim_{\substack{a \rightarrow a_0 \\ a \in E^*(S)}} \nu(a) = \nu(a_0) \quad \text{if } \nu(a_0) \neq 0$$

and

$$\lim_{\substack{a \rightarrow a_0 \\ a \in E^*(S)}} \nu(a) = 0 \pmod{2\pi} \quad \text{if } \nu(a_0) = 0.$$

(iv)

$$\lim_{\substack{a \rightarrow a_0 \\ a \in E^*(S)}} \alpha(a, h(|w(a) - w(a_0)|)) = \pi/2.$$

If the boundary of D is a smooth Jordan curve, all boundary points of D are accessible since, by a theorem of Carathéodory, any conformal mapping from $\{|z| < 1\}$ to D extends to a homeomorphism of $\{|z| \leq 1\}$ to \bar{D} . We need make no distinction between accessible boundary points of D and the boundary of D . Recall that a curve is said to be smooth if its parameterization $z(t)$ has a continuous non-zero derivative everywhere. Thus at each point of ∂D there is an inner tangent. So T^* coincides with the boundary of D . Since the boundary of D is a smooth curve, this implies that

- (a) the curve has everywhere a continuously turning tangent (and normal)
- (b) the oscillation of $z(t)$ near $z(a)$ must diminish as $t \rightarrow a$.

It is these two properties that (iii) and (iv) of Theorem 1 are characterizing. Clearly, at each point $a \in \partial D$ with $S = (0, r_a)$, properties (i)–(iv) of Theorem 1 hold. (A consideration of the smooth boundary of the unit disk illustrates the need for $h(t)$ in (iv) to tend to zero as t tends to zero.) Hence, Theorem 1 offers a geometric description of the boundary of D near almost every $a \in T^*$ that closely resembles the geometric behavior of a smooth curve. In §1.4 we will display an example that demonstrates how the smoothness property in (iv) restricts certain boundary behavior.

From the proof of this theorem the following results will be immediate:

COROLLARY 1. *For almost every $a \in T^*$,*

$$\liminf_{r \rightarrow 0} \frac{L(a, r)}{2\pi r} = \frac{1}{2}.$$

COROLLARY 2. *For almost every $a \in T^*$ and for every $\varepsilon > 0$,*

$$\left\{ r \in (0, r_a) : \frac{L(a, r)}{2\pi r} \geq \frac{1}{2} + \varepsilon \right\}$$

has density zero at 0; that is, 0 is a point of dispersion of this set [5, p. 184].

Using Corollary 2 we will then be able to show a secondary result concerning the area of D near $a \in T^*$.

THEOREM 2. *For almost every $a \in T^*$,*

$$\lim_{r \rightarrow 0} \frac{A(a, r)}{\pi r^2} = \frac{1}{2}.$$

1.4. We make a few comments and display some examples that highlight the properties of Theorems 1 and 2 and that lead to certain conclusions.

We first compare (i) and (ii) of Theorem 1 to Ostrowski's condition: We say that *Ostrowski's condition holds at $a \in T^*$* if there exist sequences $\{a'_n\}$ and $\{a''_n\}$ of accessible boundary points of D tending to $a \in T^*$ for which the corresponding sequences of complex coordinates $\{w(a'_n)\}$, $\{w(a''_n)\}$ tending to $w(a)$ satisfy

(i') $\arg(w(a'_n) - w(a)) \rightarrow \nu(a) + \pi/2$, $\arg(w(a''_n) - w(a)) \rightarrow \nu(a) - \pi/2$,

(ii')

$$\frac{|w(a'_{n+1}) - w(a)|}{|w(a'_n) - w(a)|} \rightarrow 1, \quad \frac{|w(a''_{n+1}) - w(a)|}{|w(a''_n) - w(a)|} \rightarrow 1.$$

Let $g(w)$ be a one-to-one conformal map of D onto the open unit disk. Ostrowski's condition is necessary and sufficient for $g(w)$ to be isogonal (or conformal) at a given point [6].

McMillan observes in [4, pp. 68, 73] that at each $a \in AD^*$, $g(w)$ is isogonal and as a consequence Ostrowski's condition holds. This observation is reflected in Theorem 1, parts (i) and (ii). It states that at *almost every* $a \in AD^*$ a condition slightly stronger than Ostrowski's holds. In the following two examples we illustrate how this condition restricts certain boundary behavior.

EXAMPLE 1. Let $B^* \subset T^*$ be such that the local behavior of ∂D near $a_0 \in B^*$ is similar to that shown in Figure 1. Let $B = \{w(a_0) : a_0 \in B^*\}$ and for each $a_0 \in B^*$ define $E_R(S) = \{w(a) : a \in E_R^*(S)\}$. We want to show $\Lambda(B) = 0$. Note that Ostrowski's condition holds on B^* . In fact any sequence $\{a'_n\}$ where a'_n is on the boundary over the intervals $(1/2k + 1, 1/2k)$, $k = 1, 2, \dots$, satisfies the condition. Thus the mapping is isogonal at these points. In addition, using the results of Rodin and Warschawski [8, p. 5] there is a finite non-zero angular derivative at such points. If we use property (ii) of Theorem 1, then $E_R(S)$ is on the boundary near or over the intervals $(1/2k + 1, 1/2k)$, $k = 1, 2, \dots$, and it follows that S is contained in the intervals $(1/2k + 1, 1/2k)$, $k = 1, 2, \dots$, on the




FIGURE 1




FIGURE 2

inner normal. Such a restriction violates (i) of Theorem 1 and hence Theorem 1 fails on B^* . Thus $\Lambda(B) = 0$.

EXAMPLE 2. Let $B^* \subset T^*$ be such that the local behavior of ∂D near $a_0 \in B^*$ is similar to that shown in Figure 2. The C_n closely approximate circles. Let B and $E_R(S)$ be as defined earlier. Again Ostrowski's condition holds and so the mapping is isogonal at these

points. Once again the results of Warchawski and Tsuji [9, p. 366] show that a finite non-zero angular derivative exists at these points. Using property (iv) of Theorem 1 with $h(t) = 100t$ we see that $E_R(S)$ must be contained in the C_{2k} , $k = 1, 2, \dots$. Again the density property for S is violated. Hence $\Lambda(B) = 0$ by Theorem 1.

EXAMPLE 3. So far we have demonstrated in our examples how Theorem 1 restricts certain boundary behavior where a non-zero angular derivative exists. We now construct a nontrivial example of boundary behavior that Theorem 1 does not restrict to sets of linear measure zero. We shall be working on the segment from $(0, 0)$ to $(1, 0)$ of the x axis which we denote by $[0, 1]$. Our domain D will be the half plane $\text{Im } z < 0$ and anything we join to it from the construction. On the first step of the construction we remove the middle third of the segment $[0, 1]$ and join there a rectangle R_1 of height $l_1 = 1/3$. We set $D_1 = \text{interior} (\{\text{Im } z < 0\} \cup R_1)$. At the beginning of the m th stage of the construction we are left with 2^{m-1} segments of $[0, 1]$. From the middle of each of these we remove a segment of length $d_m = 2^{-m+1}3^{-m}$ and join there a rectangle of height $l_m = (1 + 3^m)3^{-m}2^{-m-1}$. Let $R_{m,k}$, $k = 1, 2, \dots, 2^{m-1}$, denote these rectangles and set $D_m = \text{interior} (D_{m-1} \cup (\bigcup R_{m,k}))$. We let $D = \bigcup D_m$ and we make no distinction between the accessible boundary points of D and ∂D . Figure 3 displays stages of the construction near 0. Note that $\partial D \cap [0, 1]$ is a perfect set P such that $P \subset T^*$ and the Lebesgue measure of P is $1/2$. Let $B \subset P$ be such that each $p \in B$ is a point of density of P . By the Density Theorem [5, p. 187] the Lebesgue measure of B is $1/2$ and it follows that $\Lambda(B) = 1/2$. Using the results of McMillan's twist point theorem, we know a finite non-zero angular derivative exists at each point of B , with the possible exception of a set of linear measure zero. By removing such a set and relabeling B we can assume that the angular derivative exists at each point of B . At each point $p \in B$ the intersection of the circular projections of $B \cap [p, 1]$ and $B \cap [0, p]$ onto the inner normal at p is a set S_p at which Theorem 1 holds. We have constructed an example of boundary behavior that Theorem 1 does not restrict to sets of linear measure zero. In this example we also have at almost every point $a \in T^*$,

$$\lim_{r \rightarrow 0} \frac{A(a, r)}{\pi r^2} = \frac{1}{2}.$$




FIGURE 3

We finally consider Corollary 1 and Theorem 2. In [3] McMillan proved that except for a D -conformal null set of A^* ,

$$\limsup_{r \rightarrow 0} \frac{A(a, r)}{\pi r^2} \geq \frac{1}{2} \quad \text{and} \quad \limsup_{r \rightarrow 0} \frac{L(a, r)}{2\pi r} \geq \frac{1}{2}.$$

After considering a particular example, he then conjectured [3, p. 739], [4, p. 74] that except for a D -conformal null set of A^* ,

$$\liminf_{r \rightarrow 0} \frac{A(a, r)}{\pi r^2} \leq \frac{1}{2} \quad \text{and} \quad \liminf_{r \rightarrow 0} \frac{L(a, r)}{2\pi r} \leq \frac{1}{2}.$$

In this paper we not only confirm this conjecture on T^* but also evaluate these quantities.

2. Proof of Theorem 1 and of Theorem 2.

2.1. Let

$F^* = \{a_0 \in T^* : \text{for any relatively closed set } S \subset (0, r_{a_0})$
one of the properties (i)–(iv) fails}

and $F = \{w(a_0) : a_0 \in F^*\}$. For any $B^* \subset F^*$, let $B = \{w(a) : a \in B^*\}$. We must show that $\Lambda(F) = 0$.

2.2. Suppose to the contrary that $\Lambda^*(F) > 0$. Since there exists a bounded subset B of F such that $\Lambda^*(B) > 0$, we replace F by such a subset and assume without loss of generality that F is bounded.

Let ε_0 satisfy $0 < \varepsilon_0 < \pi/10$. Associate with each $a \in F^*$ rational numbers $\alpha(a)$, $\beta(a)$, $\gamma(a)$ such that

$$\begin{aligned} \pi/2 - \varepsilon_0 < \alpha(a) < \pi/2, \quad |\nu(a) - \beta(a)| < \varepsilon_0 \quad \text{and} \\ \Delta(a) = \{w(a) + \rho e^{i\nu} : 0 < \rho < \gamma(a), |\nu - \beta(a)| < \alpha(a)\} \subset D. \end{aligned}$$

It follows that there exists a $B^* \subset F^*$ such that for all $a \in B^*$, $\alpha(a) = \alpha_0$, $\beta(a) = \beta_0$, $\gamma(a) = \gamma_0$, and $\Lambda^*(B) > 0$. By replacing F^* by B^* we can assume without loss of generality that for all $a \in F^*$, $\alpha(a) = \alpha_0$, $\beta(a) = \beta_0$, and $\gamma(a) = \gamma_0$. Again associate with each $a \in F^*$ a straight $\ell(a)$ such that

$\ell(a)$ intersects the segment $\{w(a) + \rho e^{i\beta_0} : 0 < \rho < \gamma_0\}$ at right angles and the Euclidean distance from the origin to $\ell(a)$ is a rational number and one component of $\Delta(a) \setminus \ell(a)$ is triangular.

Since $\{\ell(a)\}$ is a countable set, there exists a $B^* \subset F^*$ such that for all $a \in B^*$, $\ell(a) = \ell_0$ and $\Lambda^*(B) > 0$. We thus replace F^* by B^* and assume without loss of generality that for all $a \in F^*$, $\ell(a) = \ell_0$.

For each $a \in F^*$ we define $\Delta'(a)$ to be the triangular component of $\Delta(a) \setminus \ell_0$. The set $\bigcup_{a \in F^*} \Delta'(a)$ has at most countably many components, one of which has the form $G = \bigcup_{a \in B^*} \Delta'(a)$ where $\Lambda^*(B) > 0$. Replacing F^* by B^* we again assume without loss of generality that $G = \bigcup_{a \in F^*} \Delta'(a)$ is connected. Thus, part of the boundary of G is a closed segment lying on ℓ_0 and the rest of the boundary is contained in one of the half planes determined by $\mathbb{C} \setminus \ell_0$. Without loss of generality we may assume that ℓ_0 is the x axis, $\ell_0 \cap \partial G$, is the segment from $(0, 0)$ to $(m, 0)$, which we denote by $[0, m]$, and $\partial G \setminus [0, m]$ lies in the upper half plane. Using the construction of G and the fact that, $\frac{\pi}{2} - \alpha_0 < \varepsilon_0 < \frac{\pi}{10}$, one is able to define a function $f(x)$ on $[0, m]$ such that

$$\begin{aligned} \partial G = \Gamma \cup [0, m] \quad \text{where } \Gamma = \{(x, f(x)) : x \in [0, m]\} \quad \text{and} \\ |f(x') - f(x'')| \leq \frac{1}{3}|x' - x''| \quad \text{for all } x', x'' \in [0, m]. \end{aligned}$$

It follows that ∂G is a rectifiable Jordan curve that contains the point set $F \subset \Gamma$ and that $\Lambda^*(F) > 0$. Let $\varepsilon = \Lambda^*(F)$.

We shall make no distinction between ∂G and the set of accessible boundary points of G .

We now construct a subset B^* of T^* so that

- (i) B is a closed, Λ -measurable subset of Γ for which $\Lambda^*(B \cap F) > 0$
- (ii) $\rho_0 = \text{dist}(B, [0, m]) > 0$
- (iii) If P is the projection map from $\Gamma \rightarrow [0, m]$ then $f'|_{P(B)}$ is continuous.

We have the inequalities $\Lambda^*(P(X)) \leq \Lambda^*(X) \leq 3\Lambda^*(P(X))$ for any subset of Γ . The second inequality is true since f is Lipschitz with character $1/3$. Since f is Lipschitz, the subset of $[0, m]$ where f is not differentiable has Λ^* (and thus Λ) measure zero. Thus, by Lusin's theorem, f' is continuous on some closed subset \widehat{B} of $[0, m]$ where $\Lambda^*([0, m] \setminus \widehat{B}) < \varepsilon/6$, and we may assume that $\widehat{B} \subset (0, m)$. Let B be such that $P(B) = \widehat{B}$. Then

$$\begin{aligned} \Lambda^*(B \cap F) &\geq \Lambda^*(P(B \cap F)) \\ &= \Lambda^*(P(B) \cap P(F)) = \Lambda^*(\widehat{B} \cap P(F)) \\ &\geq \Lambda^*(P(F)) - \Lambda^*(P(F) \setminus \widehat{B}). \end{aligned}$$

Now $\Lambda^*(P(F)) \geq \varepsilon/3$ and $\Lambda^*(P(F) \setminus \widehat{B}) \leq \Lambda^*([0, m] \setminus \widehat{B}) \leq \varepsilon/6$. Thus,

$$\Lambda^*(B \cap F) \geq \frac{\varepsilon}{3} - \frac{\varepsilon}{6} = \frac{\varepsilon}{6}.$$

A little thought shows that if $w \in \partial D \cap \partial G$ then there is a unique accessible boundary point a of D , with complex coordinate w , which is accessible through G . Using this observation for each $w \in B$ defines a set B^* with all the desired properties.

2.3. We have the following information on B^* and B :

- (i) For each $a \in B^*$, define $N(w(a)) = \text{Arctan}(f'(P(w(a)))) - \frac{\pi}{2}$. Since $f'|_{P(B)}$ is continuous, N is continuous on B . Also, $N(w(a))$ satisfies the same inner tangent condition as $\nu(a)$. It follows from the uniqueness of $\nu(a)$ (recall the definition of T^*) that $N(w(a)) = \nu(a)$ and as a consequence $\nu(a)$ is continuous on B^* . Thus Property (iii) of Theorem 1 holds at each point a of B^* . We now show that Property (ii) of Theorem 1 holds at each point a of B^* . In fact, let $a_0 \in B^*$.

Since f' is continuous at $P(w(a_0))$,

$$\lim_{\substack{P(w(a)) > P(w(a_0)) \\ a \rightarrow a_0, a \in B^*}} \arg(w(a) - w(a_0)) = \arctan f'(P(w(a_0))) = \nu(a_0) + \frac{\pi}{2}.$$

Similarly,

$$\begin{aligned} & \lim_{\substack{P(w(a)) < P(w(a_0)) \\ a \rightarrow a_0, a \in B^*}} \arg(w(a) - w(a_0)) \\ & = \arctan f'(P(w(a_0))) - \pi = \nu(a_0) - \frac{\pi}{2}. \end{aligned}$$

(ii) For each $w \in B$, $w = w(a)$ with $a \in B^*$, and for each $r > 0$, define $\alpha_G(w, r)$ to be the measure of the largest angle such that

$$\{w + \rho e^{i\nu} : 0 < \rho < r, |\nu - \nu(a)| < \alpha_G(w, r)\} \subset G.$$

If no such angle exists, set $\alpha_G(w, r) = 0$. Note that $\alpha_G(w, r)$ is analogous to $\alpha(a, r)$ defined in §1.3. From the construction of G , $\alpha_G(w, r)$ is positive for all $w \in B$ and $r < \rho_0 = \text{dist}(B, [0, m])$. Fix $r < \rho_0$. After drawing a picture and using the continuity of $\nu(a)$ at a_0 , one sees that $\alpha_G(w, r)$ is uppersemicontinuous at $w_0 = w(a_0)$; that is,

$$\lim_{\substack{w \rightarrow w_0 \\ w \in B}} \alpha_G(w, r) \leq \alpha_G(w_0, r).$$

We shall show that there exists a closed subset $\tilde{B} \subset B$ such that $\Lambda^*(\tilde{B} \cap F) > 0$ and for $w, w_0 \in \tilde{B}$,

$$\lim_{w \rightarrow w_0} \alpha_G(w, h(|w - w_0|)) = \pi/2.$$

From the uppersemicontinuity of α_G , for each $r < \rho_0$, $\alpha_G(w, r)$ is measurable on B . We let $\{r_n\}$ be a sequence of numbers such that $r_n \rightarrow 0$ and $r_n < \rho_0$ for all n . We define $\alpha_n(w) = \alpha_G(w, r_n)$ on B for each n . The sequence of measurable functions $\{\alpha_n(w)\}$ defined on B converges pointwise to the function $\alpha_0(w) = \pi/2$. By Egoroff's Theorem [5, p. 108] there exists a measurable subset $B_1 \subset B$ such that $\Lambda^*(B_1 \cap F) > 0$ and such that $\alpha_n(w)$ converges uniformly to $\alpha_0(w) = \pi/2$ on B_1 . Since B_1 is measurable there exists a closed set $\tilde{B} \subset B_1$, such that $\Lambda(B_1 \setminus \tilde{B})$ is sufficiently small to ensure $\Lambda^*(\tilde{B} \cap F) > 0$. We now let $w, w_0 \in \tilde{B}$ and let ε be an arbitrary positive number. Since $\{\alpha_n(w)\}$ converges uniformly on \tilde{B} to $\pi/2$, there exists $N > 0$ such that $|\pi/2 - \alpha_N(w)| < \varepsilon$ for all $w \in \tilde{B}$. Let w be sufficiently near w_0 to ensure that $h(|w - w_0|) < r_N$. Thus, for all $w \in \tilde{B}$ sufficiently near $w_0 \in \tilde{B}$, $|\pi/2 - \alpha_G(w, h(|w - w_0|))| \leq |\pi/2 - \alpha_G(w, r_N)| < \varepsilon$.

(iii) By replacing B with \tilde{B} one can assume without loss of generality that the properties listed in (i)–(ii) above hold on B^* and B and that B is closed. In addition, since

$$\alpha_G(w, h(|w - w_0|)) \leq \alpha(a, h(|w(a) - w(a_0)|)) \leq \pi/2$$

we have that

$$\lim_{\substack{a \rightarrow a_0 \\ a, a_0 \in B^*}} \alpha(a, h(|w(a) - w(a_0)|)) = \pi/2$$

and this is property (iv) of Theorem 1.

2.4. Since almost every point of $B \cap F$ is a point of density for $B \cap F$ and $\Lambda^*(B \cap F) > 0$, there exists a point $w_0 = w(a_0) \in B \cap F$ that is a point of density of the set. We fix this w_0 and consider the circular projection of Γ onto the inner normal at w_0 . (Recall that the tangent to Γ exists at w_0 .) Using the Lipschitz condition from §2.2 it can be shown [1, pp. 33, 39] that the restriction of the circular projection to that part of Γ to the right of w_0 , that is, the part of Γ from $w(a_0)$ to $(m, f(m))$, is a one-to-one map and that w_0 is a point of density of the image of B under such a map. Similarly, the restriction of the circular projection to that part of Γ to the left of w_0 is one-to-one and w_0 is a point of density of the image of B under such a map. As was done in Example 3 we define S_{w_0} to be those points on the inner normal at w_0 , whose distance from w_0 is less than ρ_0 , that are the intersection of the circular projections of that part of $B \subset \Gamma$ to the right of w_0 and that part of $B \subset \Gamma$ to the left of w_0 . Let $S = \{\rho: w_0 + \rho e^{iv(a_0)} \in S_{w_0}\}$. It follows that S is closed relative to $(0, \rho_0)$ and 0 is a point of density of S on $(0, \rho_0)$. Letting $E_R(S) = \{w(a): a \in E_R^*(S)\}$, $E_L(S) = \{w(a): a \in E_L^*(S)\}$, and $E(S) = \{w(a): a \in E^*(S)\}$, and using the one-to-one property of the restricted circular projections, one has that $E_R(S)$ is contained in that part of B to the right of w_0 , $E_L(S)$ is contained in that part of B to the left of w_0 , and $E(S)$ is contained in B . Thus from §2.3 we have an $a_0 \in F^*$ and an S that satisfies all the properties of Theorem 1. This is a contradiction. Thus F must have linear measure zero and the theorem is proved.

2.5. A result that immediately follows from (ii) of Theorem 1 is that for almost every $a \in T^*$,

$$\lim_{\substack{r \rightarrow 0 \\ r \in S}} \frac{L(a, r)}{2\pi r} = \frac{1}{2}.$$

From this Corollaries 1 and 2 can easily be established. We are now ready to prove Theorem 2. We let ε be an arbitrary positive number. Using the defining properties of the set T^* it follows that

$$(1) \quad \liminf_{r \rightarrow 0} \frac{A(a, r)}{\pi r^2} \geq \frac{1}{2} - \varepsilon$$

for every $a \in T^*$. We set

$$H = \left\{ r \in (0, r_a) : \frac{L(a, r)}{2\pi r} > \frac{1}{2} + \varepsilon \right\}.$$

For almost every $a \in T^*$, H is a measurable set that has density zero at 0. From §1.3 we have that

$$\begin{aligned} A(a, r) &= \int_0^r L(a, r') dr' \\ &= \int_{H \cap (0, r)} L(a, r') dr' + \int_{CH \cap (0, r)} L(a, r') dr', \end{aligned}$$

where CH denotes the complement of H ,

$$\leq (2\pi r)m(H \cap (0, r)) + (1/2 + \varepsilon)(\pi r^2).$$

Hence

$$(2) \quad \limsup_{r \rightarrow 0} \frac{A(a, r)}{\pi r^2} \leq 2 \lim_{r \rightarrow 0} \frac{m(H \cap (0, r))}{r} + (1/2 + \varepsilon) = 1/2 + \varepsilon.$$

Using (1) and (2) we have that the limit exists and is $1/2$.

REFERENCES

- [1] J. Marafino, *Concerning boundary behavior under conformal mappings*, Ph.D. Dissertation, University of Wisconsin-Milwaukee, 1984, pp. 33–40.
- [2] J. E. McMillan, *Boundary behavior of a conformal mapping*, *Acta Math.*, **123** (1969), 43–67.
- [3] —, *On the boundary correspondence under conformal mapping*, *Duke Math. J.*, **37** (1970), 725–739.
- [4] —, *Boundary behavior under conformal mapping*, Proceedings of the NRL Conference on Classical Function Theory, Mathematics Research Center, Naval Research Laboratory, Washington, D.C. 1970.
- [5] M. E. Monroe, *Measure and Integration*, 2nd ed., Addison-Wesley Publishing Company, (1971), 184, 187, 108–110.
- [6] A. Ostrowski, *Zur Randverzerrung bei Konformer Abbildung*, *Prace Mat. Fizycz.*, **44** (1936), 371–471.
- [7] Ch. Pommerenke, *Univalent Functions*, Vandenhoeck and Ruprecht in Göttingen, 1975.
- [8] B. Rodin and S. E. Warschawski, *Extremal length and univalent functions*, *Mathematisch Zetschrift*, **153** (1977), 1–17.
- [9] M. Tsuji, *Potential Theory in Modern Function Theory*, Maruzan Company, Tokyo, (1959), 64, 318–322.

Received September 11, 1989.

JAMES MADISON UNIVERSITY
HARRISONBURG, VA 22807

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721

R. FINN
Stanford University
Stanford, CA 94305

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Pacific Journal of Mathematics

Vol. 154, No. 1

May, 1992

Richard Arens , Pseudo regular elements in a normed ring	1
Joan Birman and William W. Menasco , Studying links via closed braids. I: A finiteness theorem	17
Etsurō Date, Michio Jimbo, Kei Miki and Tetsuji Miwa , Braid group representations arising from the generalized chiral Potts models	37
Toshihiro Hamachi , A measure theoretical proof of the Connes-Woods theorem on AT-flows	67
Allen E. Hatcher and Ulrich Oertel , Affine lamination spaces for surfaces	87
David Joyner , Simple local trace formulas for unramified p -adic groups	103
Huaxin Lin , Injective Hilbert C^* -modules	131
John Marafino , The boundary of a simply connected domain at an inner tangent point	165
Gonzalo Riera and Rubi Rodriguez , The period matrix of Bring's curve ...	179