

Pacific Journal of Mathematics

**ON THE TENSOR PRODUCT OF THETA REPRESENTATIONS
OF GL_3**

GORDAN SAVIN

ON THE TENSOR PRODUCT OF THETA REPRESENTATIONS OF GL_3

GORDAN SAVIN

Let V be the theta representation of \widetilde{GL}_3 —the two fold central extension of GL_3 . Let W be a spherical representation of GL_3 . We show that there is a nonzero GL_3 invariant trilinear form on $V \otimes V^* \otimes W$ if and only if W is a lift from SL_2 . In this case the form is unique up to a scalar.

Introduction. Let k be a global field and A its ring of adèles. Let σ be an irreducible 3 dimensional representation of the Galois group Γ of k . Assume, for simplicity, that $\sigma(\Gamma) \subset SL_3(\mathbb{C})$. Then, according to Langlands there exists an automorphic representation $\pi \subset L^2(PGL_3(k) \backslash PGL_3(\mathbb{A}))$ such that the corresponding L -functions are equal. Consider the symmetric square of the representation σ . Then, conjecturally, the corresponding L function will have a pole only if the symmetric square representation contains a copy of trivial representation. But this means that there is a quadratic form invariant under σ and therefore $\sigma(\Gamma) \subset SO_3(\mathbb{C})$. Since $SO_3(\mathbb{C}) = {}^L SL_2$, the automorphic representation π should be a lift of an automorphic representation of SL_2 . Let π_v be a local component of π . If it is spherical, π_v is the local lift of a representation of SL_2 if $\pi_v = \text{ind}_B^{\text{PGL}_3} \chi$ where χ is a character of the diagonal subgroup of PGL_3 given by

$$\chi \left(\begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} \right) = \mu \left(\frac{a}{c} \right)$$

for some unramified character $\mu: k_v^* \rightarrow \mathbb{C}^*$.

On the other hand, Patterson and Piatetski-Shapiro [PP] have constructed the symmetric square L -function corresponding to a cuspidal automorphic representation π of PGL_3 . Moreover, they showed that the residue at $s = 1$ of this L -function is

$$\int_{PGL_3(k) \backslash PGL_3(\mathbb{A})} \varphi(g) \theta(g) \theta'(g) dg$$

where $\varphi \in \pi$ and θ, θ' are “theta functions” of Kazhdan and Patterson [KP]. They are certain automorphic forms on \widetilde{GL}_3 —the two

fold central extension of GL_3 . Let F be a local field. In [FKS] we have constructed a smooth model (θ_3, V) of the local component of “theta functions”. Let (π, W) be an irreducible representation of $\mathrm{PGL}_3(F)$. From what was explained above, it is natural to ask whether there is a GL_3 invariant trilinear form on $V \otimes V^* \otimes W$. We have the following result:

THEOREM. *Let F be a local field of the characteristic $\neq 2$. Let (π, W) be a spherical representation of PGL_3 . Then there exists a GL_3 invariant trilinear form on $V \otimes V^* \otimes W$ if and only if π is the lift of a representation of SL_2 . Moreover, the form is unique up to a scalar.*

We remark that the article of Prasad [P] was perhaps the first result indicating relationship between special values of L -functions and invariant functionals.

Acknowledgment. I would like to thank Professor S. Rallis for suggesting the problem and R. Howe for the help.

Preliminaries and notation. Let P_1 (resp. P_2) be the standard $(2, 1)$ (resp. $(1, 2)$) parabolic subgroup of GL_3 . Let $P_1 = M_1U_1$ and $P_2 = M_2U_2$ be standard Levi decompositions. We shall use the letter N to denote the unipotent group of uppertriangular matrices of GL_2 and GL_3 and the letter T to denote the group of diagonal matrices of GL_2 and GL_3 . It will be clear from the context which is meant. Finally put $N_1 = N \cap M_1$, $N_2 = N \cap M_2$ and $B = TN$.

Let $P = MU$ be a parabolic subgroup and (π, V) a smooth module. Define $V(U) = \mathrm{span}\{v - \pi(u)v \mid v \in V, u \in U\}$. Then $V_U = V/V(U)$ is the module of coinvariants.

Let X be an algebraic variety over the field F . Then $S(X)$ will denote the space of locally constant, compactly supported functions on X . Obviously, $S(X_1 \times X_2) = S(X_1) \otimes S(X_2)$. Let q be an algebraic function on X . Then one can define a representation π of N on $S(X)$ by

$$\pi \left(\begin{pmatrix} 1 & n \\ & n \end{pmatrix} \right) f(x) = \phi(nq(x))f(x)$$

where ϕ is an additive character of F . It is easy to check that $S(X)_N = S(Y)$ where Y is the subvariety of X defined by $q = 0$.

We need to recall some facts about the principal series representations of $\mathrm{GL}_3(F)$. Let St denote the Steinberg representation of GL_2 .

Let λ be a multiplicative character of F^* such that $\lambda^2 = 1$. Put $\text{St}_\lambda = \text{St} \otimes \lambda(\det)$.

Let μ be a character of F^* . It defines a character $\chi_\mu: T \rightarrow \mathbf{C}^*$ by the following formula:

$$\chi_\mu \begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} = \mu \left(\frac{a}{c} \right).$$

Let $\pi(\mu) = \text{ind}_B^{\text{GL}_3} \chi_\mu$ (normalized induction). To describe the composition series of $\pi(\mu)$ we need to introduce σ_1, σ_2 representations of GL_3 defined as follows:

$$\begin{aligned} 0 \rightarrow 1 \rightarrow \text{ind}_P^{\text{GL}_3} 1 \rightarrow \sigma_1 \rightarrow 0, \\ 0 \rightarrow 1 \rightarrow \text{ind}_P^{\text{GL}_3} 1 \rightarrow \sigma_2 \rightarrow 0. \end{aligned}$$

Here the induction is not normalized! We need the following result about the principal series representations. A reader can find details in Cartier’s article [C, §III].

PROPOSITION 1. *The representations $\pi(\mu)$ are irreducible and $\pi(\mu) \cong \pi(\mu^{-1})$ unless μ is of the two following types:*

(a) $\mu = \lambda | \cdot |^{\pm 1/2}$, $\lambda^2 = 1$. *The composition series consists of $\text{ind}_P^{\text{GL}_3} \lambda(\det)$ and $\text{ind}_P^{\text{GL}_3} \text{St}_\lambda$.*

(b) $\mu = | \cdot |^{\pm 1}$. *The composition series consists of the trivial representation, the Steinberg representation, σ_1 and σ_2 .*

The central extension and theta representation. Let $(\cdot, \cdot): F^* \times F^* \rightarrow \{\pm 1\}$ be the Hilbert symbol. Let $\widetilde{\text{GL}}_n(F)$ be the 2-fold central extension of $\text{GL}_n(F)$ and $\mathfrak{s}: \text{GL}_n \rightarrow \widetilde{\text{GL}}_n$ the section as in [FKS]. The extension can be characterized in the following way:

$$\mathfrak{s}(\text{diag}(a_i))\mathfrak{s}(\text{diag}(b_i)) = \mathfrak{s}(\text{diag}(a_i b_i)) \prod_{i < j} (a_i, b_j),$$

where $\text{diag}(a_i)$ denotes the diagonal matrix with entries a_i . Moreover the section \mathfrak{s} is an isomorphism on N and we will identify N and $\mathfrak{s}(N)$. Fix a nontrivial additive character $\phi: F \rightarrow \mathbf{C}^*$. Define a function $\gamma = \gamma_\phi: F^* \rightarrow \mathbf{C}^*$ by

$$\gamma(a) = \frac{|a|^{1/2} \int \phi(ax^2) dx}{\int \phi(x^2) dx}.$$

LEMMA 1 (Weil [W1]). *The function γ has the following properties:*

(a) $\gamma(ab) = \gamma(a)\gamma(b)(a, b)$,

(b) $\overline{\gamma}_\phi = \gamma_{\overline{\phi}}$. □

DEFINITION 1. Let $C_2(F)$ be the space of locally constant functions on F^* such that

(a) $f(x) = 0$ if $|x| > c$,

(b) $f(y^2x) = f(x)$ if $|x|, |y^2x| < 1/c$,

where c is a constant depending on f .

The theta representation θ_2 of $\widetilde{\text{GL}}_2$ can be realized on the space of functions f on F^* such that $|x|^{1/4}f(x) \in C_2(F)$. The action of $\widetilde{\text{GL}}_2$ is given by the following formulae [F]:

$$\theta_2 \left(\mathbf{s} \begin{pmatrix} a & \\ & 1 \end{pmatrix} \right) f(x) = |a|^{1/2} f(ax),$$

$$\theta_2 \left(\mathbf{s} \begin{pmatrix} z & \\ & z \end{pmatrix} \right) f(x) = (x, z)\gamma(z)f(x),$$

$$\theta_2 \left(\begin{pmatrix} 1 & n \\ & 1 \end{pmatrix} \right) f(x) = \phi(nx)f(x),$$

$$\theta_2 \left(\mathbf{s} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right) f(x) = c\gamma(x)|x|^{1/2} \int_F |y|^{1/2} f(xy^2)\phi(xy) dy$$

for some constant c .

PROPOSITION 2. *Let λ be a multiplicative character of F^* such that $\lambda^2 = 1$. Then $\theta_2 \cong \theta_2 \otimes \lambda(\det)$.*

Proof. Let θ be the even Weil representation of $\widetilde{\text{SL}}_2$. Let G be the subgroup of $\widetilde{\text{GL}}_2$ consisting of the elements whose determinant is a square in F^* . It is easy to see that θ extends to G and that $\theta_2 = \text{ind}_G^{\widetilde{\text{GL}}_2} \theta$. Since $\widetilde{\text{GL}}_2/G \cong F^*/(F^*)^2$ the proposition follows.

DEFINITION 2. Let H be a group and C its center. We say that H is a Heisenberg group if H/C is abelian.

To give a characterization of θ_3 we need a simple result about Heisenberg groups (see [KP, §0.3]):

LEMMA 2. Let H be a Heisenberg group and C its center. Let δ be a character of C . Assume that δ is faithful on $[H, H] \subset C$. Then there is unique irreducible representation π_δ of H such that C acts by multiplication by δ . Moreover, $\pi_\delta \otimes \pi_{\bar{\delta}}$ is just the regular representation of H/C . \square

Let \tilde{T} be the inverse image of T in \widetilde{GL}_3 . Let Z be the center of GL_3 and \tilde{Z} the inverse image in \widetilde{GL} . It is easy to check that \tilde{Z} is the center of \widetilde{GL}_3 . The group \tilde{T} is a Heisenberg group with center $C = \tilde{Z} \cdot s(T^2)$ where T^2 is the group of diagonal matrices whose entries are squares. Define a character δ of C by

$$\delta(s(z)s(t^2)\zeta) = \gamma(z)\zeta, \quad \zeta \in \{\pm 1\}.$$

Let π_δ be the corresponding representation of \tilde{T} . Define ρ to be, as usual,

$$\rho \begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} = \left| \frac{a}{c} \right|.$$

In [FKS] we have the following theorem.

THEOREM 1. There is a unique representation (θ_3, V) of \widetilde{GL}_3 such that $V_N \cong \rho^{1/2} \otimes \pi_\delta$. The properties of θ_3 are:

- (1) $\theta_3(s(z)) = \gamma(z) \text{Id}$, $z \in Z$.
- (2) $V_{U_1} \cong \theta_2 \otimes |\det|^{1/4}$, $V_{U_2} \cong \theta_2 \otimes |\det|^{-1/4}$.
- (3) Let $V_0 = V(U_1) \cap V(U_2)$; then $V_0 \cong S(F^* \times F)$ with the action of \tilde{B} given by

$$\theta_3 \left(\mathbf{s} \begin{pmatrix} a & b & \\ & d & \\ & & 1 \end{pmatrix} \right) f(x, y) = (x, d) |ad|^{1/2} f(ax, bx + dy),$$

$$\theta_3 \left(\begin{pmatrix} 1 & u & \\ & 1 & v \\ & & 1 \end{pmatrix} \right) f(x, y) = \phi(ux + vy) f(x, y).$$

- (4) $V(U_1) + V(U_2) = V(N)$.

In particular, it follows that we have a filtration of V as a \tilde{B} module such that the quotients are $V_0, V_{U_1}(N_1), V_{U_2}(N_2)$ and V_N .

REMARK. Note that the dual representation θ_3^* is obtained by replacing ϕ by $\bar{\phi}$.

Proof of the Theorem. Let (π, W) be a representation of GL_3 . Then the existence of a nontrivial trilinear GL_3 invariant form is equivalent to the existence of a nontrivial GL_3 intertwining map from $V \otimes V^*$ to W^* . Hence we have to compute $\dim \text{Hom}_G(V \otimes V^*, W)$. Assume that $\pi = \text{ind}_B^G \chi$. Then by the Frobenius reciprocity we have $\dim \text{Hom}_G(V \otimes V^*, W) = \dim \text{Hom}_T((V \otimes V^*)_N, \rho\chi)$. In other words we have restricted the problem to computing the T -equivariant functionals on $(V \otimes V^*)_N$. From \tilde{B} filtration of V it follows that $(V \otimes V^*)_N$ has a filtration whose quotients are $(V_0 \otimes V_0^*)_N$, $(V_{U_1}(N_1) \otimes V_{U_1}^*(N_1))_{N_1}$, $(V_{U_2}(N_2) \otimes V_{U_2}^*(N_2))_{N_2}$ and $V_N \otimes V_N^*$.

Let Γ_μ be the functional on $S(F^*)$ given by

$$\Gamma_\mu(f) = \int_{F^*} f(x)\mu(x^{-1})\frac{dx}{|x|}.$$

Obviously, we have the following simple proposition:

PROPOSITION 3. *The functional Γ_μ is unique up to a nonzero constant μ -equivariant functional on $S(F^*)$ with respect to the standard action of F^* . □*

Next we need to describe F^* equivariant functionals on $C_2(F)$.

PROPOSITION 4 (see [W2]). *Let μ be a character of F^* . The functional Γ_μ extends to $C_2(F)$ if $\mu^2 \neq 1$.*

Proof. Let \mathcal{O} be the ring of integers of F and ϖ a uniformizing element. Put $q = |\varpi|^{-1}$. Assume that $\mu^2(x) = |x|^{-s}$. Let $f \in C_2(F)$. Consider the integral

$$\Lambda_s(f) = \int_{F^*} (f(x) - f(\varpi^2x))|x|^s\frac{dx}{|x|}.$$

Obviously $\Lambda_s(f)$ is defined for every s and if $\text{Re}(s) > 0$ then

$$\Lambda_s(f) = (1 - q^s)\Gamma_\mu(f).$$

This formula extends the functional Γ_μ to $C_2(F)$ if $\mu^2(x) = |x|^{-s}$ and $s \neq 0$. If μ^2 is ramified then Γ_μ extends by taking the Principal Value integral. The proposition is proved.

PROPOSITION 5. *Let χ be a character of T . The space of χ -equivariant linear functionals on $V_N \otimes V_N^*$ is at most 1-dimensional. It has the*

dimension one if and only if $\chi = \rho\lambda$

$$\lambda \begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} = \mu_1(a)\mu_2(b)\mu_3(c),$$

$$\mu_i^2 = 1 \text{ and } \mu_1 \cdot \mu_2 \cdot \mu_3 = 1.$$

Proof. It follows from Lemma 2.

Let μ be a multiplicative character of F^* . Let $\delta_{1,2}(\mu)$, $\delta_{1,3}(\mu)$ and $\delta_{2,3}(\mu)$ be the characters of T defined by

$$\begin{aligned} \delta_{12}(\mu)(t) &= \mu(a)\mu(b)^{-1}, \\ \delta_{23}(\mu)(t) &= \mu(b)\mu(c)^{-1}, \\ \delta_{13}(\mu)(t) &= \mu(a)\mu(c)^{-1} \end{aligned}$$

where $t = \text{diag}(a, b, c)$. Let $W^{T,\chi}$ denote the space of χ -equivariant functionals on a smooth T module W .

PROPOSITION 6.

- (a) $\dim(V_{U_1}(N_1) \otimes V_{U_1}^*(N_1))_{N_1}^{T,\chi} = \begin{cases} 1 & \text{if } \chi = \rho\delta_{12}(\mu), \\ 0 & \text{otherwise.} \end{cases}$
- (b) $\dim(V_{U_2}(N_2) \otimes V_{U_2}^*(N_2))_{N_2}^{T,\chi} = \begin{cases} 1 & \text{if } \chi = \rho\delta_{23}(\mu), \\ 0 & \text{otherwise.} \end{cases}$

Proof. Using property (2) of θ_3 and the description of θ_2 it is easy to check that $V_{U_1}(N_1) \cong S(F^*)$ and therefore $(V_{U_1}(N_1) \otimes V_{U_1}^*(N_1))_{N_1} \cong S(F^*)$ with the action of T given by

$$\theta_3 \otimes \theta_3^* \left(\begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} \right) f(x) = \left| \frac{a}{c} \right| \left| \frac{a}{b} \right|^{1/2} f\left(\frac{a}{b}x\right).$$

Part (a) now follows from Proposition 3. Part (b) is proved analogously.

PROPOSITION 7.

$$(V_0 \otimes V_0^*)_N^{T,\chi} = \begin{cases} 1 & \text{if } \chi = \rho\delta_{13}(\mu), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Using property (3) of θ_3 it follows that

$$(V_0 \otimes V_0^*)_{U_1} \cong S(F^* \times F)$$

with the action of TN_1 given by

$$\theta_3 \otimes \theta_3^* \left(\begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} \right) f(x, y) = \left| \frac{ab}{c^2} \right| f\left(\frac{a}{c}x, \frac{b}{c}y\right) \quad \text{and}$$

$$\theta_3 \otimes \theta_3^* \left(\begin{pmatrix} 1 & n & \\ & 1 & \\ & & 1 \end{pmatrix} \right) f(x, y) = f(x, nx + y).$$

After taking the Fourier transform in the second variable the action becomes

$$\theta_3 \otimes \theta_3^* \left(\begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} \right) f(x, y) = \left| \frac{a}{c} \right| f\left(\frac{a}{c}x, \frac{c}{b}y\right) \quad \text{and}$$

$$\theta_3 \otimes \theta_3^* \left(\begin{pmatrix} 1 & n & \\ & 1 & \\ & & 1 \end{pmatrix} \right) f(x, y) = f(x, y)\phi(nxy).$$

Therefore $(V_0 \otimes V_0^*)_N \cong S(F^*)$ with the action of T given by

$$\theta_3 \otimes \theta_3^* \left(\begin{pmatrix} a & & \\ & b & \\ & & c \end{pmatrix} \right) f(x) = \left| \frac{a}{c} \right| f\left(\frac{a}{c}x\right).$$

The proposition follows from Proposition 3.

Let us call T equivariant functionals appearing in Proposition 5 (resp. Propositions 6 and 7) of type I (resp. II and III). Since $V_N \otimes V_N^*$ is a quotient of $(V \otimes V^*)_N$, functionals of type I extend to $(V \otimes V^*)_N$. In the next several propositions we are studying extension of the functionals of type II and III to $(V \otimes V^*)_N$.

PROPOSITION 8. *The functionals of type II extend to $(V \otimes V^*)_N$ if and only if $\mu^2 \neq 1$.*

Proof. Since V_{U_1} is a quotient of V it follows that $(V_{U_1} \otimes V_{U_1}^*)_{N_1}$ is a quotient of $(V \otimes V^*)_N$. Recall that $V_{U_1} \cong \theta_2 \otimes |\det|^{1/4}$. The value of a $\rho\delta_{12}(\mu)$ equivariant functional on $(V_{U_1}(N_1) \otimes V_{U_1}^*(N_1))_{N_1}$ is given by the following integral.

$$I_\mu(f \otimes f^*) = \int_{F^*} |x|^{1/2} f(x) f^*(x) \mu(x^{-1}) \frac{dx}{|x|}.$$

If $f \in \theta_2 \otimes |\det|^{1/4}$ and $f^* \in \theta_2^* \otimes |\det|^{1/4}$ it follows from the description of θ_2 that $|x|^{1/4} f(x)$ and $|x|^{1/4} f^*(x) \in C_2(F)$. Therefore

I_μ defines a $\rho\delta_{12}(\mu)$ equivariant functional on $(V \otimes V^*)_N$ if $\mu^2 \neq 1$ by Proposition 4. It remains to deal with $\mu, \mu^2 = 1$. Let $\varphi \in V_{U_1}$ be a function given by

$$\varphi(x) = \begin{cases} |x|^{-1/4} & \text{if } |x| \leq 1 \text{ and } x \text{ is a square,} \\ 0 & \text{otherwise.} \end{cases}$$

Let

$$v = |\varpi|^2 \varphi \otimes \varphi^* - |\varpi| \theta_2 \otimes \theta_2^* \left(\begin{pmatrix} \varpi^2 & \\ & 1 \end{pmatrix} \right) \varphi \otimes \varphi^*.$$

The projection of v on $(V_{U_1} \otimes V_{U_1}^*)_{N_1}$ lies in $(V_{U_1}(N_1) \otimes V_{U_1}^*(N_1))_{N_1} \cong S(F^*)$ and is given by

$$\begin{aligned} & |\varpi|^2 \varphi(x) \varphi^*(x) - |\varpi|^3 \varphi(\varpi^2 x) \varphi^*(\varpi^2 x) \\ &= \begin{cases} -|\varpi| & \text{if } |x| = |\varpi|^{-2} \text{ and } x \text{ is a square,} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

It follows that $I_\mu(v) < 0$. On the other hand, if the functional I_μ extends to $(V \otimes V^*)_N$ then the equivariance implies $I_\mu(v) = 0$. Contradiction. Similar conclusions can be obtained for the characters $\rho\delta_{23}(\mu)$. The proposition is proved.

Let $m_{ij}(\mu)$ be the multiplicity of $\rho\delta_{ij}(\mu)$ equivariant functionals on $(V \otimes V^*)_N$. If the principal series representation $\pi(\mu)$ is irreducible then $\pi(\mu) = \text{ind}_B^G \delta_{ij}(\mu)$ for all $1 \leq i < j \leq 3$ [C]. In particular, $m_{ij}(\mu)$ is independent of i, j . Therefore, we have obtained the following corollary.

COROLLARY 1. *The functional $\rho\delta_{13}(\mu)$ of type III extends to $(V \otimes V^*)_N$ if $\mu \neq |\cdot|^{±1}, \mu^2 \neq |\cdot|^{±1}$ and $\mu^2 \neq 1$. If $\mu^2 = 1$ and $\mu \neq 1$ then it does not extend. □*

It remains to deal with $\rho\delta_{13}(1)$.

PROPOSITION 9. *The functional $\rho\delta_{13}(1)$ of type III does not extend to $(V \otimes V^*)_N$.*

Proof. The value of a $\rho\delta_{13}(\mu)$ equivariant functional on $(V_0 \otimes V_0^*)_N$ is given by the following integral:

$$I_\mu(f \otimes f^*) = \int_{F^*} \int_F f(x, y) f^*(x, y) \mu(x^{-1}) \frac{dx}{|x|} dy.$$

Let φ be a function on $F^* \times F$ given by

$$\varphi(x, y) = \begin{cases} 1 & \text{if } |x| \leq 1 \text{ and } |y| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\varphi \in V(U_1)$ and let

$$v = |\varpi| \varphi \otimes \varphi^* - \theta_3 \otimes \theta_3^* \left(\begin{pmatrix} \varpi & & \\ & 1 & \\ & & 1 \end{pmatrix} \right) \varphi \otimes \varphi^* .$$

The projection of v on $(V(U_1) \otimes V^*(U_1))_{U_1}$ lies in $(V_0 \otimes V_0^*)_{U_1} \cong S(F^* \times F)$ and is given by

$$\begin{aligned} & |\varpi| \varphi(x, y) \varphi^*(x, y) - |\varpi| \varphi(\varpi x, y) \varphi^*(\varpi x, y) \\ &= \begin{cases} -|\varpi| & \text{if } |x| = |\varpi|^{-1} \text{ and } |y| = 1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

It follows that $I_1(v) < 0$. On the other hand, if the functional I_1 extends to $(V \otimes V^*)_N$ then the equivariance implies $I_1(v) = 0$. Contradiction. The proposition is proved.

COROLLARY 2. *Let χ be a character of T . Then $\dim(V \otimes V^*)_N^{T, \chi} \leq 1$. □*

Let μ_1, μ_2, μ_3 be three characters of F^* such that $\mu_i^2 = 1$ and $\mu_1 \cdot \mu_2 \cdot \mu_3 = 1$. Let χ be a character of T defined by $\chi(\text{diag}(a, b, c)) = \mu_1(a) \mu_2(b) \mu_3(c)$. Define $\pi(\mu_1, \mu_2, \mu_3) = \text{ind}_B^G \chi$ (normalized induction). It is unitary irreducible representation. We are now ready to formulate our main result:

THEOREM. *Let (π, W) be a quotient of a principal series representation of GL_3 . Then the space of GL_3 invariant trilinear forms on $V \otimes V^* \otimes W$ is 0 or 1 dimensional. The dimension is 0 unless π is one of the following:*

- (a) $\pi(\mu_1, \mu_2, \mu_3), \mu_i^2 = 1$ and $\mu_1 \mu_2 \mu_3 = 1$,
- (b) $\pi(\mu), \mu^2 \neq |\cdot|^{\pm 1}$ and $\mu \neq |\cdot|^{\pm 1}$,
- (c) *trivial representation,*
- (d) $\text{ind}_P^{\text{GL}_3} \mu, \mu^2 = 1$,
- (e) $\text{ind}_P^{\text{GL}_3} \text{St}_\mu, \mu^2 = 1$,
- (f) σ_1, σ_2 and St .

In cases (a)–(d) the dimension is 1. In cases (e) and (f) the dimension is ≤ 1 .

Proof. Clearly the dimension is 0 unless π is one of the representations in (a)–(f). Since representations in (a) and (b) are irreducible these two cases follow from Corollary 2. The trace $\text{tr}: V \times V^* \rightarrow \mathbf{C}$

is a GL_3 invariant trilinear form for $\pi = 1$. We can similarly deal with the representations in (d). Indeed, $V_U \otimes V_U^*$ is a quotient of $(V \otimes V^*)_U$. Since $V_U \cong \theta_2 \otimes |\det|^{1/4}$ and $\theta_2 \cong \theta_2 \otimes \mu(\det)$ by Proposition 2 we can define an appropriate P -equivariant functional on $(V \otimes V^*)_U$ defining a map from $V \otimes V^*$ into $\text{ind}_P^G \mu$. The theorem is proved.

COROLLARY. *Let (π, W) be a spherical representation of GL_3 . Then there exists a GL_3 invariant trilinear form on $V \otimes V^* \otimes W$ if and only if π is the lift of a representation of SL_2 . In this case the form is unique up to a scalar.*

Proof. Note that there is only one nontrivial unramified character μ of F^* such that $\mu^2 = 1$. Therefore if $\pi(\mu_1, \mu_2, \mu_3)$ is spherical then $\pi(\mu_1, \mu_2, \mu_3) \cong \pi(\mu, 1, \mu^{-1})$ for some unramified character μ , $\mu^2 = 1$.

A final remark. Recently Bump and Ginsburg [BG] have generalized the work of Patterson and Piatetski-Shapiro to construct an integral representation of the symmetric square L -function corresponding to a cuspidal automorphic representation π of PGL_n . As in the case $n = 3$, the residue at $s = 1$ of the L -function is

$$\int_{PGL_n(k) \backslash PGL_n(\mathbb{A})} \varphi(g)\theta(g)\theta'(g) dg$$

where $\varphi \in \pi$ and θ, θ' are “theta functions” of \widetilde{GL}_n —the two fold central extension of GL_n . The result of Bump and Ginzburg suggests the following generalization of our result:

CONJECTURE. *Let F be a local field of the characteristic $\neq 2$ and let (θ, V) be the theta representation of \widetilde{GL}_n . Let (π, W) be a spherical representation of PGL_n . Then there exists a GL_n invariant trilinear form on $V \otimes V^* \otimes W$ if and only if π is the lift of a representation of $Sp(2m)$ if $n = 2m + 1$ or π is the lift of a representation of $SO(2m)$ if $n = 2m$.*

REFERENCES

[BG] D. Bump and D. Ginzburg, *Symmetric square L-functions on $GL(r)$* , preprint.
 [C] P. Cartier, *Representations of p -adic groups*, Proceedings of Symposia in Pure Mathematics, XXXIII (1979), 111–156.

- [F] Y. Flicker, *Explicit realization of a higher metaplectic representation*, Indag. Math., Vol. I, (1990), 417–434.
- [FKS] Y. Flicker, D. Kazhdan and G. Savin, *Explicit realization of a metaplectic representation*, J. Analyse Math., **55** (1990), 17–39.
- [KP] D. Kazhdan and S. J. Patterson, *Metaplectic forms*, Publ. Math. IHES, **59** (1984), 35–142.
- [PP] S. J. Patterson and I. Piatetski-Shapiro, *The symmetric square L -function attached to a cuspid automorphic representation of GL_3* , Math. Ann., **283** (1989), 551–572.
- [P] D. Prasad, *Trilinear forms for $GL(2)$ of a local field and ε -factors*, Comp. Math., **75** (1990), 1–40.
- [W1] A. Weil, *Sur certains groupes d'opérateurs unitaires*, Acta Math., **111** (1964), 143–211.
- [W2] —, Séminaire Bourbaki, No. 312.

Received March 12, 1991. Partially supported by NSF.

YALE UNIVERSITY
NEW HAVEN, CT 06520

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

NICHOLAŞ ERCOLANI
University of Arizona
Tucson, AZ 85721

R. FINN
Stanford University
Stanford, CA 94305

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1992 by Pacific Journal of Mathematics

Pacific Journal of Mathematics

Vol. 154, No. 2

June, 1992

Manuel (Rodriguez) de León, J. A. Oubiña, P. R. Rodrigues and Modesto R. Salgado , Almost s -tangent manifolds of higher order	201
Martin Engman , New spectral characterization theorems for S^2	215
Yuval Zvi Flicker , The adjoint representation L -function for $GL(n)$	231
Enrique Alberto Gonzalez-Velasco and Lee Kenneth Jones , On the range of an unbounded partly atomic vector-valued measure	245
Takayuki Hibi , Face number inequalities for matroid complexes and Cohen-Macaulay types of Stanley-Reisner rings of distributive lattices	253
Hervé Jacquet and Stephen James Rallis , Kloosterman integrals for skew symmetric matrices	265
Shulim Kaliman , Two remarks on polynomials in two variables	285
Kirk Lancaster , Qualitative behavior of solutions of elliptic free boundary problems	297
Feng Luo , Actions of finite groups on knot complements	317
James Joseph Madden and Charles Madison Stanton , One-dimensional Nash groups	331
Christopher K. McCord , Estimating Nielsen numbers on infrasolvmanifolds	345
Gordan Savin , On the tensor product of theta representations of GL_3	369
Gerold Wagner , On means of distances on the surface of a sphere. II. (Upper bounds)	381