

*Pacific
Journal of
Mathematics*

HOMOMORPHISMS OF BUNCE-DEDDENS ALGEBRAS

CORNEL PASNICU

Volume 155 No. 1

September 1992

HOMOMORPHISMS OF BUNCE-DEDDENS ALGEBRAS

CORNEL PASNICU

The homomorphisms of a Bunce-Deddens algebra A are described. Necessary and sufficient conditions for an automorphism of the canonical UHF-subalgebra of A to have an extension to an automorphism of A are given.

The Bunce-Deddens algebras were introduced in [5]. They are interesting particular examples of inductive limits of the form $\varinjlim C(X_i, F_i)$ (where the F_i 's are finite dimensional C^* -algebras), whose study was suggested in [9]. In this paper we analyse the homomorphisms and the automorphisms of the Bunce-Deddens algebras, since their good knowledge could spread some light in the above general problem raised by E. G. Effros.

A Bunce-Deddens algebra A is a certain C^* -inductive limit $\varinjlim C(\mathbb{T}, M_{n(i)})$ (see [5]). It contains a canonical UHF-algebra B , namely the C^* -subalgebra generated by the constant functions in the algebras $C(\mathbb{T}, M_{n(i)})$. Necessary and sufficient conditions for an automorphism of B to have an extension to an automorphism of A are given (Theorem 2). A key fact proved in this paper is that B is dense in A with respect to the norm given by the unique trace of A (see Proposition 2). It is also shown that the centralizer of $\{\Phi \in \text{Aut}(A): \Phi(B) = B\}$ in $\text{Aut}(A)$ is trivial (Proposition 5) and the same thing about the centralizer of $\{\Phi \in \text{Aut}(B): (\exists) \tilde{\Phi} \in \text{Aut}(A) \text{ such that } \tilde{\Phi}|_B = \Phi\}$ in $\text{Aut}(B)$ (Proposition 4).

We also describe the endomorphisms of the Bunce-Deddens algebras, showing that they are approximately inner in a weak sense (see Theorem 1 for a more general case), but not necessarily approximately inner, since they don't always induce the identity in K_1 (see Proposition 3).

The author is grateful to M. Dădărlat for useful discussions.

Thanks are due also to the referee for his suggestions on the first version of this paper.

1. In this paper we shall consider only unital C^* -algebras.

For a compact space X and C^* -algebra A we shall identify

$C(X, A) = C(X) \otimes A$ in the canonical way and we shall consider the embedding $A \subset C(X, A)$, where, each element in A is seen as a constant function on X .

By a homomorphism of C^* -algebras we shall mean a unital $*$ -homomorphism and by an automorphism of a C^* -algebra, a $*$ -automorphism. Let $\text{Hom}(A, B)$ be the homomorphisms $A \rightarrow B$, and $\text{Aut}(A)$ the automorphisms of A . $\text{ad } u \in \text{Aut}(A)$ will denote the map $\text{ad } u(x) = uxu^*$, $x \in A$, where u is a unitary in A . By a trace of A we shall mean a central state of A .

A Bunce-Deddens algebra A will be the C^* -inductive limit of a system:

$$C(\mathbf{T}, M_{n(1)}) \xrightarrow{\Phi_1} C(\mathbf{T}, M_{n(2)}) \xrightarrow{\Phi_2} \dots$$

where $(n(i))_i$ is a strictly increasing sequence of positive integers with $n(k)$ dividing $n(k+1)$ for all $k \geq 1$ and where each homomorphism Φ_k is given by:

$$\Phi_k(a) = \begin{bmatrix} a & & & & \\ & a & & & \\ & & 0 & & \\ & & \cdot & \cdot & \\ 0 & & & \cdot & \\ & & & & a \end{bmatrix}, \quad a \in M_{n(k)},$$

$$\Phi_k \left(\begin{bmatrix} 0 & 0 & 0 & 0 & z \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & \cdot & 0 & 0 \\ & & \cdot & \cdot & \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix} \right) = \begin{bmatrix} 0 & 0 & 0 & 0 & z \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & \cdot & 0 & 0 \\ & & \cdot & \cdot & \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

(see [5]). Here $z \in C(\mathbf{T})$ is the map given by $\mathbf{T} \ni t \mapsto t \in \mathbf{C}$. We shall simply denote by $S \in A$ the unitary represented in each $C(\mathbf{T}, M_{n(i)})$ by the matrix:

$$\begin{bmatrix} 0 & 0 & 0 & 0 & z \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & \cdot & 0 & 0 \\ & & \cdot & \cdot & \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

Note that A is simple [5], has a unique trace ([4], see also [1]) and is the C^* -algebra generated by B and S (see e.g. [5]).

We shall say that (A, B) is a *canonical pair* if

$$A = \varinjlim (C(\mathbb{T}, M_{n(i)}), \Phi_i)$$

is a Bunce-Deddens algebra (as above) and $B \subset A$ is the UHF-algebra given by $B = \varinjlim (M_{n(i)}, \Phi_i|_{M_{n(i)}})$.

For a C^* -algebra A , we shall denote by $U(A)$ the unitary group of A . We denote $U(n) := U(M_n)$ (of course, by M_n we mean the $n \times n$ complex matrices). $K_1(A)$ will denote the K_1 -group ([12], [2], [8]) and if $\Phi \in \text{Hom}(A, B)$, $K_1(\Phi): K_1(A) \rightarrow K_1(B)$ denotes the natural group homomorphism.

For a space X , we shall denote by $\text{Vect}(X)$ the isomorphism classes of complex vector bundles on X . We say, that $\text{Vect}(X)$ is *torsion free* if any $E \in \text{Vect}(X)$ such that $E \oplus E \oplus \dots \oplus E$ (n -times) is a trivial vector bundle for some n , is (isomorphic to) the trivial bundle.

In this paper we shall consider only C^* -inductive limits with unital injective bonding homomorphisms.

2. We begin with a general result, which will be used in the sequel. It shows that any two homomorphisms from a UHF-algebra to a more general C^* -inductive limit are approximately inner equivalent:

PROPOSITION 1. *Consider two homomorphisms $\Phi, \Psi: A \rightarrow B = \varinjlim B_i$. Here A is a UHF-algebra and each B_i is a direct sum of C^* -algebras of the form $C(X, M_n)$, where each X is a compact connected space such that $\text{Vect}(X)$ is torsion free.*

Then, there is a sequence $(u_n)_{n \geq 1}$ in $U(B)$ such that:

$$\Phi(x) = \lim_n u_n \Psi(x) u_n^*, \quad x \in A.$$

Proof. Suppose that $A = \varinjlim A_i$, where each A_i is a full matrix algebra. For any fixed i , arguing as in [3, Lemma 2.3], we find v_i, w_i in $U(B)$ and $j = j(i)$ such that:

$$v_i \Phi(A_i) v_i^*, w_i \Psi(A_i) w_i^* \subset B_j.$$

Using ([6]; see also [7, Corollary 2.2]) for each component (in B_j) of $v_i \Phi(\cdot) v_i^*, w_i \Psi(\cdot) w_i^*: A_i \rightarrow B_j$, we obtain finally $u_i \in U(B)$ such that:

$$\Phi(x) = u_i \Psi(x) u_i^*, \quad x \in A_i.$$

Since for any $p \geq q$ and any $x \in A_q$ we have:

$$\Phi(x) = u_q \Psi(x) u_q^* = u_p \Psi(x) u_p^*$$

one easily obtains:

$$\Phi(x) = \lim_n u_n \Psi(x) u_n^*, \quad x \in A.$$

NOTATIONS. For a C^* -algebra A with a unique trace τ , we shall denote by $L^2(A)$ the separate completion of A with respect to the seminorm $A \ni a \mapsto \tau(a^*a)^{1/2} \in \mathbf{R}_+$. The induced norm on $L^2(A)$ will be denoted by $\|\cdot\|_\tau$. Note that $(L^2(A), \|\cdot\|_\tau)$ is a Hilbert space. When $(x_n)_{n \geq 1}$ is a sequence in $L^2(A)$ with $\|x_n - x\|_\tau \rightarrow 0$ for some $x \in L^2(A)$, we shall write $\tau\text{-}\lim_n x_n = x$.

The following proposition will be important in the sequel:

PROPOSITION 2. *Let (A, B) be a canonical pair (see §1). Then B is dense in $(L^2(A), \|\cdot\|_\tau)$ (where τ is the trace of A).*

Proof. Consider $A = \varinjlim (C(\mathbf{T}, M_{n(i)}), \Phi_i)$ as in §1. Since A is simple and is generated as C^* -algebra by B and S (see §1), it is enough to prove that $S \in \overline{B}^{\|\cdot\|_\tau}$.

For each $m \in \mathbf{N}$, one has:

$$S = b_m + e_{1, n(m)}^{(m)} S^{n(m)}$$

where

$$b_m := \sum_{i=1}^{n(m)-1} e_{i+1, i}^{(m)}$$

and $(e_{i,j}^{(m)})_{i,j=1}^{n(m)}$ is the canonical system of matrix units in $M_{n(m)} \subset C(\mathbf{T}, M_{n(m)})$.

We have:

$$\begin{aligned} \|S - b_m\|_\tau^2 &= \tau((S^{n(m)})^* e_{n(m), 1}^{(m)} e_{1, n(m)}^{(m)} S^{n(m)}) \\ &= \tau(e_{n(m), n(m)}^{(m)}) = \frac{1}{n(m)} \rightarrow 0 \quad \text{as } m \rightarrow \infty \end{aligned}$$

and each $b_m \in B$. Hence $S \in \overline{B}^{\|\cdot\|_\tau}$, and the proof is completed.

The following corollary was obtained in [1] (and in the particular case when the Bunce-Deddens algebra is of type 2^∞ in [5]). Our proof is simpler and shorter.

COROLLARY. *Let (A, B) be a canonical pair. Then $B' \cap A = C \cdot 1_A$.*

Proof. Let τ be the trace of A . Take an element x in $B' \cap A$. By the above proposition we deduce that it belongs to the center of A (the maps $(A, \|\cdot\|_\tau) \ni a \mapsto ax \in (A, \|\cdot\|_\tau)$ and $(A, \|\cdot\|_\tau) \ni a \mapsto xa \in (A, \|\cdot\|_\tau)$ are continuous) which is trivial since A is a simple C^* -algebra.

The following result gives a description of the homomorphisms between two Bunce-Deddens algebras. Observe first that if A and B are Bunce-Deddens algebras such that $\text{Hom}(A, B) \neq \emptyset$, then $A \subset B$ (see [5, Theorem 2 and the proof of Theorem 4]).

THEOREM 1. *Let (A, D) be a canonical pair and B a Bunce-Deddens algebra such that $A \subset B$. Let τ be the trace of B .*

If $\Phi \in \text{Hom}(A, B)$ then there is a sequence $(u_n)_{n \geq 1}$ in $U(B)$ such that:

- (a) $\Phi(x) = \tau\text{-}\lim_n u_n x u_n^*$, $x \in A$, and
- (b) $\Phi(x) = \lim_n u_n x u_n^*$, $x \in D$.

Proof. Proposition 1 gives a sequence $(u_n)_{n \geq 1}$ in $U(B)$ such that (b) holds. The fact that (a) is satisfied for the same sequence $(u_n)_{n \geq 1}$ follows using Proposition 2 and also the fact that for any $x \in A$ and $n \in \mathbb{N}$ we have:

$$\|u_n x u_n^*\|_\tau = \|\Phi(x)\|_\tau = \|x\|_\tau$$

by the unicity of the trace on a Bunce-Deddens algebra.

Having the above result, one could suspect that any endomorphism of a Bunce-Deddens algebra is approximately inner. The answer follows from:

PROPOSITION 3. *Let (A, B) be a canonical pair. Then there is a symmetry Φ of A such that $K_1(\Phi) = -\text{id}_{K_1(A)}$ (and hence Φ is not approximately inner).*

The proof follows from the following lemma (see also [2, 10.11.5]):

LEMMA 1. *Let (A, B) be a canonical pair. Then, there is a symmetry Φ of A such that $\Phi(S) = S^*$ and $\Phi(B) = B$.*

Proof. Suppose that $A = \varinjlim (C(\mathbf{T}, M_{m(i)}), \Phi_i)$ as in §1. For each n , take:

$$u_n := \begin{bmatrix} & & & & 1 \\ & 0 & & & \\ & & 1 & & \\ & & \cdot & \cdot & \\ & & \cdot & \cdot & 0 \\ 1 & & & & \end{bmatrix} \in U(m(n)) \subset U(C(\mathbf{T}, M_{m(n)})).$$

Observe that $u_n = u_n^*$ and that each diagram:

$$\begin{array}{ccc} M_{m(n)} & \xrightarrow{\Phi_n|_{M_{m(n)}}} & M_{m(n+1)} \\ \downarrow \text{ad } u_n & & \downarrow \text{ad } u_{n+1} \\ M_{m(n)} & \xrightarrow{\Phi_n|_{M_{m(n)}}} & M_{m(n+1)} \end{array}$$

commutes. Hence we obtain an automorphism Φ of B such that:

$$\Phi(x) = \lim_n u_n x u_n, \quad x \in B.$$

Let τ be the trace of A . We shall prove that:

$$\lim_n \|u_n S u_n - S^*\|_\tau = 0$$

which, by Theorem 2, will imply that Φ extends to an automorphism of A , also denoted by Φ , such that $\Phi(S) = S^*$ (don't forget that $u_n = u_n^*$).

Since for any arbitrary fixed n we have:

$$S = \begin{bmatrix} 0 & 0 & & 0 & 0 & z \\ 1 & 0 & & 0 & 0 & 0 \\ & & \ddots & & & \\ 0 & 0 & & 1 & 0 & 0 \\ 0 & 0 & & 0 & 1 & 0 \end{bmatrix} \in C(\mathbf{T}, M_{m(n)})$$

(see §1) we get:

$$u_n S u_n = \begin{bmatrix} 0 & 1 & 0 & & 0 & 0 \\ 0 & 0 & 1 & & 0 & 0 \\ & & & \ddots & & \\ 0 & 0 & 0 & & 0 & 1 \\ z & 0 & 0 & & 0 & 0 \end{bmatrix}.$$

Then one easily obtains:

$$\begin{aligned} \|u_n S u_n - S^*\|_\tau^2 &= \tau \left(\left[\begin{array}{cccc} |z - \bar{z}|^2 & & & 0 \\ & 0 & & \\ & & 0 & \\ & & & \ddots \\ 0 & & & & 0 \end{array} \right] \right) \\ &\leq 4 \cdot \tau \left(\left[\begin{array}{cccc} 1 & & & \\ & 0 & & 0 \\ & & 0 & \\ & & & \ddots \\ & & 0 & & 0 \end{array} \right] \right) \\ &= \frac{4}{m(n)} \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Since $\Phi^2|_B = \text{id}_B$ ($u_n = u_n^*$ for each n), $\Phi^2(S) = S$ and A is the C^* -algebra generated by B and S , it follows that Φ is a symmetry of A .

Proof of Proposition 3. Let Φ be the symmetry of A given by the above lemma. Suppose that $K_1(\Phi) = \text{id}_{K_1(A)}$. Then $[\Phi(S)] = [S^*] = [S]$ in $K_1(A)$ and hence $2[S] = 0$. But it is known that $K_1(A) = \mathbf{Z}$ and that $[S]$ is a generator (see [11] and [10]). It follows that $[S] = 0$, a contradiction.

Now we are interested in knowing under which conditions an automorphism of B extends to an automorphism of A ; here (A, B) will be a canonical pair. The answer to this natural problem is given by:

THEOREM 2. *Consider $\Phi \in \text{Aut}(B)$ and let $(u_n)_{n \geq 1}$ be a sequence in $U(B)$ such that $\Phi(x) = \lim_n u_n x u_n^*$, $x \in B$. Let τ be the trace of A . Then:*

$$\begin{aligned} &\Phi \text{ extends to an automorphism of } A \\ &\Leftrightarrow \tau\text{-}\lim_n u_n S u_n^*, \tau\text{-}\lim_n u_n^* S u_n \in A \end{aligned}$$

and when Φ extends, it has a unique extension $\tilde{\Phi} \in \text{Aut}(A)$, where:

$$\tilde{\Phi}(x) = \tau\text{-}\lim_n u_n x u_n^*$$

and

$$\tilde{\Phi}^{-1}(x) = \tau\text{-}\lim_n u_n^* x u_n$$

for any $x \in A$.

In the proof of this theorem we shall use the following:

LEMMA 2. Let (A, B) be a canonical pair and D a C^* -algebra with a unique trace. If $\Phi, \Psi \in \text{Hom}(A, D)$ are such that:

$$\Phi|_B = \Psi|_B$$

then:

$$\Phi = \Psi.$$

Proof. Since A and D have unique traces, denoted by τ respectively σ , one obtains:

$$\|\Phi(x)\|_\sigma = \|\Psi(x)\|_\sigma \leq \|x\|_\tau, \quad x \in A.$$

Hence, using Proposition 2 and the fact that $\Phi|_B = \Psi|_B$, it follows that $\Phi = \Psi$.

Proof of Theorem 2. Observe first that the unicity of the extension (when it exists) follows from the above lemma.

“ \Rightarrow ” Let $\tilde{\Phi} \in \text{Aut}(A)$ be such that $\tilde{\Phi}|_B = \Phi$. Then, by the proof of Theorem 1 and the above remark, it follows that:

$$\tilde{\Phi}(x) = \tau\text{-}\lim_n u_n x u_n^*, \quad x \in A.$$

Hence $\tau\text{-}\lim_n u_n S u_n^* = \tilde{\Phi}(S) \in A$.

The other relation is obtained working with Φ^{-1} .

“ \Leftarrow ” If B is seen in its GNS representation in $B(L^2(B))$ associated with the (unique) trace of B , we have:

$$\Phi(x) = U x U^*, \quad x \in B,$$

where $U \in U(B(L^2(B)))$ is given by $U(b) := \Phi(b)$, $b \in B$. Since $L^2(B) = L^2(A)$ (by Proposition 2), we have $U \in U(B(L^2(A)))$ and we can define $\tilde{\Phi} \in \text{Hom}(A, B(L^2(A)))$ by:

$$\tilde{\Phi}(x) = U x U^*, \quad x \in A.$$

Here A is seen in its GNS representation in $B(L^2(A))$ associated with the (unique) trace of A . Obviously $\tilde{\Phi}|_B = \Phi$.

By the proof of Proposition 2 there is a sequence $(b_k)_{k \geq 1}$ in B such that $\|S - b_k\|_\tau \rightarrow 0$ and $\|b_k\| = 1, k \geq 1$. Then, since $x_n \xrightarrow{\|\cdot\|_\tau} 0$ in A means $x_n \xrightarrow{\text{so}} 0$ in $B(L^2(A))$ when $\{\|x_n\|\}$ is bounded, we have:

$$\begin{aligned} \tilde{\Phi}(S) &= USU^* = \text{so-}\lim_k Ub_kU^* \\ &= \text{so-}\lim_k \Phi(b_k) = \text{so-}\lim_k (\text{so-}\lim_n u_n b_k u_n^*). \end{aligned}$$

On the other hand observe that:

$$\tau\text{-}\lim_n u_n x u_n^* \text{ exists in } L^2(A) \text{ for any } x \in A$$

(use the fact that the limit already exists for any $x \in B$, use Proposition 2 and the equality $\|u_n x u_n^*\|_\tau = \|x\|_\tau$, true for $x \in A$ and $n \in \mathbb{N}$). Therefore, we may write:

$$\begin{aligned} &\|\tau\text{-}\lim_n u_n b_k u_n^* - \tau\text{-}\lim_n u_n S u_n^*\|_\tau \\ &= \lim_n \|u_n (b_k - S) u_n^*\|_\tau = \|b_k - S\|_\tau \end{aligned}$$

and hence:

$$\tau\text{-}\lim_k (\tau\text{-}\lim_n u_n b_k u_n^*) = \tau\text{-}\lim_n u_n S u_n^* \text{ (in } L^2(A)\text{)}.$$

But, by hypothesis, $\tau\text{-}\lim_n u_n S u_n^* \in A$. Using again that $x_n \xrightarrow{\|\cdot\|_\tau} 0$ in A means $x_n \xrightarrow{\text{so}} 0$ in $B(L^2(A))$ when $\{\|x_n\|\}$ is bounded, we have:

$$\tilde{\Phi}(S) = \text{so-}\lim_k (\text{so-}\lim_n u_n b_k u_n^*) = \text{so-}\lim_n u_n S u_n^* \in A.$$

But A is the C^* -algebra generated by B and S . Hence $\tilde{\Phi}(A) \subset A$ and, as in “ \Rightarrow ”, we deduce:

$$\tilde{\Phi}(x) = \tau\text{-}\lim_n u_n x u_n^*, \quad x \in A.$$

The proof ends if we repeat the above arguments for Φ^{-1} , where $\Phi^{-1}(x) = \lim_n u_n^* x u_n, x \in B$, in this way we get $\tilde{\Phi}^{-1} \in \text{Aut}(A)$.

Question. Does any automorphism of B extend to an automorphism of A , whenever (A, B) is a canonical pair?

Our feeling is that the answer is negative.

REMARK. If we replace the above B with a certain C^* -subalgebra of A , it is easy to see that the answer to the corresponding question is negative. Let $A = \varinjlim (C(\mathbb{T}, M_{n(k)}), \Phi_k)$ be a Bunce-Deddens

algebra as in §1, where $n(k) = 2^k$, $k \geq 1$. Let D be the C^* -algebra which is the closure in A of the constant diagonal functions in $C(\mathbf{T}, M_{2^k})$, $k \geq 1$. Observe that there are canonical isomorphisms $D \cong C_{\text{red}}^*(G) \cong C(\widehat{G})$, where $G := \{z \in \mathbf{T} \mid z^{2^k} = 1 \text{ for some integer } k \geq 1\}$ and hence \widehat{G} is the group of the dyadic integers. It is not difficult to see that there are automorphisms of D which do not preserve the trace (induced from A) and, hence, cannot be extended to A .

Let again (A, B) be a canonical pair. Denote

$$H = \{\Phi \in \text{Aut}(B) : (\exists) \tilde{\Phi} \in \text{Aut}(A) \text{ such that } \tilde{\Phi}|_B = \Phi\}$$

and $G = \text{Aut}(B)$. We shall prove that the centralizer of H in G is trivial:

PROPOSITION 4. $\{\Phi \in G : \Phi \circ \Psi = \Psi \circ \Phi \text{ for any } \Psi \in H\} = \{\text{id}_B\}$.

Proof. Fix $\Phi \in G$ which commutes with every element of H . Since for any $u \in U(B)$, $\text{ad } u \in G$ belongs also to H , we have:

$$\Phi \circ \text{ad } u = \text{ad } u \circ \Phi \Leftrightarrow \Phi(u)^* u$$

commutes with

$$\Phi(B) = B \Leftrightarrow \Phi(u)^* u \in \mathbf{T} \cdot 1_B$$

(since B is simple and hence its center is trivial).

Therefore, for any $u \in U(B)$ we have:

$$\Phi(u) = \gamma(u)u$$

where $\gamma: U(B) \rightarrow \mathbf{T}$ is a continuous map.

Let τ be the (unique) trace of B . Then, we obtain:

$$\tau(u) = \tau(\Phi(u)) = \gamma(u)\tau(u), \quad u \in U(B).$$

But it is not difficult to see that $\{u \in U(B) : \tau(u) \neq 0\}$ is dense in $U(B)$. Therefore:

$$\gamma(u) = 1, \quad u \in U(B)$$

which implies that:

$$\Phi(u) = u, \quad u \in U(B)$$

and hence:

$$\Phi = \text{id}_B.$$

Also, we can prove the following:

PROPOSITION 5. *Let (A, B) be a canonical pair. Then, the centralizer of $\{\Phi \in \text{Aut}(A): \Phi(B) = B\}$ in $\text{Aut}(A)$ is trivial.*

Proof. Fix $\Phi \in \text{Aut}(A)$ which commutes with every element in $\{\Psi \in \text{Aut}(A): \Psi(B) = B\}$. For any $u \in U(B)$, $\text{ad } u \in \text{Aut}(A)$ and $\text{ad } u(B) = B$. Hence:

$$\Phi \circ \text{ad } u = \text{ad } u \circ \Phi, \quad u \in U(B).$$

Since A is simple, we deduce (as in the proof of the above proposition) that:

$$\Phi|_B = \text{id}_B.$$

By Lemma 2, it follows that:

$$\Phi = \text{id}_A.$$

REFERENCES

- [1] R. J. Archbold, *An averaging process for C^* -algebras related to weighted shifts*, Proc. London Math. Soc., (3), **35** (1977), 541–554.
- [2] B. Blackadar, *K -theory for Operator Algebras*, Springer MSRI series, 1986.
- [3] O. Bratteli, *Inductive limits of finite dimensional C^* -algebras*, Trans. Amer. Math. Soc., **171** (1972), 195–234.
- [4] J. Bunce and J. Deddens, *C^* -algebras generated by weighted shifts*, Indiana Univ. Math. J., **23** (1973), 257–271.
- [5] ———, *A family of simple C^* -algebras related to weighted shift operators*, J. Funct. Anal., **19** (1975), 13–24.
- [6] M. Dădărlat and V. Deaconu, *On some homomorphisms $\Phi: C(X) \otimes F_1 \rightarrow C(Y) \otimes F_2$* , preprint INCREST, 1986.
- [7] M. Dădărlat and C. Pasnicu, *Inductive limits of $C(X)$ -modules and continuous fields of AF -algebras*, J. Funct. Anal., **85** (1989), 103–116.
- [8] E. G. Effros, *Dimensions and C^* -algebras*, CBMS Regional Conference Series in Mathematics, No. 46, Amer. Math. Soc., Providence, R.I., 1981.
- [9] ———, *On the structure of C^* -algebras: Some old and some new problems*, in *Operator Algebras and Applications*, Proc. Sympos. Pure Math., Amer. Math. Soc., Providence, RI, 1982.
- [10] D. E. Evans, *Gauge actions on O_A* , J. Operator Theory, **7** (1982), 79–100.
- [11] P. G. Ghatage and W. J. Phillips, *C^* -algebras generated by weighted shifts II*, Indiana Univ. Math. J., **30** (1981), 539–345.
- [12] J. L. Taylor, *Banach algebras and topology in Algebras in Analysis* (ed. J. H. Williamson), Academic Press, 1975, 118–186.

Received November 20, 1990 and in revised form July 15, 1991.

INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY
P.O. BOX 1-764
RO-70700, BUCHAREST, ROMANIA

PACIFIC JOURNAL OF MATHEMATICS
EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@henri.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

PACIFIC JOURNAL OF MATHEMATICS

Volume 155 No. 1 September 1992

Characterization of modular correspondences by geometric properties	1
ALLAN RUSSELL ADLER	
Representations of convex nondentable sets	29
SPIROS ARGYROS and IRENE DELIYANNI	
Isomorphisms of spaces of continuous affine functions	71
CHO-HO CHU and HENRY BRUCE COHEN	
Universal classes of Orlicz function spaces	87
FRANCISCO LUIS HERNÁNDEZ RODRÍGUEZ and CESAR RUIZ	
Asymptotic behavior of the curvature of the Bergman metric of the thin domains	99
KANG-TAE KIM	
Quadratic central polynomials with derivation and involution	111
CHARLES PHILIP LANSKI	
Nonsplit ring spectra and products of β -elements in the stable homotopy of Moore spaces	129
JIN KUN LIN	
Orientation and string structures on loop space	143
DENNIS MCLAUGHLIN	
Homomorphisms of Bunce-Deddens algebras	157
CORNEL PASNICU	
Certain C^* -algebras with real rank zero and their corona and multiplier algebras. Part I	169
SHUANG ZHANG	
Correction to: "On the density of twistor elementary states"	199
MICHAEL G. EASTWOOD and A. M. PILATO	



0030-8730(1992)155:1;1-M