

*Pacific
Journal of
Mathematics*

COBCAT AND SINGULAR BORDISM

A. K. DAS AND S. S. KHARE

Volume 155 No. 2

October 1992

COBCAT AND SINGULAR BORDISM

A. K. DAS AND S. S. KHARE

Dold proved that a homomorphism $\phi: H^n(\text{BO}) \rightarrow \mathbb{Z}_2$ corresponds to a manifold M^n if and only if $\phi(\text{Sq}^p u + v_p \cdot u) = 0$, $\forall p \geq 0$ and $\forall u \in H^{n-p}(\text{BO})$, v_p being the Wu class. The object of the present work is to have a singular analogue of this result and to study the bordism classification of singular manifolds in BO .

1. Introduction. Singh [1] has developed the notion of cobcat for a manifold M^n and has classified, upto bordism, all manifolds M^n with $\text{cobcat}(M^n) \leq 3$. $\text{Cobcat}(M^n)$ was defined to be the smallest positive integer k such that the number $\langle W_{i_1} \cdots W_{i_p}, [M^n] \rangle = 0$ for all partitions $i_1 + \cdots + i_p = n$ with $k \leq p \leq n$.

Here we develop the notion of cobcat for a singular manifold (M^n, f) in a space X and discuss the bordism classification of all singular manifolds (M^n, f) in BO with $\text{cobcat}(M^n, f) \leq 3$, $n = 2^r$.

Here all the manifolds are to be unoriented, smooth and closed, and all the homology and cohomology coefficients are to be in \mathbb{Z}_2 . The space X is such that for each n , $H_n(X)$ and hence $H^n(X)$ is a finite dimensional vector space over \mathbb{Z}_2 .

We are thankful to Dr. H. K. Mukharjee of NEHU (India) for giving helpful suggestions. The first author is also grateful to NBHM for its financial support during the course of this work.

2. Preliminaries. Consider the set $N_n(X)$ of bordism classes of n -dimensional singular manifolds (M^n, f) in X , $f: M^n \rightarrow X$ being a continuous map. We know that $N_n(X)$ is an abelian group under the operation "disjoint union"

$$[M_1^n, f_1] + [M_2^n, f_2] = [M_1^n \sqcup M_2^n, f_1 \sqcup f_2],$$

where $f_1 \sqcup f_2: M_1^n \sqcup M_2^n \rightarrow X$ is given by

$$f_1 \sqcup f_2(x) = \begin{cases} f_1(x) & \text{if } x \in M_1^n, \\ f_2(x) & \text{if } x \in M_2^n. \end{cases}$$

Further, we have

$$N_*(X) = \bigoplus_{n \geq 0} N_n(X).$$

We know that for a point, $N_*(pt) = N_*$, the unoriented bordism ring, and there is a N_* -module structure in $N_*(X)$ given by

$$[M^n, f] \times [N^m] = [M^n \times N^m, f\pi],$$

where $\pi: M^n \times N^m \rightarrow M^n$ is the projection.

For a singular manifold (M^n, f) in X let $\tau: M^n \rightarrow \text{BO}$ be the classifying map of the tangent bundle over M^n . Then there is defined a homomorphism $t: H^n(\text{BO} \times X) \rightarrow \mathbb{Z}_2$ given by

$$t(w \otimes x) = \langle (\tau, f)^*(w \otimes x), [M^n] \rangle = \langle \tau^*(w)f^*(x), [M^n] \rangle,$$

where $w \otimes x \in H^n(\text{BO} \times X) = \bigoplus_{i=0}^n H^{n-i}(\text{BO}) \otimes H^i(X)$ and $(\tau, f): M^n \rightarrow \text{BO} \times X$ is given by $(\tau, f)(z) = (\tau(z), f(z))$.

The number $\langle W_{i_1} \cdots W_{i_r} f^*(x_{n-p}), [M^n] \rangle$ is called the Stiefel-Whitney number of (M^n, f) associated to the cohomology class $x_{n-p} \in H^{n-p}(X)$ corresponding to the partition $i_1 + \cdots + i_r = p$. Moreover, this number is as usual bordism invariant [2].

Analogous to [1], given a singular manifold (M^n, f) in X there is associated a Poincaré algebra P^* given as follows:

Let $J = \{z \in H^*(\text{BO} \times X): \text{either } \dim z > n, \text{ or for all } z' \in H^{n-\dim z}(\text{BO} \times X), \langle (\tau, f)^*z(\tau, f)^*z', [M^n] \rangle = 0\}$.

It is easy to see that J is an ideal of the graded algebra $H^*(\text{BO} \times X)$. Set

$$P^* = \frac{H^*(\text{BO} \times X)}{J}, \text{ the quotient algebra.}$$

Let $q: H^*(\text{BO} \times X) \rightarrow P^*$ be the quotient map. Clearly, $P^* = 0$ if and only if (M^n, f) bounds. Let $z \in H^*(\text{BO} \times X)$; we say, “ $z = 0$ in P^* ” if $q(z) = 0$.

As in [1], we have the following proposition, whose verification is a routine matter.

PROPOSITION 2.1. *If (M^n, f) is not a boundary then*

- (a) P^* is an n -dimensional graded algebra with Poincaré duality,
- (b) the Steenrod algebra acts on P^* with the action given by

$$\text{Sq}^i(q(z)) = q(\text{Sq}^i(z)),$$

- (c) if $z \in H^n(\text{BO} \times X)$ then $q(z) = 0$ if and only if

$$t(z) = \langle (\tau, f)^*z, [M^n] \rangle = 0. \quad \square$$

It is easy to see that for all $p \geq 0$, and for all $z \in H^{n-p}(\text{BO} \times X)$,

$$t(\text{Sq}^p(z) + (v_p \otimes 1)z) = 0,$$

where $v_p \in H^p(\text{BO})$ is the Wu class. So, in view of the above proposition, we have $\text{Sq}^p(z) = (v_p \otimes 1)z$ in P^* .

3. Extension of Dold's and Milnor's results to singular case. Dold [3] has proved the following

Result 3.1. For each integer $n \geq 0$, if $\phi: H^n(\mathbf{BO}) \rightarrow \mathbb{Z}_2$ is a homomorphism then there is an n -dimensional closed manifold M^n with $\phi(w) = \langle \tau^*(w), [M^n] \rangle$ for all $w \in H^n(\mathbf{BO})$ if and only if $\phi(\text{Sq}^p(u) + v_p \cdot u) = 0$ for all $u \in H^{n-p}(\mathbf{BO})$ and for all $p \geq 0$, $v_p \in H^p(\mathbf{BO})$ being the Wu class. \square

Here we shall extend this result to the singular case as follows:

THEOREM 3.2. For each $n \geq 0$, if $h: H^n(\mathbf{BO} \times X) \rightarrow \mathbb{Z}_2$ is a homomorphism then there is an n -dimensional singular manifold (M^n, f) with $h(w \otimes x) = \langle \tau^*(w)f^*(x), [M^n] \rangle$ for all $w \otimes x \in H^n(\mathbf{BO} \times X)$ if and only if $h(\text{Sq}^p(u \otimes y) + (v_p \cdot u) \otimes y) = 0$ for all $u \otimes y \in H^{n-p}(\mathbf{BO} \times X)$ and for all $p \geq 0$, $v_p \in H^p(\mathbf{BO})$ being the Wu class.

Proof. It is easy to see that the condition is necessary. We prove that the condition is sufficient also. Let $\{c_{m,i}\}_{i \in I_m}$ denote a basis for the vector space $H_m(X)$ over \mathbb{Z}_2 , $m \geq 0$. Let $c^{m,i} \in H^m(X)$ be the cohomology class dual to $c_{m,i}$ i.e. $\langle c^{m,i}, c_{m,j} \rangle = \delta_{ij}$. Note that $\{c^{m,i}\}$ forms a basis for $H^m(X)$. Now, for each $c_{m,j}$ we can choose a singular manifold (M_j^m, f_j^m) with $(f_j^m)_*([M_j^m]) = c_{m,j}$, [2]. Thus, we have

$$\langle (f_j^m)^* c^{m,i}, [M_j^m] \rangle = \delta_{ij}.$$

Now,

$$H^n(\mathbf{BO} \times X) = \bigoplus_{l=0}^n H^l(\mathbf{BO}) \otimes H^{n-l}(X).$$

Define,

$$h_i^0: H^0(\mathbf{BO}) \rightarrow \mathbb{Z}_2$$

by

$$h_i^0(1) = h(1 \otimes c^{n,i}).$$

Clearly, h_i^0 satisfies the condition given in Result (3.1) and so there exists a manifold V_i^0 such that

$$h_i^0(1) = \langle \tau^*(1), [V_i^0] \rangle = \langle 1, [V_i^0] \rangle,$$

for each $i \in I_n$.

Define $h_i^1: H^1(\text{BO}) \rightarrow \mathbb{Z}_2$ by

$$\begin{aligned} h_i^1(w) &= h(w \otimes c^{n-1, i}) \\ &\quad + \sum_{j \in I_n} \langle \tau^*(w)(f_j^n \pi)^*(c^{n-1, i}), [M_j^n \times V_j^0] \rangle \\ &= h(w \otimes c^{n-1, i}) \\ &\quad + \sum_j \langle \tau^*w \cdot ((f_j^n)^* c^{n-1, i} \otimes 1), [M_j^n \times V_j^0] \rangle. \end{aligned}$$

Now,

$$\begin{aligned} h_i^1(\text{Sq}^1(1) + v_1 \cdot 1) &= h(v_1 \otimes c^{n-1, i}) + \sum_j \langle \tau^*v_1 \cdot ((f_j^n)^* c^{n-1, i} \otimes 1), [M_j^n \times V_j^0] \rangle \\ &= h(\text{Sq}^1(1 \otimes c^{n-1, i})) + \sum_j \langle \text{Sq}^1((f_j^n)^* c^{n-1, i} \otimes 1), [M_j^n \times V_j^0] \rangle \\ &= h(1 \otimes \text{Sq}^1 c^{n-1, i}) + \sum_j \langle (f_j^n)^* \text{Sq}^1 c^{n-1, i}, [M_j^n] \rangle \langle 1, [V_j^0] \rangle. \end{aligned}$$

Since, $\text{Sq}^1 c^{n-1, i} \in H^n(X)$, there is a subset $K_n \subset I_n$ such that

$$\text{Sq}^1 c^{n-1, i} = \sum_{k \in K_n} c^{n, k}.$$

Therefore,

$$\begin{aligned} h_i^1(\text{Sq}^1(1) + v_1 \cdot 1) &= \sum_{k \in K_n} h(1 \otimes c^{n, k}) + \sum_j \sum_{k \in K_n} \langle (f_j^n)^* c^{n, k}, [M_j^n] \rangle h_j^0(1) \\ &= \sum_k h_k^0(1) + \sum_k h_k^0(1), \quad \text{since } \langle (f_j^n)^* c^{n, k}, [M_j^n] \rangle = \delta_{kj} \\ &= 0. \end{aligned}$$

So, by Result (3.1), there exists a manifold V_i^1 such that

$$h_i^1(w) = \langle \tau^*w, [V_i^1] \rangle,$$

for each $i \in I_{n-1}$.

Now, using induction, we define $h_i^l: H^l(\text{BO}) \rightarrow \mathbb{Z}_2$ by

$$\begin{aligned} h_i^l(w) &= h(w \otimes c^{n-l, i}) \\ &\quad + \sum_{\substack{j \in I_{n-m} \\ 0 \leq m < l}} \langle \tau^*w \cdot ((f_j^{n-m})^* c^{n-l, i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle \end{aligned}$$

where $1 \leq l \leq n$, and each V_j^m is given by h_j^m ($m < l$).

Now, it is enough to show that h_i^l satisfies the condition given in (3.1). For if it is so, then there exists a family $\{V_i^l\}_{i \in I_{n-l}}$ of manifolds such that for each $i \in I_{n-l}$,

$$h_i^l(w) = \langle \tau^* w, [V_i^l] \rangle \quad (1 \leq l \leq n).$$

Also, we already have a family $\{V_i^0\}_{i \in I_n}$ of manifolds such that

$$h_i^0(1) = \langle 1, [V_i^0] \rangle,$$

for each $i \in I_n$. It is then easy to see that the given homomorphism $h: H^n(\mathbf{BO} \times X) \rightarrow \mathbb{Z}_2$ corresponds to the singular manifold (M^n, f) given by

$$M^n = \bigsqcup M_i^{n-l} \times V_i^l \quad \text{and} \quad f = \bigsqcup (f_i^{n-l} \pi),$$

where the disjoint union \bigsqcup is taken over all $i \in I_{n-l}$ and all $0 \leq l \leq n$, and $\pi: M_i^{n-l} \times V_i^l \rightarrow M_i^{n-l}$ denotes the projection map.

Note that, for each $p \geq 0$ and for each $u \in H^{l-p}(\mathbf{BO})$,

$$h_i^l(\text{Sq}^p u + v_p \cdot u) = \text{I} + \text{II} + \text{III},$$

where

$$\begin{aligned} \text{I} &= h((\text{Sq}^p u + v_p \cdot u) \otimes c^{n-l, i}) \\ &= h(\text{Sq}^p u \otimes c^{n-l, i} + (v_p \cdot u) \otimes c^{n-l, i}) \\ &= h \left(\sum_{\substack{r+s=p \\ r \neq p}} \text{Sq}^r u \otimes \text{Sq}^s c^{n-l, i} \right), \quad \text{by hypothesis on } h, \\ &= \sum_{\substack{r+s=p \\ r \neq p}} h(\text{Sq}^r u \otimes \text{Sq}^s c^{n-l, i}), \\ \text{II} &= \sum_{\substack{j \in I_{n-m} \\ 0 \leq m < l}} \langle \tau^*(\text{Sq}^p u)((f_j^{n-m})^* c^{n-l, i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle \\ &= \sum_{j, m} \langle \text{Sq}^p(\tau^* u)((f_j^{n-m})^* c^{n-l, i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle, \quad \text{and} \\ \text{III} &= \sum_{j, m} \langle \tau^*(v_p \cdot u)((f_j^{n-m})^* c^{n-l, i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle \\ &= \sum_{j, m} \langle \text{Sq}^p(\tau^* u((f_j^{n-m})^* c^{n-l, i} \otimes 1)), [M_j^{n-m} \times V_j^m] \rangle. \end{aligned}$$

So,

$$\begin{aligned} \text{II} + \text{III} &= \sum_{j,m} \sum_{\substack{r+s=p \\ r \neq p}} \langle \text{Sq}^r(\tau^*u)(\text{Sq}^s(f_j^{n-m})^*c^{n-l,i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle \\ &= \sum_{j,m} \sum_{\substack{r+s=p \\ r \neq p}} \langle \text{Sq}^r(\tau^*u)((f_j^{n-m})^* \text{Sq}^s c^{n-l,i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle. \end{aligned}$$

Now, since $\text{Sq}^s c^{n-l,i} \in H^{n-l+s}(X)$, there is a subset K_{n-l+s} of I_{n-l+s} such that

$$\text{Sq}^s c^{n-l,i} = \sum_{k \in K_{n-l+s}} c^{n-l+s,k}, \quad (1 \leq s \leq p).$$

So,

$$\begin{aligned} (1) \quad & h(\text{Sq}^r u \otimes \text{Sq}^s c^{n-l,i}) \\ &= \sum_{k \in K_{n-l+s}} h(\text{Sq}^r u \otimes c^{n-l+s,k}) \\ &= \sum_k \left\{ \sum_{\substack{j,m \\ (m < l-s)}} \langle \tau^*(\text{Sq}^r u)((f_j^{n-m})^*c^{n-l+s,k} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle \right. \\ & \qquad \qquad \qquad \left. + \langle \tau^*(\text{Sq}^r u), [V_k^{l-s}] \rangle \right\}, \\ & \qquad \qquad \qquad \text{noting that } l-s < l \\ &= \sum_{\substack{j,m \\ (m < l-s)}} \langle \text{Sq}^r(\tau^*u)((f_j^{n-m})^* \text{Sq}^s c^{n-l,i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle \\ & \quad + \sum_k \langle \text{Sq}^r(\tau^*u), [V_k^{l-s}] \rangle. \end{aligned}$$

Also,

$$\begin{aligned} (*) \quad & \sum_{\substack{j,m \\ (l-s \leq m < l)}} \langle \text{Sq}^r(\tau^*u)((f_j^{n-m})^* \text{Sq}^s c^{n-l,i} \otimes 1), [M_j^{n-m} \times V_j^m] \rangle \\ &= \sum_j \langle \text{Sq}^r(\tau^*u)((f_j^{n-l+s})^* \text{Sq}^s c^{n-l,i} \otimes 1), [M_j^{n-l+s} \times V_j^{l-s}] \rangle, \end{aligned}$$

by dimensional consideration, since $m \geq l-s$. Further, we note that u is a polynomial in Stiefel-Whitney classes of BO, so that $\text{Sq}^r(\tau^*u)$ is

a polynomial in Stiefel-Whitney classes of $M_j^{n-l+s} \times V_j^{l-s}$. Therefore the above expression (*) becomes equal to

$$\begin{aligned} (2) \quad & \sum_j \langle (f_j^{n-l+s})^* \text{Sq}^s c^{n-l,i}, [M_j^{n-l+s}] \langle \text{Sq}^r(\tau^* u), [V_j^{l-s}] \rangle \rangle \\ &= \sum_j \sum_{k \in K_{n-l+s}} \langle (f_j^{n-l+s})^* c^{n-l+s,k}, [M_j^{n-l+s}] \langle \text{Sq}^r(\tau^* u), [V_j^{l-s}] \rangle \rangle \\ &= \sum_k \langle \text{Sq}^r(\tau^* u), [V_k^{l-s}] \rangle. \end{aligned}$$

Hence, combining I with (1) and II + III with (2), it follows that

$$h_i^l(\text{Sq}^p u + v_p \cdot u) = \text{I} + \text{II} + \text{III} = 0.$$

That is, h_i^l satisfies the condition given in (3.1). □

Now, consider the universal bundle $\gamma: \text{EO} \rightarrow \text{BO}$ and the cartesian product $\gamma \times \gamma$ over $\text{BO} \times \text{BO}$. Let $\mu: \text{BO} \times \text{BO} \rightarrow \text{BO}$ be the classifying map of $\gamma \times \gamma$. μ has the property that

$$\mu^*(W_i) = \sum_{k=0}^i W_k \otimes W_{i-k}.$$

The product of two singular manifolds (M^m, f) and (M^n, g) in BO is given by $(M^m \times M^n, \mu \circ (f \times g))$, and this product induces a multiplication in $N_*(\text{BO})$ given by

$$[M^m, f] \times [M^n, g] = [M^m \times M^n, \mu \circ (f \times g)],$$

which makes $N_*(\text{BO})$ an algebra over \mathbb{Z}_2 .

Analogous to [4], we have

LEMMA 3.3. *The Stiefel-Whitney numbers*

$$\langle W_{2i_1} \cdots W_{2i_r} (\mu \circ (g \times g))^* (W_{2i_{r+1}} \cdots W_{2i_{r+s}}), [N \times N] \rangle$$

of the product $(N, g) \times (N, g)$ in BO are equal to

$$\langle W_{i_1} \cdots W_{i_r} g^* (W_{i_{r+1}} \cdots W_{i_{r+s}}), [N] \rangle,$$

while the numbers

$$\langle W_{j_1} \cdots W_{j_p} (\mu \circ (g \times g))^* (W_{j_{p+1}} \cdots W_{j_{p+q}}), [N \times N] \rangle$$

are zero if some j_h is odd.

Proof. Routine verification. □

THEOREM 3.4. *Let (M^{2n}, f) be a singular manifold in BO, such that*

$$\langle W_{j_1} \cdots W_{j_p} f^*(W_{j_{p+1}} \cdots W_{j_{p+q}}), [M^{2n}] \rangle = 0,$$

whenever some j_h is odd. Then

$$[M^{2n}, f] = [(N^n, g) \times (N^n, g)] \text{ in } N_{2n}(\text{BO}).$$

Proof. We shall construct a singular manifold (N^n, g) in BO whose Stiefel-Whitney numbers

$$\langle W_{i_1} \cdots W_{i_r} g^*(W_{i_{r+1}} \cdots W_{i_{r+s}}), [N^n] \rangle$$

are equal to

$$\langle W_{2i_1} \cdots W_{2i_r} f^*(W_{2i_{r+1}} \cdots W_{2i_{r+s}}), [M^{2n}] \rangle.$$

This will imply that (M^{2n}, f) is cobordant to $(N^n, g) \times (N^n, g)$, by (3.3).

Let $R^n \subset H^n(\text{BO} \times \text{BO})$ be the vector space generated by all elements of the form $\text{Sq}^p(x \otimes y) + (v_p \cdot x) \otimes y$. The Stiefel-Whitney numbers of each manifold (N^n, g) determine a homomorphism

$$h_N: H^n(\text{BO} \times \text{BO}) \rightarrow \mathbb{Z}_2$$

given by $h_N(x \otimes y) = \langle (\tau, g)^*(x \otimes y), [N] \rangle$, and by Theorem (3.2) we know that a given homomorphism $H^n(\text{BO} \times \text{BO}) \rightarrow \mathbb{Z}_2$ corresponds to a singular manifold in BO if and only if it annihilates R^n .

Define the “doubling homomorphism”

$$d: H^*(\text{BO} \times \text{BO}) \rightarrow H^*(\text{BO} \times \text{BO})$$

by

$$d(W_i \otimes W_j) = W_{2i} \otimes W_{2j}.$$

Let (M^{2n}, f) satisfy the hypothesis of Theorem (3.4). Then we shall show that $h_M \circ d: H^n(\text{BO} \times \text{BO}) \rightarrow \mathbb{Z}_2$ annihilates R^n . This will prove the existence of the required manifold (N^n, g) .

Let $I \subset H^*(\text{BO} \times \text{BO})$ denote the ideal generated by the family $\{W_i \otimes 1, 1 \otimes W_i\}_{i \text{ odd}}$. Note that

$$\begin{aligned} \text{Sq}^{2i} d(W_j \otimes 1) &= (\text{Sq}^{2i} W_{2j}) \otimes 1 \\ &= \left(\sum_{k=0}^{2i} \binom{2j-2i+k-1}{k} W_{2i-k} W_{2j+k} \right) \otimes 1, \end{aligned}$$

where $\binom{p}{q}$ denotes the binomial coefficients reduced modulo 2. Therefore using the fact that

$$\binom{2j - 2i + 2l - 1}{2l} = \binom{j - i + l - 1}{l}$$

we get

$$\begin{aligned} \text{Sq}^{2i} d(W_j \otimes 1) &\equiv \left(\sum_{l=0}^i \binom{j - i + l - 1}{l} W_{2i-2l} W_{2j+2l} \right) \otimes 1 \pmod{I} \\ &\equiv d(\text{Sq}^i(W_j \otimes 1)) \pmod{I}. \end{aligned}$$

Similarly, $\text{Sq}^{2i} d(1 \otimes W_j) \equiv d(\text{Sq}^i(1 \otimes W_j)) \pmod{I}$. Further, if

$$\begin{aligned} \text{Sq}^{2i} d(x \otimes y) &\equiv d(\text{Sq}^i(x \otimes y)) \pmod{I} \quad \text{and} \\ \text{Sq}^{2i} d(x \otimes y) &\equiv d(\text{Sq}^i(x' \otimes y')) \pmod{I}, \end{aligned}$$

then

$$\begin{aligned} \text{Sq}^{2i} d((x \otimes y)(x' \otimes y')) &\equiv \sum_{p+q=i} (\text{Sq}^{2p} d(x \otimes y))(\text{Sq}^{2q} d(x' \otimes y')) \pmod{I} \\ &\equiv \sum_{p+q=i} (d(\text{Sq}^p(x \otimes y)))(d(\text{Sq}^q(x' \otimes y'))) \pmod{I} \\ &\equiv d(\text{Sq}^i((x \otimes y)(x' \otimes y'))) \pmod{I}. \end{aligned}$$

Hence, by induction, it follows that

$$\text{Sq}^{2i} d(x \otimes y) \equiv d(\text{Sq}^i(x \otimes y)) \pmod{I},$$

for each $x \otimes y \in H^*(\text{BO} \times \text{BO})$.

It is simple to verify that I is closed under Steenrod squaring operation. Applying induction on p , one gets

$$d(v_p \otimes 1) \equiv (v_{2p} \otimes 1) \pmod{I}.$$

Now, consider the manifold (M^{2n}, f) . By the hypothesis on (M^{2n}, f) we have

$$h_M(I^{2n}) = 0, \quad \text{where } I^{2n} = I \cap H^{2n}(\text{BO} \times \text{BO}).$$

Therefore, for any generator $\text{Sq}^p(x \otimes y) + (v_p \otimes 1)(x \otimes y)$ of R^n we have, using the congruences established above,

$$\begin{aligned} (h_M \circ d)(\text{Sq}^p(x \otimes y) + (v_p \otimes 1)(x \otimes y)) &= h_M(\text{Sq}^{2p}(d(x \otimes y)) + (v_{2p} \otimes 1)d(x \otimes y) + (\text{terms in } I^{2n})) \\ &= 0. \end{aligned}$$

That is, $h_M \circ d$ annihilates R^n and so by Theorem (3.2) there exists a singular manifold (N^n, g) in BO such that $h_M \circ d = h_N$. Hence, the theorem follows. \square

4. Cobcat and singular bordism in BO. Analogous to [1], we define the cobcat for a singular manifold (M^n, f) in X as follows

DEFINITION. Cobcat(M^n, f) is the smallest positive integer k such that for each $m, 0 \leq m \leq n$, the number

$$\langle W_{i_1} \cdots W_{i_p} f^*(x_{j_1} \cdots x_{j_q}), [M^n] \rangle = 0$$

for all partitions $i_1 + \cdots + i_p$ of m and for all partitions $j_1 + \cdots + j_q$ of $n - m$, with $k \leq p + q \leq n$ ($x_{j_h} \in H^{j_h}(X)$ for all j_h). If no such k exists define cobcat(M^n, f) = $n + 1$.

- REMARK 4.1.** (a) Cobcat(M^n) \leq cobcat(M^n, f),
 (b) cobcat(M^n, f) = 1 if and only if (M^n, f) bounds,
 (c) cobcat(M^n, f) \leq nil(Im(τ, f)*) \leq cat(τ, f) \leq cat(M^n),
 (d)

$$\text{cobcat}(M_1^n \sqcup M_2^n, f_1 \sqcup f_2) \leq \max\{\text{cobcat}(M_1^n, f_1), \text{cobcat}(M_2^n, f_2)\}.$$

Now we shall discuss the singular version of some results proved in [1]. Let P^* be the Poincaré algebra associated to the singular manifold (M^n, f) in X . As in [1], an element z of any graded algebra A^* will be called k -decomposable if it is zero or is the sum of the products $z_1 \cdot z_2 \cdot \cdots \cdot z_p$ where $z_i \in A^*$ with $\dim z_i > 0$ for each i , and $p \geq k$.

PROPOSITION 4.2. Let cobcat(M^n, f) $\leq k$.

- (a) If $z \in P^*$ is k -decomposable, then z is zero.
 (b) If $z \in P^*$ is $(k - 1)$ -decomposable and $\dim z < n$ then z is zero.

Proof. Note that any k -decomposable element z of $H^*(\text{BO} \times X) \cong H^*(\text{BO}) \otimes H^*(X)$ can be written as a sum of the products $z_1 \cdot z_2 \cdot \cdots \cdot z_p$, where each z_i is of the type $W_j \otimes 1$ or $1 \otimes x_j$, and $p \geq k$. Also for any $z' \in H^{n - \dim z}(\text{BO} \times X)$ we have

$$\langle (\tau, f)^* z_1 (\tau, f)^* z_2 \cdots (\tau, f)^* z_p (\tau, f)^* z', [M^n] \rangle = 0,$$

since cobcat(M^n, f) $\leq k$ and $p \geq k$. Hence (a) follows, using the fact that any k -decomposable element of P^* can be obtained from a k -decomposable element of $H^*(\text{BO} \times X)$.

As in ([1], 1.2), (b) follows from (a). \square

From now on, the ambient space X will be taken to be the universal base space BO and (M^n, f) will denote a singular manifold in BO with $\text{cobcat}(M^n, f) \leq 3$.

LEMMA 4.3. (a) *If $z(W_i \otimes 1) = 0$ and $z(1 \otimes W_i) = 0$ in P^* , where $z \in H^{n-i}(\text{BO} \times \text{BO})$ and $0 < i < n$, then $z = 0$ in P^* .*

(b) *For $j > 0$,*

$$W_{2j+1} \otimes 1 = \begin{cases} \text{Sq}^1(W_{2j} \otimes 1) & \text{if } 2j + 1 < n, \\ 0 & \text{if } 2j + 1 = n, \end{cases}$$

and

$$1 \otimes W_{2j+1} = \text{Sq}^1(1 \otimes W_{2j}) \quad \text{if } 2j + 1 < n$$

in P^* .

Proof. (a) By the hypothesis, the last proposition and the fact that $\text{cobcat}(M^n, f) \leq 3$, we have

$$\langle (\tau, f)^* z(\tau, f)^* z', [M^n] \rangle = \langle (\tau, f)^*(z \cdot z'), [M^n] \rangle = 0$$

for all $z' \in H^i(\text{BO} \times \text{BO})$. So $z = 0$ in P^* .

(b) Note that

$$\text{Sq}^1(W_{2j} \otimes 1) = (W_1 \otimes 1)(W_{2j} \otimes 1) + W_{2j+1} \otimes 1.$$

If $2j + 1 < n$ then $(W_1 \otimes 1)(W_{2j} \otimes 1)$, being decomposable, is zero in P^* . If $2j + 1 = n$, then $(W_1 \otimes 1)(W_{2j} \otimes 1) = \text{Sq}^1(W_{2j} \otimes 1)$ in P^* . For the last part of (b) one has

$$\text{Sq}^1(1 \otimes W_{2j}) = (1 \otimes W_1)(1 \otimes W_{2j}) + 1 \otimes W_{2j+1}. \quad \square$$

LEMMA 4.4. *If n is even and $n > 2$, then $W_i \otimes 1 = 0$ and $1 \otimes W_i = 0$ in P^* for all odd i .*

Proof. By Lemma (4.3), it is enough to show that

(a) $(W_i \otimes 1)(W_{n-i} \otimes 1) = 0$,

(b) $(W_i \otimes 1)(1 \otimes W_{n-i}) = 0$,

(c) $(1 \otimes W_i)(1 \otimes W_{n-i}) = 0$ in P^* for all odd i .

For (a), let $i = 1$; then

$$(W_1 \otimes 1)(W_{n-1} \otimes 1) = \text{Sq}^1(W_{n-1} \otimes 1) = \text{Sq}^1 \text{Sq}^1(W_{n-2} \otimes 1) = 0$$

in P^* , using (4.3) and the fact that $\text{Sq}^1 \text{Sq}^1 = 0$. Now let $i = 2j + 1$,

$j > 0$ and $n - i = 2k + 1, k > 0$; then by (4.3)

$$\begin{aligned} (W_i \otimes 1)(W_{n-i} \otimes 1) &= \text{Sq}^1(W_{2j} \otimes 1) \text{Sq}^1(W_{2k} \otimes 1) \\ &= \text{Sq}^1((W_{2j} \otimes 1) \text{Sq}^1(W_{2k} \otimes 1)) \\ &= (W_1 \otimes 1)(W_{2j} \otimes 1) \text{Sq}^1(W_{2k} \otimes 1) = 0 \end{aligned}$$

in P^* , as it is 3-decomposable. Hence (a) follows. For (b), using the same technique as in (a), we have $(W_i \otimes 1)(1 \otimes W_{n-i}) = 0$ in P^* for all odd $i < n - 1$. However, for $i = n - 1$, we have by (4.3)

$$\begin{aligned} (W_{n-1} \otimes 1)(1 \otimes W_1) &= \text{Sq}^1(W_{n-2} \otimes 1)(1 \otimes W_1) \\ &= \text{Sq}^1((W_{n-2} \otimes 1)(1 \otimes W_1)) + (W_{n-2} \otimes 1) \text{Sq}^1(1 \otimes W_1) \\ &= (W_1 \otimes 1)(W_{n-2} \otimes 1)(1 \otimes W_1) \\ &\quad + (W_{n-2} \otimes 1)(1 \otimes W_1)(1 \otimes W_1) = 0 \end{aligned}$$

in P^* , as it is 3-decomposable. Thus (b) follows. Now, (c) can be proved by the same technique used in (a) and (b) above. \square

PROPOSITION 4.5. *If (M^n, f) is a non-bounding n -dimensional singular manifold in BO with $\text{cobcat}(M^n, f) \leq 3$, where n is even and $n > 2$, then (M^n, f) is cobordant to a product $(N, g) \times (N, g)$ in BO, where (N, g) is also non-bounding and $\text{cobcat}(N, g) \leq 3$.*

Proof. By Theorem (3.4) and Lemma (4.4), there exists a singular manifold (N, g) is BO of dimension $n/2$ such that (M^n, f) is cobordant to the product $(N, g) \times (N, g)$. Also,

$$\begin{aligned} \langle W_{i_1} \cdots W_{i_p} g^*(W_{i_{p+1}} \cdots W_{i_{p+q}}), [N] \rangle \\ = \langle W_{2i_1} \cdots W_{2i_p} f^*(W_{2i_{p+1}} \cdots W_{2i_{p+q}}), [M] \rangle, \end{aligned}$$

where $i_1 + \cdots + i_{p+q} = n/2$ is a partition of $n/2$. Hence the proposition follows. \square

In the above proposition one can observe that if the underlying manifold M^n in (M^n, f) were a boundary then N in (N, g) would also be a boundary. Further, using induction, one can easily get the following

COROLLARY 4.6. *Let (M^n, f) be an n -dimensional singular manifold in BO with $\text{cobcat}(M^n, f) \leq 3$. Let $n = 2^r \cdot m$ where either m is odd and $m \geq 3$, or $m = 2$. Then either (M^n, f) is a boundary or else (M^n, f) is cobordant to $(N, g)^{2^r}$, where (N, g) is a non-bounding m -dimensional singular manifold in BO with $\text{cobcat}(N, g) \leq 3$. \square*

Finally, in view of the above results and the fact that $N_2(\text{BO})$ is generated by the bordism classes $[(RP^1, \tau_1)^2]$, $[RP^2, \tau_1]$ and $[RP^2, c]$, where $\tau_1: RP^i \rightarrow \text{BO}$ is the classifying map of the canonical line bundle over RP^i ($i = 1, 2$) and $c: RP^2 \rightarrow \text{BO}$ is the constant map, we can make the following remarks.

REMARK 4.7. Let (M^n, f) be as in (4.6) and $n = 2^r$, $r \geq 1$. Then either (M^n, f) is a boundary or else (M^n, f) is cobordant to $(N, g)^{2^{r-1}}$, where (N, g) is a 2-dimensional singular manifold generated by $(RP^1, \tau_1)^2$, (RP^2, τ_1) and/or (RP^2, c) . \square

REMARK 4.8. In Remark (4.7) if, in addition, the underlying manifold M^n in (M^n, f) were a boundary then (N, g) would be equal to exactly one of the following

- (i) $(RP^1, \tau_1)^2$,
- (ii) $(RP^2, c) \sqcup (RP^2, \tau_1)$, or
- (iii) $(RP^1, \tau_1)^2 \sqcup (RP^2, c) \sqcup (RP^2, \tau_1)$,

where $(M_1, f_1) \sqcup (M_2, f_2) = (M_1 \sqcup M_2, f_1 \sqcup f_2)$. \square

REFERENCES

- [1] H. Singh, *Lusternik-Schnirelmann category and cobordism*, Proc. Amer. Math. Soc., **102** (1) (1988), 183–190.
- [2] P. E. Conner, *Differentiable Periodic Maps*, Lecture Notes in Mathematics, vol. 738, Springer-Verlag, Berlin-Heidelberg-New York 1978.
- [3] R. E. Stong, *Notes on Cobordism Theory*, Math. Notes, Princeton Univ. Press, Princeton, N.J., 1968.
- [4] J. W. Milnor, *On the Stiefel-Whitney numbers of complex manifolds and spin manifolds*, Topology, **3** (1965), 223–230.

Received February 19, 1991 and in revised form July 8, 1991.

NORTH EASTERN HILL UNIVERSITY
SHILLONG, 793003
MEGHALAYA, INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@henri.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1992 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 155 No. 2 October 1992

Invariant subspaces and harmonic conjugation on compact abelian groups	201
NAKHLE HABIB ASMAR, EARL ROBERT BERKSON and THOMAS ALASTAIR GILLESPIE	
Cobcat and singular bordism	215
A. K. DAS and S. S. KHARE	
Poles of Eisenstein series on SL_n induced from maximal parabolics	229
PAUL FEIT	
On the postulation of 0-dimensional subschemes on a smooth quadric	251
SALVATORE GIUFFRIDA, RENATO MAGGIONI and ALFIO RAGUSA	
The Euler class for “piecewise” groups	283
PETER ABRAHAM GREENBERG	
L^p -Fourier transforms on nilpotent Lie groups and solvable Lie groups acting on Siegel domains	295
JUNKO INOUE	
Fuchsian moduli on a Riemann surface—its Poisson structure and Poincaré-Lefschetz duality	319
KATSUNORI IWASAKI	
Interpolation between H^p spaces and noncommutative generalizations. I	341
GILLES PISIER	
The braid index of generalized cables	369
ROBERT FONES WILLIAMS and ROBERT FONES WILLIAMS	
BMO and Hankel operators on Bergman spaces	377
KEHE ZHU	



0030-8730(1992)155:2;1-L