

*Pacific  
Journal of  
Mathematics*

**LOCAL REAL ANALYTIC BOUNDARY REGULARITY OF AN  
INTEGRAL SOLUTION OPERATOR OF THE  $\bar{\partial}$ -EQUATION ON  
CONVEX DOMAINS**

ZHENHUA CHEN

LOCAL REAL ANALYTIC BOUNDARY REGULARITY  
OF AN INTEGRAL SOLUTION OPERATOR  
OF THE  $\bar{\partial}$ -EQUATION ON CONVEX DOMAINS

ZHENHUA CHEN

**In this paper we show that a well known integral solution operator of the  $\bar{\partial}$ -equation on a convex domain  $\Omega$  locally preserves the real analyticity of  $\bar{\partial}$ -closed  $(0, 1)$  forms at boundary points near which  $\partial\Omega$  is totally convex in the complex tangential directions.**

**1. Introduction.** The real analytic boundary regularity of the canonical solution or Kohn's solution of the  $\bar{\partial}$ -equation was studied by many researchers [2], [3], [7], [8], [9], [10], [14], [15], [16]. In general, the canonical solution is not explicit. An interesting question is, whether we can find some explicit or computable solution of the  $\bar{\partial}$ -equation that is real analytically regular up to the boundary of pseudoconvex domains. In this paper we prove the following local real analytic boundary regularity of a well known integral solution operator of the  $\bar{\partial}$ -equation for totally convex domains (terminology will be defined in §2). The real analytic boundary regularity of the canonical solution of such domains has not been proved yet. The recent work of Boas and Straube [1] gives the global  $C^\infty$  boundary regularity of the canonical solution for convex domains.

**THEOREM.** *Suppose  $\Omega$  is a bounded convex domain in  $\mathbb{C}^n$  with  $C^2$  boundary. If  $p$  is a real analytic boundary point of  $\Omega$ , and  $\partial\Omega$  is totally convex at  $p$  in the complex tangential directions, then the Henkin operator  $T$  locally preserves the real analyticity of  $\bar{\partial}$ -closed  $(0, 1)$  forms up to the boundary point  $p$ .*

It had been conjectured for some time that global analytic hypoellipticity of the  $\bar{\partial}$ -Neumann problem held for weakly pseudoconvex domains, and this remains open. But local results on finite type domains came as a surprise comparing with  $C^\infty$  results. A recent counterexample of Christ and Geller [6] shows that local real analytic boundary regularity of the  $\bar{\partial}$ -equation does not hold for general pseudoconvex domains of finite type. We would like to point out that the domain

used by Christ and Geller in their counterexample does not satisfy the total convexity, which is the key assumption in our theorem, at the boundary point  $z = 0$ .

**2. Preliminaries.** Let  $h$  be a function on a domain  $\Omega \subseteq \mathbb{C}^n$  and  $p$  be a boundary point of  $\Omega$ . We say that  $h$  is *real analytic* at  $p$  if  $h$  has a convergent power series expansion in a neighborhood  $U$  of  $p$  in  $\mathbb{C}^n$ . We say that a  $(0, 1)$  form is a real analytic at  $p$  if its coefficients are all real analytic at  $p$ . Let  $E$  be a subset of  $\overline{\Omega}$ . We define

$C^\omega(E) =$  all functions that are real analytic at every point of  $E$ .

Suppose  $\Omega$  is a bounded convex domain in  $\mathbb{C}^n$  with  $C^2$  boundary. Following the standard  $\bar{\partial}$ -construction we can obtain an explicit solution of the  $\bar{\partial}$ -equation  $\bar{\partial}u = f$  on  $\Omega$  as follows. Let

$$\delta(z) = \begin{cases} -\text{dist}(z, \partial\Omega), & z \in \Omega, \\ \text{dist}(z, \partial\Omega), & z \notin \Omega, \end{cases}$$

$$\Phi(\zeta, z) = \sum_{j=1}^n \frac{\partial\delta}{\partial\zeta_j}(\zeta)(\zeta_j - z_j),$$

$$B = B(\zeta, z) = \frac{\partial_\zeta(|\zeta - z|^2)}{|\zeta - z|^2}, \quad W(\zeta, z) = \sum_{j=1}^n \frac{(\partial\delta/\partial\zeta_j)(\zeta)}{\Phi(\zeta, z)} d\zeta_j,$$

$$\widehat{W} = \widehat{W}(\lambda, \zeta, z) = \lambda W(\zeta, z) + (1 - \lambda)B(\zeta, z),$$

$$\Omega(B(\zeta, z)) = C_n B \wedge (\bar{\partial}_\zeta B)^{n-1}, \quad \Omega(\widehat{W}) = C_n \widehat{W} \wedge (\bar{\partial}_{\zeta, \lambda} \widehat{W})^{n-1},$$

where

$$\bar{\partial}_{\zeta, \lambda} = \bar{\partial}_\zeta + d_\lambda, \quad C_n = \frac{1}{(2\pi i)^n}.$$

Let

$$T(h)(z) = \int_{\partial\Omega \times I} h(\zeta) \wedge \Omega(\widehat{W}(\lambda, \zeta, z)) - \int_{\Omega} h(\zeta) \wedge \Omega(B(\zeta, z)).$$

It is well known that for any given  $\bar{\partial}$ -closed  $(0, 1)$  form  $f$  with coefficients in  $C^1(\overline{\Omega})$ ,  $u(z) = T(f)(z)$  is a solution of the  $\bar{\partial}$ -equation  $\bar{\partial}u = f$  in  $\Omega$  (see, for example, [13]). We call  $T$  the *Henkin operator*.

We say that the Henkin operator  $T$  *globally preserves the real analyticity* of  $\bar{\partial}$ -closed  $(0, 1)$  forms up to the boundary of  $\Omega$  if for any  $\bar{\partial}$ -closed  $(0, 1)$  form  $f$  with coefficients in  $C^\omega(\overline{\Omega})$  it follows that  $T(f) \in C^\omega(\overline{\Omega})$ . We say that the Henkin operator  $T$  *locally preserves the real analyticity* of  $\bar{\partial}$ -closed  $(0, 1)$  forms up to the boundary point

$p$  if for any  $\bar{\partial}$ -closed  $(0, 1)$  form  $f$  with coefficients in  $C^1(\bar{\Omega})$  and real analytic at  $p$  it follows that  $T(f)$  is also real analytic at  $p$ . It is clear that if the Henkin operator  $T$  locally preserves the real analyticity of  $\bar{\partial}$  closed  $(0, 1)$  forms up to the boundary of  $\Omega$  then  $T$  globally preserves the real analyticity of  $\bar{\partial}$ -closed  $(0, 1)$  forms up to the boundary of  $\Omega$ .

The following total convexity was used by Range [12] for studying the Carathéodory metric and holomorphic mappings. Here we use it as the basic assumption for our local real analytic boundary regularity result.

**DEFINITION.** Let  $\Omega$  be a convex domain in  $\mathbb{C}^n$  and  $p$  a  $C^1$  boundary point of  $\Omega$ . We say that  $\partial\Omega$  is *totally convex at  $p$  in the complex tangential directions* if there exists a neighborhood  $U$  of  $p$  in  $\mathbb{C}^n$  such that

$$\bar{\Omega} \cap (H_\zeta(\partial\Omega) + \{\zeta\}) = \{\zeta\},$$

for any  $\zeta \in \partial\Omega \cap U$ , where  $H_\zeta(\partial\Omega)$  is the complex tangent space of  $\partial\Omega$  at  $\zeta$ .

**3. Real analytic boundary regularity of Cauchy-Fantappie transform.** The following “integration by parts” lemma was proved in [5]:

**LEMMA 1.** *Let  $\Omega$  be a bounded convex domain in  $\mathbb{C}^n$  with  $C^2$  boundary, and  $p$  be a boundary point of  $\Omega$ . If there exists a neighborhood  $U$  of  $p$  in  $\mathbb{C}^n$  such that  $\partial\bar{\partial}/\partial\zeta_n \neq 0$  in  $U$  and*

$$\begin{aligned} a_n(\zeta, z) &\equiv \frac{\partial\Phi}{\partial\zeta_n}(\zeta, z) \\ &\quad - \frac{\partial\bar{\delta}}{\partial\zeta_n}(\zeta) \left( \frac{\partial\delta}{\partial\bar{\zeta}_n}(\zeta) \right)^{-1} \frac{\partial\Phi}{\partial\bar{\zeta}_n}(\zeta, z) \neq 0, \quad \text{in } U \times U, \end{aligned}$$

then for any positive integer  $k$  and any function  $h \in C^\infty(\bar{\Omega})$  with support in  $U \cup \bar{\Omega}$  we have

$$\begin{aligned} &\int_{\partial\Omega} \frac{h(\zeta)}{\Phi^k(\zeta, z)} d\zeta_1 \wedge \cdots \wedge d\zeta_n \wedge d\bar{\zeta}_1 \wedge \cdots \wedge d\bar{\zeta}_{n-1} \\ &= \frac{(-1)^{k-1}}{(k-1)!} \int_{\partial\Omega} (L_n)^{k+1}(h)(\zeta) \Phi(\zeta, z) \log \Phi(\zeta, z) d\zeta_1 \\ &\quad \wedge \cdots \wedge d\zeta_n \wedge d\bar{\zeta}_1 \wedge \cdots \wedge d\bar{\zeta}_{n-1} \\ &\quad + \frac{(-1)^{n+k-1}}{(k-1)!} \int_{\partial\Omega} (L_n)^k(h)(\zeta) d\zeta_1 \wedge \cdots \wedge d\zeta_n \wedge d\bar{\zeta}_1 \wedge \cdots \wedge d\bar{\zeta}_{n-1}, \end{aligned}$$

for any  $z \in U \cap \Omega$ , where

$$\begin{aligned} L_n &= \left( \frac{\partial \delta}{\partial \bar{\zeta}_n} \right)^{-1} \left( \frac{\partial \delta}{\partial \bar{\zeta}_n} \right) \frac{\partial}{\partial \bar{\zeta}_n} - \frac{\partial}{\partial \zeta_n} \\ &\quad + \left( \frac{\partial \delta}{\partial \bar{\zeta}_n} \right)^{-1} \left( \frac{\partial \delta}{\partial \bar{\zeta}_n} \right) \frac{\partial (a_n^{-1})}{\partial \bar{\zeta}_n} - \frac{\partial (a_n^{-1})}{\partial \zeta_n} \end{aligned}$$

is a first order differential operator.

Notice that  $H_\zeta(\partial\Omega) + \{\zeta\} = \{z \in \mathbb{C}^n : \Phi(\zeta, z) = 0\}$ . By the continuity of the function  $\Phi(\zeta, z)$ , we have

**LEMMA 2.** *Let  $\Omega$  be a bounded convex domain in  $\mathbb{C}^n$  with  $C^1$  boundary. If  $\partial\Omega$  is totally convex at a boundary point  $p$  of  $\Omega$  in the complex tangential directions, then there exists a neighborhood  $V$  of  $p$  in  $\mathbb{C}^n$  such that for any neighborhood  $U \subset\subset V$  of  $p$  in  $\mathbb{C}^n$  there exists a constant  $m > 0$  such that*

$$|\Phi(\zeta, z)| \geq m \quad \text{for any } \zeta \in \partial\Omega \cap (\mathbb{C}^n \setminus V) \text{ and any } z \in \bar{\Omega} \cap \bar{U}.$$

Since cut-off functions are not real analytic, when we localize our results we need the following result of Ehrenpreis (for a proof see [15]):

**LEMMA 3.** *Given  $U \subset\subset V \subset\subset \mathbb{C}^n$  there exist constants  $C, M > 0$  such that for any integer  $k$  we can find  $\psi_k \in C^\infty(\mathbb{C}^n)$  such that  $\psi_k \equiv 1$  in  $U$ ,  $\text{Supp } \psi_k \subset V$  and*

$$\left| \frac{\partial^{|\alpha|+|\beta|}}{\partial z^\alpha \partial \bar{z}^\beta} \psi_k(z) \right| \leq CM^{|\alpha|+|\beta|} k^{|\alpha|+|\beta|},$$

for any  $z$  in  $\mathbb{C}^n$ , and any  $\alpha, \beta : |\alpha| + |\beta| \leq k$ .

We now prove a local real analytic boundary regularity result about the Cauchy-Fantappie transform which may be of independent interest. A similar result was obtained by S.-C. Chen [4]. In [4] S.-C. Chen used an estimate of  $\Phi(\zeta, z)$ . Such an estimate can be obtained by using a fundamental theorem of Lojasiewicz about analytic varieties [11]. In our proof we do not use the fundamental theorem of Lojasiewicz.

**PROPOSITION 4.** *Suppose  $\Omega$  is a bounded convex domain in  $\mathbb{C}^n$  with  $C^2$  boundary. If  $p$  is a real analytic boundary point of  $\Omega$ , and  $\partial\Omega$*

is totally convex at  $p$  in the complex tangential directions, then the Cauchy-Fantappie transform  $C$

$$C(h)(z) = \int_{\partial\Omega} h(\zeta)\Omega(W(\zeta, z))$$

locally preserves the real analyticity up to the boundary point  $p$ .

*Proof.* Let  $h$  be any function that is in  $C^1(\overline{\Omega})$  and real analytic at  $p$ . We show that  $C(h)$  is real analytic at  $p$ . It suffices to show that for some neighborhood  $U$  of  $p$  in  $\mathbb{C}^n$  there exist constants  $C, M > 0$  such that

$$\left| \frac{\partial^{|\alpha|+|\beta|}}{\partial z^\alpha \partial \bar{z}^\beta} C(h)(z) \right| \leq CM^{|\alpha|+|\beta|} (|\alpha| + |\beta|)!,$$

for any  $z$  in  $U \cup \Omega$ , and any multi-indices  $\alpha, \beta$ . (For our convenience, constants  $C$  and  $M$  may take different values at different occurrences.)

Since  $\Phi(\zeta, z)$  is a holomorphic function of  $z$  in  $\Omega$ . When  $\beta > 0$  we have

$$\frac{\partial^{|\alpha|+|\beta|}}{\partial z^\alpha \partial \bar{z}^\beta} C(h)(z) = 0,$$

for any  $z \in \Omega$ . Hence we only need to estimate

$$\frac{\partial^{|\alpha|+|\beta|}}{\partial z^\alpha \partial \bar{z}^\beta} C(h)$$

with  $\beta = 0$ .

Let  $V_1$  be a neighborhood of  $p$  in  $\mathbb{C}^n$  such that  $h$  is real analytic on  $V_1$ . Since  $\delta$  is a defining function of  $\Omega$ , there exist a neighborhood  $V_2 \subseteq V_1$  of  $p$  in  $\mathbb{C}^n$  and some  $j$  such that  $\partial\delta/\partial\zeta_j \neq 0$  in  $V_2$  and

$$a_j(\zeta, z) = \frac{\partial\Phi}{\partial\zeta_j}(\zeta, z) - \frac{\partial\delta}{\partial\zeta_j}(\zeta) \left( \frac{\partial\delta}{\partial\zeta_j}(\zeta) \right)^{-1} \frac{\partial\Phi}{\partial\bar{\zeta}_j}(\zeta, z) \neq 0,$$

for any  $\zeta, z \in V_2$ . By Lemma 2, there exists a neighborhood  $V \subset\subset V_2$  of  $p$  in  $\mathbb{C}^n$  such that for any neighborhood  $U \subset\subset V$  of  $p$  in  $\mathbb{C}^n$  there exists a positive constant  $m$  such that

$$|\Phi(\zeta, z)| \geq m, \quad \text{for any } \zeta \in \partial\Omega \cap (\mathbb{C}^n \setminus V) \text{ and any } z \in U \cap \Omega.$$

By Lemma 3, there exist constants  $C, M > 0$  such that for any integer  $k$  we can find  $\psi_k \in C^\infty(\mathbb{C}^n)$  such that  $\psi_k \equiv 1$  in  $V$ ,  $\text{Supp } \psi_k \subset V_2$  and

$$\left| \frac{\partial^{|\alpha|+|\beta|}}{\partial z^\alpha \partial \bar{z}^\beta} \psi_k(z) \right| \leq CM^{|\alpha|+|\beta|} k^{|\alpha|+|\beta|},$$

for any  $z$  in  $\mathbb{C}^n$ , and any  $\alpha, \beta: |\alpha| + |\beta| \leq k$ .

Let  $U$  be a neighborhood of  $p$  in  $\mathbb{C}^n$  such that  $U \subset\subset V$ . We estimate

$$\frac{\partial^{|\alpha|}}{\partial z^\alpha} C(h)$$

on  $U \cap \Omega$ .

For any multi-index  $\alpha$ , we have

$$C(h) = C(\psi_{|\alpha|+n+1}h) + C((1 - \psi_{|\alpha|+n+1})h).$$

We first estimate  $C((1 - \psi_{|\alpha|+n+1})h)$ :

$$\begin{aligned} & \left| \frac{\partial^{|\alpha|}}{\partial z^\alpha} C((1 - \psi_{|\alpha|+n+1})h)(z) \right| \\ &= \left| \frac{\partial^{|\alpha|}}{\partial z^\alpha} \int_{\partial\Omega} (1 - \psi_{|\alpha|+n+1}(\zeta))h(\zeta)\Omega(W(\zeta, z)) \right| \\ &= \left| \frac{\partial^{|\alpha|}}{\partial z^\alpha} \int_{\partial\Omega} \frac{(1 - \psi_{|\alpha|+n+1}(\zeta))h(\zeta)}{\Phi^n(\zeta, z)} \Omega \left( \sum_{j=1}^n \frac{\partial\delta}{\partial\zeta_j} d\zeta_j \right) \right| \\ &= \left| \int_{\partial\Omega} \frac{\partial^{|\alpha|}}{\partial z^\alpha} \left( \frac{(1 - \psi_{|\alpha|+n+1}(\zeta))h(\zeta)}{\Phi^n(\zeta, z)} \right) \Omega \left( \sum_{j=1}^n \frac{\partial\delta}{\partial\zeta_j} d\zeta_j \right) \right| \\ &= \left| \int_{\partial\Omega} \frac{n(n+1)\cdots(n+|\alpha|-1) \left( \frac{\partial\delta}{\partial\zeta} \right)^\alpha (1 - \psi_{|\alpha|+n+1}(\zeta))h(\zeta)}{\Phi^{n+|\alpha|}(\zeta, z)} \right. \\ & \qquad \qquad \qquad \left. \times \Omega \left( \sum_{j=1}^n \frac{\partial\delta}{\partial\zeta_j} d\zeta_j \right) \right|. \end{aligned}$$

Since  $1 - \psi_{|\alpha|+n+1} \equiv 0$  in  $V$

$$\begin{aligned} &= \left| \int_{\partial\Omega \cap (\mathbb{C}^n \setminus V)} \frac{n(n+1)\cdots(n+|\alpha|-1) \left( \frac{\partial\delta}{\partial\zeta} \right)^\alpha (1 - \psi_{|\alpha|+n+1}(\zeta))h(\zeta)}{\Phi^{n+|\alpha|}(\zeta, z)} \right. \\ & \qquad \qquad \qquad \left. \times \Omega \left( \sum_{j=1}^n \frac{\partial\delta}{\partial\zeta_j} d\zeta_j \right) \right| \\ &\leq CM^{|\alpha|}|\alpha|!. \end{aligned}$$

for any  $z$  in  $U \cap \Omega$ .

We now estimate  $C(\psi_{|\alpha|+n+1}h)$ : Without loss of generality, we assume  $j = n$ . Set

$$\Omega \left( \sum_{j=1}^n \frac{\partial \delta}{\partial \zeta_j} d\zeta_j \right) = R(\zeta) d\zeta_1 \wedge \cdots \wedge d\zeta_n \wedge d\bar{\zeta}_1 \wedge \cdots \wedge d\bar{\zeta}_{n-1}.$$

Then

$$\begin{aligned} & \frac{\partial^{|\alpha|}}{\partial z^\alpha} C(\psi_{|\alpha|+n+1}h)(z) \\ &= \int_{\partial\Omega} \frac{n(n+1) \cdots (n+|\alpha|-1) \left( \frac{\partial \delta}{\partial \zeta} \right)^\alpha \psi_{|\alpha|+n+1} h R}{\Phi^{n+|\alpha|}(\zeta, z)} d\zeta_1 \\ & \quad \wedge \cdots \wedge d\zeta_n \wedge d\bar{\zeta}_1 \wedge \cdots \wedge d\bar{\zeta}_{n-1}. \end{aligned}$$

Notice that  $\psi_{|\alpha|+n+1}$  is supported in  $V \cap \bar{\Omega}$ . By Lemma 1:

$$\begin{aligned} &= \frac{(-1)^{|\alpha|+n}}{(|\alpha|+n)!} \int_{\partial\Omega} (L_n)^{n+|\alpha|+1} \left( n(n+1) \cdots (n+|\alpha|-1) \right. \\ & \quad \times \left. \left( \frac{\partial \delta}{\partial \zeta} \right)^\alpha \psi_{|\alpha|+n+1} h R \right) \Phi \log \Phi \\ & \quad \times d\zeta_1 \wedge \cdots \wedge d\zeta_n \wedge d\bar{\zeta}_1 \wedge \cdots \wedge d\bar{\zeta}_{n-1} \\ &+ \frac{(-1)^{|\alpha|}}{(|\alpha|+n)!} \int_{\partial\Omega} (L_n)^{n+|\alpha|} \left( n(n+1) \cdots (n+|\alpha|-1) \right. \\ & \quad \times \left. \left( \frac{\partial \delta}{\partial \zeta} \right)^\alpha \psi_{|\alpha|+n+1} h R \right) \\ & \quad \times d\zeta_1 \wedge \cdots \wedge d\zeta_n \wedge d\bar{\zeta}_1 \wedge \cdots \wedge d\bar{\zeta}_{n-1}, \end{aligned}$$

for any  $z \in V \cap \Omega$ .

Since  $h$ ,  $\delta$  and  $R$  are real analytic, and

$$|\Phi(\zeta, z) \log \Phi(\zeta, z)| \leq M,$$

for any  $\zeta \in V \cap \partial\Omega$  and any  $z \in V \cap \Omega$ , we have

$$\left| \frac{\partial^{|\alpha|}}{\partial z^\alpha} C(\psi_{|\alpha|+n+1}h)(z) \right| \leq CM^{|\alpha|} |\alpha|!,$$

for any  $z$  in  $V \cap \Omega$ .

Hence

$$\left| \frac{\partial^{|\alpha|}}{\partial z^\alpha} C(h)(z) \right| \leq CM^{|\alpha|} |\alpha|,$$

for any  $z$  in  $U \cap \Omega$ .

This completes the proof.

**4. Proof of the theorem.** Let  $f$  be a  $\bar{\partial}$ -closed  $(0, 1)$  form with coefficients in  $C^1(\bar{\Omega})$  and real analytic at the boundary point  $p$ . We show that  $T(f)$  is also real analytic at  $p$ .

Let  $U$  be a small ball with center  $p$  such that  $f$  is real analytic in  $U$ . By the Cauchy-Kovalevsky theorem, there exists a solution  $u_p$  in  $C^\omega(U)$  of the equation  $\bar{\partial}u_p = f$  in  $U$ .

Let  $V \subset\subset U$  be a neighborhood of  $p$  in  $\mathbb{C}^n$ , and  $\psi \in C^\infty(\mathbb{C}^n)$  such that  $\psi \equiv 1$  in  $V$  and  $\psi \equiv 0$  in  $\mathbb{C}^n \setminus U$ . Then

$$T(f) = T(\psi f) + T((1 - \psi)f).$$

Since the kernel of  $T$  is real analytic in the  $z$  variable when the variable  $\zeta$  is restricted to the support of  $(1 - \psi)f$ , by Lemma 2,  $T((1 - \psi)f)$  is real analytic at  $p$ .

Notice that

$$T(\psi f) = T(\psi \bar{\partial}u_p) = T(\bar{\partial}(\psi Tu_p)) - T((\bar{\partial}\psi)u_p).$$

By a standard homotopy argument, we have

$$T(\bar{\partial}(\psi u_p)) = \psi u_p - C(\psi u_p).$$

Hence

$$T(\psi f) = \psi u_p - C(\psi u_p) - T((\bar{\partial}\psi)u_p).$$

Clearly,  $\psi u_p$  is real analytic at  $p$ . By Proposition 4,  $C(\psi u_p)$  is real analytic at  $p$ . Since the kernel of  $T$  is real analytic in the  $z$  variable when the variable  $\zeta$  is restricted to the support of  $(\bar{\partial}\psi)u_p$ , by Lemma 2,  $T((\bar{\partial}\psi)u_p)$  is real analytic at  $p$ . Hence  $T(\psi f)$  is real analytic at  $p$ . Since both  $T(\psi f)$  and  $T((1 - \psi)f)$  are real analytic at  $p$  it follows that  $T(f)$  is real analytic at  $p$ .

This completes the proof.

#### REFERENCES

- [1] H. P. Boas and E. J. Straube, *Sobolev estimates for the  $\bar{\partial}$ -Neumann operator on convex domains in  $\mathbb{C}^n$* , preprint.
- [2] S.-C. Chen, *Global analytic hypoellipticity of the  $\bar{\partial}$ -Neumann problem on circular domains*, *Invent. Math.*, **92** (1988), 173–185.

- [3] —, *Global real analyticity of solutions of the  $\bar{\partial}$ -Neumann problem on Reinhardt domains*, Indiana Univ. Math. J., **37** (1988), 421–430.
- [4] —, *Real analytic boundary regularity of the Cauchy transform on convex domains*, Proc. Amer. Math. Soc., **108** (1990), 423–432.
- [5] Z. Chen, *The local  $C^\infty$  boundary regularity of an integral solution operator of the  $\bar{\partial}$ -equation on convex domains*, preprint.
- [6] M. Christ and D. Geller, *Counterexamples to analytic hypoellipticity for domains of finite type*, preprint.
- [7] M. Derridj and D. S. Tartakoff, *On the global real analyticity for the  $\bar{\partial}$ -Neumann problem*, Comm. Partial Differential Equations, **5** (1976), 401–435.
- [8] —, *Local analyticity for  $\square_b$  and the  $\bar{\partial}$ -Neumann problem at certain weakly pseudoconvex points*, preprint.
- [9] —, *Local analyticity for the  $\bar{\partial}$ -Neumann problem for  $\square_b$  in some model domains without maximal estimates*, preprint.
- [10] G. Komatsu, *Global analytic hypoellipticity of the  $\bar{\partial}$ -Neumann problem*, Tôhoku Math. J., **28** (1976), 145–156.
- [11] S. Lojasiewicz, *Sur la problème de la division*, Studia Math., **18** (1959), 87–136.
- [12] R. M. Range, *The Caratheodory metric and holomorphic maps on a class of weakly pseudoconvex domains*, Pacific J. Math., **78** (1978), 173–188.
- [13] —, *Holomorphic Functions and Integral Representations in Several Complex Variables*, Springer-Verlag, New York, 1986.
- [14] D. S. Tartakoff, *Local analytic hypoellipticity for  $\square_b$  on non-degenerate Cauchy-Riemann manifolds*, Proc. Nat. Acad. Sci. U.S.A., **75** (1978), 3027–3028.
- [15] —, *On the local real analyticity of solutions on  $\square_b$  and the  $\bar{\partial}$ -Neumann problem*, Acta Math., **145** (1980), 117–204.
- [16] F. Trèves, *Analytic hypo-ellipticity of a class of pseudodifferential operators with double characteristics and application to the  $\bar{\partial}$ -Neumann problem*, Comm. Partial Differential Equations, **3** (1978), 475–642.

Received April 2, 1991. Much of the work in this paper is contained in the author's Washington University Ph.D. Thesis. The author is grateful to Steven G. Krantz for his advice and encouragement.

UNIVERSITY OF MICHIGAN  
ANN ARBOR, MI 48109



# PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982)      F. WOLF (1904–1989)

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555  
vsv@math.ucla.edu

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112  
clemens@math.utah.edu

F. MICHAEL CHRIST  
University of California  
Los Angeles, CA 90024-1555  
christ@math.ucla.edu

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093  
tenright@ucsd.edu

NICHOLAS ERCOLANI  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

R. FINN  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720  
vfr@math.berkeley.edu

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

C. C. MOORE  
University of California  
Berkeley, CA 94720

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106  
mgscharl@henri.ucsb.edu

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

# PACIFIC JOURNAL OF MATHEMATICS

Volume 156 No. 1 November 1992

---

Enveloping algebras of Lie groups with discrete series	1
NGUYEN HUU ANH and VUONG MANH SON	
Asymptotic behavior of eigenvalues for a class of pseudodifferential operators on $\mathbf{R}^n$	19
JUNICHI ARAMAKI	
A hybrid of theorems of Vinogradov and Piatetski-Shapiro	45
ANTAL BALOG and JOHN BENJAMIN FRIEDLANDER	
Chaos in terms of the map $x \rightarrow \omega(x, f)$	63
ANDREW MICHAEL BRUCKNER and JACK GARY CEDER	
Local real analytic boundary regularity of an integral solution operator of the $\bar{\partial}$ -equation on convex domains	97
ZHENHUA CHEN	
On some properties of exhaustion maps between bounded domains	107
CHI KEUNG CHEUNG	
A generalization of maximal functions on compact semisimple Lie groups	119
HENDRA GUNAWAN	
Stability of nonsingular group orbits	135
CLARK DEAN HORTON	
Bordism and regular homotopy of low-dimensional immersions	155
JOHN FORBES HUGHES	
On six-connected finite $H$ -spaces	185
JAMES PEICHENG LIN and FRANK WILLIAMS	



0030-8730(1992)156:1;1-L