

*Pacific
Journal of
Mathematics*

**ASYMPTOTICALLY FREE FAMILIES OF RANDOM
UNITARIES IN SYMMETRIC GROUPS**

ALEXANDRU MIHAI NICA

Volume 157 No. 2

February 1993

ASYMPTOTICALLY FREE FAMILIES OF RANDOM UNITARIES IN SYMMETRIC GROUPS

ALEXANDRU NICA

We prove that independent, Haar distributed families of random unitaries in symmetric groups are asymptotically free.

1. Introduction. In this note we prove that independent, Haar distributed families of random unitaries in symmetric groups are asymptotically free.

If G_n is a closed subgroup of the unitary group $U(n)$, then by a random unitary in G_n we understand a measurable function $f: X \rightarrow G_n$, where (X, \mathcal{F}, P) is a (fixed) probability space. A random unitary in G_n has a distribution, which is a probability measure on G_n , and we can define the notion of independence for a family of random unitaries, exactly as it is done for usual real-valued random variables. A family $(f_\omega)_{\omega \in \Omega}$ of random unitaries in G_n will be called, following [1], standard-independent if it is independent and if the distribution of every f_ω is the Haar measure on G_n .

Now, a family $(f_\omega)_{\omega \in \Omega}$ of random unitaries in G_n can be also viewed as a family of unitaries in the non-commutative probability space (\mathfrak{M}_n, τ_n) , where \mathfrak{M}_n is the unital $*$ -algebra of measurable functions $X \rightarrow \text{Mat}_n(\mathbb{C})$, having bounded entries, and τ_n is the trace-state of \mathfrak{M}_n obtained by integrating the normalized trace of $\text{Mat}_n(\mathbb{C})$. From this point of view, the concept analogous to independence to be considered is the property of $(f_\omega)_{\omega \in \Omega}$ of being or not being free (see [2]). This property can be expressed in terms of a naturally defined "non-commutative distribution" of $(f_\omega)_{\omega \in \Omega}$, which is a state on the group algebra of the free group on Ω generators.

Hence, we are in a situation when both concepts of independence and freeness can be considered. It seems to be a deep phenomenon that, as found by D. Voiculescu in [1] for several important examples of such situations, one can hope independence to give rise (at least in good cases) to asymptotic freeness.

In our particular framework, the problem-type reflecting this phenomenon can be stated as follows: "For every $n \geq 1$, let $(f_n, \omega)_{\omega \in \Omega}$

be a standard-independent family of random unitaries in the closed subgroup G_n of $U(n)$. Is it true, for a reasonable series $(G_n)_{n=1}^\infty$, that the families $(f_{n,\omega})_{\omega \in \Omega}$ are asymptotically free for $n \rightarrow \infty$?" In [1] it is proved that this is true for $G_n = U(n)$, and it is found that the limit non-commutative distribution of the families $(f_{n,\omega})_{\omega \in \Omega}$ is, roughly speaking, the free product of Ω copies of the Haar measure on the circle.

The goal of this paper is to prove that the same holds if every G_n is a semidirect product $A_n^n \rtimes_{\alpha_n} S_n$, with S_n the group of permutations of $\{1, 2, \dots, n\}$, A_n a closed subgroup of the circle, and $\alpha_n: S_n \rightarrow \text{Aut}(A_n^n)$ the natural action,

$$\alpha_n(t)(z_1, z_2, \dots, z_n) = (z_{t^{-1}(1)}, z_{t^{-1}(2)}, \dots, z_{t^{-1}(n)}).$$

The paper is subdivided into sections as follows: in §2 we fix the notations and review the concepts related to non-commutative probability spaces that we need. In §3 we state precisely the problem-type concerning asymptotic freeness for standard-independent families of random unitaries. In §4 we prove the main theorem of the paper, namely that asymptotic freeness holds for the series $(S_n)_{n=1}^\infty$ of symmetric groups, and in §5 we extend the result to the above-mentioned case of semidirect products.

We are deeply indebted to Dan Voiculescu for his constant support during the preparation of this work. We are thankful for the good atmosphere of the Operator Algebra theme year at the C. R. M., Montreal, where the paper was written.

2. Basic definitions. In this section we fix the notations and briefly review the basic concepts about non-commutative random variables that we need (for a more detailed exposition, see [2]).

2.1. By a non-commutative probability space we shall understand a pair (\mathfrak{A}, σ) , with \mathfrak{A} a unital $*$ -algebra and σ a state of \mathfrak{A} (i.e. $\sigma: \mathfrak{A} \rightarrow \mathbb{C}$ linear, $\sigma(a^*a) \geq 0$ for any $a \in \mathfrak{A}$, $\sigma(1) = 1$).

2.2. **-distributions for families of unitaries.* For a non-void set Ω , let $F(\Omega)$ be the free group having a free family of generators indexed by Ω , and let $\mathbb{C}[F(\Omega)]$ be its group algebra. It is handy to view $\mathbb{C}[F(\Omega)]$ as an algebra of non-commutative trigonometric polynomials, i.e., as having a linear basis consisting of 1 and the monomials $X_{\omega_1}^{\alpha_1} X_{\omega_2}^{\alpha_2} \cdots X_{\omega_m}^{\alpha_m}$, with $\omega_1 \neq \omega_2 \neq \cdots \neq \omega_m$ in Ω and $\alpha_1, \alpha_2, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$. (Note: By $\omega_1 \neq \omega_2 \neq \cdots \neq \omega_m$ we shall always mean that $\omega_i \neq \omega_{i+1}$, $1 \leq i \leq m-1$.) Of course, if $\text{card } \Omega = 1$, then $\mathbb{C}[F(\Omega)] = \mathbb{C}[\mathbb{Z}]$ is the algebra of usual trigonometric polynomials.

$\mathbb{C}[F(\Omega)]$ has a natural $*$ -operation, uniquely determined by the condition that every X_ω ($\omega \in \Omega$) be unitary, and has the following universality property: for any unital $*$ -algebra \mathfrak{A} , and for any family $(u_\omega)_{\omega \in \Omega}$ of unitaries of \mathfrak{A} , there exists a unique homomorphism of unital $*$ -algebras, $\Pi: \mathbb{C}[F(\Omega)] \rightarrow \mathfrak{A}$, such that $\Pi(X_\omega) = u_\omega$, for every ω in Ω .

DEFINITION. Let (\mathfrak{A}, σ) be a non-commutative probability space. The $*$ -distribution of a unitary $u \in \mathfrak{A}$ will be the state $\mu_u: \mathbb{C}[\mathbb{Z}] \rightarrow \mathbb{C}$ obtained by composing σ with the unique homomorphism $\mathbb{C}[\mathbb{Z}] \rightarrow \mathfrak{A}$ sending X into u . More generally, the $*$ -distribution of a family $(u_\omega)_{\omega \in \Omega}$ of unitaries in \mathfrak{A} will be the state $\mu: \mathbb{C}[F(\Omega)] \rightarrow \mathbb{C}$ obtained by composing σ with the unique homomorphism $\mathbb{C}[F(\Omega)] \rightarrow \mathfrak{A}$ sending X_ω into u_ω for every ω .

2.3. Example (Haar distribution on $\mathbb{C}[\mathbb{Z}]$). Consider the algebra $C(\mathbb{T})$ of continuous functions on the circle, with pointwise operations, let $\sigma: C(\mathbb{T}) \rightarrow \mathbb{C}$ be integration with respect to Haar measure, and let $\text{id} \in C(\mathbb{T})$ be the unitary $\text{id}(z) = z$. The $*$ -distribution $\mu_{\text{id}}: \mathbb{C}[\mathbb{Z}] \rightarrow \mathbb{C}$ will be called the Haar distribution on $\mathbb{C}[\mathbb{Z}]$; it is clearly determined by the property that $\mu_{\text{id}}(X^k) = 0$ for any $k \neq 0$.

2.4. Free families. Let (\mathfrak{A}, σ) be a non-commutative probability space. A family $(u_\omega)_{\omega \in \Omega}$ of unitaries in \mathfrak{A} is called free if for any $\omega_1 \neq \omega_2 \neq \dots \neq \omega_m$ in Ω and p_1, p_2, \dots, p_m in $\mathbb{C}[\mathbb{Z}]$ such that $\sigma(p_j(u_{\omega_j})) = 0$, $1 \leq j \leq m$, we also have that $\sigma(p_1(u_{\omega_1}) \cdots p_m(u_{\omega_m})) = 0$.

Remark that the fact whether the family $(u_\omega)_{\omega \in \Omega}$ is free or not depends only on its $*$ -distribution μ on $\mathbb{C}[F(\Omega)]$. More precisely, it is easy to check that $(u_\omega)_{\omega \in \Omega}$ is free in (\mathfrak{A}, σ) if and only if $(X_\omega)_{\omega \in \Omega}$ is free in $(\mathbb{C}[F(\Omega)], \mu)$.

2.5. Asymptotically free families. Let, for every $n \geq 1$, $(u_{n,\omega})_{\omega \in \Omega}$ be a family of unitaries in the non-commutative probability space $(\mathfrak{A}_n, \sigma_n)$, and let μ_n be its $*$ -distribution. The families $(u_{n,\omega})_{\omega \in \Omega}$ are said to converge in distribution (for $n \rightarrow \infty$) to the state μ on $\mathbb{C}[F(\Omega)]$ if $\mu_n(p) \xrightarrow{n \rightarrow \infty} \mu(p)$, for any p in $\mathbb{C}[F(\Omega)]$. If, moreover, the limit state μ has the property that the family $(X_\omega)_{\omega \in \Omega}$ is free in $(\mathbb{C}[F(\Omega)], \mu)$, then the families $(u_{n,\omega})_{\omega \in \Omega}$ are said to be asymptotically free for $n \rightarrow \infty$.

2.6. Random matrices. For (X, \mathcal{F}, P) a probability space and n a positive integer, we shall work with the unital $*$ -algebra \mathfrak{M}_n of

measurable functions $f: X \rightarrow \text{Mat}_n(\mathbb{C})$ having the property that for any $1 \leq i, j \leq n$, the entry $f_{i,j}: X \rightarrow \mathbb{C}$ is bounded. This is only a subalgebra of what one usually calls the algebra of $n \times n$ random matrices on (X, \mathcal{F}, P) (see the definitions preceding Theorem 2.2 of [1]), but it will be sufficient for our purposes. On \mathfrak{M}_n we have a canonical trace state, defined by:

$$(1) \quad \tau_n(f) = \frac{1}{n} \int_X \text{Tr} f(x) dP(x) = \frac{1}{n} \sum_{j=1}^n \int_X f_{j,j} dP.$$

2.7. Random unitaries. Let (X, \mathcal{F}, P) and n be as above, and assume that we are also given a closed subgroup G of $U(n)$. A measurable function $f: X \rightarrow G$ (which is in particular a unitary in the algebra \mathfrak{M}_n defined at 2.6) will be called a random unitary in G .

As a unitary in the non-commutative probability space (\mathfrak{M}_n, τ_n) , a random unitary f in G has of course a *-distribution in the sense of 2.2. But in this case, we also have the distribution of f defined in the classical sense, which is the probability measure on the Borel σ -algebra of G , given by the formula: $\lambda_f(A) = P(f^{-1}(A))$. More generally, a finite family $(f_\omega)_{\omega \in \Omega}$ of random unitaries in G has a joint distribution, which is the probability measure λ on the compact group G^Ω , given by

$$\lambda(A) = P(\{x \in X | (f_\omega(x))_{\omega \in \Omega} \in A\}), \quad A \subseteq G^\Omega \text{ Borel set.}$$

If $\lambda = \prod_{\omega \in \Omega} \lambda_{f_\omega}$, the family $(f_\omega)_{\omega \in \Omega}$ is called independent. If this happens and, moreover, each λ_{f_ω} coincides with the Haar measure on G , the family $(f_\omega)_{\omega \in \Omega}$ will be called standard-independent (see 3.7 of [1]).

An arbitrary family (finite or not) $(f_\omega)_{\omega \in \Omega}$ of random unitaries in G will be called standard-independent if so is $(f_\omega)_{\omega \in \Omega_0}$ for any finite subset Ω_0 of Ω . As it is easily checked, this is equivalent to the fact that

$$(2) \quad \int_X \varphi_1(f_{\omega_1}(x)) \cdots \varphi_m(f_{\omega_m}(x)) dP(x) = \prod_{q=1}^m \int_G \varphi_q(t) dt,$$

for any $\omega_1, \dots, \omega_m$ in Ω such that $\omega_i \neq \omega_j$ when $i \neq j$, and for any $\varphi_1, \dots, \varphi_m$ in $C(G)$; on the right side of (2), integration is done with respect to the Haar measure of G .

3. The setting of the problem.

3.1. The enunciation of the problem can be done as follows. For every $n \geq 1$, let $(f_{n,\omega})_{\omega \in \Omega}$ be a standard-independent family of random unitaries in the closed subgroup G_n of $U(n)$. Is it true, for reasonable series $(G_n)_{n=1}^\infty$, that:

(i) the families $(f_{n,\omega})_{\omega \in \Omega}$, regarded in the non-commutative probability spaces (\mathfrak{M}_n, τ_n) , are asymptotically free, and

(ii) For every $\omega \in \Omega$, the unitaries $f_{n,\omega}$ converge in distribution (for $n \rightarrow \infty$) to the Haar distribution on $\mathbb{C}[\mathbb{Z}]$?

3.2. *Reformulation.* For concrete computations, it is useful to remark that (i) and (ii) of 3.1 together are equivalent to the following assertion: For any $\omega_1 \neq \omega_2 \neq \dots \neq \omega_m$ in Ω , and any $\alpha_1, \alpha_2, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$, we have:

$$(3) \quad \lim_{n \rightarrow \infty} \tau_n(f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m}) = 0.$$

Indeed, (i) + (ii) mean that the *-distributions of the families $(f_{n,\omega})_{\omega \in \Omega}$ converge, for $n \rightarrow \infty$, to a state μ on $\mathbb{C}[F(\Omega)]$ with the following properties:

(j) $(X_\omega)_{\omega \in \Omega}$ is a free family in $(\mathbb{C}[F(\Omega)], \mu)$;

(jj) $\mu(X_\omega^k) = 0$ for any ω in Ω and k in $\mathbb{Z} \setminus \{0\}$.

But (j) and (jj) together are clearly equivalent to

(jjj) $\mu(X_{\omega_1}^{\alpha_1} \cdots X_{\omega_m}^{\alpha_m}) = 0$, for any $\omega_1 \neq \omega_2 \neq \dots \neq \omega_m$ in Ω and $\alpha_1, \alpha_2, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$

(in particular (j) + (jj) determine μ completely).

So, if (i) and (ii) hold, we have for any $\omega_1 \neq \omega_2 \neq \dots \neq \omega_m$ in Ω and any $\alpha_1, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$:

$$\lim_{n \rightarrow \infty} \tau_n(f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m}) = \mu(X_{\omega_1}^{\alpha_1} \cdots X_{\omega_m}^{\alpha_m}) = 0;$$

conversely, if (3) holds, then the *-distributions of the families $(f_{n,\omega})_{\omega \in \Omega}$ converge for $n \rightarrow \infty$ to a state on $\mathbb{C}[F(\Omega)]$ satisfying (jjj), and we have (i) + (ii).

3.3. **REMARK.** The expression $\tau_n(f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m})$ appearing in (3) depends in fact only on G_n (and not on the probability space (X, \mathcal{F}, P) we started with). Indeed, if $f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m} = f$, then for any $1 \leq j \leq n$, the entry $f_{j,j}$ of f is a polynomial in the entries of $f_{n,\omega_1}, \dots, f_{n,\omega_m}$, and $\int_X f_{j,j} dP$ is seen not to depend on (X, \mathcal{F}, P) because of (2) of 2.7.

3.4. *Reformulation for G finite.* Let us assume that the subgroups $G_n \subseteq U(n)$ considered at 3.1 are finite.

Let $(f_{n,\omega})_{\omega \in \Omega}$ be a family of random unitaries in G_n . Since $C(G_n)$ is the linear span of characteristic functions of one-point sets, and Haar measure on G_n is the normalized counting measure, the relation (2) of 2.7 (i.e. the standard-independence of $(f_{n,\omega})_{\omega \in \Omega}$) is seen to be equivalent to

$$(4) \quad P(f_{n,\omega_1}^{-1}(t_1) \cap \cdots \cap f_{n,\omega_m}^{-1}(t_m)) = \frac{1}{(\text{card } G_n)^m},$$

for any $\omega_1, \dots, \omega_m$ in Ω such that $\omega_i \neq \omega_j$ when $i \neq j$, and for any t_1, \dots, t_m in G_n .

Further, let $(f_{n,\omega})_{\omega \in \Omega}$ be a standard-independent family of random unitaries in G_n , and let us compute $\tau_n(f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m})$ for some $\omega_1 \neq \omega_2 \neq \cdots \neq \omega_m$ in Ω and $\alpha_1, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$. When q runs from 1 to m , ω_q describes a subset $\omega'_1, \dots, \omega'_k$ of Ω , with $k \leq m$. In other words, we have written $\omega_q = \omega'_q$, ($1 \leq q \leq m$), with $\omega'_i \neq \omega'_j$ for $i \neq j$. The hypothesis $\omega_1 \neq \omega_2 \neq \cdots \neq \omega_m$ becomes $c_1 \neq c_2 \neq \cdots \neq c_m$.

We claim that

$$(5) \quad \tau_n(f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m}) = \frac{1}{n(\text{card } G_n)^k} \sum_{t_1, \dots, t_k \in G_n} \text{Tr}(t_{c_1}^{\alpha_1} \cdots t_{c_m}^{\alpha_m}).$$

Indeed, for any t_1, \dots, t_k in G_n , the function $f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m}$ is constant and equal to $t_{c_1}^{\alpha_1} \cdots t_{c_m}^{\alpha_m}$ on the set $f_{n,\omega'_1}^{-1}(t_1) \cap \cdots \cap f_{n,\omega'_k}^{-1}(t_k)$, which has measure $1/(\text{card } G_n)^k$, by (4). The sets of this form realise a partition of X (when t_1, \dots, t_k describe G_n); decomposing the integral which appears in the formula (1) of 2.6 after this partition, we get (5).

We conclude that a sufficient condition for having an affirmative answer to (i) and (ii) of 3.1 (for any indexing set Ω) is, in this case

$$(6) \quad \lim_{n \rightarrow \infty} \frac{1}{n(\text{card } G_n)^k} \sum_{t_1, \dots, t_k \in G_n} \text{Tr}(t_{c_1}^{\alpha_1} \cdots t_{c_m}^{\alpha_m}) = 0,$$

for any $k \geq 1$, $c_1 \neq c_2 \neq \cdots \neq c_m$ exhausting $\{1, \dots, k\}$ and $\alpha_1, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$. Clearly, this condition is also necessary (take $\Omega = \{1, \dots, k\}$ and $\omega_1 = c_1, \dots, \omega_m = c_m$).

4. Asymptotic freeness in the case of the symmetric groups.

4.1. *Statement of the result.* We view the symmetric group S_n as a subgroup of $U(n)$, by identifying every t in S_n with the corresponding permutation matrix (the entry (i, j) equals 1 if $t(j) = i$, and 0 otherwise). We shall prove that:

THEOREM. *The assertions (i) and (ii) of 3.1 are true for the series $(S_n)_{n=1}^\infty$.*

As we saw at 3.4, this comes to

$$(7) \quad \lim_{n \rightarrow \infty} \frac{1}{n(n!)^k} \sum_{t_1, \dots, t_k \in S_n} \text{Tr}(t_{c_1}^{\alpha_1} \dots t_{c_m}^{\alpha_m}) = 0,$$

for any $k \geq 1$, $c_1 \neq c_2 \neq \dots \neq c_m$ exhausting $\{1, \dots, k\}$ and $\alpha_1, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$.

The numbers: $k, c_1, \dots, c_m, \alpha_1, \dots, \alpha_m$ will be fixed for the rest of this section. In fact, we shall also fix for the rest of the section an integer n , not too small (for instance such that $n > 2(|\alpha_1| + \dots + |\alpha_m|)$), and prove the inequality:

$$(8) \quad \frac{1}{n(n!)^k} \sum_{t_1, \dots, t_k \in S_n} \text{Tr}(t_{c_1}^{\alpha_1} \dots t_{c_m}^{\alpha_m}) \leq \frac{1}{n}(\alpha + 2)^\alpha,$$

with $\alpha = |\alpha_1| + \dots + |\alpha_m|$. Clearly, (8) implies (7) and hence the proposition.

4.2. REMARK. In some particular cases, the left side of (8) can be computed precisely. We give here some examples (we omit the proofs, since they are not part of the main stream of this paper).

1°. Assume that there exists $1 \leq j \leq k$ with the following property: there is only one q ($1 \leq q \leq m$) with $c_q = j$, and for that q we have $\alpha_q = \pm 1$. Then the left side of (8) is exactly $1/n$.

2°. Assume that for every $1 \leq j \leq k$ there is only one q ($1 \leq q \leq m$) with $c_q = j$ (but instead there is no condition on the exponents). After a change of indices, the non-commutative monomial $t_{c_1}^{\alpha_1} \dots t_{c_m}^{\alpha_m}$ becomes $t_1^{\alpha_1} \dots t_k^{\alpha_k}$. We have

$$\frac{1}{n(n!)^k} \sum_{t_1, \dots, t_k \in S_n} \text{Tr}(t_1^{\alpha_1} \dots t_k^{\alpha_k}) = \frac{1}{n} + \frac{\prod_{j=1}^k (\Delta(\alpha_j) - 1)}{n(n-1)^{k-1}},$$

where $\Delta(\alpha_j)$ denotes the number of positive divisors of α_j .

3°. Assume that the non-commutative monomial $t_{c_1}^{\alpha_1} \dots t_{c_m}^{\alpha_m}$ is the commutator of two permutations. We have:

$$\frac{1}{n(n!)^2} \sum_{t_1, t_2 \in S_n} \text{Tr}(t_1 t_2 t_1^{-1} t_2^{-2}) = \frac{1}{n-1}.$$

It would be interesting to find such precise evaluations in the general case.

4.3. NOTATIONS. (a) Besides $k, c_1, \dots, c_m, \alpha_1, \dots, \alpha_m$ and $\alpha = |\alpha_1| + \dots + |\alpha_m|$ which were fixed at 4.1, we also fix the following partition of $\{1, 2, \dots, \alpha\}$ into integer valued intervals:

$$\begin{aligned}
 I_1 &= \{1, \dots, |\alpha_1|\}, \\
 I_2 &= \{|\alpha_1| + 1, \dots, |\alpha_1| + |\alpha_2|\}, \\
 &\dots\dots\dots \\
 I_m &= \{|\alpha_1| + \dots + |\alpha_{m-1}| + 1, \dots, |\alpha_1| + \dots + |\alpha_{m-1}| + |\alpha_m|\}.
 \end{aligned}$$

(b) *Relations.* By a relation we shall mean a subset of $\{1, \dots, n\}^2$. A relation R will be called injective if the two projections on the components are injective when restricted to R . For any t in S_n , the relation R_t associated to t will be $\{(i, j) | t(i) = j\}$.

It is easy to see that for a given relation R , there exist permutations t such that $R \subseteq R_t$ if and only if R is injective; if this happens, the number of permutations t such that $R \subseteq R_t$ equals $(n - (\text{card } R))!$.

(c) *Cycles.* By a cycle we shall understand a sequence $\xi = (u_1, \dots, u_\alpha, u_{\alpha+1})$ of numbers in $\{1, \dots, n\}$, such that $u_{\alpha+1} = u_1$. To a cycle ξ we shall associate k relations, $R_1(\xi), \dots, R_k(\xi)$. It will be useful at 4.7 to have the construction made for any sequence $\eta = (u_1, \dots, u_\beta)$ of elements of $\{1, \dots, n\}$, with $2 \leq \beta \leq \alpha + 1$. So, having such a sequence η , we define for any $1 \leq j \leq k$ a relation $R_j(\eta)$ as follows: we take all the numbers $1 \leq a \leq \beta - 1$ which belong to intervals I_q ($1 \leq q \leq m$) having $c_q = j$; and for any such $a \in I_q$ we take into $R_j(\eta)$ the couple:

$$\begin{cases} (u_{a+1}, u_a), & \text{if } \alpha_q > 0; \\ (u_a, u_{a+1}), & \text{if } \alpha_q < 0. \end{cases}$$

A sequence $\eta = (u_1, \dots, u_\beta)$ will be called injective if the relations $R_1(\eta), \dots, R_k(\eta)$ are so.

Having a cycle ξ and a k -tuple (t_1, \dots, t_k) in S_n^k , we shall write $\xi \prec (t_1, \dots, t_k)$ if $R_j(\xi) \subseteq R_{t_j}$, for all $1 \leq j \leq k$.

Now, for $\xi = (u_1, \dots, u_\alpha, u_{\alpha+1})$ a cycle, t a permutation and $1 \leq j \leq k$, we clearly have: $R_j(\xi) \subseteq R_t \Leftrightarrow t^{\text{sign } \alpha_q}(u_{a+1}) = u_a$, for any $1 \leq q \leq m$ such that $c_q = j$, and for any $a \in I_q$. This gives the criterion

$$\begin{aligned}
 (9) \quad \xi &= (u_1, \dots, u_\alpha, u_{\alpha+1}) \prec (t_1, \dots, t_k) \\
 &\Leftrightarrow t_{c_q}^{\text{sign } \alpha_q}(u_{a+1}) = u_a, \quad \forall 1 \leq q \leq m, a \in I_q.
 \end{aligned}$$

4.4. LEMMA. For any t_1, \dots, t_k in S_n we have:

$$\text{Tr}(t_{c_1}^{\alpha_1} \cdots t_{c_m}^{\alpha_m}) = \text{card}\{\xi \text{ cycle} \mid \xi \prec (t_1, \dots, t_k)\}.$$

Proof. Let $t_{c_q}^{\text{sign} \alpha_q} \stackrel{\text{def}}{=} s_q = (s_q; u, v)_{1 \leq u, v \leq n} \in U(n)$. Then:

$$\begin{aligned} \text{Tr}(t_{c_1}^{\alpha_1} \cdots t_{c_m}^{\alpha_m}) &= \text{Tr}(s_1^{|\alpha_1|} \cdots s_m^{|\alpha_m|}) \\ &= \sum_{\xi=(u_1, \dots, u_{\alpha+1}), \text{ cycle}} \left(\prod_{q=1}^m \prod_{a \in I_q} s_q; u_a, u_{a+1} \right). \end{aligned}$$

Since $s_q; u, v$ equals 1 if $t_{c_q}^{\text{sign} \alpha_q}(v) = u$, and 0 otherwise, every term of the last sum is 0 or 1, and it is 1 if and only if $t_{c_q}^{\text{sign} \alpha_q}(u_{a+1}) = u_a$ for every $1 \leq q \leq m$ and $a \in I_q$. Comparing with (9) we obtain the desired equality. \square

4.5. LEMMA. For any injective cycle ξ we have

$$(10) \quad \begin{aligned} \text{card}\{(t_1, \dots, t_k) \in S_n^k \mid \xi \prec (t_1, \dots, t_k)\} \\ \leq (n!)^k (n/2)^{-\sum_{j=1}^k \text{card } R_j(\xi)}. \end{aligned}$$

Proof. We have:

$$\{(t_1, \dots, t_k) \in S_n^k \mid \xi \prec (t_1, \dots, t_k)\} = \prod_{j=1}^k \{t \in S_n \mid R_t \supseteq R_j(\xi)\},$$

so that the cardinal to be majorized is

$$\begin{aligned} &\prod_{j=1}^k \text{card}\{t \in S_n \mid R_t \supseteq R_j(\xi)\} \\ &= \prod_{j=1}^k (n - (\text{card } R_j(\xi)))! \quad (\text{by 4.3b}) \\ &= (n!)^k n^{-\sum_{j=1}^k \text{card } R_j(\xi)} \\ &\quad \cdot \prod_{j=1}^k \left(\frac{n}{n} \cdot \frac{n}{n-1} \cdots \frac{n}{n - (\text{card } R_j(\xi)) + 1} \right). \end{aligned}$$

Now, from 4.3c) it is clear that $\sum_{j=1}^k \text{card } R_j(\xi) \leq \sum_{q=1}^m \text{card } I_q = \alpha$. This implies that $\frac{n}{n}, \frac{n}{n-1}, \dots, \frac{n}{n - (\text{card } R_j(\xi)) + 1}$ are not greater than $\frac{n}{n-\alpha+1} < 2$ (we assumed in 4.1 that n is not too small); majorizing all these factors with 2 in the last expression, we get (10). \square

4.6. *A reduction of the problem.* Applying Lemma 4.4, changing the order of summation and applying after that Lemma 4.5 we get

$$\begin{aligned} & \frac{1}{n(n!)^k} \sum_{t_1, \dots, t_k \in S_n} \text{Tr}(t_{c_1}^{\alpha_1} \dots t_{c_m}^{\alpha_m}) \\ &= \frac{1}{n(n!)^k} \sum_{t_1, \dots, t_k \in S_n} \text{card}\{\xi \text{ cycle} \mid \xi \prec (t_1, \dots, t_k)\} \\ &= \frac{1}{n(n!)^k} \sum_{\xi \text{ cycle}} \text{card}\{(t_1, \dots, t_k) \in S_n^k \mid \xi \prec (t_1, \dots, t_k)\} \\ &\leq \frac{1}{n(n!)^k} \sum_{\xi \text{ injective cycle}} (n!)^k (n/2)^{-\sum_{j=1}^k \text{card } R_j(\xi)}. \end{aligned}$$

The sum $\sum_{j=1}^k \text{card } R_j(\xi)$ takes values not greater than α , as remarked in the proof of 4.5, and not less than k (since obviously every $R_j(\xi)$ is non-void). Hence the last expression equals:

$$\frac{1}{n} \sum_{l=k}^{\alpha} (2/n)^l \text{card} \left\{ \xi \text{ injective cycle} \mid \sum_{j=1}^k \text{card } R_j(\xi) = l \right\},$$

and a simple computation shows that (8) will follow if we can prove that:

$$(11) \quad \text{card} \left\{ \xi \mid \xi \text{ injective cycle}, \sum_{j=1}^k \text{card } R_j(\xi) = l \right\} \leq \binom{\alpha}{l} \alpha^{\alpha-l} n^l,$$

for every $k \leq l \leq \alpha$.

4.7. *The tree of injective cycles.* In order to estimate the cardinals of sets of injective cycles needed in (11), it is convenient to have all the injective cycles placed together in a rooted tree.

Let T be a rooted tree, let V be its set of vertices, and let v_0 be its root. For any v in V , the length of the (unique) path connecting v and v_0 will be called the level of v , and denoted by $L(v)$. The vertices of level $L(v) + 1$ which can be connected with v by a path of length 1 will be called the successors of v ; their number, denoted by $D(v)$, will be called the degree of v .

The rooted tree T will be called $(1, n)$ -regular if

- (i) it has a maximal value of the levels, $L_{\max} \geq 2$;
- (ii) any vertex v with $L(v) \neq L_{\max}$ has $D(v) = 1$ or $n - \alpha \leq D(v) \leq n$ (of course, for $L(v) = L_{\max}$ we have $D(v) = 0$).

Clearly, if T is a $(1, n)$ -regular rooted tree, then for any vertex v of T , the subtree T_v of T generated by v is also $(1, n)$ -regular.

By a labeling of the $(1, n)$ -regular tree T we shall understand a function $e: V \setminus \{v_0\} \rightarrow \{1, \dots, n\}$, having the property that whenever $v \in V$ has $n - \alpha \leq D(v) \leq n$, the restriction of e to the set of the successors of v is one-to-one.

PROPOSITION. *One can construct a rooted $(1, n)$ -regular tree T , with $L_{\max} = \alpha + 1$, and a labeling $e: V \setminus \{v_0\} \rightarrow \{1, \dots, n\}$ such that*

1°. *For any $2 \leq \beta \leq \alpha + 1$, there is a canonical bijection from $\{v \in V \mid L(v) = \beta\}$ onto the injective sequences of β numbers in $\{1, \dots, n\}$ (see 4.3c), given by the following rule: for any v in V with $L(v) = \beta$, we take the unique path $v_0, v_1, \dots, v_\beta = v$ connecting v to the root, and we associate to it the sequence $(e(v_1), \dots, e(v_\beta))$.*

2°. *For any v in V with $n - \alpha \leq D(v) \leq n$ and $L(v) \leq \alpha - 1$, at most α of the successors of v have degree 1.*

3°. *Let ξ be an injective cycle, and let v be the unique vertex with $L(v) = \alpha + 1$ associated to ξ at 1°. Then, denoting the path between v_0 and v by $v_0, v_1, \dots, v_{\alpha+1} = v$, we have:*

$$\text{card}\{1 \leq \beta \leq \alpha \mid D(v_\beta) = 1\} = \alpha - \sum_{j=1}^k \text{card } R_j(\xi).$$

Proof. We shall construct the levels of the tree inductively, and define the labeling at the same time, taking care that 1° holds.

The level 0 contains only one vertex, the root, which is not labeled. The level 1 contains n vertices, labeled from 1 to n . The level 2 contains n^2 vertices, and more precisely, every vertex of the level 1 has n successors, labeled from 1 to n . It is clear that the rule described at 1° gives a bijection between the vertices of the level 2 and the sequences of two numbers in $\{1, \dots, n\}$ (which are all injective). If $\alpha = 1$, then this is T , and 1°, 2°, 3° are easily checked. For the rest of the proof, we shall suppose that $\alpha \geq 2$.

Now, let us assume that for some $2 \leq \beta \leq \alpha$ we have constructed the tree and the labeling up to the level β , such that 1° is satisfied. For constructing the level $\beta + 1$, what we have to do is provide an algorithm which decides, for a given vertex v with $L(v) = \beta$, what $D(v)$ should be, and which indicates the labels of the successors of v . Let q ($1 \leq q \leq m$) be such that $\beta \in I_q$, and consider $j = c_q \in \{1, \dots, k\}$, $\alpha_q \in \mathbb{Z} \setminus \{0\}$. If $\alpha_q > 0$, the algorithm sounds like this:

“Take v with $L(v) = \beta$, consider the path $v_0, v_1, \dots, v_\beta = v$ connecting v to the root, consider the sequence $\eta = (e(v_1), \dots, e(v_\beta))$ and the relation $R_j(\eta)$ (defined at 4.3c). If there is $u \in \{1, \dots, n\}$ such that $(u, e(v_\beta)) \in R_j(\eta)$, then put $D(v) = 1$, and the label of the unique successor of v is u . Otherwise, $n - \alpha \leq D(v) \leq n$, and the labels of the successors of v are $\{1, \dots, n\} \setminus \pi_1(R_j(\eta))$.” If $\alpha_q < 0$, the algorithm is the same, but we replace “ $(u, e(v_\beta))$ ” with “ $(e(v_\beta), u)$ ”.

It is easy to check that the level $\beta + 1$ constructed in this manner has the property of 1° . Hence the construction can be reiterated up to $\beta = \alpha$, giving us a labeled $(1, n)$ -regular rooted tree T , with maximal value of the levels $\alpha + 1$, and satisfying 1° .

To prove 2° , we need the following

LEMMA. *Let v' be in V such that $L(v') = \beta' \leq \alpha$ and $D(v') = 1$. Let $v_0, v_1, \dots, v_{\beta'} = v'$ be the path connecting v' with the root. Then there exists $1 \leq \gamma \leq \beta' - 1$ such that $e(v') = e(v_\gamma)$.*

Proof of the Lemma. We denote by q, q' , respectively, the numbers in $\{1, \dots, m\}$ such that $\beta' - 1 \in I_q, \beta' \in I_{q'}$ (note that $L(v') = \beta', D(v') = 1$ imply $\beta' \geq 3$, so that q makes sense; clearly, $q = q'$ or $q = q' - 1$).

Let us assume that $e(v') \neq e(v_\gamma)$ for every $1 \leq \gamma \leq \beta' - 1$, and obtain a contradiction. To make a choice, suppose that $\alpha_{q'} > 0$. From $D(v') = 1$ and the way we constructed the tree, we infer that there exists $1 \leq \gamma \leq \beta' - 1$, belonging to an interval I_p , such that $c_p = c_{q'}$ and:

$$\begin{cases} e(v_\gamma) = e(v'), & \text{if } \alpha_p > 0, \\ e(v_{\gamma+1}) = e(v'), & \text{if } \alpha_p < 0. \end{cases}$$

Because of the assumptions we made, the only possibility is that $\alpha_p < 0$ and $\gamma = \beta' - 1$ (hence $p = q$). This gives $c_q = c_{q'}$ and $\alpha_q < 0$. Further, $\alpha_q < 0 < \alpha_{q'}$ implies $q \neq q'$, and hence $q = q' - 1$; so we get $c_q = c_{q-1}$, a contradiction. If $\alpha_{q'} < 0$, we proceed in the same manner.

The proof of 2° is now immediate. Take v in V with $L(v) = \beta \leq \alpha - 1, n - \alpha \leq D(v) \leq n$, and let $v_0, v_1, \dots, v_\beta = v$ be the unique path connecting v with the root. If a successor v' of v has $D(v') = 1$, then, by the lemma, $e(v') \in \{e(v_1), \dots, e(v_\beta)\}$ which has at most $\alpha - 1$ elements. Since the labeling is one-to-one on the set of successors of v , we obtain 2° .

Finally, let ξ , v and $v_0, v_1, \dots, v_{\alpha+1} = v$ be as in 3°. From the construction of the tree it is clear that, for any $2 \leq \beta \leq \alpha$:

$$\sum_{j=1}^k \text{card } R_j(e(v_1), \dots, e(v_\beta), e(v_{\beta+1})) - \sum_{j=1}^k \text{card } R_j(e(v_1), \dots, e(v_\beta)) = \begin{cases} 0, & \text{if } D(v_\beta) = 1; \\ 1, & \text{if } n - \alpha \leq D(v_\beta) \leq n. \end{cases}$$

Hence the sum:

$$\text{card}\{1 \leq \gamma \leq \beta \mid D(v_\gamma) = 1\} + \sum_{j=1}^k \text{card } R_j(e(v_1), \dots, e(v_\beta), e(v_{\beta+1})),$$

considered for $1 \leq \beta \leq \alpha$, increases with 1 when β increases with 1. Since for $\beta = 1$:

$$\text{card}\{1 \leq \gamma \leq 1 \mid D(v_\gamma) = 1\} + \sum_{j=1}^k \text{card } R_j(e(v_1), e(v_2)) = 0 + 1 = 1,$$

by putting $\beta = \alpha$ we get assertion 3°. \square

4.8. Remarks on $(1, n)$ -regular rooted trees.

4.8.1. LEMMA. Let T be a rooted tree with maximal value of the levels $L_{\max} = \beta + 1$, and such that every vertex v with $L(v) \leq \beta$ has $D(v) \leq n$. Then, for any $0 \leq \gamma \leq \beta + 1$, T has at most n^γ vertices of level γ .

The proof of 4.8.1 is clear, by induction on γ .

4.8.2. LEMMA. Let T be a $(1, n)$ -regular rooted tree, with maximal value of the levels $\beta + 1$, and with the following property: for any vertex v with $L(v) \leq \beta - 1$ and $n - \alpha \leq D(v) \leq n$, at most α of the successors of v have degree 1. Then for any $1 \leq \gamma \leq \beta$, there are no more than $\alpha n^{\gamma-1}$ vertices of degree 1 on the level γ .

Proof. The case $\gamma = 1$ is clear. If $\gamma \geq 2$, denote by $N_\delta^{(i)}$ the number of vertices of degree i on the level δ , $i \in \{1, \dots, n\}$, $\delta \in \{\gamma - 1, \gamma\}$. We have

$$N_\gamma^{(1)} \leq \alpha \sum_{i=n-\alpha}^n N_{\gamma-1}^{(i)} + N_{\gamma-1}^{(1)} \leq \alpha \sum_{i=1}^n N_{\gamma-1}^{(i)} \stackrel{4.8.1}{\leq} \alpha n^{\gamma-1}. \quad \square$$

4.8.3. LEMMA. *Let T be a $(1, n)$ -regular rooted tree with the property stated at 4.8.2. Fix $1 \leq \beta_1 \leq \dots \leq \beta_h \leq \beta$, and let N be the number of vertices with $L(v) = \beta$ and with the following property: if $v_0, v_1, \dots, v_\beta = v$ is the path connecting v with the root, then $D(v_{\beta_1}) = \dots = D(v_{\beta_h}) = 1$. Then $N \leq \alpha^h n^{\beta-h}$.*

Proof. We first take the case $h = 1$. Let $V_{\beta_1}^{(1)}$ be the set of vertices of degree 1 on the level β_1 , and for any v in $V_{\beta_1}^{(1)}$, let T_v be the subtree of T generated by v . Clearly, $N = \sum_{v \in V_{\beta_1}^{(1)}} N_v$, with N_v the number of vertices of T_v having level $\beta - \beta_1$ (in T_v). 4.8.1 gives us that $N_v \leq n^{\beta-\beta_1}$ ($v \in V_{\beta_1}^{(1)}$), and 4.8.2 that $\text{card } V_{\beta_1}^{(1)} \leq \alpha n^{\beta_1-1}$; hence $N \leq \alpha n^{\beta-1}$.

We now make induction on h . Assume the lemma proved for all the possible choices $1 \leq \beta_1 < \dots < \beta_h \leq \beta$ (β natural) and let us prove it for a system $1 \leq \beta_1 < \dots < \beta_h < \beta_{h+1} \leq \beta$. Defining $V_{\beta_1}^{(1)}$ as in the preceding paragraph, we have again the formula $N = \sum_{v \in V_{\beta_1}^{(1)}} N_v$, where this time N_v is the number to be majorized with respect to the tree T_v and the system $1 \leq \beta_2 - \beta_1 < \dots < \beta_{h+1} - \beta_1 \leq \beta - \beta_1$. Hence, by 4.8.2 and the induction hypothesis:

$$N \leq (\alpha n^{\beta_1-1})(\alpha^h n^{\beta-\beta_1-h}) = \alpha^{h+1} n^{\beta-(h+1)}. \quad \square$$

4.8.4. PROPOSITION. *Let T be a $(1, n)$ -regular rooted tree, with the property stated at 4.8.2, and with the maximal value of the levels $\alpha + 1$. Let $0 \leq h \leq \alpha$ be a fixed integer. For any vertex v with $L(v) = \alpha$, we consider the path $v_0, v_1, \dots, v_\alpha = v$ connecting v to the root. Then*

$$\text{card}\{v | L(v) = \alpha, \text{card}\{1 \leq \beta \leq \alpha | D(v_\beta) = 1\} \geq h\} \leq \binom{\alpha}{h} \alpha^h n^{\alpha-h}.$$

Proof. For $h = 0$ we have to prove that the number of vertices of level α is not greater than n^α (which is in 4.8.1). If $h \geq 1$, then for any v having $\text{card}\{1 \leq \beta \leq \alpha | D(v_\beta) = 1\} \geq h$ we choose a system $1 \leq \beta_1 < \dots < \beta_h \leq \alpha$ such that $D(v_{\beta_1}) = \dots = D(v_{\beta_h}) = 1$; after that, we sum after all the possible choices of $1 \leq \beta_1 < \dots < \beta_h \leq \alpha$, and apply 4.8.3. \square

4.9. End of the proof of Theorem 4.1. We were left to prove (11) of 4.6. Let us fix l ($k \leq l \leq \alpha$), and denote $\alpha - l$ by h . We consider

the tree T of the injective sequences, constructed at 4.7. To every injective cycle ξ having $\sum_{j=1}^k \text{card } R_j(\xi) = l$ we associate the unique vertex v of this tree such that: $L(v) = \alpha$, and $(e(v_1), \dots, e(v_\alpha))$ are the first α components of ξ (as usual, $v_0, v_1, \dots, v_\alpha = v$ is the path connecting v to the root). Taking into account point 3° of Proposition 4.7, we see that $\xi \rightarrow v$ is a one-to-one mapping into the set of vertices $\{v | L(v) = \alpha, \text{card}\{1 \leq \beta \leq \alpha | D(v_\beta) = 1\} = h\}$. Hence:

$$\begin{aligned} & \text{card} \left\{ \xi \text{ injective cycle} \mid \sum_{j=1}^k \text{card } R_j(\xi) = l \right\} \\ & \leq \text{card}\{v | L(v) = \alpha, \text{card}\{1 \leq \beta \leq \alpha | D(v_\beta) = 1\} = h\} \\ & \leq \text{card}\{v | L(v) = \alpha, \text{card}\{1 \leq \beta \leq \alpha | D(v_\beta) = 1\} \geq h\} \\ & \leq \binom{\alpha}{h} \alpha^h n^{\alpha-h} = \binom{\alpha}{l} \alpha^{\alpha-l} n^l. \quad \square \end{aligned}$$

5. Asymptotic freeness in the case of Weyl groups. For any positive integer n , let A_n be a closed subgroup of the circle (i.e., $A_n = \mathbb{T}$ or $A_n = \mathbb{Z}/r_n\mathbb{Z}$ for some r_n), and let G_n be the semidirect product $A_n^n \rtimes_{\alpha_n} S_n$, where the action α_n of S_n on A_n^n is

$$\alpha_n(t)(z_1, \dots, z_n) = (z_{t^{-1}(1)}, \dots, z_{t^{-1}(n)}).$$

We can view G_n as a subgroup of $U(n)$, by identifying $g = ((z_1, \dots, z_n), t) \in G_n$ with the matrix having the (i, j) entry equal to z_i , if $t(j) = i$, and to 0, if $t(j) \neq i$. We have

THEOREM. *The assertions (i) and (ii) of 3.1 are true for the series $(G_n)_{n=1}^\infty$.*

To see this, we only need to take $\pi_n: G_n \rightarrow S_n$ the projection, and make the obvious remark that the inequality (12) appearing in the next lemma is valid:

LEMMA. *For $n \geq 1$, let G_n and H_n be closed subgroups of $U(n)$, and let π_n be a continuous homomorphism of G_n onto H_n , with the property that*

$$(12) \quad \text{Tr } \pi_n(g) \geq |\text{Tr } g|, \quad \text{for any } g \text{ in } G_n.$$

In this situation, if (i) and (ii) of 3.1 are valid for the series $(H_n)_{n=1}^\infty$, then they are also valid for the series $(G_n)_{n=1}^\infty$.

Proof. Consider, for every $n \geq 1$, a standard-independent family $(f_{n,\omega})_{\omega \in \Omega}$ of random unitaries in G_n .

Note first that $(\pi_n \circ f_{n,\omega})_{\omega \in \Omega}$ is standard-independent in H_n . Indeed, for any $\omega_1, \dots, \omega_m$ in Ω such that $\omega_i \neq \omega_j$ when $i \neq j$, and for any $\varphi_1, \dots, \varphi_m$ in $C(H_n)$:

$$\begin{aligned} & \int_X \varphi_1((\pi_n \circ f_{n,\omega_1})(x)) \cdots \varphi_m((\pi_n \circ f_{n,\omega_m})(x)) dP(x) \\ &= \int_X (\varphi_1 \circ \pi_n)(f_{n,\omega_1}(x)) \cdots (\varphi_m \circ \pi_n)(f_{n,\omega_m}(x)) dP(x) \\ &\stackrel{(2) \text{ of 2.7}}{=} \prod_{q=1}^m \int_{G_n} (\varphi_q \circ \pi_n)(g) dg \\ &= \prod_{q=1}^m \int_{H_n} \varphi_q(h) dh. \end{aligned}$$

On the other hand, for any $\omega_1 \neq \omega_2 \neq \dots \neq \omega_m$ in Ω and $\alpha_1, \dots, \alpha_m$ in $\mathbb{Z} \setminus \{0\}$ we have

$$\begin{aligned} & |\tau_n(f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m})| \\ &\leq \frac{1}{n} \int_X |\text{Tr}((f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m})(x))| dP(x) \\ &\stackrel{(12)}{\leq} \frac{1}{n} \int_X \text{Tr}((\pi_n \circ (f_{n,\omega_1}^{\alpha_1} \cdots f_{n,\omega_m}^{\alpha_m}))(x)) dP(x) \\ &= \frac{1}{n} \int_X \text{Tr}(((\pi_n \circ f_{n,\omega_1})^{\alpha_1} \cdots (\pi_n \circ f_{n,\omega_m})^{\alpha_m})(x)) dP(x) \\ &= \tau_n((\pi_n \circ f_{n,\omega_1})^{\alpha_1} \cdots (\pi_n \circ f_{n,\omega_m})^{\alpha_m}). \end{aligned}$$

Taking into account the considerations of 3.2, the last inequalities clearly finish the proof. \square

REFERENCES

- [1] D. Voiculescu, *Limit laws for random matrices and free products*, Invent. Math., **104** (1991), 201–220.
- [2] ———, *Free non-commutative random variables, random matrices and the II_1 factors of free groups*, in Quantum Probability and Related Topics (L. Accardi, editor), 1991, pp. 473–487.

Received April 30, 1991.

UNIVERSITY OF CALIFORNIA
BERKELEY, CA 94720

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982) F. WOLF (1904–1989)

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@henri.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Julie Speckart, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1993 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 157 No. 2 February 1993

Strong integral summability and the Stone-Čech compactification of the half-line	201
JEFF CONNOR and MARY ANNE SWARDSON	
The endlich Baer splitting property	225
THEODORE GERARD FATICONI	
The formal group of the Jacobian of an algebraic curve	241
MARGARET N. FREIJE	
Concordances of metrics of positive scalar curvature	257
PAWEL GAJER	
Explicit construction of certain split extensions of number fields and constructing cyclic classfields	269
STANLEY JOSEPH GURAK	
Asymptotically free families of random unitaries in symmetric groups	295
ALEXANDRU MIHAI NICA	
On purifiable subgroups and the intersection problem	311
TAKASHI OKUYAMA	
On the incidence cycles of a curve: some geometric interpretations	325
LUCIANA RAMELLA	
On some explicit formulas in the theory of Weil representation	335
R. RANGA RAO	
An analytic family of uniformly bounded representations of a free product of discrete groups	373
JANUSZ WYSOCZAŃSKI	
Errata: "Dentability, trees, and Dunford-Pettis operators on L_1 "	389
MARIA GIRARDI and ZHIBAO HU	
Errata: "Poincaré cobordism exact sequences and characterisation"	395
HIMADRI KUMAR MUKERJEE	



0030-8730(1993)157:2;1-I