

*Pacific
Journal of
Mathematics*

**ERRATA: “DENTABILITY, TREES, AND DUNFORD-PETTIS
OPERATORS ON L_1 ”**

MARIA GIRARDI AND ZHIBAO HU

Volume 157 No. 2

February 1993

ERRATA
CORRECTION TO
DENTABILITY, TREES, AND
DUNFORD-PETTIS OPERATORS ON L_1

MARIA GIRARDI AND ZHIBAO HU

Volume 148 (1991), 59–79

A Banach space has the complete continuity property if all its bounded subsets are midpoint Bocce dentable. We show that a lemma used in the original proposed proof of this result is false; however, we give a proof to show that the result is indeed true.

1. Introduction. Throughout this paper, \mathfrak{X} denotes an arbitrary Banach space, \mathfrak{X}^* the dual space of \mathfrak{X} , $B(\mathfrak{X})$ the closed unit ball of \mathfrak{X} , and $S(\mathfrak{X})$ the unit sphere of \mathfrak{X} . The triple (Ω, Σ, μ) refers to the Lebesgue measure space on $[0, 1]$, Σ^+ to the sets in Σ with positive measure, and L_1 to $L_1(\Omega, \Sigma, \mu)$. The σ -field generated by a partition π of $[0, 1]$ is $\sigma(\pi)$. The conditional expectation of $f \in L_1$ given a σ -field \mathcal{B} is $E(f|\mathcal{B})$.

A Banach space \mathfrak{X} has the *complete continuity property* (CCP) if each bounded linear operator from L_1 into \mathfrak{X} is *Dunford-Pettis* (i.e. carries weakly convergent sequences onto norm convergent sequences). Since a representable operator is Dunford-Pettis, the CCP is a weakening of the Radon-Nikodým property (RNP). Recall that a Banach space has the RNP if and only if all its bounded subsets are dentable. A subset D of \mathfrak{X} is *dentable* if for each $\varepsilon > 0$ there is x in D such that $x \notin \overline{\text{co}}(\{y \in D: \|x - y\| \geq \varepsilon\})$. Midpoint Bocce dentability is a weakening of dentability. The subset D is *midpoint Bocce dentable* if for each $\varepsilon > 0$ there is a finite subset F of D such that for each x^* in $B(\mathfrak{X}^*)$ there is x in F satisfying:

$$\text{if } x = \frac{1}{2}z_1 + \frac{1}{2}z_2 \text{ with } z_i \in D \text{ then } |x^*(x - z_1)| \equiv |x^*(x - z_2)| < \varepsilon.$$

The following theorem is presented in [G1].

THEOREM 1. *\mathfrak{X} has the CCP if all bounded subsets of \mathfrak{X} are midpoint Bocce dentable.*

Our purpose in writing this note is to show that Lemma 2.9 in [G1] (which was used in [G1] to prove Theorem 1) is false and to provide a proof of the theorem. Lemma 2.9 asserts that if A is in Σ^+ and f in $L_\infty(\mu)$ is not constant a.e. on A , then there is an increasing sequence $\{\pi_n\}$ of positive finite measurable partitions of A such that $\sigma(\bigcup \pi_n) = \Sigma \cap A$ and for each n

$$\mu \left(\bigcup \left\{ E : E \in \pi_n \text{ and } \frac{\int_E f d\mu}{\mu(E)} \geq \frac{\int_A f d\mu}{\mu(A)} \right\} \right) = \frac{\mu(A)}{2}.$$

Example 2 shows that Lemma 2.9 is false.

EXAMPLE 2. Let $f = 3\chi_{[0, \frac{1}{4}]} - \chi_{[\frac{1}{4}, 1]}$. Then $\int_\Omega f d\mu = 0$. Suppose that $\{\pi_n\}$ is an increasing sequence of positive finite measurable partitions of $[0, 1]$ such that for each n

$$\mu \left(\bigcup \left\{ E : E \in \pi_n \text{ and } \frac{\int_E f d\mu}{\mu(E)} \geq 0 \right\} \right) = \frac{1}{2}.$$

Then $\sigma(\bigcup \pi_n) \neq \Sigma$.

Proof. Consider the martingale $\{f_n\}$ given by

$$f_n(\cdot) = E(f | \sigma(\pi_n)) = \sum_{E \in \pi_n} \frac{\int_E f d\mu}{\mu(E)} \chi_E(\cdot).$$

For each $n \in \mathbb{N}$ put

$$P_n = \bigcup \left\{ E : E \in \pi_n \text{ and } \int_E f d\mu \geq 0 \right\} \quad \text{and} \quad Q_n = P_n \cap \left(\frac{1}{4}, 1\right].$$

Since $\mu(P_n) = \frac{1}{2}$, we have that $\mu(Q_n) \geq \frac{1}{4}$. Thus

$$\begin{aligned} \int_\Omega |f_n - f| d\mu &\geq \int_{Q_n} |f_n - f| d\mu \geq \int_{Q_n} (f_n - -1) d\mu \\ &\geq \int_{Q_n} 1 d\mu = \mu(Q_n) \geq \frac{1}{4}. \end{aligned}$$

We know that such a martingale $E(f | \sigma(\pi_n))$ converges in L_1 norm to $E(f | \sigma(\bigcup \pi_n))$. But $E(f | \Sigma) = f$. Thus $\sigma(\bigcup \pi_n) \neq \Sigma$. \square

The error in the proof of Lemma 2.9 occurred in assuming that if A is in Σ^+ and $\{\pi_n\}$ is an increasing sequence of positive measurable partitions of A such that for each n and each E in π_n the $\mu(E) \leq \varepsilon_n$ with $\lim_n \varepsilon_n = 0$, then $\sigma(\bigcup \pi_n) = \Sigma \cap A$. This seemingly sound assertion is not true as shown by the following counterexample.

EXAMPLE 3. For $n \in \mathbb{N}$ and $1 \leq i \leq 2^n$, define

$$E_i^n = \left[\frac{i-1}{2^{n+1}}, \frac{i}{2^{n+1}} \right) \cup \left[\frac{1}{2} + \frac{i-1}{2^{n+1}}, \frac{1}{2} + \frac{i}{2^{n+1}} \right)$$

and

$$\pi_n = \{E_i^n : 1 \leq i \leq 2^n\}.$$

Clearly $\{\pi_n\}$ is an increasing sequence of positive measurable partitions of $[0, 1]$ such that $\mu(E) = 2^{-n}$ for each n and each $E \in \pi_n$. Let $f = \chi_{[0, \frac{1}{2}]}$. An easy computation shows that $E(f|\sigma(\pi_n)) = \frac{1}{2}\chi_{[0, 1]}$. We know that such a martingale $E(f|\sigma(\pi_n))$ converges in L_1 norm to $E(f|\sigma(\bigcup \pi_n))$. But $E(f|\Sigma) = f$. Thus $\sigma(\bigcup \pi_n) \neq \Sigma$. \square

2. **Proof of theorem.** Our proof of Theorem 1 uses the following observations. For f in L_1 and A in Σ , the average value and the Bocce oscillation of f on A respectively are

$$m_A(f) \equiv \frac{\int_A f d\mu}{\mu(A)}$$

and

$$\text{Bocce-osc } f|_A \equiv \frac{\int_A |f - m_A(f)| d\mu}{\mu(A)}$$

observing the convention that $0/0$ is 0.

LEMMA 4. Fix A in Σ and f in L_1 . There is a subset E of A with $2\mu(E) = \mu(A)$ and

$$\frac{1}{2} \text{ Bocce-osc } f|_A \leq |m_E(f) - m_A(f)|.$$

Furthermore, for each subset E of A with $2\mu(E) = \mu(A)$,

$$|m_E(f) - m_A(f)| \leq \text{Bocce-osc } f|_A.$$

Proof. Without loss of generality, $A = \Omega$ and $\int_{\Omega} f d\mu = 0$ and $\int_{\Omega} |f| d\mu = 1$. With this normalization, $\text{Bocce-osc } f|_A = 1$ and $|m_E(f) - m_A(f)| = |m_E(f)|$. Let $P = [f \geq 0]$ and $N = [f < 0]$.

The first claim now reads that $\frac{1}{2} \leq 2|\int_E f d\mu|$ for some subset E of measure one half. Wlog $\mu(P) \geq \frac{1}{2}$. Partition P into 2 sets, P_1 and P_2 , of equal measure such that $\int_{P_2} f d\mu \leq \int_{P_1} f d\mu$. Note that

$$\begin{aligned} 1 &= \int_{\Omega} |f| d\mu = \int_P f d\mu + \int_N -f d\mu \\ &= 2 \int_P f d\mu = 2 \left[\int_{P_1} f d\mu + \int_{P_2} f d\mu \right] \leq 4 \int_{P_1} f d\mu. \end{aligned}$$

Since $\mu(P_1) \leq \frac{1}{2} \leq \mu(P)$, we can find a set E such that $P_1 \subset E \subset P$ and $\mu(E) = \frac{1}{2}$. For such a set E

$$\frac{1}{4} \leq \int_{P_1} f d\mu \leq \int_E f d\mu,$$

as needed.

Normalized, the second claim reads that for each subset E of measure $\frac{1}{2}$

$$2 \left| \int_E f d\mu \right| \leq 1.$$

Fix a subset E of measure $\frac{1}{2}$. Wlog $\int_{E \cap N} -f d\mu \leq \int_{E \cap P} f d\mu$. So

$$\begin{aligned} \left| \int_E f d\mu \right| &= \left| \int_{E \cap P} f d\mu + \int_{E \cap N} f d\mu \right| \\ &\leq \left| \int_{E \cap P} f d\mu \right| \leq \int_P |f| d\mu = \frac{1}{2}, \end{aligned}$$

as needed. □

A subset K of L_1 satisfies the *Bocce criterion* if for each $\varepsilon > 0$ and B in Σ^+ there is a finite collection \mathcal{F} of subsets of B each with positive measure such that for each f in K there is an A in \mathcal{F} satisfying

$$(*) \quad \text{Bocce-osc } f|_A < \varepsilon.$$

Lemma 4 provides an equivalent formulation of the Bocce criterion; namely we can replace condition $(*)$ by the condition

$$(**) \quad \begin{aligned} &\text{if the subset } E \text{ of } A \text{ has half the measure of } A, \\ &\text{then } |m_E(f) - m_A(f)| < \varepsilon. \end{aligned}$$

We now attack the proof of Theorem 1. Our proof follows mainly the proof in [G1].

Proof of Theorem 1. Let all bounded subsets of \mathfrak{X} be midpoint Bocce dentable. Fix a bounded linear operator T from L_1 into \mathfrak{X} . It suffices to show that the subset $T^*(B(\mathfrak{X}^*))$ of L_1 satisfies the Bocce criterion (this is a necessary and sufficient condition for T to be Dunford-Pettis [G2]). To this end, fix $\varepsilon > 0$ and B in Σ^+ .

Consider the vector measure F from Σ into \mathfrak{X} given by $F(E) = T(\chi_E)$. For $x^* \in \mathfrak{X}^*$

$$m_E(T^*x^*) = \frac{x^*F(E)}{\mu(E)}$$

since $\int_E (T^*x^*) d\mu = x^*T(\chi_E) = x^*F(E)$.

Since the subset $\{\frac{F(E)}{\mu(E)} : E \subset B \text{ and } E \in \Sigma^+\}$ of \mathfrak{X} is bounded, it is midpoint Bocce dentable. Accordingly, there is a finite collection \mathcal{F} of subsets of B each in Σ^+ such that for each $x^* \in B(\mathfrak{X}^*)$ there is a set A in \mathcal{F} such that if

$$\frac{F(A)}{\mu(A)} = \frac{1}{2} \frac{F(E_1)}{\mu(E_1)} + \frac{1}{2} \frac{F(E_2)}{\mu(E_2)}$$

for some subsets E_i of B with $E_i \in \Sigma^+$, then

$$\left| \frac{x^*F(E_1)}{\mu(E_1)} - \frac{x^*F(A)}{\mu(A)} \right| \equiv \left| \frac{x^*F(E_2)}{\mu(E_2)} - \frac{x^*F(A)}{\mu(A)} \right| < \varepsilon.$$

Fix $x^* \in B(\mathfrak{X}^*)$ and find the associated A in \mathcal{F} .

At this point we turn to our new formulation of the Bocce criterion (whereas [G1] used the old formulation and Lemma 2.9).

This $A \in \mathcal{F}$ satisfies the condition (**). For consider a subset E of A with $\mu(E) = \frac{1}{2}\mu(A)$. Since

$$\frac{F(A)}{\mu(A)} = \frac{1}{2} \frac{F(E)}{\mu(E)} + \frac{1}{2} \frac{F(A \setminus E)}{\mu(A \setminus E)}$$

we have that

$$|m_E(T^*x^*) - m_A(T^*x^*)| \equiv \left| \frac{x^*F(E)}{\mu(E)} - \frac{x^*F(A)}{\mu(A)} \right| < \varepsilon.$$

Thus $T^*(B(\mathfrak{X}^*))$ satisfies the Bocce criterion, as needed. □

3. Closing comments. A relatively weakly compact subset of L_1 is relatively norm compact if and only if it satisfies the Bocce criterion [G2]. Thus our new formulation of the Bocce criterion provides another (perhaps at times more useful) method for testing for norm compactness in L_1 .

Fix A in Σ^+ and f in L_1 . Put

$$M_A(f) = \sup \{ |m_E(f) - m_A(f)| : E \subset A \text{ and } 2\mu(E) = \mu(A) \}.$$

This supremum is obtained. For just normalize so that $A = \Omega$ and $\int_\Omega f d\mu = 0$ and $\int_\Omega |f| d\mu = 1$. As Ralph Howard pointed out, next find disjoint subsets E_1 and E_2 of measure $\frac{1}{2}$ and $a \in \mathbb{R}$ such that

$$E_1 \subset [f \leq a] \text{ and } E_2 \subset [f \geq a].$$

Then $M_A(f)$ will be the larger of $|m_{E_1}(f)|$ and $|m_{E_2}(f)|$.

Basically, our Lemma 4 says that

$$\frac{1}{2} \text{Bocce-osc } f|_A \leq M_A(f) \leq \text{Bocce-osc } f|_A.$$

These bounds are the best possible.

For the second inequality, consider the function defined on $A \equiv [0, 1]$ by

$$f = \chi_{[0, \frac{1}{2})} - \chi_{[\frac{1}{2}, 1]}.$$

Straightforward calculations show that $m_{[0, \frac{1}{2}]}(f) = 1$ and that $\text{Bocce-osc } f|_A = 1$. Thus

$$M_A(f) = \text{Bocce-osc } f|_A.$$

As for the first inequality, consider the family of functions defined on $A \equiv [0, 1]$ by

$$f_\delta = \frac{\delta - 1}{\delta} \chi_{[0, \delta)} + \chi_{[\delta, 1]}$$

for $0 < \delta < \frac{1}{2}$. Straightforward calculations show that

$$M_A(f_\delta) = \frac{1}{2(1 - \delta)} \text{Bocce-osc } f_\delta|_A.$$

Actually $M_A(f) = \frac{1}{2} \text{Bocce-osc } f|_A$ if and only if f is the zero function on A .

REFERENCES

- [B] R.D. Bourgin, *Geometric Aspects of Convex Sets With the Radon-Nikodým Property*, Lecture Notes in Math., vol. 933, Springer-Verlag, Berlin and New York, 1983.
- [DU] J. Diestel and J. J. Uhl, Jr., *Vector Measures*, Math. Surveys, no. 15, Amer. Math. Soc., Providence, R.I., 1977.
- [G1] Maria Girardi, *Dentability, trees, and Dunford-Pettis operators on L_1* , Pacific J. Math., **148** (1991), 59–79.
- [G2] ———, *Compactness in L_1 , Dunford-Pettis operators, geometry of Banach spaces*, Proc. Amer. Math. Soc., **111** (1991), 767–777.

Received April 24, 1992.

UNIVERSITY OF SOUTH CAROLINA
COLUMBIA, SC 29208

AND

MIAMI UNIVERSITY
OXFORD, OH 45056

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982) F. WOLF (1904–1989)

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@henri.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Julie Speckart, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1993 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 157 No. 2 February 1993

Strong integral summability and the Stone-Čech compactification of the half-line	201
JEFF CONNOR and MARY ANNE SWARDSON	
The endlich Baer splitting property	225
THEODORE GERARD FATICONI	
The formal group of the Jacobian of an algebraic curve	241
MARGARET N. FREIJE	
Concordances of metrics of positive scalar curvature	257
PAWEL GAJER	
Explicit construction of certain split extensions of number fields and constructing cyclic classfields	269
STANLEY JOSEPH GURAK	
Asymptotically free families of random unitaries in symmetric groups	295
ALEXANDRU MIHAI NICA	
On purifiable subgroups and the intersection problem	311
TAKASHI OKUYAMA	
On the incidence cycles of a curve: some geometric interpretations	325
LUCIANA RAMELLA	
On some explicit formulas in the theory of Weil representation	335
R. RANGA RAO	
An analytic family of uniformly bounded representations of a free product of discrete groups	373
JANUSZ WYSOCZAŃSKI	
Errata: "Dentability, trees, and Dunford-Pettis operators on L_1 "	389
MARIA GIRARDI and ZHIBAO HU	
Errata: "Poincaré cobordism exact sequences and characterisation"	395
HIMADRI KUMAR MUKERJEE	



0030-8730(1993)157:2;1-I