

*Pacific
Journal of
Mathematics*

**L^n SOLUTIONS OF THE STATIONARY AND NONSTATIONARY
NAVIER-STOKES EQUATIONS IN R^n**

ZHI MIN CHEN

Volume 158 No. 2

April 1993

L^n SOLUTIONS OF THE STATIONARY AND NONSTATIONARY NAVIER-STOKES EQUATIONS IN R^n

ZHI-MIN CHEN

It is shown that the Navier-Stokes equations in the whole space R^n ($n \geq 3$) admit a unique small stationary solution which may be formed as a limit of a nonstationary solution as $t \rightarrow \infty$ in L^n -norms.

0. Introduction. As is well known, the existence of solutions to the exterior stationary Navier-Stokes equations was studied by Finn [2, 3], and small solutions from Finn [2, 3] may be formed as limits of nonstationary solutions as time $t \rightarrow \infty$ in local or global L^2 -norms (cf. Heywood [9, 10], Galdi and Rionero [6], Miyakawa and Sohr [16], Borchers and Miyakawa [1]) and in the norms of other function spaces (cf. Heywood [11], Musuda [14]). However, it is still unknown even in the case of whole spaces whether or not

$$(0.1) \quad \|v(t) - w\|_n + t^{1/2} \|Dv(t) - Dw\|_n + t^{1/2} \|v(t) - w\|_\infty \rightarrow 0$$

as $t \rightarrow \infty$,

provided that w and v are, respectively, the solutions to the stationary Navier-Stokes equations

$$(0.2) \quad -\Delta w + (w \cdot D)w + d\bar{p} = f, \quad D \cdot w = 0 \quad \text{in } R^n$$

and the nonstationary Navier-Stokes equations

$$(0.3) \quad v_t - \Delta v + (v \cdot D)v + D\bar{p} = f, \quad D \cdot v = 0 \quad \text{in } R^n \times (0, \infty),$$

$$v(0) = v_0 \quad \text{in } R^n.$$

Here and in what follows, $n \geq 3$ denotes the space dimension, \bar{p} and $\bar{\bar{p}}$ represent the pressures associated with w and v , respectively, D = the gradient, $f = f(x)$ is a prescribed function, the dot \cdot denotes the scalar product in R^n , and $\|\cdot\|_r$ denotes the norm of the Lebesgue space $L^r = L^r(R^n; R^n)$.

The purpose of the paper is to show that (0.2) and (0.3) admit small regular solutions w and $v(t)$ in L^n , respectively, such that (0.1) is valid. The problem above is, as usual, said to be a stability problem

for w , which has been studied by Kozono and Ozawa [13] in the case of bounded domains. From our view point, the global existence results of Kato [12] may be regarded as the stability theorems around the rest flow $w \equiv 0$.

In this paper we shall use the following spaces.

- C_0^∞ = the set of compactly supported solenoidal $u \in C^\infty(\mathbb{R}^n; \mathbb{R}^n)$,
- J^r = the completion of C_0^∞ in L^r for $1 < r < \infty$,
- $W^{k,r}$ = the Sobolev space $W^{k,r}(\mathbb{R}^n; \mathbb{R}^n)$ for $1 < r < \infty$ and $k = 1, 2$,
- $\widehat{W}^{1,r} = \{u \in L^{nr/(n-r)}; Du \in L^r(\mathbb{R}^n; \mathbb{R}^{n^2})\}$ for $1 < r < n$,
- $\widehat{W}^{2,r} = \{u \in W^{1, nr/(n-r)}; D^2u \in L^r(\mathbb{R}^n; \mathbb{R}^{n^3})\}$ for $1 < r < n/2$,

where $D^2 =$ the Hessian matrix $[D_i D_j]_{n \times n}$ with $D_k = \partial/\partial x_k$. Moreover, we denote by P the linear bounded projection from L^r onto J^r for $1 < r < \infty$ (cf. [15] for details), by A the Stokes operator $-PA$ associated with the domain $W^{2,r} \cap J^r$ for $1 < r < \infty$, by (\cdot, \cdot) the duality pairing between L^r and $(L^r)^*$ for $1 \leq r < \infty$, and we set

$$\|u\|_{-1,r} = \sup\{|(u, v)|; v \in C_0^\infty, \|Dv\|_{r/(r-1)} = 1\} \quad \text{for } 1 < r < \infty.$$

Our main results read as follows.

THEOREM 0.1. *For $n \geq 3$ there is a small $0 < d < 1$ such that (0.1) admits a unique solution*

$$w \in J^n \cap \widehat{W}^{1, 2n/3} \cap \widehat{W}^{1, 2n/5} \quad \text{with } \|Dw\|_{n/2} \leq d$$

satisfying

$$\begin{aligned} \|Dw\|_{n/2} + \|w\|_n &\leq C\|f\|_{-1, n/2}, \\ \|Dw\|_{2n/5} + \|Dw\|_{2n/3} + \|w\|_{2n} + \|w\|_{2n/3} \\ &\leq C(\|f\|_{-1, 2n/5} + \|f\|_{-1, 2n/3}) \end{aligned}$$

with C independent of f and w , provided that

$$f \in C_0^\infty \quad \text{and} \quad \|f\|_{-1, n/2} \leq d^2.$$

THEOREM 0.2. *Let $n \geq 3$, $f \in C_0^\infty$, $v_0 \in J^n$, and let $\|v_0\|_n$ and $\|f\|_{-1, 2n/5} + \|f\|_{-1, 2n/3}$ be sufficiently small. Then (0.3) admits a unique solution*

$$v \in BC([0, \infty); J^n) \quad \text{and} \quad t^{1/2}D(v(t) - w) \in BC([0, \infty); L^n(\mathbb{R}^n; \mathbb{R}^{n^2}))$$

such that (0.1) is valid, where w is the solution of (0.2) from Theorem 0.1 and BC denotes the class of bounded and continuous functions.

Since there is no boundary to worry about in the whole space, our proof largely depends on the fact that P commutes with D , and also based on the theory of analytic semigroups in various L^r spaces. Such an approach is developed from Fujita and Kato [5] and Kato [12].

In §1 we prove Theorem 0.1. In §2 we obtain resolvent estimates for the perturbed operator $Au + P(u \cdot D)w + P(w \cdot D)u$ and therefore deduce decay estimates for the analytic semigroups generated by the perturbed operator. Theorem 0.2 is proved in §3.

1. Proof of Theorem 0.1. From the Sobolev inequality

$$(1.1) \quad C^{-1} \|u\|_{nr/(n-2r)} \leq \|Du\|_{nr/(n-r)} \leq C \|D^2u\|_r \quad \text{for } 1 < r < n/2,$$

the Calderon-Zygmund inequality (cf. [7])

$$\|D^2u\|_r \leq C \|\Delta u\|_r \quad \text{for } 1 < r < \infty,$$

the density of $\{Au; u \in C_0^\infty\}$ in J^r for $1 < r < n/2$, and the fact that P commutes with Δ , it follows that the Stokes operator A can be extended to a bounded and invertible operator from $J^{nr/(n-2r)} \cap \widehat{W}^{2,r}$ onto J^r for $1 < r < n/2$. Consequently, we set the operator

$$T: J^n \cap \widehat{W}^{1,2n/5} \cap \widehat{W}^{1,2n/3} \rightarrow \widehat{W}^{2,r} \quad \text{for } n/3 < r < n/2$$

such that

$$Tw = T_f w = A^{-1}(f - P(w \cdot D)w).$$

It is easy to see that to seek solutions of (0.2) means to seek fixed points of T .

Let $2n/5 \leq r \leq 2n/3$, $w \in J^n \cap \widehat{W}^{1,2n/5} \cap \widehat{W}^{1,2n/3}$, $v \in C_0^\infty$. Then by the divergence condition $D \cdot w = 0$, we have

$$\begin{aligned} (DTw, Dv) &= (f, v) - ((w \cdot D)w, v) \\ &= (f, v) + (w, (w \cdot D)v) \\ &\leq (f, v) + \|w\|_n \|w\|_{nr/(n-r)} \|Dv\|_{r/(r-1)}. \end{aligned}$$

Combining this with the inequality (cf. [17, 18])

$$\|DTw\|_r \leq C \sup\{|(DTw, Dv)|; v \in C_0^\infty, \|Dv\|_{r/(r-1)} = 1\}$$

with $C = C(n)$, we have, by (1.1),

$$\|DTw\|_r \leq C(n)(\|f\|_{-1,r} + \|Dw\|_{n/2} \|Dw\|_r),$$

and, similarly, for $u, w \in J^n \cap \widehat{W}^{1,2n/5} \cap \widehat{W}^{1,2n/3}$

$$\|DTw - DTu\|_r \leq C(n)(\|Dw\|_{n/2} + \|Du\|_{n/2})\|Dw - Du\|_r.$$

Consequently, there is a small positive d such that T is a contraction mapping from the complete metric space

$$\{w \in J^n \cap \widehat{W}^{1,2n/5} \cap \widehat{W}^{1,2n/3}; \|Dw\|_{n/2} \leq d\}$$

into itself provided that $f \in C_0^\infty$ with $\|f\|_{-1,n/2} < d^2$. We thus obtain the desired assertion by making use of the contraction mapping principle and (1.1). The proof is complete.

2. $L^p - L^q$ estimates. In the remainder of the paper we denote by w the solution of (0.2) given in Theorem 0.1, and by C the various constants which are always independent of the quantities $u, v, w, f, a, t,$ and z . Moreover we set

$$\begin{aligned} S &= \{z \in \mathbb{C}; -3\pi/4 < \arg z < 3\pi/4\}, \\ Lu &= Au + Bu; \quad Bu = P(u \cdot D)w + P(w \cdot D)u, \\ L^*u &= Au + B^*u; \quad B^*u = -P(w \cdot D)u + \sum_{i=1}^n Pu^i Dw^i \end{aligned}$$

for $u = (u^1, \dots, u^n)$ and $w = (w^1, \dots, w^n)$.

In arriving at $L^p - L^q$ estimates, we begin with the resolvent estimates for L and L^* .

LEMMA 2.1. *Let $z \in S$ and $u \in C_0^\infty$. Then we have*

$$(2.1) \quad |z| \|(L + z)^{-1}u\|_r \leq C\|u\|_r \quad \text{for } 1 < r < \infty,$$

$$(2.2) \quad |z| \|(L^* + z)^{-1}u\|_r \leq C\|u\|_r \quad \text{for } 1 < r < \infty,$$

$$(2.3) \quad |z|^{1/2} \|D(L + z)^{-1}u\|_r \leq C\|u\|_r \quad \text{for } 1 < r < n,$$

$$(2.4) \quad |z|^{1/2} \|D(L^* + z)^{-1}u\|_r \leq C\|u\|_r \quad \text{for } 1 < r < \infty,$$

provided that $\|Dw\|_{n/2}$ is sufficiently small;

$$(2.5) \quad |z|^{3/4} \|(L + z)^{-1}u\|_\infty \leq C\|u\|_{2n},$$

$$(2.6) \quad |z|^{1/2} \|D(L + z)^{-1}u\|_n \leq C(\|u\|_n + |z|^{-1/4} \|u\|_{2n}),$$

provided that $\|w\|_{2n}^{1/2} \|w\|_{2n/3}^{1/2}$ is sufficiently small.

Proof. Let us recall the well-known resolvent estimates for the Stokes operator (cf. [15])

$$(2.7) \quad \begin{aligned} |z| \|(A + z)^{-1}u\|_r + |z|^{1/2} \|D(A + z)^{-1}u\|_r \\ + \|D^2(A + z)^{-1}u\|_r \leq C\|u\|_r \end{aligned}$$

for $z \in S$, $1 < r < \infty$ and $u \in J^r$, and the Gagliardo-Nirenberg inequality (cf. [4])

$$(2.8) \quad \|u\|_q \leq C \|u\|_r^{1-h} \|Du\|_p^h$$

for $1 < r, p \leq q \leq \infty, 0 \leq h < 1, -n/q = h(1 - n/p) - (1 - h)n/r, u \in C_0^\infty$. Let us suppose $z \in S$ and $u \in J^r \cap W^{1,r}$ for $1 < r < \infty$.

Step 1. We prove (2.1) and (2.2). From (2.7), (1.1), the Hölder inequality and the boundedness of P in L^r -spaces it follows that for $1 < r < n/2, p = nr/(n - r)$ and $q = nr/(n - 2r)$,

$$\begin{aligned} \|B(A + z)^{-1}u\|_r &\leq C \|w\|_n \|D(A + z)^{-1}u\|_p \\ &\quad + C \|Dw\|_{n/2} \|(A + z)^{-1}u\|_q \\ &\leq C \|Dw\|_{n/2} \|D^2(A + z)^{-1}u\|_r \\ &\leq C \|Dw\|_{n/2} \|u\|_r \\ &\leq (1/2) \|u\|_r, \quad \text{by setting } C \|Dw\|_{n/2} < 1/2. \end{aligned}$$

This is together with (2.7) and the identity

$$L + z = (1 + B(A + z)^{-1})(A + z)$$

implies

$$|z| \|(L + z)^{-1}u\|_r \leq C \|u\|_r \quad \text{for } 1 < r < n/2.$$

Similarly, we have

$$|z| \|(L^* + z)^{-1}u\|_r \leq C \|u\|_r \quad \text{for } 1 < r < n/2.$$

This yields for $n < r < \infty, v \in L^{r'}$ with $r' = r/(r - 1)$,

$$((L + z)^{-1}u, v) = (u, (L^* + z)^{-1}Pv) \leq C |z|^{-1} \|u\|_r \|v\|_{r'}$$

and hence the validity of (2.1) with $n < r < \infty$. Thus (2.1) with $n/2 \leq r \leq n$ follows immediately from the Marcinkiewicz interpolation theorem (cf. [7]). (2.2) is verified in the same way.

Step 2. We prove (2.3). Observing that $1 < r < n$ and applying the condition $D \cdot u = D \cdot w = 0$ and the fact that D commutes with P yields

$$(2.9) \quad (A + z)^{-1}Bu = \sum_{i=1}^n D_i (A + z)^{-1}P(u^i w + w^i u),$$

we have, by (2.7) and (1.1),

$$\begin{aligned} \|(A + z)^{-1}Bu\|_r &\leq C |z|^{-1/2} \|w\|_n \|u\|_{nr/(n-r)} \\ &\leq C \|Dw\|_{n/2} \|Du\|_r |z|^{-1/2} \\ &\leq 2^{-1} |z|^{-1/2} \|Du\|_r, \quad \text{by setting } C \|Dw\|_{n/2} < 1/2 \end{aligned}$$

and

$$(2.10) \quad \begin{aligned} \|D(A+z)^{-1}Bu\|_r &\leq C\|w\|_n\|u\|_{nr/(n-r)} \\ &\leq C\|Dw\|_{n/2}\|Du\|_r \\ &\leq (1/2)\|Du\|_r, \quad \text{by setting } C\|Dw\|_{n/2} < 1/2. \end{aligned}$$

Consequently, we have

$$(2.11) \quad \|((A+z)^{-1}B)^k u\|_r \leq 2^{-k}|z|^{-1/2}\|Du\|_r, \quad \text{for integer } k > 0,$$

and so

$$D(L+z)^{-1}u = \sum_{k=0}^{\infty} D((A+z)^{-1}B)^k (A+z)^{-1}u \quad \text{in } L^r.$$

Applying (2.10) to the preceding identity repeatedly and using (2.7), we have

$$\|D(L+z)^{-1}u\|_r \leq 2\|D(A+z)^{-1}u\|_r \leq C|z|^{-1/2}\|u\|_r$$

as required.

Step 3. We prove (2.4). Observing that

$$\begin{aligned} (D_i(L^*+z)^{-1}u, v) &= -(u, (L+z)^{-1}D_iPv) \\ &\leq \|u\|_r\|(L+z)^{-1}D_iPv\|_{r'} \end{aligned}$$

for $i = 1, \dots, n$, $1 < r < \infty$, $r' = r/(r-1)$ and $v \in W^{1,r'}$, we need only to show the estimate

$$(2.12) \quad \|(L+z)^{-1}Du\|_r \leq C|z|^{-1/2}\|u\|_r, \quad \text{for } 1 < r < \infty.$$

Indeed, taking (2.9), (1.1) and (2.7) into account, we have for $n \leq r < \infty$,

$$\begin{aligned} \|(A+z)^{-1}Bu\|_r &\leq C \sum_{i=1}^n \|DD_i(A+z)^{-1}P(uw^i + wu^i)\|_{nr/(n+r)} \\ &\leq C\|Dw\|_{n/2}\|u\|_r \leq (1/2)\|u\|_r, \end{aligned}$$

by setting $C\|Dw\|_{n/2} < 1/2$, and hence for $n \leq r < \infty$,

$$\begin{aligned} \|(L+z)^{-1}Du\|_r &= \|(1 + (A+z)^{-1}B)^{-1}D(A+z)^{-1}u\|_r \\ &\leq 2\|D(A+z)^{-1}u\|_r \\ &\leq C|z|^{-1/2}\|u\|_r \end{aligned}$$

which arrives at (2.12) for $n \leq r < \infty$. Moreover (2.12) with $1 < r < n$ is verified as follows:

$$\begin{aligned} \|(L+z)^{-1}Du\|_r &= \|(1+(A+z)^{-1}B)^{-1}D(A+z)^{-1}u\|_r \\ &\leq \|D(A+z)^{-1}u\|_r + |z|^{-1/2}\|D^2(A+z)^{-1}u\|_r \\ &\leq C|z|^{-1/2}\|u\|_r, \end{aligned}$$

where we have used (2.11) and (2.7).

Step 4. We prove (2.5). By (2.8) and (2.7), we obtain

$$(2.13) \quad \|(A+z)^{-1}u\|_\infty \leq C\|(A+z)^{-1}u\|_{2n}^{1/2}\|D(A+z)^{-1}u\|_{2n}^{1/2} \\ \leq C|z|^{-3/4}\|u\|_{2n},$$

and, by (2.7), (2.8), (1.1) and (2.9),

$$(2.14) \quad \|(A+z)^{-1}Bu\|_\infty \leq C\|(A+z)^{-1}Bu\|_{2n}^{1/2}\|D(A+z)^{-1}Bu\|_{2n}^{1/2} \\ \leq C\|D(A+z)^{-1}Bu\|_{2n/3}^{1/2}\|D(A+z)^{-1}Bu\|_{2n}^{1/2} \\ \leq C\sum_{i=1}^n\|u^iw+w^iu\|_{2n/3}^{1/2}\|u^iw+w^iu\|_{2n}^{1/2} \\ \leq C\|w\|_{2n/3}^{1/2}\|w\|_{2n}^{1/2}\|u\|_\infty \\ \leq (1/2)\|u\|_\infty, \quad \text{by setting } C\|w\|_{2n/3}^{1/2}\|w\|_{2n}^{1/2} < 1/2.$$

We thus obtain

$$\begin{aligned} \|(L+z)^{-1}u\|_\infty &= \|(1+(A+z)^{-1}B)^{-1}(A+z)^{-1}u\|_\infty \\ &\leq 2\|(A+z)^{-1}u\|_\infty \leq C|z|^{-3/4}\|u\|_{2n} \end{aligned}$$

and hence the validity of (2.5).

Step 5. We prove (2.6). By (1.1), (2.9) and (2.7),

$$\begin{aligned} \|(A+z)^{-1}Bu\|_n &\leq C\|D(A+z)^{-1}Bu\|_{n/2} \\ &\leq C\|w\|_{2n}^{1/2}\|w\|_{2n/3}^{1/2}\|u\|_n \\ &\leq (1/2)\|u\|_n, \quad \text{by setting } C\|w\|_{2n}^{1/2}\|w\|_{2n/3}^{1/2} < 1/2 \end{aligned}$$

and, by (2.9), (2.7) and (1.1),

$$\begin{aligned} \|D(A+z)^{-1}Bu\|_n &\leq C\|w\|_n\|u\|_\infty \\ &\leq C\|w\|_{2n/3}^{1/2}\|w\|_{2n}^{1/2}\|u\|_\infty \\ &\leq (1/2)\|u\|_\infty, \quad \text{by setting } C\|w\|_{2n/3}^{1/2}\|w\|_{2n}^{1/2} < 1/2. \end{aligned}$$

Hence, it is easy to see that

$$\begin{aligned}
\|D(L+z)^{-1}u\|_n &\leq \sum_{k=0}^{\infty} \|D((A+z)^{-1}B)^k(A+z)^{-1}u\|_n \\
&\leq \|D(A+z)^{-1}u\|_n + \sum_{k=0}^{\infty} \|((A+z)^{-1}B)^k(A+z)^{-1}u\|_{\infty} \\
&\leq \|D(A+z)^{-1}u\|_n + \|(A+z)^{-1}u\|_{\infty}, \quad \text{by (2.14),} \\
&\leq C(|z|^{1/2}\|u\|_n + |z|^{-3/4}\|u\|_{2n}), \quad \text{by (2.7), (2.13).}
\end{aligned}$$

The proof is complete.

As an immediate consequence of (2.1) and (2.2), we conclude that L and L^* generate strongly continuous analytic semigroups e^{-tL} and e^{-tL^*} in J^r with $1 < r < \infty$, respectively, provided $\|Dw\|_{n/2}$ is sufficiently small. What is more, we can now proceed to the proof of the following $L^p - L^r$ estimates.

THEOREM 2.1. *Let $t > 0$, $1 < q \leq n$, $v \in J^q$ and $u \in C_0^{\infty}$. Then we have*

$$(2.15) \quad \|e^{-tL}u\|_p \leq Ct^{-(n/r-n/p)/2}\|u\|_r \quad \text{for } 1 < r \leq p < \infty,$$

provided that $\|Dw\|_{n/2}$ is sufficiently small;

$$(2.16) \quad \|e^{-tL}u\|_{\infty} + \|De^{-tL}u\|_n \leq Ct^{-n/2r}\|u\|_r \quad \text{for } 1 < r \leq n,$$

$$(2.17) \quad t^{n/2q}(t^{-1/2}\|e^{-tL}v\|_n + \|e^{-tL}v\|_{\infty} + \|De^{-tL}v\|_n) \rightarrow 0$$

as $t \rightarrow \infty$,

provided that $\|w\|_{2n}^{1/2}\|w\|_{2n/3}^{1/2}$ is sufficiently small.

Proof. By making use of the semigroup property of e^{-tL} , Lemma 2.1, and the Dunford integral (cf. [8]) via a standard calculation, we have

$$(2.18) \quad \|e^{-tL^*}u\|_r + t^{1/2}\|De^{-tL^*}u\|_r \leq C\|u\|_r \quad \text{for } 1 < r < \infty,$$

$$(2.19) \quad \|e^{-tL}u\|_{\infty} + \|De^{-tL}u\|_n \\ \leq Ct^{-1/2}\|e^{-(t/2)L}u\|_n + Ct^{-1/4}\|e^{-(t/2)L}u\|_{2n}$$

under the assumptions of Theorem 2.1.

It follows from (2.8) and (2.18) that

$$\begin{aligned} \|e^{-tL^*} u\|_p &\leq C \|e^{-tL^*} u\|_r^{1-n/r+n/p} \|De^{-tL^*} u\|_r^{n/r-n/p} \\ &\leq C t^{-(n/r-n/p)/2} \|u\|_r \end{aligned}$$

for $1 < r \leq p < \infty$ and $n/r - n/p < 1$. Combining this with the semigroup property of e^{-tL^*} , we have for $1 < r \leq p < \infty$ and $a \in L^{p/(p-1)}$,

$$\|e^{-tL^*} Pa\|_{r/(r-1)} \leq C t^{-(n/r-n/p)/2} \|a\|_{p/(p-1)},$$

and hence

$$(e^{-tL} u, a) = (u, e^{-tL^*} Pa) \leq C t^{-(n/r-n/p)} \|u\|_r \|a\|_{p/(p-1)}$$

This gives (2.15). (2.16) follows from (2.19) and (2.15).

To prove (2.17), we note for $a \in J^q \cap J^r$ with $1 < r < q$,

$$\begin{aligned} t^{n/2q} (t^{-1/2} \|e^{-tL} v\|_n + \|e^{-tL} v\|_\infty + \|De^{-tL} v\|_n) \\ \leq C \|v - a\|_q + C t^{-(n/r-n/q)/2} \|a\|_r, \end{aligned}$$

where we have used (2.15) and (2.16). Hence the density of $J^q \cap J^r$ in J^q implies (2.17). The proof is complete.

3. Proof of Theorem 0.2. From Theorem 0.1 we can suppose that $\|Dw\|_{n/2} + \|w\|_{2n}^{1/2} \|w\|_{2n/3}^{1/2}$ is small such that (2.15)–(2.17) holds.

By using the projection P to (0.2)–(0.3), and setting $u(t) = v(t) - w$ and $a = v_0 - w$, then (0.2)–(0.3) leads to the evolution equation

$$(3.1) \quad (d/dt)u + Lu = -P(u \cdot D)u \quad (t > 0), \quad u(0) = a$$

in J^n . Hence, our goal now remains to show that (3.1) has a unique solution u belonging to the space

$$U \equiv \{u \in BC([0, \infty); J^n); t^{1/2} Du(t) \in BC([0, \infty); L^n(R^n; R^{n^2}))\}$$

such that

$$Hu(t) \equiv \|u(t)\|_n + t^{1/2} \|u(t)\|_\infty + t^{1/2} \|Du(t)\|_\infty \rightarrow 0 \quad \text{as } t \rightarrow \infty$$

provided that $a \in J^n$ with $\|a\|_n$ small enough.

Let us impose the following notation.

$$\|u\| = \sup_{t>0} Hu(t),$$

$$W = \{u \in U; \|u\| < \infty, Hu(t) \rightarrow 0 \text{ as } t \rightarrow \infty\},$$

$$Mu(t) = u_0(t) - \int_0^t e^{-(t-s)L} P(u \cdot D)u(s) ds; \quad u_0(t) = e^{-tL} a.$$

Observing that for $u \in C_0^\infty$,

$$\begin{aligned} & t^{1/2}(\|e^{-tL}P(u \cdot D)u\|_\infty + \|De^{-tL}P(u \cdot D)u\|_n) + \|e^{-tL}P(u \cdot D)u\|_n \\ & \leq Ct^{-1/4}\|(u \cdot D)u\|_{2n/3}, \quad \text{by (2.15)-(2.16),} \\ & \leq Ct^{-1/4}\|u\|_{2n}\|Du\|_n, \end{aligned}$$

and

$$\|u\|_{2n} \leq \|u\|_n^{1/2}\|u\|_\infty^{1/2},$$

we have for $u \in W$,

$$\begin{aligned} (3.2) \quad & \|Mu(t)\|_n + t^{1/2}\|Mu(t)\|_\infty + t^{1/2}\|DMu(t)\|_n \\ & \leq C\|a\|_n + C \int_0^t (t-s)^{-1/4}\|u(s)\|_{2n}\|Du(s)\|_n ds \\ & \quad + C \int_0^t t^{1/2}(t-s)^{-3/4}\|u(s)\|_{2n}\|Du(s)\|_n ds, \\ & \hspace{15em} \text{by (2.15)-(2.16),} \\ & \leq C\|a\|_n + C\|u\|^2, \end{aligned}$$

and what is more, by using (2.17) and the property

$$Hu_0(t) + Hu(t) \rightarrow 0 \quad \text{as } t \rightarrow \infty$$

via a calculation similar to (3.2), we have

$$H(Mu)(t) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Moreover, by a standard calculation from [19] or [12], we have $Mu \in U$ for $u \in W$, and so $M: W \rightarrow W$ and

$$\|Mu\| \leq C\|a\|_n + C\|u\|^2.$$

Additionally, similar to (3.2), we obtain for $u_1, u_2 \in W$,

$$\|Mu_1 - Mu_2\| \leq C(\|u_1\| + \|u_2\|)\|u_1 - u_2\|.$$

From contraction mapping principle it follows that M has a fixed point u in W provided $\|a\|_n$ is sufficiently small. As in [12, 5], we find that the fixed point u is the desired solution which exists uniquely in U . The proof is complete.

REFERENCES

- [1] W. Borchers and T. Miyakawa, L^2 decay for Navier-Stokes flows in unbounded domains, with application to exterior stationary flows, to appear.
- [2] R. Finn, On the exterior stationary problem for the Navier-Stokes equations, and associated perturbation problems, Arch. Rational Mech. Anal., **19** (1965), 363-406.

- [3] ———, *Mathematical questions relating to viscous fluid flow in an exterior domain*, Rocky Mountain J. Math., **3** (1973), 107–140.
- [4] A. Friedman, *Partial Differential Equations*, Academic Press, New York, 1969.
- [5] H. Fujita and T. Kato, *On the Navier-Stokes initial value problems. I*, Arch. Rational Mech. Anal., **16** (1964), 269–315.
- [6] G. P. Galdi and S. Rionero, *Weighted Energy Methods in Fluid Dynamics and Elasticity*, Lecture Notes in Math., vol. 1134, Springer-Verlag, Berlin and New York, 1985.
- [7] G. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, New York, 1983.
- [8] J. A. Goldstein, *Semigroups of Linear Operators and Applications*, Oxford University Press, New York, 1985.
- [9] J. G. Heywood, *On stationary solutions of the Navier-Stokes equations as limits of nonstationary solutions*, Arch. Rational Mech. Anal., **37** (1970), 48–60.
- [10] ———, *The exterior nonstationary problem for the Navier-Stokes equations*, Acta Math., **129** (1972), 11–34.
- [11] ———, *The Navier-Stokes equations: on the existence, regularity and decay of solutions*, Indiana Univ. Math. J., **29** (1980), 639–681.
- [12] T. Kato, *Strong L^q -solutions of the Navier-Stokes equations in R^n , with applications to weak solutions*, Math. Z., **187** (1984), 471–480.
- [13] H. Kozono and T. Ozawa, *Stability in L^r for the Navier-Stokes flow in an n -dimensional bounded domain*, J. Math. Anal. Appl., **152** (1990), 35–45.
- [14] K. Masuda, *On the stability of incompressible viscous fluid motions past objects*, J. Math. Soc., Japan, **27** (1975), 294–327.
- [15] M. McCracken, *The resolvent problem for the Stokes equations on halfspace in L_p* , SIAM J. Math. Anal., **12** (1981), 201–228.
- [16] T. Miyakawa and H. Sohr, *On energy inequality, smoothness and large time behavior in L^2 for weak solutions of the Navier-Stokes equations in exterior domains*, Math. Z., **199** (1988), 455–478.
- [17] J. Schnute and M. Shinbrot, *The Cauchy problem for the Navier-Stokes equations*, Quart. J. Math. Oxford, **24** (1973), 457–473.
- [18] M. Shinbrot, *The Cauchy problem for the Navier-Stokes equations: a correction*, Quart. J. Math. Oxford, **27** (1976), 135–137.
- [19] F. Weissler, *The Navier-Stokes initial value problem in L^p* , Arch. Rational Mech. Anal., **74** (1980), 219–230.

Received July 11, 1991 and in revised form March 13, 1992.

TIANJIN UNIVERSITY
TIANJIN 3000 72
PEOPLE'S REPUBLIC OF CHINA

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982) F. WOLF (1904–1989)

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@henri.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Julie Speckart, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 75 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$200.00 a year (10 issues). Special rate: \$100.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

This publication was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$,
the American Mathematical Society's $\mathcal{T}\mathcal{E}\mathcal{X}$ macro system.
Copyright © 1993 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 158 No. 2 April 1993

- On the extension of Lipschitz functions from boundaries of subvarieties to strongly pseudoconvex domains 201
K. ADACHI and HIROSHI KAJIMOTO
- On a nonlinear equation related to the geometry of the diffeomorphism group 223
DAVID DAI-WAI BAO, JACQUES LAFONTAINE and TUDOR S. RATIU
- Fixed points of boundary-preserving maps of surfaces 243
ROBERT F. BROWN and BRIAN SANDERSON
- On orthomorphisms between von Neumann preduals and a problem of Araki 265
L. J. BUNCE and JOHN DAVID MAITLAND WRIGHT
- Primitive subalgebras of complex Lie algebras. I. Primitive subalgebras of the classical complex Lie algebras 273
I. V. CHEKALOV
- L^n solutions of the stationary and nonstationary Navier-Stokes equations in R^n 293
ZHI MIN CHEN
- Some applications of Bell's theorem to weakly pseudoconvex domains 305
XIAO JUN HUANG
- On isotropic submanifolds and evolution of quasicauistics 317
STANISŁAW JANECZKO
- Currents, metrics and Moishezon manifolds 335
SHANYU JI
- Stationary surfaces in Minkowski spaces. I. A representation formula 353
JIANGFAN LI
- The dual pair $(U(1), U(1))$ over a p -adic field 365
COURTNEY HUGHES MOEN
- Any knot complement covers at most one knot complement 387
SHICHENG WANG and YING QING WU



0030-8730(1993)158:2;1-H