

*Pacific
Journal of
Mathematics*

ON AMBIENTAL BORDISM

CARLOS BIASI

Volume 163 No. 1

March 1994

ON AMBIENTAL BORDISM

CARLOS BIASI

Let M^m be a closed and oriented submanifold of a closed or oriented manifold N^n , such that $[M, i] = 0 \in \Omega_m(N)$, where $i: M \rightarrow N$ is the inclusion and $\Omega_m(N)$ is the m th oriented bordism group of N . If $n = m + 2$ or $m \leq 3$ or $m \leq 4$ and $n \neq 7$ then M bounds in N .

Introduction. Let us consider M^m a closed submanifold of N^n . In this paper, we study the possibility that there exists submanifold $W^{m+1} \subset N^n$ such that $\partial W = M$. If $M = S^m$ and $N = S^{m+2}$, such that a submanifold W is called a Seifert surface knot S^m . In [5], Sato showed that every connected closed and oriented submanifold M^m of S^{m+2} is a boundary of an oriented surface of S^{m+2} .

In [4], Hirsch studies the following problem: If a compact connected and oriented manifold M^m bounds, does there exist embedding from M^m into \mathbb{R}^n which is a boundary in \mathbb{R}^n ?

The answer is yes, if $n \geq 2m$.

The difference between the two problems is that, in our case, the embedding from M into N is fixed.

There is an obvious necessary condition for the existence of W , when M and N are oriented manifolds.

Let $\Omega_m(N)$ be the m th oriented bordism group of N [2]. If $i: M \rightarrow N$ is the inclusion map, we can define an element $[M, i]$ in $\Omega_m(N)$ and see that $[M, i] = 0$ if M bounds in N .

Generally, the converse is not true, but sometimes the vanishing of $[M, i]$ guarantees the existence of W , for example if the codimension $n - m$ is large.

We prove the following theorem.

THEOREM 5.2. *Let us suppose that $M^m \subset N^n$, $n > m + 1$, is such that $[M, i] = 0$ in $\Omega_m(N)$. Then M bounds in N if one of the following conditions occurs:*

- (a) $n = m + 2$,
- (b) $m \leq 3$,
- (c) $m \leq 4$ and $n \neq 7$.

In his Doctoral thesis [1] the author proved that, when $n = 2m + 1$, and M and N are closed and oriented, a submanifold $M \subset N$ bounds in N if, and only if, $[M, i] = 0 \in \Omega_m(N)$.

1. A more general problem of ambient bordism. Let

$$G \subset O(n - m - 1), \quad n > m + 1,$$

be a closed transformation group and let $\gamma_G \rightarrow BG$ be the classifying fiber bundle of $(n - m - 1)$ -vector bundles which have a G -structure.

Let us consider MG the Thom space of γ_G . We have:

$$\pi_i(MG) = \begin{cases} 0, & i < n - m - 1, \\ \mathbb{Z}, & i = n - m - 1 \text{ and } G \subset \text{SO}(n - m - 1), \\ \mathbb{Z}_2, & i = n - m - 1 \text{ and } G \not\subset \text{SO}(n - m - 1). \end{cases}$$

Let us consider now N^n to be a closed connected manifold which we assume to be oriented if $G \subset \text{SO}(n - m - 1)$. (If $G \not\subset \text{SO}(n - m - 1)$ we drop the orientability hypothesis.)

Let $M^m \subset N^n$ be a closed submanifold and let us suppose that the normal fiber bundle ν_M of M in N has a cross section s , nowhere zero, such that $\nu_M = \{s\} \oplus \xi$, where $\{s\}$ is a subbundle generated by s and ξ is a $(n - m - 1)$ -vector bundle endowed with a G -structure.

We shall say that a submanifold $W \subset N$ satisfies condition (*) if it has the properties:

- (i) $\partial W = M$ and s is the inward-pointing vector field on ∂W .
- (ii) the normal fiber bundle ν_W has a G -structure which agrees with the given G -structure of ξ over M . (Observe that $\xi = \nu_W|_M$.)

2. Primary obstruction to the existence of W . Let V be a closed tabular neighborhood of M in N , $A = \partial V$ and $X = N - \overset{\circ}{V}$. We can think s a function $s: M \rightarrow A$. Then $s(M)$ is a submanifold of A , whose normal fiber bundle is isomorphic to ξ . By the Thom construction there exists a function $f: A \rightarrow MG$ such that, if ∞ is the point at infinity to MG , then f is differentiable on $A - f^{-1}(\infty)$, f is transversal to BG and $f^{-1}(BG) = (M)$ [6].

We shall take $\pi_{m-n-1}(MG)$ as the cohomology coefficient group. Let $e \in H^{n-m-1}(MG)$ be the fundamental class of the space MG . We know that $f^*(e) = \alpha$, where α is the dual class of $s_*(\mu_M)$ and μ_M is the fundamental class of M .

If $f: A \rightarrow MG$ extends to a map $\bar{f}: X \rightarrow MG$, then we can suppose, up to homotopy, that \bar{f} is differentiable in $X - \bar{f}^{-1}(\infty)$ and that \bar{f} is transversal to BG . Taking $W_1 = \bar{f}^{-1}(BG)$ we obtain a submanifold of X whose boundary is $s(M)$.

Let us observe that this submanifold can be extended to a submanifold W which satisfies condition $(*)$.

We conclude then that there exists W , satisfying $(*)$, if and only if f extends to X .

The class $\delta f^*(e)$ is the obstruction to the extension of f to the $(n - m)$ -skeleton of X , where $\delta: H^{n-m-1}(A) \rightarrow H^{n-m}(X, A)$ is the coboundary operator.

Consider the commutative diagram:

$$\begin{array}{ccc} H^{n-m-1}(A) & \xrightarrow{\delta} & H^{n-m}(X, A) \\ \downarrow D & & \downarrow D \\ H_m(A) & \xrightarrow{s_*} & H_m(X) \cong H_m(N - M). \end{array}$$

We conclude that the primary obstruction to the extension of f , up to duality, is the element $s_*(\mu_M) \in H_m(N - M)$ (regarding s as function from M into $N - M$).

Hence, we have:

PROPOSITION 2.1. *f extended to the $(n - m)$ -skeleton of X if, and only if, $s_*(\mu_M) = 0$ in $H_m(N - M)$.*

Assuming that $s_*(\mu_M) = 0$, let us consider two cases:

1. $G = O(n - m - 1)$.

Here, f extends up to the $(n - m + 1)$ -skeleton of X , because $\pi_{n-m}(MG) = 0$ and, if $n - m = 2$, then f extends to all of X since $MO(1)$ is a $K(\mathbb{Z}_2, 1)$ space.

2. $G = SO(n - m - 1)$.

Since $\pi_{n-m+i}(MG) = 0$, $i = 0, 1, 2$, f extends up to the $(n - m + 3)$ -skeleton of X . Hence, if $\dim M \leq 3$, f extends.

On the other hand, if $n - m = 2$ or 3 then MG is a $K(\mathbb{Z}, 1)$ or $K(\mathbb{Z}, 2)$, respectively. In any case, f extends globally.

3. Oriented ambiental bordism. From now on, all manifolds and submanifolds will be considered to be oriented.

THEOREM 3.1. *Let us suppose that:*

(a) $H_j(X) = 0$, $0 < j < m - 3$.

(b) *The canonical homomorphism $\pi_{n-1}(\text{MSO}(n - m - 1)) \xrightarrow{\varphi} \Omega_m$ is injective.*

There exists W satisfying $()$ if, and only if, $s_*(\mu_M) = 0 \in H_m(X)$ and M is a boundary.*

Proof. Let us use the notation $\pi_i = \pi_i(\text{MSO}(n - m - 1))$. If $s_*(\mu_M) = 0$, then f extends to the $(n - m)$ -skeleton of X .

From hypothesis (a) and Lefschetz duality, we conclude that

$$H^j(X, A, \pi_{j-1}) = 0, \quad n - m < j < n.$$

Let D be an open disk of $X - A$. Since X is orientable, $H^j(X - D, A, \pi_{j-1}) \cong H^j(X, A, \pi_{j-1}) = 0$, $n - m < j < n$. Hence, there exists an extension $\bar{f}: X - D \rightarrow Y$ of $f: A \rightarrow Y$, where $Y = \text{MSO}(n - m - 1)$.

Let us consider $S = \partial D$ and $h = \bar{f}|_{\partial D}: S \rightarrow Y$. We may suppose that h is transversal to $\text{BSO}(n - m - 1)$ and let

$$M^m = h^{-1}(\text{BSO}(n - m - 1)).$$

Consider $\bar{W} = \bar{f}^{-1}(\text{BSO}(n - m - 1))$, a bordism between M_1 and $s(M)$. Since $s(M)$ is a boundary, M_1 also is.

We have also that $\psi([h]) = [M_1] = 0$ and since ψ is a monomorphism, h is homotopic to a constant map and so h extends over D .

The converse is straightforward. \square

4. On the existence of normal vector fields homologous to zero in $N - M$. In the next section, we show that in certain situations it is possible to obtain a cross-section $s: M \rightarrow S(\nu_M)$ such that $s_*(\mu_M) = 0 \in H_m(N - M)$, where $S(\nu_M) \rightarrow M$ is the normal sphere bundle of M in N .

PROPOSITION 4.1. *The Euler class of the normal bundle of M^m in N^n is zero if and only if $i_*(\mu_M) \subset \text{im } j_*$, where μ_M is the fundamental class of M and $i: M \rightarrow N$, $j: N - M \rightarrow N$ are inclusion maps.*

Proof. Let us consider $e \in H^{n-m}(M, \mathbb{Z})$, the Euler class of the normal bundle ν_M , and let $D_A: H^{n-m}(M; \mathbb{Z}) \rightarrow H_m(N, N - M; \mathbb{Z})$ be the Alexander duality. We have that $D_A(e) = \alpha_*(\mu_M)$ where α_* is induced by the inclusion map $\alpha: (N, N - M)$.

Using the exact sequence of pair $(N, N - M)$ it follows that $\alpha_*(\mu_M) = 0$ if, and only if, $i_*(\mu_M) \subset \text{im } j_*$. \square

COROLLARY 4.2. *If $M^m \subset N^n$ is homologous to zero, $n - m = 2$ or $n \geq 2m$, then M has a normal vector field that is nowhere zero.*

Proof. By Proposition 4.1 the Euler class of ν_M is zero. Then there is a nowhere zero normal vector field on the $(n - m)$ -skeleton

of M , which can be extended to all M , because $n - m \geq m$ or $\pi_i(R^2 - 0) = 0$, $i > 1$ in the case $n - m = 2$. \square

Let $\pi: E \rightarrow M^m$ be a differentiable $SO(n + 1)$ -bundle with fiber S^n and base M^m (and oriented manifold).

If $s: M \rightarrow E$ is a cross-section, let θ_s be the Poincaré dual to $\bar{s}_*(\mu_M)$, where $\bar{s} = -s$ is the opposite cross-section to s .

Having fixed a cross-section $s_0: M \rightarrow E$, the following diagrams are commutative:

$$\begin{array}{ccccc}
 [M, E] & & & & \\
 \downarrow \xi & \searrow \varphi & & & \\
 H^n(M) & \xrightarrow{\pi^*} & H^n(E) & & \\
 \downarrow D & & \downarrow D & & \\
 H_{m-n}(M) & \xrightarrow{\Delta} & H_m(E) & \xrightarrow{\pi_*} & H_m(M)
 \end{array}$$

where $[M, E]$ is the set of homotopy classes of cross-sections, $\xi([s]) = \bar{s}^*(\theta_{\bar{s}_0})$; $\varphi([s]) = \theta_{\bar{s}_0} - \theta_{\bar{s}}$, is Poincaré duality and last line is a portion of the generalized Gysin sequence.

We define $\psi: [M, E] \rightarrow H_m(E)$ by $\psi([s]) = s_{s_*}(\mu_M) - s_*(\mu_M)$ and observe that $\psi = D \circ \xi$.

If $m \leq n + 1$ or $n = 1$, then the function ξ is onto and so the image of ψ is the kernel of π_* .

This fact will be applied in the proof of Proposition 4.3 below, where the fiber bundle to be considered is $S(\nu_M) \rightarrow M$.

PROPOSITION 4.3. *Let $M^m \subset N^n$, $n = m + 2$ or $n \geq 2m$, be an oriented submanifold homologous to zero in an oriented manifold N . Then there exists a cross-section $r: M \rightarrow S(\nu_M)$ such that its image is homologous to zero in $H_m(N - m)$.*

Proof. Let $s_0: M \rightarrow S(\nu_M)$ be a cross-section that is nowhere zero (Corollary 4.2) and let us consider the commutative diagrams:

$$\begin{array}{ccccc}
 & s_{0*} \nearrow & H_m(S(\nu_M)) & \xrightarrow{\pi_*} & H_m(M) \\
 H_m(M) & & \downarrow l_* & & \downarrow i_* \\
 & s_* \searrow & H_m(N - M) & \xrightarrow{j_*} & H_m(N)
 \end{array}$$

where $s_* = l_*(s_{0*})$ and l_* is induced by the inclusion $S(\nu_M) \rightarrow (N - M)$.

We have $j_*s_*(\mu_M) = i_*\pi_*s_0(\mu_M) = 0$ implying that $s_*(\mu_M)$ belongs to the kernel of j_* which is the image of $\partial: H_{m+1}(N, N-M) \rightarrow H_m(N-M)$.

Let us consider the following commutative diagram:

$$\begin{array}{ccc} H_{m+1}(D(\nu_M), S(\nu_M)) & \xrightarrow{\partial} & H_m(S(\nu_M)) \\ \downarrow exc & & \downarrow j_* \\ H_{m+1}(N, N-M) & \xrightarrow{\partial} & H_m(N-M). \end{array}$$

It follows that there exists an element $\mu \in H_m(S(\nu_M))$ such that $\mu \in \text{Ker } \pi_*$ and $j_* = s_*(\mu_M)$.

Since $\text{im } \psi = \text{ker } \pi_*$, there exists a cross-section $r: M \rightarrow S(\nu_M)$ such that $\psi([r]) = \mu$.

But $\psi([r]) = s_0(\mu_M) \rightarrow r_*(\mu_M)$ so $j_*r_*(\mu_M) = 0$ in $H_m(N-M)$. Hence, the image of $r: M \rightarrow S(\nu_M)$ is homologous to zero in $N-M$.

5. A theorem on ambient bordism. Let us consider $\Omega_j(N)$ to be the j th bordism group of N .

If $H_j(N) = 0$, $0 < j < m-3$, it is possible using the bordism spectral sequence [2] to show that the function $\Omega_m(N) \rightarrow H_m(N) \oplus \Omega_m$, which associates to each pair $[M, f]$ the element $\mu([M, f]) + [M]$, is an isomorphism, where μ is the canonical homomorphism.

In the proof of Theorem 5.2, we are going to use the following lemma, which has been proved in [1] (the proof, if $q > m$, is due to Thom [6]).

LEMMA 5.1. *The homomorphism $\varphi: \pi_{q+m}(\text{MSO}(q)) \rightarrow \Omega_m$, $q \geq m$, is an isomorphism.*

THEOREM 5.2. *Let us suppose $M^m \subset N^n$, $n > m+1$, is such that $[M, i] = 0$ in $\Omega_m(N)$. Then M bounds in N if one of the following conditions occurs:*

- (a) $n = m+2$,
- (b) $m \leq 3$,
- (c) $m \leq 4$ and $n \neq 7$.

Proof. Any one of the conditions (a), (b) and (c), based on previous results, imply that normal bundle ν_M has a cross-section nowhere zero such that, considering s as a function from M into $N-M$, $s_*(\mu_M) = 0 \in H_m(N-M)$.

If (a) or (b) occurs, the theorem follows from case 2, already discussed in §2

If $n = 4$ and $n \geq 8$, we apply Theorem 3.1.

REMARK 1. If $n = m + 2$ or $m \leq 3$, then $[M, i] = 0 \in \Omega_m(N)$ if, and only if, M is homologous to zero in N .

REMARK 2. When $m = 4$ and $n \neq 7$, although we shall prove that $[M, i] = 0$ implies the existence of a normal section nowhere zero (Th. 5.3) we are not able to prove that there exists a normal vector field homologous to zero in $N - M$, which in this case would be sufficient to prove the conclusion of Theorem 5.2.

THEOREM 5.3. *Let us suppose $M^4 \subset N^7$. If $[M, i] = 0$ in $\Omega_4(N)$ then ν_M has a cross-section which is nowhere zero.*

Proof. There exists $W \subset N \times I$ such that $\partial W = M \times 0 \subset N \times I$ [1].

Let ν_W and ν_M be the normal fiber bundles of W in $N \times I$ and of M in N , respectively. We can also suppose that $\nu_W|_{M \times 0} = \nu_M$.

Let us consider $\overline{W} \subset N \times \mathbb{R}$ to be the double of W and let $i: \overline{W} \rightarrow N \times \mathbb{R}$ and $j: N \times \mathbb{R} \rightarrow \overline{W} \rightarrow N \times \mathbb{R}$ be inclusion maps.

Since $i_*(\mu_{\overline{W}}) \subset \text{im } j_*$, then \overline{W} has a normal vector field which is nowhere zero in $N \times \mathbb{R}$ up to the 3-skeleton of \overline{W} .

Hence, there exists a 2-dimensional oriented vector bundle ξ over M such that $\nu_M|M^{(3)} = \xi \otimes \mathcal{E}^1$.

Let us consider e to be the Euler class of ξ in $H^2(M^{(3)})$ and let $\bar{e} \in H^2(M)$ be such that $io^*(\bar{e}) = e$, where $i: M^{(3)} \rightarrow M$ is the inclusion map.

Let $\bar{\xi}$ be a 2-dimensional vector bundle over M such that its Euler class is \bar{e} . Let us observe that $\bar{\xi}|_{M^{(3)}} = \xi$.

Let $f, g: M \rightarrow \text{BSO}(3)$ be classifying maps $\bar{\xi} \oplus \mathcal{E}^1$ and ν_M , respectively.

Since the Euler classes of $\bar{\xi} \oplus \mathcal{E}^1$ and of ν_M are equal, then their second Stiefel-Whitney classes are equal.

Let \tilde{p}_1 be the Pontryagin class of the classifying fiber bundle $\tilde{\gamma} \rightarrow \text{BSO}(3)$ and let \bar{e} be the Euler class of $\tilde{\gamma}$. Since $f^*(\tilde{p}_1) = g^*(\tilde{p}_1)$. Hence, the vector bundles $\bar{\xi} \oplus \mathcal{E}^1$ and ν_M are equivalent [3]. \square

REFERENCES

- [1] C. Biasi, *L-equivalent and ambient bordism of submanifolds*, (Doctoral Thesis), ICMSC-USP, São Carlos (1981).
- [2] P. E. Conner and E. E. Floyd, *Differentiable Periodic Maps*, Springer-Verlag (1964).
- [3] A. Dold and H. Whitney, *Classification of oriented sphere bundles over a 4-complex*, Ann. of Math., **69** (1959), 667–677.
- [4] M. Hirsch, *On embedding of bounding manifolds in euclidean space*, Ann. of Math., (2), **74** (1961), 494–497.
- [5] N. Sato, *Cobordism of semiboundary links*, Topology Appl., **18** (1984), 225–234.
- [6] R. Thom, *Quelques propriétés globales de variétés différentiable*, Comment. Math. Helv., **28** (1954), 17–86.

Received November 13, 1991 and in revised form November 19, 1992.

ICMSC-USP

DEPARTAMENTO DE MATEMÁTICA

RUA DR. CARLOS BOTELHO, 1465

CAIXA POSTAL 668

13560-970-SÃO CARLOS-SP-BRASIL

CONTENTS

| | |
|---|-----|
| P. Ahern and W. Cohn, Weighted maximal functions and derivatives of invariant Poisson integrals of potentials | 1 |
| V. Ancona, T. Peternell, and J. A. Wiśniewski, Fano bundles and splitting theorems on projective spaces and quadrics | 17 |
| G. M. Beffa, A transverse structure for the Lie-Poisson bracket on the dual of the Virasoro algebra | 43 |
| C. Biasi, On ambient bordism | 73 |
| P. Debes and M. D. Fried, Nonrigid constructions in Galois theory | 81 |
| K. Dykema, Interpolated free group factors | 123 |
| S.-A. Kim and D. Minda, Two-point distortion theorems for univalent functions | 137 |
| X. Mo, Value distribution of the Gauss map and the total curvature of complete minimal surface in R^m | 159 |
| Z. Shen, On complete Riemannian manifolds with collapsed ends | 175 |
| F. Zheng, Curvature characterization of certain bounded domains of holomorphy | 183 |
| S. Zhu, The classification of complete locally conformally flat manifolds of non-negative Ricci curvature | 189 |

PACIFIC JOURNAL OF MATHEMATICS

Volume 163 No. 1 March 1994

| | |
|--|-----|
| Weighted maximal functions and derivatives of invariant Poisson integrals of potentials | 1 |
| PATRICK ROBERT AHERN and WILLIAM S. COHN | |
| Fano bundles and splitting theorems on projective spaces and quadrics | 17 |
| VINCENZO ANCONA, THOMAS PETERNELL and JAROSLAW WISNIEWSKI | |
| A transverse structure for the Lie-Poisson bracket on the dual of the Virasoro algebra | 43 |
| GLORIA MARI BEFFA | |
| On ambient bordism | 73 |
| CARLOS BIASI | |
| Nonrigid constructions in Galois theory | 81 |
| PIERRE DÈBES and MICHAEL FRIED | |
| Interpolated free group factors | 123 |
| KENNETH JAY DYKEMA | |
| Two-point distortion theorems for univalent functions | 137 |
| SEONG-A KIM and C. DAVID (CARL) MINDA | |
| Value distribution of the Gauss map and the total curvature of complete minimal surface in R^m | 159 |
| XIAOKANG MO | |
| On complete Riemannian manifolds with collapsed ends | 175 |
| ZHONGMIN SHEN | |
| Curvature characterization of certain bounded domains of holomorphy | 183 |
| FANGYANG ZHENG | |
| The classification of complete locally conformally flat manifolds of nonnegative Ricci curvature | 189 |
| SHUN-HUI ZHU | |



0030-8730(1994)163:1;1-J