

*Pacific
Journal of
Mathematics*

MÖBIUS-INVARIANT HILBERT SPACES IN POLYDISCS

H. TURGAY KAPTANOGLU

MÖBIUS-INVARIANT HILBERT SPACES IN POLYDISCS

H. TURGAY KAPTANOĞLU

We define the Dirichlet space \mathcal{D} on the unit polydisc \mathbb{U}^n of \mathbb{C}^n . \mathcal{D} is a semi-Hilbert space of holomorphic functions, contains the holomorphic polynomials densely, is invariant under compositions with the biholomorphic automorphisms of \mathbb{U}^n , and its semi-norm is preserved under such compositions. We show that \mathcal{D} is unique with these properties. We also prove \mathcal{D} is unique if we assume that the semi-norm of a function in \mathcal{D} composed with an automorphism is only equivalent in the metric sense to the semi-norm of the original function. Members of a subclass of \mathcal{D} given by a norm can be written as potentials of \mathcal{L}^2 -functions on the n -torus \mathbb{T}^n . We prove that the functions in this subclass satisfy strong-type inequalities and have tangential limits almost everywhere on $\partial\mathbb{U}^n$. We also make capacity estimates on the size of the exceptional sets on $\partial\mathbb{U}^n$.

1. Introduction. *Möbius-invariant spaces.* Let \mathbb{U} be the open unit disc in \mathbb{C} and \mathbb{T} be the unit circle bounding it. The open unit polydisc \mathbb{U}^n and the torus \mathbb{T}^n in \mathbb{C}^n are the cartesian products of n unit discs and n unit circles, respectively. \mathbb{T}^n is the distinguished boundary of \mathbb{U}^n and forms only a small part of the topological boundary $\partial\mathbb{U}^n$ of \mathbb{U}^n . We denote by \mathcal{M} the group of all biholomorphic automorphisms of \mathbb{U}^n (the Möbius group). The subgroup of linear automorphisms in \mathcal{M} is denoted by \mathcal{U} . The space of holomorphic functions with domain \mathbb{U}^n will be called $\mathcal{H}(\mathbb{U}^n)$ and will carry the topology of uniform convergence on compact subsets of \mathbb{U}^n .

A semi-inner product on a complex vector space \mathcal{H} is a sesquilinear functional on $\mathcal{H} \times \mathcal{H}$ with all the properties of an inner product except that it is possible to have $\langle\langle a, a \rangle\rangle = 0$ when $a \neq 0$. $\|a\| = \sqrt{\langle\langle a, a \rangle\rangle}$ is the associated semi-norm. We assume $\langle\langle \cdot, \cdot \rangle\rangle$ is not identically zero.

DEFINITION 1.1. \mathcal{H} is called a Hilbert space of holomorphic functions on \mathbb{U}^n if

- (i) \mathcal{H} is a linear subspace of $\mathcal{H}(\mathbb{U}^n)$,
- (ii) the semi-inner product $\langle\langle \cdot, \cdot \rangle\rangle$ of \mathcal{H} is complete,
- (iii) \mathcal{H} contains all (holomorphic) polynomials,
- (iv) polynomials are dense in \mathcal{H} in the topology of the semi-norm $\|\cdot\|$ of \mathcal{H} .

A space \mathcal{H} of functions on U^n is \mathcal{M} -invariant if $f \circ \Psi \in \mathcal{H}$ whenever $f \in \mathcal{H}$ and $\Psi \in \mathcal{M}$. An \mathcal{M} -invariant Hilbert space \mathcal{H} of holomorphic functions on U^n will be called an \mathcal{M} -space for brevity. \mathcal{U} -invariance and \mathcal{U} -space have similar definitions.

\mathbb{N} , \mathbb{Z}_+ , \mathbb{Z} , \mathbb{R} denote the set of nonnegative integers, positive integers, integers, and real numbers, respectively. A multi-index $\alpha = (\alpha_1, \dots, \alpha_n)$ is a point in \mathbb{N}^n . \sum_α indicates a summation with α running over all the points in \mathbb{N}^n , and $\sum'_{\alpha \in \mathbb{I}}$ is a summation where we consider only those α in the index set \mathbb{I} with all positive components. Let also $D_j = \partial/\partial z_j$ and $\bar{D}_j = \partial/\partial \bar{z}_j$. The following abbreviated notations will be used:

$$\begin{aligned} |\alpha| &= \alpha_1 + \dots + \alpha_n, & z^\alpha &= z_1^{\alpha_1} \dots z_n^{\alpha_n}, \\ \alpha! &= \alpha_1! \dots \alpha_n!, & D^\alpha &= D_1^{\alpha_1} \dots D_n^{\alpha_n}. \end{aligned}$$

The Dirichlet space $\mathcal{D}(U^n)$ is the class of $f(z) = \sum_\alpha f_\alpha z^\alpha \in \mathcal{H}(U^n)$ with

$$(1.1) \quad \|f\|_{\mathcal{D}}^2 = \sum_\alpha \alpha_1 \dots \alpha_n |f_\alpha|^2 = \sum_{k=0}^\infty \sum_{|\alpha|=k} \alpha_1 \dots \alpha_n |f_\alpha|^2 < \infty.$$

Equivalently, $\mathcal{D}(U^n)$ is the class of those $f \in \mathcal{H}(U^n)$ with

$$\|f\|_{\mathcal{D}}^2 = \int_{U^n} |D_1 \dots D_n f|^2 d\mu_n < \infty,$$

where μ_n is the Lebesgue measure on U^n normalized so that $\mu_n(U^n) = 1$. The semi-norm $\|\cdot\|_{\mathcal{D}}$ is obtained from the semi-inner product

$$\langle\langle f, g \rangle\rangle_{\mathcal{D}} = \sum_\alpha \alpha_1 \dots \alpha_n f_\alpha \bar{g}_\alpha = \int_{U^n} (D_1 \dots D_n f) \overline{(D_1 \dots D_n g)} d\mu_n.$$

Main results. In this work, we first prove two theorems which show that the Dirichlet space is unique among \mathcal{M} -spaces that have certain properties.

THEOREM A. *Let \mathcal{H} be an \mathcal{M} -space and suppose that*

$$(1.2) \quad \|f\| = \|f \circ \Psi\| \quad (f \in \mathcal{H}, \Psi \in \mathcal{M}).$$

Then

$$\|f\| = C \|f\|_{\mathcal{D}} \quad (f \in \mathcal{H}),$$

where $C = \|z_1 \dots z_n\|^2$. Thus \mathcal{H} is $\mathcal{D}(U^n)$.

Note that the assumption on the semi-norm is equivalent to $\langle\langle f \circ \Psi, g \circ \Psi \rangle\rangle = \langle\langle f, g \rangle\rangle$, and the conclusion implies that $\langle\langle f, g \rangle\rangle = C \langle\langle f, g \rangle\rangle_{\mathcal{D}}$, for all $f, g \in \mathcal{H}$ and $\Psi \in \mathcal{M}$.

For the second theorem, we need a strengthening of condition (ii) of Definition 1.1. To derive it, we write the Taylor series expansion at 0 of an $f \in \mathcal{H}$ by separating the higher and lower dimensional terms. For example, when $n = 2$, using (z, w) for (z_1, z_2) and (k, l) for (α_1, α_2) , we write

$$f(z, w) = f_{00} + \sum_{k=1}^{\infty} f_{k0} z^k + \sum_{l=1}^{\infty} f_{0l} w^l + \sum_{k, l=1}^{\infty} f_{kl} z^k w^l.$$

Since we assume conditions (iii) and (iv) of Definition 1.1, we can define a norm on \mathcal{H} by

$$\| \| f \| \|^2 = |f_{00}|^2 + \sum_{k=1}^{\infty} |f_{k0}|^2 \|z^k\|_1^2 + \sum_{l=1}^{\infty} |f_{0l}|^2 \|w^l\|_1^2 + \sum_{k, l=1}^{\infty} |f_{kl}|^2 \|z^k w^l\|^2,$$

where $\| \cdot \|$ is still the semi-norm of $\mathcal{H} \subset \mathcal{H}(\mathbb{U}^n)$ and $\| \cdot \|_1$ denotes the semi-norm of a similar $\mathcal{H}' \subset \mathcal{H}(\mathbb{U})$. We already know (see [1]) that $\| \cdot \|_1$ is equivalent to $\| \cdot \|_{\mathcal{D}}$ in \mathbb{U} . Since our proof of Theorem B is by induction on the dimension of the polydisc, the above definition of $\| \| \cdot \| \|$ makes sense. Now we make the alternate assumption

(ii)' \mathcal{H} is complete in the norm $\| \| \cdot \| \|$.

A similar condition was assumed in [1], whereas [4] assumes (ii).

THEOREM B. *Let \mathcal{H} be an \mathcal{M} -space in the sense modified by (ii)' and assume that there is a positive constant $\delta < 1$ such that*

$$(1.3) \quad \delta \| f \| \leq \| f \circ \Psi \| \leq \frac{1}{\delta} \| f \| \quad (f \in \mathcal{H}, \Psi \in \mathcal{M}).$$

Then there exists positive constants K_1 and K_2 such that

$$K_1 \| f \|_{\mathcal{D}} \leq \| f \| \leq K_2 \| f \|_{\mathcal{D}} \quad (f \in \mathcal{H}).$$

Thus $\mathcal{D}(\mathbb{U}^n)$ is unique again.

The proofs of these theorems will be presented in §2.

Next, we consider a subspace of the Dirichlet space, one that is defined by a genuine norm similar to $\| \| \cdot \| \|$. This space is not \mathcal{M} -invariant any more, but the stronger conditions on it allow us to prove

that it has *tangential* limits as we approach $\partial\mathbb{U}^n$. In fact, tangential limits exist for a wider class of functions which are potentials of certain functions in $\mathcal{L}^2(\mathbb{T}^n)$. The precise definitions and theorems are stated in §3. Theorems C and D at the end of that section are the major results in this direction.

In earlier work, Arazy and Fisher [1] proved, under slightly different hypotheses, the analogs of Theorem A and Theorem B in \mathbb{U} . Zhu [5] found the equivalent of Theorem A for the unit ball in \mathbb{C}^n when $n \geq 2$. Nagel, Rudin and Shapiro [3] obtained the unit-disc versions of Theorems C and D.

After the submission of the manuscript, we were informed by a referee that in the preprint *Invariant Hilbert Spaces of Analytic Functions on Bounded Symmetric Domains* by J. Arazy and S. D. Fisher, results analogous to Theorems A and B were established for all irreducible bounded symmetric domains.

NOTATION. λ_n is the Lebesgue measure on \mathbb{T}^n both normalized to have mass 1; i.e., it is the Haar measure on the compact abelian group \mathbb{T}^n . If $p \in [1, \infty)$, its *conjugate* is $q = p/(p - 1)$. The \mathcal{L}^p - and ℓ^p -spaces will have their usual meaning. z_j will usually be an element of \mathbb{U} and ζ_j of \mathbb{T} . Apart from the usual big \mathcal{O} notation, we will use $u \sim v$ to mean both $u = \mathcal{O}(v)$ and $v = \mathcal{O}(u)$, and $u \approx v$ to mean u/v has a finite positive limit.

The *Poisson integral* of an $f \in \mathcal{L}^1(\mathbb{T}^n)$ is

$$P[f](z) = \int_{\mathbb{T}^n} f(\zeta) \prod_{j=1}^n \frac{1 - |z_j|^2}{|1 - z_j \bar{\zeta}_j|^2} d\lambda_n(\zeta) \quad (z \in \mathbb{U}^n),$$

and its *Cauchy integral* is

$$C[f](z) = \int_{\mathbb{T}^n} f(\zeta) \prod_{j=1}^n \frac{1}{1 - z_j \bar{\zeta}_j} d\lambda_n(\zeta) \quad (z \in \mathbb{U}^n),$$

where the products are called the *Poisson kernel* $P(z, \zeta)$ and the *Cauchy kernel* $C(z, \zeta)$ for \mathbb{U}^n , respectively. These transforms have the following invariance properties: If $f \in \mathcal{L}^1(\lambda_n)$, $\Psi \in \mathcal{M}$, and $U \in \mathcal{U}$, then

$$P[f \circ \Psi] = P[f] \circ \Psi \quad \text{and} \quad C[f \circ U] = C[f] \circ U.$$

The automorphisms of \mathbb{U}^n for $n \geq 2$ are generated by the following three subgroups: *rotations* in each variable separately

$$R_\theta(z) = (e^{i\theta_1} z_1, \dots, e^{i\theta_n} z_n),$$

Möbius transformations in each variable separately

$$\Phi_w(z) = (\phi_{w_1}(z_1), \dots, \phi_{w_n}(z_n)),$$

and the coordinate permutations. Here $\theta \in [-\pi, \pi]^n$ and $w \in \mathbb{U}^n$ are fixed, Möbius transformations are in the form

$$(1.4) \quad \phi_w(z) = \frac{w - z}{1 - \bar{w}z} \quad (w \in \mathbb{U}, z \in \bar{\mathbb{U}}),$$

and the coordinate permutations are nothing but the $n!$ members of the symmetric group \mathcal{S}_n on n objects. Thus an arbitrary $\Psi \in \mathcal{M}$ can be written in the form

$$\Psi(z) = \left(e^{i\theta_1} \phi_{w_1}(z_{\sigma(1)}), \dots, e^{i\theta_n} \phi_{w_n}(z_{\sigma(n)}) \right),$$

for some $w \in \mathbb{U}^n$ and $\theta \in [-\pi, \pi]^n$, and $\sigma \in \mathcal{S}_n$. \mathcal{U} is generated by $\sigma \in \mathcal{S}_n$ and the rotations R_θ . Each Möbius transformation Φ_w is an involution (its inverse is itself) exchanging 0 and w . \mathcal{M} acts transitively on \mathbb{U}^n : if $a, b \in \mathbb{U}^n$, then $\Phi_b \circ \Phi_a \in \mathcal{M}$ moves a to b (and b to a). Finally, \mathcal{M}^* denotes the component of the identity in \mathcal{M} ; i.e., \mathcal{M}^* is \mathcal{M} without the action of \mathcal{S}_n .

2. Uniqueness of the Dirichlet space. We start by showing that $\mathcal{D}(\mathbb{U}^n)$ has all the properties of a Hilbert space in the sense of Definition 1.1. Clearly the polynomials are in $\mathcal{D}(\mathbb{U}^n)$ and $\|z^\alpha\|^2 = \alpha_1 \cdots \alpha_n$ for all $\alpha \in \mathbb{N}^n$. A quick look at (1.1) shows that the polynomials are dense in $\mathcal{D}(\mathbb{U}^n)$ with respect to $\|\cdot\|_{\mathcal{D}}$. Again from (1.1), identifying g by $\{g_\alpha\}$, we see that $\mathcal{D}(\mathbb{U}^n)$ is a weighted ℓ^2 -space, hence every Cauchy sequence $\{f_j\}$ in $\mathcal{D}(\mathbb{U}^n)$ converges in $\|\cdot\|_{\mathcal{D}}$ to some $f \in \mathcal{D}(\mathbb{U}^n)$ represented by $\{f_\alpha\}$ for $\alpha \in \mathbb{Z}_+^n$. To show that f is holomorphic, let $f_m(z) = \sum_{k=1}^m \sum'_{|\alpha|=k} f_\alpha z^\alpha$ and pick $\varepsilon > 0$. For any $0 < r < 1$ and positive integers $m > l > n$,

$$\begin{aligned} \sup_{z \in r\bar{\mathbb{U}}^n} |(f_m - f_l)(z)| &= \sup_{z \in r\bar{\mathbb{U}}^n} \left| \sum_{k=l+1}^m \sum'_{|\alpha|=k} f_\alpha z^\alpha \right| \leq \sum_{k=l+1}^m \sum'_{|\alpha|=k} |f_\alpha| r^{|\alpha|} \\ &\leq \left(\sum_{k=l+1}^m \sum'_{|\alpha|=k} |f_\alpha|^2 \right)^{1/2} \left(\sum_{k=l+1}^m \sum'_{|\alpha|=k} r^{2|\alpha|} \right)^{1/2} \\ &< \left(\sum_{k=l+1}^m \sum_{|\alpha|=k} \alpha_1 \cdots \alpha_n |f_\alpha|^2 \right)^{1/2} \left(\sum_{k=l+1}^m k^n r^{2k} \right)^{1/2}. \end{aligned}$$

The first factor is less than ε when l and m are large enough because $f \in \mathcal{D}(\mathbb{U}^n)$, and the second factor is bounded as $l, m \rightarrow \infty$. Hence

$f(z) = \sum'_\alpha f_\alpha z^\alpha$ is uniformly convergent on compact subsets of \mathbb{U}^n , and this proves $f \in \mathcal{H}(\mathbb{U}^n)$. Note that we need not know f_α if $\alpha \in \mathbb{N}^n \setminus \mathbb{Z}_+^n$. These coefficients of f can be taken arbitrarily as long as f remains holomorphic.

LEMMA 2.1. \mathcal{M} is generated by \mathcal{S}_n , rotations $R_\omega(z) = (e^{i\omega} z_1, z_2, \dots, z_n)$ with $\omega \in [-\pi, \pi]$, and Möbius transformations of the form $\Phi_t(z) = (\phi_t(z_1), z_2, \dots, z_n)$ with $0 \leq t < 1$.

PROPOSITION 2.2. $\mathcal{D}(\mathbb{U}^n)$ is \mathcal{M} -invariant.

Proof. The integral form of the Dirichlet semi-norm uses the measure μ_n which is invariant under rotations and permutations. Thus $\mathcal{D}(\mathbb{U}^n)$ is \mathcal{U} -invariant. To prove invariance under Möbius transformations, in view of Lemma 2.1, it suffices to consider

$$w = \Phi_r(z) = \left(\frac{r - z_1}{1 - rz_1}, z_2, \dots, z_n \right).$$

Then $D_1^z(f \circ \Phi_r) = (D_1^w f)dw_1/dz_1$ and

$$\begin{aligned} |D_1^z D_2 \cdots D_n(f \circ \Phi_r)|^2 &= |D_1^w D_2 \cdots D_n f|^2 \frac{(r^2 - 1)^2}{|1 - rz_1|^4} \\ &= |D_1^w D_2 \cdots D_n f|^2 J_{\Re} \Phi_r(z), \end{aligned}$$

since $dw_1/dz_1 = (r^2 - 1)/(1 - rz_1)^2$, where $J_{\Re} \Phi_r$ is the real Jacobian of Φ_r . Therefore

$$\begin{aligned} \|f \circ \Phi_r\|_{\mathcal{D}}^2 &= \int_{\mathbb{U}^n} |D_1^z D_2 \cdots D_n(f \circ \Phi_r)(z)|^2 d\mu_n(z) \\ &= \int_{\mathbb{U}^n} |D_1^w D_2 \cdots D_n f(w)|^2 \frac{(r^2 - 1)^2}{|1 - rz_1|^4} \frac{1}{J_{\Re} \Phi_r(z)} d\mu_n(w) \\ &= \int_{\mathbb{U}^n} |D_1^w D_2 \cdots D_n f(w)|^2 d\mu_n(w) = \|f\|_{\mathcal{D}}^2. \quad \square \end{aligned}$$

Note that when $n \geq 2$, $\mathcal{D}(\mathbb{U}^n)$ does not put any conditions on the infinitely many power series coefficients of f , those with at least one $\alpha_j = 0$, i.e., those in $\mathbb{N}^n \setminus \mathbb{Z}_+^n$. Thus if each term in the Taylor expansion of some $f \in \mathcal{H}(\mathbb{U}^n)$ depends on fewer than n variables, then $\|f\|_{\mathcal{D}} = 0$ and $f \in \mathcal{D}$. The Dirichlet space can also be thought

of as a quotient space of holomorphic functions satisfying (1.1) where the functions whose Taylor series differ by terms depending on at most $n - 1$ variables are identified. Trivially any holomorphic function f of fewer than n variables or any constant f has $\|f\|_{\mathcal{D}} = 0$ and is in $\mathcal{D}(\mathbb{U}^n)$. For comparison, when $n = 1$, only constants (a one-dimensional subspace) have zero Dirichlet semi-norm.

We can define a *modified* Dirichlet space $\tilde{\mathcal{D}}(\mathbb{U}^n)$ similar to $\mathcal{D}(\mathbb{U}^n)$ by considering a norm instead of a semi-norm. This requires some control on *all* the power series coefficients of $f \in \mathcal{H}(\mathbb{U}^n)$. For simplicity let's look at the case $n = 2$. With notation as before, let

$$\begin{aligned} \|f\|_{\tilde{\mathcal{D}}}^2 &= |f_{00}|^2 + \sum_{k=1}^{\infty} k|f_{k0}|^2 + \sum_{l=1}^{\infty} l|f_{0l}|^2 + \sum_{k,l=1}^{\infty} kl|f_{kl}|^2 \\ &= |f(0, 0)|^2 + \int_{\mathbb{U}} |D_1 f(z, 0)|^2 d\mu_1(z) \\ &\quad + \int_{\mathbb{U}} |D_2 f(0, w)|^2 d\mu_1(w) \\ &\quad + \int_{\mathbb{U}^2} |D_1 D_2 f(z, w)|^2 d\mu_2(z, w). \end{aligned}$$

This norm is \mathcal{U} -invariant, but not \mathcal{M} -invariant; in fact, none of its first three terms is preserved under compositions with Φ_r .

Proof of Theorem A. First, $\langle\langle z^\alpha, z^\beta \rangle\rangle = 0$ if $\alpha \neq \beta$. To see this, assume, without loss of generality, $\alpha_1 \neq \beta_1$. Let ω be an irrational multiple of π and consider the rotation $R_\omega(z) = (e^{i\omega} z_1, z_2, \dots, z_n)$. By the \mathcal{M} -invariance of $\langle\langle \cdot, \cdot \rangle\rangle$,

$$\begin{aligned} \langle\langle z^\alpha, z^\beta \rangle\rangle &= \langle\langle z^\alpha \circ R_\omega, z^\beta \circ R_\omega \rangle\rangle \\ &= \langle\langle e^{i\alpha_1 \omega} z_1^{\alpha_1} z_2^{\alpha_2} \dots z_n^{\alpha_n}, e^{i\beta_1 \omega} z_1^{\beta_1} z_2^{\beta_2} \dots z_n^{\beta_n} \rangle\rangle \\ &= e^{i(\alpha_1 - \beta_1)\omega} \langle\langle z^\alpha, z^\beta \rangle\rangle, \end{aligned}$$

and the desired orthogonality result follows.

Put $C_\alpha = \langle\langle z^\alpha, z^\alpha \rangle\rangle$. Note that C_α is defined only for $\alpha \in \mathbb{N}^n$. If β is another multi-index and $\beta = \sigma(\alpha)$ for some $\sigma \in \mathcal{S}_n$, then by the \mathcal{M} -invariance of $\langle\langle \cdot, \cdot \rangle\rangle$ again, $C_\alpha = C_\beta$.

Now let $0 < r_j < 1$, $\Psi = (\phi_{r_1}, \dots, \phi_{r_n})$, and consider $f(z) = \prod_{j=1}^n (1 - r_j z_j) \in \mathcal{H}$. Then since z^α is orthogonal to z^β for $\alpha \neq \beta$,

we get

$$\begin{aligned} \langle\langle f, f \rangle\rangle &= \left\| \prod_{j=1}^n (1 - r_j z_j) \right\|^2 = \left\| 1 - \sum_{j=1}^n r_j z_j + \dots + (-1)^m \prod_{j=1}^n r_j z_j \right\|^2 \\ &= C_{(0, \dots, 0)} + \left(\sum_{j=1}^n r_j^2 \right) C_{(1, 0, \dots, 0)} \\ &\quad + \left(\sum_{l>j=1}^n r_j^2 r_l^2 \right) C_{(1, 1, 0, \dots, 0)} + \dots + \left(\prod_{j=1}^n r_j^2 \right) C_{(1, \dots, 1)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} (f \circ \Psi)(z) &= \prod_{j=1}^n \left(1 - r_j \frac{r_j - z_j}{1 - r_j z_j} \right) = \prod_{j=1}^n \frac{1 - r_j^2}{1 - r_j z_j} \\ &= \prod_{j=1}^n \left[(1 - r_j^2) \sum_{\alpha_j=0}^{\infty} r_j^{\alpha_j} z_j^{\alpha_j} \right] = \left(\prod_{j=1}^n (1 - r_j^2) \right) \left(\sum_{\alpha} r^{\alpha} z^{\alpha} \right); \end{aligned}$$

now the density of the polynomials and the axioms of a Hilbert space imply

$$\langle\langle f \circ \Psi, f \circ \Psi \rangle\rangle = \left(\prod_{j=1}^n (1 - r_j^2) \right)^2 \sum_{\alpha} r^{2\alpha} C_{\alpha}.$$

Putting $x_j = r_j^2$ and using the \mathcal{M} -invariance of the semi-norm gives

$$\begin{aligned} (2.1) \quad C_{(0, \dots, 0)} &+ \left(\sum_{j=1}^n x_j \right) C_{(1, 0, \dots, 0)} + \left(\sum_{l>j=1}^n x_j x_l \right) C_{(1, 1, 0, \dots, 0)} \\ &+ \dots + \left(\prod_{j=1}^n x_j \right) C_{(1, \dots, 1)} = \left(\prod_{j=1}^n (1 - x_j) \right)^2 \sum_{\alpha} x^{\alpha} C_{\alpha}. \end{aligned}$$

The constant terms $(C_{(0, \dots, 0)})$ cancel, and if we set the coefficients of $x_1, x_1 x_2, \dots$, and $x_1 x_2 \dots x_n$ on either side equal to each other, we obtain, respectively,

$$\begin{aligned} C_{(1, 0, \dots, 0)} &= C_{(1, 0, \dots, 0)} - 2C_{(0, \dots, 0)}, \\ C_{(1, 1, 0, \dots, 0)} &= C_{(1, 1, 0, \dots, 0)} - 2^2 C_{(1, 0, \dots, 0)}, \\ &\vdots \\ C_{(1, \dots, 1)} &= C_{(1, \dots, 1)} - 2^n C_{(1, \dots, 1, 0)}. \end{aligned}$$

These imply $C_{(0, \dots, 0)} = 0$, $C_{(1, 0, \dots, 0)} = 0$, \dots , $C_{(1, \dots, 1, 0)} = 0$. After the elimination of the terms that are zero, (2.1) simplifies to

$$\begin{aligned} \sum_{\alpha} x^{\alpha} C_{\alpha} &= \frac{x_1 \cdots x_n C_{(1, \dots, 1)}}{(1-x_1)^2 \cdots (1-x_n)^2} \\ &= C_{(1, \dots, 1)} \left(\sum_{\alpha_1=1}^{\infty} \alpha_1 x_1^{\alpha_1} \right) \cdots \left(\sum_{\alpha_n=1}^{\infty} \alpha_n x_n^{\alpha_n} \right); \end{aligned}$$

and this implies

$$C_{\alpha} = \alpha_1 \cdots \alpha_n C_{(1, \dots, 1)} \quad (\alpha \in \mathbb{Z}_+^n).$$

Thus the norm of a monomial of fewer than n variables is zero. Since the polynomials are dense in \mathcal{H} , the same result is true for any $f \in \mathcal{H}$ whose Taylor expansion consists of monomials depending on fewer than n variables. But $C_{(1, \dots, 1)} \neq 0$, because otherwise, since the polynomials are dense in \mathcal{H} , $\langle \cdot, \cdot \rangle$ would be identically zero contrary to hypothesis. Then renaming $C_{(1, \dots, 1)} = C$ completes the proof. \square

Proof of Theorem B. We will only show how the two-variable case is obtained from the one-variable case. This then can be adapted to prove by induction the case for arbitrary \mathbb{U}^n . Unless explicitly stated, subscripted C 's will denote positive constants that are independent of any parameters.

Step 1. We begin by introducing two other semi-inner products on \mathcal{H} . For $f, g \in \mathcal{H}$, let

$$[f, g] = \int_{\mathbb{T}^2} \langle f \circ R_{\theta}, g \circ R_{\theta} \rangle d\lambda_2(\theta)$$

and

$$\langle f, g \rangle = m \langle f \circ \Phi, g \circ \Phi \rangle,$$

where m is an invariant mean (see [2]) on the abelian subgroup

$$\mathcal{N} = \{ \Phi = (\phi_s, \phi_t) : 0 \leq s, t < 1 \}$$

of \mathcal{M} . To actually make \mathcal{N} abelian, in this proof we change our definition of a Möbius transformation so that $\phi_w(z)$ is the negative of what is given in (1.4). The earlier definition was adopted to make ϕ_w an involution, since it simplified calculations involving ϕ_w^{-1} . The required boundedness condition for the existence of this nonunique

mean is furnished by (1.3). Rotations and \mathcal{N} , along with \mathcal{S}_2 , suffice to generate \mathcal{M} , by Lemma 2.1.

Now $[\cdot, \cdot]$ is rotation-invariant

$$(2.2) \quad [f, g] = [f \circ R_\theta, g \circ R_\theta] \quad (\theta \in [-\pi, \pi]^n),$$

and $\langle \cdot, \cdot \rangle$ is \mathcal{N} -invariant

$$(2.3) \quad \langle f, g \rangle = \langle f \circ \Phi, g \circ \Phi \rangle \quad (\Phi \in \mathcal{N}).$$

Moreover, (1.3) implies

$$(2.4) \quad \delta^2 \|f\|^2 \leq [f, f] \leq \frac{1}{\delta^2} \|f\|^2 \quad (f \in \mathcal{H}),$$

and

$$(2.5) \quad \delta^2 \|f\|^2 \leq \langle f, f \rangle \leq \frac{1}{\delta^2} \|f\|^2 \quad (f \in \mathcal{H});$$

and combining these two, we further obtain

$$(2.6) \quad \delta^4 [f, f] \leq \langle f, f \rangle \leq \frac{1}{\delta^4} [f, f] \quad (f \in \mathcal{H}).$$

(2.4) and (2.5) show that the semi-norms associated to $[\cdot, \cdot]$ and $\langle \cdot, \cdot \rangle$ are both equivalent to $\|\cdot\|$.

As in the proof of Theorem A, the rotation-invariance of $[\cdot, \cdot]$ gives the orthogonality condition

$$(2.7) \quad [z^{k_1} w^{l_1}, z^{k_2} w^{l_2}] = 0 \quad ((k_1, l_1) \neq (k_2, l_2)),$$

which leads to

$$[f, f] = \sum_{k, l=0}^{\infty} |f_{kl}|^2 [z^k w^l, z^k w^l] \quad (f \in \mathcal{H}).$$

Therefore, to prove the theorem, it suffices to show that

$$K_1 kl \leq [z^k w^l, z^k w^l] \leq K_2 kl \quad ((k, l) \in \mathbb{N}^2)$$

or, equivalently,

$$K_3 kl \leq \langle z^k w^l, z^k w^l \rangle \leq K_4 kl \quad ((k, l) \in \mathbb{N}^2)$$

for some positive constants K_1 , K_2 , K_3 , and K_4 . Clearly $K_2 \geq K_1$ and $K_4 \geq K_3$.

Step 2. Claim:

$$(2.8) \quad \langle z^k w^l, z^k w^l \rangle \neq 0 \quad \text{if } k \geq 1 \text{ and } l \geq 1.$$

Suppose it is zero for some (N, M) ; then $[z^N w^M, z^N w^M] = 0$ also. Then for $0 \leq s, t < 1$, if we use (2.3), (2.6) and power series expansion

$$\begin{aligned} 0 &= \langle z^N w^M, z^N w^M \rangle \\ &= \left\langle \left(\frac{z-s}{1-sz} \right)^N \left(\frac{w-t}{1-wt} \right)^M, \left(\frac{z-s}{1-sz} \right)^N \left(\frac{w-t}{1-wt} \right)^M \right\rangle \\ &\geq \delta^8 \left[\left(\frac{z-s}{1-sz} \right)^N \left(\frac{w-t}{1-wt} \right)^M, \left(\frac{z-s}{1-sz} \right)^N \left(\frac{w-t}{1-wt} \right)^M \right] \\ &= \left[\sum_{k=0}^{\infty} c_{kN}(s) z^k \sum_{l=0}^{\infty} c'_{lM}(t) w^l, \sum_{k=0}^{\infty} c_{kN}(s) z^k \sum_{l=0}^{\infty} c'_{lM}(t) w^l \right] \\ &= \sum_{k,l=0}^{\infty} |c_{kN}(s)|^2 |c'_{lM}(t)|^2 [z^k w^l, z^k w^l]. \end{aligned}$$

A tedious computation shows that the coefficients $c_{kN}(s) \neq 0$ for any k, N , and s as given above; the same is obviously true for $c'_{lM}(t)$. Thus $[z^k w^l, z^k w^l] = 0$. This means that every element in \mathcal{H} has zero norm and contradicts our basic assumption that $\langle \cdot, \cdot \rangle$ is not identically zero. Hence the claim is proved.

The one-variable result can be stated as

$$(2.9) \quad C_1 k \leq \langle z^k, z^k \rangle_1 \leq C_2 k \quad (k \in \mathbb{N}).$$

It is a consequence of condition (ii)' of Definition 1.1 and of (2.8) that the subspace of \mathcal{H} consisting of functions whose Taylor series expansion at 0 depend only on z is closed. Then (2.9) implies that, for fixed $M \in \mathbb{N}$,

$$(2.10) \quad C_3 k \leq \langle z^k w^M, z^k w^M \rangle \leq C_4 k \quad (k \in \mathbb{N}),$$

and we have a similar equation when the power of z is held constant. Of course, the constants C_3 and C_4 are different for different M . It is our aim to find their explicit dependence on M . If we had only finitely many M , we could pick C_3 and C_4 independently of M and the proof would be over. In the sequel, whenever we have only finitely many n or M , we will use this fact without further reference.

Step 3 (upper bound). Let $M \in \mathbb{N}$ be fixed and $k, j \in \mathbb{N}$. Put

$$\begin{aligned} \alpha_{kM}^{jM} &= \langle z^k w^M, z^j w^M \rangle, & \beta_{kM} &= \langle z^k w^M, z^k w^M \rangle, \\ b_{kM} &= [z^k w^M, z^k w^M]. \end{aligned}$$

By (2.3), $\alpha_{kM}^{jM} = \langle \phi_s^k w^M, \phi_s^j w^M \rangle$ for any $s \in [0, 1)$. Differentiate both sides of this equality with respect to s and set $s = 0$. Then take $k = j + 1$ and add the resulting expressions from $j = 0$ to $j = N \geq 1$; and finally divide both sides by $N + 1$. The result is

$$(2.11) \quad \frac{\beta_{N+1,M}}{N+1} = \frac{2S_{NM}}{N(N+1)} - \frac{\alpha_{N+2,M}^{NM}}{N},$$

where $S_{NM} = \frac{1}{2}\beta_{0M} + \sum_{k=1}^N \beta_{kM}$. In particular, $\alpha_{N+2,M}^{NM}$ is real. Now using (2.3) and (2.5), and letting $s^2 = \frac{N}{N+1}$, we obtain

$$\begin{aligned} \frac{1}{\delta^4}\beta_{1M} &= \frac{1}{\delta^4}\langle \phi_s w^M, \phi_s w^M \rangle \geq [\phi_s w^M, \phi_s w^M] \\ &= s^2 b_{0M} + (1 - s^2)^2 \sum_{k=0}^{\infty} s^{2k} b_{k+1,M} \\ &\geq \frac{\delta^4}{(N+1)^2} \sum_{k=0}^N \left(\frac{N}{N+1}\right)^k \beta_{k+1,M} \geq \frac{\delta^4}{(N+1)^2 e} \sum_{k=0}^N \beta_{k+1,M}, \end{aligned}$$

which implies

$$S_{NM} \leq \left(\frac{e\beta_{1M}}{\delta^8} + \frac{\beta_{0M}}{2}\right)N^2.$$

Using (2.10) twice on the right side gives

$$(2.12) \quad S_{NM} \leq C_5 M N^2.$$

It is this inequality and its pair (2.15) below that allow us to pass from one variable to several variables.

As a special case, when $M = 0$, we get $S_{N0} = 0$ for all $N \geq 1$. Symmetric nature of the calculation shows also $S_{0N} = 0$ for all $N \geq 1$. It follows that

$$(2.13) \quad \beta_{N0} = \langle z^N, z^N \rangle = 0 \quad \text{and} \quad \beta_{0N} = \langle w^N, w^N \rangle = 0 \quad (N \in \mathbb{Z}_+).$$

The inequality

$$\delta^8(\beta_{NM} + \beta_{N+2,M}) \leq \langle (z^N + z^{N+2})w^M, (z^N + z^{N+2})w^M \rangle$$

is a direct consequence of (2.6) and (2.7). Using this, after some routine calculation, (2.11) can be written as

$$\begin{aligned} 2\frac{\beta_{N+1,M}}{N+1} &\leq \frac{4S_{NM}}{N(N+1)} + \frac{2\beta_{N+2,M}}{N(N+2)} + (1 - \delta^8) \left(\frac{\beta_{NM}}{N} + \frac{\beta_{N+2,M}}{N+2}\right) \\ &\leq 6C_5 M + (1 - \delta^8) \left(\frac{\beta_{NM}}{N} + \frac{\beta_{N+2,M}}{N+2}\right), \end{aligned}$$

which is equivalent to $2\gamma_{N+1, M} \leq (1 - \delta^8)(\gamma_{NM} + \gamma_{N+2, M})$ if we let

$$\gamma_{NM} = \frac{\beta_{NM}}{N} - \frac{3C_5M}{\delta^8}.$$

A result in [1] shows $\gamma_{N+1, M} \leq |\gamma_{NM}|$ for positive N . Then using (2.8), we get

$$\frac{\beta_{NM}}{N} - \frac{3C_5M}{\delta^8} \leq \left| \frac{\beta_{1M}}{1} - \frac{3C_5M}{\delta^8} \right| \leq C_6M.$$

Multiplying both sides by M , we conclude

$$\beta_{NM} \leq \left(C_6 + \frac{3C_5}{\delta^8} \right) NM = K_4 NM,$$

which holds for $N \geq 1$ and $M \geq 0$, and for $M \geq 1$ and $N \geq 0$, by the symmetry of the computation.

Step 4 (lower bound). If we combine the result of Step 1 with (2.6), we also get

$$(2.14) \quad b_{kM} \leq \frac{K_4}{\delta^4} kM.$$

We use (2.6), (2.7), (2.3), (2.14), and take $s^2 = \frac{N}{N+1}$ to calculate

$$\begin{aligned} \delta^8 s^2 b_{1M} &\leq \delta^8 [(1 + sz)w^M, (1 + sz)w^M] \\ &= \delta^4 \langle (1 + s\phi_s)w^M, (1 + s\phi_s)w^M \rangle \\ &\leq (1 - s^2)^2 \left[\frac{w^M}{1 - sz}, \frac{w^M}{1 - sz} \right] = \left(\frac{1}{N + 1} \right)^2 \sum_{k=0}^{\infty} s^{2k} b_{kM} \\ &\leq \frac{1}{(N + 1)^2} \left(\sum_{k=0}^{mN} b_{kM} + \frac{K_2}{\delta^4} M \sum_{k=mN+1}^{\infty} k \left(\frac{N}{N + 1} \right)^k \right), \end{aligned}$$

where m will be determined shortly. After approximating the second sum by an integral, we have

$$\delta^8 \left(\frac{N}{N + 1} \right) b_{1M} \leq \frac{1}{(N + 1)^2} \sum_{k=0}^{mN} b_{kM} + \frac{K_2}{\delta^4} M(m + 1)e^{-m}.$$

Because of (2.10), M in the last term can be replaced with $C_8 b_{1M}$. Now choose m so large that $K_2 \delta^{-4} (m + 1) e^{-m} \leq \delta^8 / 3$. Using (2.10) once again and some simplification yields

$$(2.15) \quad \sum_{k=0}^N b_{kM} \geq C_9 M N^2,$$

which is the reverse inequality for (2.12). Combining (2.15) with (2.11), we obtain

$$(2.16) \quad \delta^4 C_9 M \leq \frac{\beta_{N+1, M}}{N+1} + \left(\frac{\beta_{NM}}{N}\right)^{1/2} \left(\frac{\beta_{N+2, M}}{N+2}\right)^{1/2}.$$

Now let

$$\phi_s^N(z) = \left(\frac{z-s}{1-sz}\right)^N = \sum_{k=0}^{\infty} c_{kN}(s) z^k$$

and consider

$$\begin{aligned} b_{NM} &\geq \delta^4 \beta_{NM} = \delta^4 \langle \phi_s^N w^M, \phi_s^N w^M \rangle \geq \delta^8 [\phi_s^N w^M, \phi_s^N w^M] \\ &= \delta^8 \sum_{k=1}^{\infty} |c_{kN}(s)|^2 b_{kM} \geq \delta^8 |c_{N+1, N}(s)|^2 b_{N+1, M}. \end{aligned}$$

A calculation in [1] shows that $|c_{N+1, N}(s)|^2 \geq 1/2$ for all $s \in [0, 1)$ and $N \geq 1$. Thus there is a constant C_{10} such that

$$(2.17) \quad \beta_{N+1, M} \leq C_{10} \beta_{NM}.$$

Now (2.16) and (2.17) together imply

$$\beta_{NM} \geq \left(\frac{\delta^4 C_9}{4C_{10}}\right) NM = K_3 NM,$$

for $N \geq 1$ and $M \geq 0$, and for $M \geq 1$ and $N \geq 0$.

Step 5. The only term we have not yet accounted for is $\beta_{00} = \langle 1, 1 \rangle$. Since it represents a one-dimensional subspace of \mathcal{H} , we now know $\mathcal{H} = \mathcal{D}(\mathbb{U}^2)$. To complete the proof, we will also show $\beta_{00} = b_{00} = 0$. To obtain a contradiction, suppose $b_{00} = [1, 1] \neq 0$. Let $f \in \mathcal{H}$, Φ be a Möbius transformation, and denote the power series coefficients of f and $f \circ \Phi$ by f_{kl} and f'_{kl} , respectively. Because of (2.13)

$$\|f\|^2 \geq \delta^2 [f, f] = \delta^2 \left(b_{00} |f(0, 0)|^2 + \sum_{k, l=1}^{\infty} |f_{kl}|^2 b_{kl} \right)$$

and

$$[f \circ \Phi, f \circ \Phi] = b_{00} |f(\Phi(0, 0))|^2 + \sum_{k, l=1}^{\infty} |f'_{kl}|^2 b_{kl}.$$

We have

$$\begin{aligned} \sum_{k, l=1}^{\infty} |f'_{kl}|^2 b_{kl} &\geq K_1 \sum_{k, l=1}^{\infty} |f_{kl}|^2 kl = K_1 \|f \circ \Phi\|_{\mathcal{D}}^2 = K_1 \|f\|_{\mathcal{D}}^2 \\ &= K_1 \sum_{k, l=1}^{\infty} |f_{kl}|^2 kl \geq \frac{K_1}{K_2} \sum_{k, l=1}^{\infty} |f_{kl}|^2 b_{kl}. \end{aligned}$$

Hence

$$\begin{aligned} b_{00}|f(0, 0)|^2 + \sum_{k,l=1}^{\infty} |f_{kl}|^2 b_{kl} &= [f, f] \geq \delta^8 [f \circ \Phi, f \circ \Phi] \\ &= \delta^8 \left(b_{00} |f(\Phi(0, 0))|^2 + \sum_{k,l=1}^{\infty} |f'_{kl}|^2 b_{kl} \right) \\ &= \delta^8 \left(b_{00} |f(\Phi(0, 0))|^2 + \frac{K_1}{K_2} \sum_{k,l=1}^{\infty} |f_{kl}|^2 b_{kl} \right), \end{aligned}$$

from which we obtain

$$b_{00}|f(0, 0)|^2 + \left(1 - \delta^8 \frac{K_1}{K_2}\right) \sum_{k,l=1}^{\infty} |f_{kl}|^2 b_{kl} \geq \delta^8 b_{00} |f(\Phi(0, 0))|^2.$$

The left hand side of this equation is finite since it is equivalent to $\|f\|^2$. Since $\Phi(0, 0)$ can be any point in \mathbb{U}^2 , it follows that every element of \mathcal{H} , i.e., of $\mathcal{D}(\mathbb{U}^2)$, is bounded. But a Dirichlet space contains unbounded elements. In \mathbb{U} , this is seen most easily by the Area Theorem; in \mathbb{U}^2 , we take an unbounded function depending only on one variable. Therefore $b_{00} = 0$ and we are done. \square

COROLLARY 2.3. *Theorem B is true even if (1.3) holds only for $\Psi \in \mathcal{M}^*$. Theorem A is true even if (1.2) holds only for $\Psi \in \mathcal{M}^*$.*

Proof. The proof of Theorem B uses coordinate permutations nowhere. Theorem A is a consequence of Theorem B. \square

3. Boundary behavior. Dirichlet-type spaces. This section requires some new notions that were studied in \mathbb{U} in [3] and [4]. For each $\delta = (\delta_1, \dots, \delta_n)$ with each $0 \leq \delta_j \leq 1$, we define the *Dirichlet-type spaces* $\mathcal{D}_\delta(\mathbb{U}^n)$ to consist of those $f(z) = \sum_\alpha f_\alpha z^\alpha \in \mathcal{H}(\mathbb{U}^n)$ that satisfy

$$\|f\|_{\mathcal{D}_\delta} = \sum_\alpha \alpha_1^{2\delta_1} \cdots \alpha_n^{2\delta_n} |f_\alpha|^2 < \infty.$$

This definition makes sense even if some $\delta_j = 0$ if we interpret $0^0 = 1$. In fact, if all the $\delta_j = 0$, then $\mathcal{D}_\delta(\mathbb{U}^n) = \mathcal{H}^2(\mathbb{U}^n)$. The space corresponding to $\delta_1 = \dots = \delta_n = 1$ consists of functions f with $D_1 \cdots D_n f \in \mathcal{H}^2(\mathbb{U}^n)$. When $\delta_1 = \dots = \delta_n = 1/2$, we have the Dirichlet space. For $n = 1$, all Dirichlet-type spaces are contained in $\mathcal{H}^2(\mathbb{U})$, but this is not true if $n > 1$.

Some subclasses of $\mathcal{D}_\delta(\mathbb{U}^n)$ have certain integral representations: If $F \in \mathcal{L}^2(\mathbb{T}^n)$, for $0 \leq \delta_j < 1$ and $z \in \mathbb{U}^n$, set

$$(3.1) \quad f(z) = \int_{\mathbb{T}^n} F(\zeta) \prod_{j=1}^n \frac{1}{(1 - z_j \bar{\zeta}_j)^{1-\delta_j}} d\lambda_n(\zeta).$$

The product is the Cauchy kernel each of whose factors is raised to a fractional power. Omitting j , each factor can be expanded as

$$(1 - z \bar{\zeta})^{\delta-1} = \sum_{\alpha=0}^{\infty} b_\alpha z^\alpha \bar{\zeta}^\alpha$$

where

$$b_\alpha = \frac{\Gamma(1 - \delta + \alpha)}{\Gamma(1 - \delta)\Gamma(1 + \alpha)} \sim \frac{1}{\alpha^\delta}.$$

In particular $b_0 = 1$. Let c_α be the $(\alpha_1, \dots, \alpha_n)$ th Fourier coefficient of F ; i.e.,

$$c_\alpha = \int_{\mathbb{T}^n} \bar{\zeta}^\alpha F(\zeta) d\lambda_n(\zeta).$$

Setting $f_\alpha = b_{\alpha_1} \cdots b_{\alpha_n} c_\alpha$, we get

$$\begin{aligned} f(z) &= \int_{\mathbb{T}^n} \prod_{j=1}^n \sum_{\alpha_j=0}^{\infty} b_{\alpha_j} z_j^{\alpha_j} \bar{\zeta}_j^{\alpha_j} F(\zeta) d\lambda_n(\zeta) \\ &= \sum_{\alpha} b_{\alpha_1} \cdots b_{\alpha_n} z^\alpha \int_{\mathbb{T}^n} \bar{\zeta}^\alpha F(\zeta) d\lambda_n(\zeta) \\ &= \sum_{\alpha} b_{\alpha_1} \cdots b_{\alpha_n} z^\alpha c_\alpha = \sum_{\alpha} f_\alpha z^\alpha. \end{aligned}$$

Now

$$\|f\|_{\mathcal{D}_\delta} = \sum_{\alpha} \alpha_1^{2\delta_1} \cdots \alpha_n^{2\delta_n} |b_{\alpha_1}|^2 \cdots |b_{\alpha_n}|^2 |c_\alpha|^2 \sim \sum_{\alpha} |c_\alpha|^2 < \infty.$$

Hence $f \in \mathcal{D}_\delta(\mathbb{U}^n)$; i.e., any f given by (3.1) is in a Dirichlet-type space.

But not all $f \in \mathcal{D}_\delta(\mathbb{U}^n)$ have integral representations as in (3.1), because a Dirichlet-type space does not control all the power series coefficients of its members. However, we can define a space $\tilde{\mathcal{D}}_\delta(\mathbb{U}^n)$ similar to $\mathcal{D}_\delta(\mathbb{U}^n)$ in which an integral representation is possible. Let's concentrate on the case $n = 2$ again for simplicity. With obvious

notation, $f \in \tilde{\mathcal{D}}_\delta(\mathbb{U}^2)$ if and only if

$$(3.2) \quad \|f\|_{\tilde{\mathcal{D}}_\delta}^2 = |f_{00}|^2 + \sum_{k=1}^\infty k^{2\delta_1} |f_{k0}|^2 + \sum_{l=1}^\infty l^{2\delta_2} |f_{0l}|^2 + \sum_{k,l=1}^\infty k^{2\delta_1} l^{2\delta_2} |f_{kl}|^2 < \infty.$$

Given $f(z) = \sum_{kl} f_{kl} z^k w^l \in \tilde{\mathcal{D}}_\delta(\mathbb{U}^2)$, let $c_{kl} = f_{kl}/b_k b_l$ if $(k, l) \in \mathbb{N}^2$ (recall that $b_0 = 1$), and let $c_{kl} = 0$ otherwise. Then, using (3.2),

$$\sum_{(k,l) \in \mathbb{Z}^2} |c_{kl}|^2 = |c_{00}|^2 + \sum_{k=1}^\infty |c_{k0}|^2 + \sum_{l=1}^\infty |c_{0l}|^2 + \sum_{k,l=1}^\infty |c_{kl}|^2 \sim \|f\|_{\tilde{\mathcal{D}}_\delta}^2 < \infty.$$

Thus there is an $F \in \mathcal{L}^2(\mathbb{T}^2)$ such that $\hat{F}(k, l) = c_{kl}$. Therefore

$$\begin{aligned} f(z) &= \sum_{k,l=0}^\infty f_{kl} z^k w^l = \sum_{k,l=0}^\infty c_{kl} b_k b_l z^k w^l \\ &= \sum_{k,l=0}^\infty b_k b_l z^k w^l \int_{\mathbb{T}^2} \bar{\zeta}_1^k \bar{\zeta}_2^l F(\zeta) d\lambda_2(\zeta) \\ &= \int_{\mathbb{T}^2} \frac{F(\zeta) d\lambda_2(\zeta)}{(1 - \bar{\zeta}_1 z)^{1-\delta_1} (1 - \bar{\zeta}_2 w)^{1-\delta_2}}. \end{aligned}$$

Clearly F is not unique. In fact, c_{kl} can be defined arbitrarily for $(k, l) \notin \mathbb{N}^2$ as long as we retain $\sum_{(k,l) \in \mathbb{Z}^2} |c_{kl}|^2 < \infty$.

Kernels and potentials. From now on, we will also use $e^{i\theta_j}$ for $\zeta_j \in \mathbb{T}$, $e^{i\varphi_j}$ for $\eta_j \in \mathbb{T}$, and $r_j e^{i\theta_j} = r_j \zeta_j$ for $z_j \in \mathbb{U}$. The point $(1, \dots, 1) \in \mathbb{T}^n$ corresponding to $\theta_1 = \dots = \theta_n = 0$ will act like the origin in \mathbb{R}^n . Now the Poisson kernel takes the more familiar form

$$P_r(\theta) = \prod_{j=1}^n \frac{1 - r_j^2}{1 - 2r_j \cos \theta_j + r_j^2},$$

and it is considered as a function of θ indexed by r . So the \mathcal{L}^p -norm of a Poisson integral will be obtained by an integration on the θ -variable and will still depend on r .

A kernel K is a nonnegative \mathcal{L}^1 -function on \mathbb{T}^n which is even and decreasing in each $|\theta_j|$ when the other variables are kept fixed. We will also have $K(1, \dots, 1) = \infty$ and normalize as $\|K\|_1 = 1$. A potential is the convolution of an \mathcal{L}^p -function F on \mathbb{T}^n with a kernel. Thus (3.1) defines $f(z) \in \tilde{\mathcal{D}}_\delta(\mathbb{U}^n)$ as a potential. The Poisson integral is simply the convolution with the Poisson kernel.

Let's define the *Bessel kernels* on the torus. For $0 < \delta_j \leq 1$, let

$$G_\delta(\zeta) = \prod_{j=1}^n g_{\delta_j}(\theta_j) = \prod_{j=1}^n \left(1 + \frac{1}{2} \sum_{\alpha_j \neq 0} |\alpha_j|^{-\delta_j} \zeta_j^{\alpha_j} \right) \\ = \prod_{j=1}^n \left(1 + \sum_{\alpha_j=1}^{\infty} \alpha_j^{-\delta_j} \cos(\alpha_j \theta_j) \right),$$

where each g_{δ_j} is a Bessel kernel on the unit circle. $g_\delta(0) = \infty$, g_δ is a decreasing function of $|\omega|$ for $\omega \neq 0$, $g_\delta(\omega) > 0$, and $g_\delta(-\omega) = g_\delta(\omega)$. Each $g_{\delta_j} \in \mathcal{L}^1(\mathbb{T})$, so $G_\delta \in \mathcal{L}^1(\mathbb{T}^n)$, and $\|G_\delta\|_1 = \prod_{j=1}^n \|g_{\delta_j}\|_1 = 1$. When $0 < \delta < 1$,

$$g_\delta(\omega) \approx \left| \sin \frac{\omega}{2} \right|^{\delta-1} \quad \text{as } \omega \rightarrow 0.$$

Also $g_1(\omega) = 1 - \log|2 \sin(\omega/2)|$. $P_r[g_\delta] = P_r * g_\delta$ is the harmonic extension of g_δ to \mathbb{U} . As $r \rightarrow 1$, it satisfies the following:

$$(3.3) \quad \|P_r * g_\delta\|_q \sim (1-r)^{\delta-1/p} \quad (\delta p < 1),$$

$$(3.4) \quad \|P_r * g_\delta\|_q \sim \left(\log \frac{1}{1-r} \right)^{1/q} \quad (\delta p = 1, p > 1),$$

$$\|P_r * g_1\|_\infty \sim \log \frac{1}{1-r} \quad (\delta = 1, p = 1).$$

$P_r[G_\delta]$ possesses these properties in each variable separately.

On the unit circle, for $0 < \delta < 1$, the *modified Bessel kernels* are

$$\tilde{g}_\delta(\omega) = (1 - e^{i\omega})^{\delta-1} \quad \text{and} \quad \tilde{g}_1(\omega) = \log \frac{1}{1 - e^{i\omega}}.$$

On the torus, let $\tilde{G}_\delta(\zeta) = \prod_{j=1}^n \tilde{g}_{\delta_j}(\theta_j)$. These functions are not positive, so they are not properly kernels, but they are dominated by the Bessel kernels: There are constants $C_\delta > 0$ such that $|\tilde{G}_\delta| \leq C_\delta G_\delta$. If each δ_j is less than 1,

$$P_r[\tilde{G}_\delta](\theta) = \prod_{j=1}^n P_{r_j}[\tilde{g}_{\delta_j}](\theta_j) = \prod_{j=1}^n (1 - z_j)^{\delta_j-1},$$

with a logarithmic term if some $\delta_k = 1$. For $F \in \mathcal{L}^p(\mathbb{T}^n)$, the map that takes F to $G_\delta * F$ is one-to-one, and the Cauchy integral of $\tilde{G}_\delta * F$ is the same as its Poisson integral:

$$P_r[\tilde{G}_\delta * F](\theta) = (P_r * \tilde{G}_\delta * F)(\theta) \\ = \int_{\mathbb{T}^n} F(\zeta) \prod_{j=1}^n \frac{1}{(1 - z_j \bar{\zeta}_j)^{1-\delta_j}} d\lambda_n(\zeta) = f(z).$$

Thus we have obtained the integral in (3.1). From now on, F and f will always be related as in this equation. $\delta_j = 1$ does not give rise to a Cauchy-type integral; so we will not pay any attention to this case any more.

Tangential limits. Define the *tangential approach regions* to the unit circle:

$$(3.5) \quad A_{\gamma,c}(\varphi) = \left\{ re^{i\theta} : 1 - r > c \left| \sin \frac{\theta - \varphi}{2} \right|^\gamma \right\},$$

$$E_{\gamma,c}(\varphi) = \left\{ re^{i\theta} : 1 - r > \exp \left(-c \left| \sin \frac{\theta - \varphi}{2} \right|^{-\gamma} \right) \right\}.$$

$A_{\gamma,c}(\varphi)$ has (polynomial) order of contact γ , and $E_{\gamma,c}(\varphi)$ makes exponential contact, with \mathbb{T} . A function f defined in \mathbb{U} has $A_\gamma(E_\gamma)$ -limit L at $e^{i\varphi}$ if $f(z) \rightarrow L$ as $z \rightarrow e^{i\varphi}$ within $A_{\gamma,c}(E_{\gamma,c})$ for every $c > 0$. In [3], it was shown that Poisson integrals of the modified Bessel potentials have A_γ -limits a.e. on \mathbb{T} if $\delta p < 1$ for $\gamma = \frac{1}{1-\delta p}$, and E_γ -limits a.e. on \mathbb{T} if $\delta p = 1$ for $\gamma = q - 1$.

Let $Q = Q(\eta, s)$ be the cube centered at $\eta \in \mathbb{T}^n$ with sides $s = (s_1, \dots, s_n)$, where each s_j has the same order as $\max\{s_j : 1 \leq j \leq n\} \rightarrow 0$. Its volume is $\lambda_n(Q) = s_1 \cdots s_n$. If $F \in \mathcal{L}^p(\mathbb{T}^n)$, its Hardy-Littlewood maximal function is

$$(M_p F)(\eta) = \sup_{0 < s_1, \dots, s_n \leq 1} \left(\frac{1}{\lambda_n(Q)} \int_Q |f|^p d\lambda_n \right)^{1/p}.$$

M_1 is of weak type $(1, 1)$; and since $M_p F = (M_1 |F|^p)^{1/p}$, M_p is of weak type (p, p) . Thus there are C_p such that

$$\lambda_n(\{M_p F > t\}) \leq \frac{C_p}{t^p} \|F\|_p^p \quad (F \in \mathcal{L}^p(\mathbb{T}^n), t \in (0, \infty)).$$

The proofs of the following assertions are similar to the proofs given in [3] for $n = 1$ and will be omitted. Some of them are valid in more general situations. The first result is obtained using the straightforward inequality

$$\int_{\mathbb{T}^n} |F| G_\delta d\lambda_n \leq (M_1 F)(1, \dots, 1) \int_{\mathbb{T}^n} G_\delta d\lambda_n = (M_1 F)(1, \dots, 1),$$

which holds for any $F \in \mathcal{L}^1(\mathbb{T}^n)$, and whose proof is also in [3].

THEOREM 3.1. *There is a $C_p < \infty$ such that for $F \in \mathcal{L}^p(\mathbb{T}^n)$ and $\zeta, \eta \in \mathbb{T}^n$,*

$$|(G_\delta * F)(\zeta)| \leq C_p \left[(M_p F)(\eta) \left(\prod_{j=1}^n |\zeta_j - \eta_j|^{1/p} \right) \|G_\delta\|_q + (M_1 F)(\eta) \right].$$

Convolution of two kernels is a kernel; so Theorem 3.1 holds with $P_r * G_\delta = P_r[G_\delta]$ in place of G_δ , which has the desired properties $\|P_r * G_\delta\|_1 = \|P_r\|_1 \|G_\delta\|_1 = 1$ and $\|P_r * G_\delta\|_q = \prod_{j=1}^n \|P_{r_j} * g_{\delta_j}\|_q$. The Hölder inequality gives $M_1 F \leq M_p F$. In addition, $|\zeta_j - \eta_j|$ can be replaced by $|\theta_j - \varphi_j|$, or even by $|\sin((\theta_j - \varphi_j)/2)|$, since they are all of the same order as $\zeta \rightarrow \eta$. Lastly, we can put \tilde{G}_δ in place of G_δ on the left side of the inequality since the latter dominates the former. Hence Theorem 3.1 yields

THEOREM 3.2. *If $F \in \mathcal{L}^p(\mathbb{T}^n)$, then, using $z = re^{i\theta}$, for all $e^{i\theta}$, $e^{i\varphi} \in \mathbb{T}^n$,*

$$(3.6) \quad |f(z)| \leq C_p (M_p F)(\varphi) \left[1 + \prod_{j=1}^n \left| \sin \frac{\theta_j - \varphi_j}{2} \right|^{1/p} \|P_{r_j} * g_{\delta_j}\|_q \right].$$

For given φ , any bound on the product on the right side gives a bound on $f(z)$. This leads us to the tangential approach regions to \mathbb{T}^n . So fix an $\eta \in \mathbb{T}^n$. As $z \rightarrow \zeta$, all $r_j \rightarrow 1$; and because of (3.3) and (3.4), $\|P_r * G_\delta\|_q \sim \prod_{j=1}^n b_j$, where $b_j = (1 - r_j)^{\delta_j - 1/p}$ or $b_j = (\log(1/(1 - r_j)))^{1/q}$ depending on whether $\delta_j p < 1$ or $\delta_j p = 1$, respectively. In other words, an approach region should be determined by

$$\prod_{j=1}^n b_j \left| \sin \frac{\theta_j - \varphi_j}{2} \right|^{1/p} < c.$$

So define $B_{\gamma,c}(\eta)$ by

$$(3.7) \quad B_{\gamma,c}(\eta) = B_{\gamma,c}(\varphi) = \left\{ z \in \mathbb{U}^n : \prod_{j=1}^n b_j^{-1/\gamma_j} \left| \sin \frac{\theta_j - \varphi_j}{2} \right| < c \right\}.$$

Each of the factors in the above product is related to one of the regions in (3.5). In particular, points in a cartesian product of one-dimensional approach regions such as $B_{\gamma_1,c_1}(\varphi_1) \times \dots \times B_{\gamma_n,c_n}(\varphi_n)$, where each $B_{\gamma_j,c_j}(\varphi_j)$ is either $A_{\gamma_j,c_j}(\varphi_j)$ or $E_{\gamma_j,c_j}(\varphi_j)$, satisfy the criterion for being in $B_{\gamma,c}(\eta)$. Hence an approach region can make exponential contact with \mathbb{T}^n in one (complex) direction and polynomial contact in another. For $\eta \in \mathbb{T}^n$, the *maximal functions* associated to these approach regions are defined as

$$(M_{\tilde{G}_\delta, \gamma, c f})(\eta) = \sup \{ |f(z)| : z \in B_{\gamma,c}(\eta) \}.$$

A function f defined in \mathbb{U}^n is said to have B_γ -limit L at $\eta \in \mathbb{T}^n$ if $f(z) \rightarrow L$ as $z \rightarrow \eta$ within $B_{\gamma,c}(\eta)$ for every $c > 0$.

THEOREM 3.3. *If $F \in \mathcal{L}^p(\mathbb{T}^n)$ and $t \in (0, \infty)$, there is a $C = C(p, c)$ such that*

$$\lambda_n(\{M_{G_\delta, \gamma, c} f > t\}) < \frac{C}{t^p} \|F\|_p^p,$$

where

$$(3.8) \quad \gamma_j = \frac{1}{1 - \delta_j p} \quad \text{if } \delta_j p < 1 \quad \text{and} \quad \gamma_j = q - 1 \quad \text{if } \delta_j p = 1.$$

This follows from the weak-type- (p, p) estimate for M_p . The weak-type estimate gives rise to a convergence theorem via classical arguments. This is the content of part (i) of Theorem C. The case $p = 1$ of part (ii) of that theorem also follows from Theorem 3.3. Henceforth, p, δ , and γ will always be related by (3.8).

Capacities. For $E \subset \mathbb{T}^n$, let $T(G_\delta, p, E)$ be the set of all nonnegative $F \in \mathcal{L}^p(\mathbb{T}^n)$ such that $(G_\delta * F)(\zeta) \geq 1$ for all $\zeta \in E$. The p -capacity of E is

$$\Sigma_{G_\delta}(E) = \inf\{\|F\|_p^p : F \in T(G_\delta, p, E)\}.$$

$\Sigma_{G_\delta}(E) = 0$ implies $\lambda_n(E) = 0$. The functions $G_\delta * F$ are defined Σ_{G_δ} -almost everywhere. If $F \in T(P_r * G_\delta, p, E)$, then $\Sigma_{P_r}(E) \leq \Sigma_{P_r * G_\delta}(E)$. For $\eta \in \mathbb{T}^n$ and fixed $\rho > 1$, let

$$\Gamma(\eta) = \{z \in \mathbb{U}^n : |z_j - \eta_j| < \rho(1 - |z_j|), 1 \leq j \leq n\},$$

and set $S(E) = \mathbb{U}^n \setminus \bigcup_{\eta \notin E} \Gamma(\eta)$. $\Gamma(\eta)$ is the cartesian product of n sets each of which is asymptotic, as $z_j \rightarrow \eta_j$, to an angle-shaped approach region in \mathbb{U} with vertex at η_j . For $\eta \in \mathbb{T}^n$, the *nontangential maximal function* is

$$(Nf)(\eta) = \sup\{|f(z)| : z \in \Gamma(\eta)\}.$$

For $W \subset \mathbb{U}^n$, $J_c^\gamma(W)$ is the set of $\eta \in \mathbb{T}^n$ for which W intersects $B_{\gamma, c}(\eta)$.

LEMMA 3.4. *There exists a constant $b = b(n) > 0$ such that if $F \geq 0$ on \mathbb{T}^n , $F \geq 1$ on $E \in \mathbb{T}^n$, and $z \in S(E)$, then $P_r[F](z) > b$.*

We will use this lemma with $G_\delta * F$ in place of F . It leads to the following lower estimate for capacities.

PROPOSITION 3.5. *If $F \in \mathcal{L}^1(\mathbb{T}^n)$ and $0 < t < \infty$, then $\{M_{G_\delta, \gamma, cf} > t\}$ is contained in $J_c^\gamma(S(\{Nf > t\}))$. Thus, there is a $C = C(p, c)$ such that if $t \in (0, \infty)$ and $F \in \mathcal{L}^1(\mathbb{T}^n)$, we have*

$$\lambda_n(\{M_{G_\delta, \gamma, cf} > t\}) \leq C \Sigma_{G_\delta}(\{Nf > t\}).$$

THEOREM 3.6. *For $1 < p < \infty$, there is a constant $C_p < \infty$ such that if $F \in \mathcal{L}^p(\mathbb{T}^n)$ and $F \geq 0$, then*

$$\int_0^\infty \Sigma_{G_\delta}(\{G_\delta * F > t\}) d(t^n) \leq C_p \|F\|_p^p.$$

Combining Lemma 3.4, Proposition 3.5, and Theorem 3.6 with the fact that G_δ dominates \tilde{G}_δ and that $N(G_\delta * F) = G_\delta * NF$, we obtain the strong-type estimates in part (ii) of Theorem C.

THEOREM C. *Let $1 \leq p < \infty$, $F \in \mathcal{L}^p(\mathbb{T}^n)$, $0 < \delta_j < 1$, define f as in (3.1), pick γ_j as in (3.8), and for $\zeta \in \mathbb{T}^n$, construct $B_{\gamma, c}(\zeta)$ as in (3.7).*

- (i) *The B_γ -limit of f exists a.e. $[\lambda_n]$ on \mathbb{T}^n*
- (ii) *There are positive constants C_p such that*

$$\begin{aligned} \|M_{G_\delta, \gamma, cf}\|_p &\leq C_p \|F\|_p && (1 < p < \infty), \\ \lambda_n(\{M_{G_\delta, \gamma, cf} > t\}) &\leq \frac{C_1}{t} \|F\|_1 && (p = 1, 0 < t < \infty). \end{aligned}$$

If $\zeta \in \partial\mathbb{U}^n \setminus \mathbb{T}^n$, then only one component of ζ , say the n th, has $|\zeta_n| = 1$. Then the first $n - 1$ factors in the product in (3.6) are bounded as $z \rightarrow \zeta$. So in this case, it suffices to apply the one-variable result in the n th variable. The approach regions are restricted only in the n th component as in (3.5), and (z_1, \dots, z_{n-1}) can approach $(\zeta_1, \dots, \zeta_{n-1}) \in \partial\mathbb{U}^n$ in any manner whatsoever. Theorem C remains valid except that in part (ii), we would use one-dimensional norm and Lebesgue measure.

When $p = 2$ and all the $\delta_j = 1/2$, this theorem takes care of $\tilde{\mathcal{D}}(\mathbb{U}^n)$, but cannot deal with $\mathcal{D}(\mathbb{U}^n)$. Thus the functions in the modified Dirichlet space have tangential ($B_{(1, \dots, 1)}$ -) limits at almost every boundary point of the unit polydisc. When $n = 1$, since $\mathcal{H}^2(\mathbb{U})$ includes all Dirichlet-type spaces, elements of $\mathcal{D}(\mathbb{U})$ have nontangential limits a.e. on \mathbb{T} .

Now we will look at the size of the exceptional sets. From Lemma 3.4, part (ii) of Theorem C, and the first part of Proposition 3.5, we obtain

LEMMA 3.7. *If $1 < p < \infty$, then for some $C = C(p, c)$*

$$\Sigma_{P_r}(J_c^\gamma(S(E))) \leq C \Sigma_{P_r * G_\delta}(E).$$

Hence for $F \in \mathcal{L}^p(\mathbb{T}^n)$,

$$\Sigma_{P_r}(\{M_{G_\delta, \gamma, c} f > t\}) \leq C \Sigma_{P_r * G_\delta}(\{Nf > t\}).$$

THEOREM 3.8. *If $1 < p < \infty$ and $F \in \mathcal{L}^p(\mathbb{T}^n)$, then*

$$\int_0^\infty \Sigma_{P_r}(\{M_{G_\delta, \gamma, c} f > t\}) d(t^n) \leq C \|F\|_p^p,$$

and thus

$$\Sigma_{P_r}(\{M_{G_\delta, \gamma, c} f > t\}) \leq C \frac{\|F\|_p^p}{t^p}.$$

This theorem is an analog of Theorem C in the language of capacities and proved similarly.

THEOREM D. *Let $1 < p < \infty$, $F \in \mathcal{L}^p(\mathbb{T}^n)$, and f be as in (3.1).*

(i) *There is a set $E_1 \subset \mathbb{T}^n$ with $\Sigma_{P_r * G_\delta}(E_1) = 0$ such that the nontangential limit of f exists at every point of $\mathbb{T}^n \setminus E_1$.*

(ii) *There is a set $E_2 \subset \mathbb{T}^n$ with $\Sigma_{P_r}(E_2) = 0$ such that the B_γ -limit of f exists at every point of $\mathbb{T}^n \setminus E_2$.*

This result is a consequence of the basic properties of capacities and Theorem 3.10. For points on $\partial\mathbb{U}^n \setminus \mathbb{T}^n$, the one-variable result can again be used to reach a similar conclusion. Hence if $p = 2$ and all the $\delta_j = 1/2$, the points on $\partial\mathbb{U}^n$ where the modified Dirichlet space does not have nontangential limits have zero capacity in some sense.

REFERENCES

- [1] J. Arazy and S. D. Fisher, *The uniqueness of the Dirichlet space among Möbius-invariant Hilbert spaces*, Illinois J. Math., **29** (1985), 449–462.
- [2] E. Hewitt and K. Ross, *Abstract Harmonic Analysis*, I, 2nd ed., Grundlehren Math. Wiss., vol. 115, Springer, Berlin, 1979.
- [3] A. Nagel, W. Rudin and J. H. Shapiro, *Tangential boundary behavior of functions in Dirichlet-type spaces*, Ann. of Math., **116** (1982), 331–360.
- [4] W. Rudin, *The variation of holomorphic functions on tangential boundary curves*, Michigan Math. J., **32** (1985), 41–45.

- [5] K. Zhu, *Möbius-invariant Hilbert spaces of holomorphic functions in the unit ball of \mathbb{C}^n* , Trans. Amer. Math. Soc., **323** (1991), 823–842.

Received May 11, 1992. Part of this work represents a part of the author's Ph.D. thesis completed under the supervision of Walter Rudin. The author wishes to thank Walter Rudin and Frank Forelli for several useful discussions. Supported under TBAG-ÇG2 by TÜB.İTAK.

MATHEMATICS DEPARTMENT
UNIVERSITY OF WISCONSIN
MADISON, WI 53706

Current address: Mathematics Department
Middle East Technical University
Ankara 06531 Turkey

E-mail address: mathtk@vm.cc.metu.edu.tr

CONTENTS

D. Bisch, A note on intermediate subfactors	201
M. J. Carro and J. Soria, Tent spaces over general approach regions and pointwise estimates	217
J. A. Charris, M. E. H. Ismail, and S. Monsalve, On sieved orthogonal polynomials X: general blocks of recurrence relations	237
K. S. Chou and T. Y. H. Wan, Asymptotic radial symmetry for solutions of $\Delta u + e^u = 0$ in a punctured disc	269
M. E. Fogel, Knots with algebraic unknotting number one	277
K. N. Jones, The structure of closed nonpositively curved Euclidean cone 3-manifolds	297
M. Kaneda, On the Frobenius morphism of flag schemes	315
H. T. Kaptanoğlu, Möbius-invariant Hilbert spaces in polydiscs	337
K. Kuribayashi, The cohomology ring of the spaces of loops on Lie groups and homogeneous spaces	361
A. Thompson, A note on Murasugi sums	393

PACIFIC JOURNAL OF MATHEMATICS

Volume 163 No. 2 April 1994

A note on intermediate subfactors	201
DIETMAR BISCH	
Tent spaces over general approach regions and pointwise estimates	217
MARÍA J. CARRO and JAVIER SORIA	
On sieved orthogonal polynomials. X. General blocks of recurrence relations	237
JAIRO A. CHARRIS, MOURAD ISMAIL and SERGIO MONSALVE	
Asymptotic radial symmetry for solutions of $\Delta u + e^u = 0$ in a punctured disc	269
KAI SENG (KAISING) CHOU (TSO) and TOM YAU-HENG WAN	
Knots with algebraic unknotting number one	277
MICAHA ELTON FOGEL	
The structure of closed non-positively curved Euclidean cone 3-manifolds	297
KERRY NELSON JONES	
On the Frobenius morphism of flag schemes	315
MASAHARU KANEDA	
Möbius-invariant Hilbert spaces in polydiscs	337
H. TURGAY KAPTANOGLU	
The cohomology ring of the spaces of loops on Lie groups and homogeneous spaces	361
KATSUHIKO KURIBAYASHI	
A note on Murasugi sums	393
ABIGAIL A. THOMPSON	