

*Pacific
Journal of
Mathematics*

THE L^p THEORY OF STANDARD HOMOMORPHISMS

FEREIDOUN GHAHRAMANI AND SANDY GRABINER

Volume 168 No. 1

March 1995

THE L^p THEORY OF STANDARD HOMOMORPHISMS

F. GHAHRAMANI AND S. GRABINER

Suppose that $\phi : L^1(\omega_1) \rightarrow L^1(\omega_2)$ is a continuous nonzero homomorphism between weighted convolution algebras on R^+ , and let ϕ also designate the extension of this map to the corresponding measure algebras $M(\omega_1)$ and $M(\omega_2)$. For $1 < p < \infty$, we prove: (a) the semigroup $\mu_t = \phi(\delta_t)$ acts as a strongly continuous semigroup on $L^p(\omega_2)$; (b) Whenever $L^1(\omega_1) * f$ is dense in $L^1(\omega_1)$, then $L^p(\omega_2) * \phi(f)$ is dense in $L^p(\omega_2)$; (c) Each h in $L^p(\omega_2)$ can be factored as $h = \phi(f) * g$; (d) ϕ is continuous from the strong operator topology of $M(\omega_1)$ acting on $L^1(\omega_1)$ to the strong operator topology of $M(\omega_2)$ acting on $L^p(\omega_2)$.

1. Introduction. In this paper we show that the L^p analogue of a number of questions we have studied ([10], [8], [11], [7]) involving homomorphisms and semigroups on weighted L^1 spaces on $R^+ = [0, \infty)$ all have positive answers when $1 < p < \infty$. If $\omega(t) > 0$ is a Borel function on R^+ which is locally bounded and locally bounded away from 0 and if $1 \leq p < \infty$, we let $L^p(\omega)$ be the Banach space of (equivalence classes of) measurable functions on R^+ with $f\omega$ in $L^p(R^+)$, with the inherited norm

$$\|f\| = \|f\|_{\omega,p} = \|f\omega\|_p = \left(\int_0^\infty |f(t)\omega(t)|^p dt \right)^{1/p}.$$

We are particularly interested in the case that $L^1(\omega)$ is a Banach algebra and all $L^p(\omega)$ are $L^1(\omega)$ -modules under the usual convolution multiplication $f * g(x) = \int_0^x f(x-t)g(t) dt$. Therefore we will usually assume that $\omega(t)$ is an *algebra weight*, that is $\omega(t)$ satisfies:

- (1) $\omega(x+y) \leq \omega(x)\omega(y)$;
- (2) $\omega(x)$ is right continuous;
- (3) $\omega(0) = 1$.

(1), (2), and (3) are just normalizations and are essentially equivalent to $L^1(\omega)$ being an algebra in which case $L^p(\omega)$ is an $L^1(\omega)$ -module [9], where the module action is convolution. The most

important cases are the classical case $\omega(t) \equiv 1$, so that $L^1(\omega) = L^1(\mathbb{R}^+)$, and the case $\lim_{t \rightarrow \infty} \omega(t)^{\frac{1}{t}} = 0$, so that $L^1(\omega)$ is a radical Banach algebra.

When $\omega(t)$ is an algebra weight, the space $M(\omega)$ of locally finite Borel measures satisfying $\|\mu\| = \|\mu\|_\omega = \int_{\mathbb{R}^+} \omega(t) d|\mu|(t) < \infty$ is also a Banach algebra under convolution and each $L^p(\omega)$ is an $M(\omega)$ Banach module. We will usually identify the measure μ in $M(\omega)$ with the linear operator of convolution by μ on $L^p(\omega)$, so that $M(\omega)$ has a strong operator topology for its action on each $L^p(\omega)$. Particularly important is the fact that convolution by the point mass δ_a is right translation by a , so that the set of all δ_t for $t \geq 0$ is identified with the strongly continuous right translation semigroup on $L^p(\omega)$ for $1 \leq p < \infty$.

Suppose now that $\phi : L^1(\omega_1) \rightarrow L^1(\omega_2)$ is a continuous algebra homomorphism, with $\omega_1(t)$ and $\omega_2(t)$ algebra weights. This homomorphism has a unique extension to a homomorphism, which we also call ϕ , from $M(\omega_1)$ to $M(\omega_2)$ [10, Theorem 3.4, p. 596], so that $\mu_t = \phi(\delta_t)$ is a semigroup in $M(\omega_2)$. The following definition lists the properties we would like to prove for ϕ .

DEFINITION (1.1). We say the above homomorphism ϕ is *standard for p* , where $1 \leq p < \infty$ is fixed, if the following properties all hold.

(a) The semigroup $\mu_t = \phi(\delta_t)$ is strongly continuous on $L^p(\omega_2)$; that is $\lim_{t \rightarrow 0} \mu_t * g = g$ for all g in $L^p(\omega_2)$.

(b) Whenever $L^1(\omega_1) * f$ is dense in $L^1(\omega_1)$, then $L^p(\omega_2) * \phi(f)$ is dense in $L^p(\omega_2)$.

(c) For each h in $L^p(\omega_2)$, we can write $h = \phi(f) * g$ for some $f \in L^1(\omega_1)$ and $g \in L^p(\omega_2)$.

(d) Whenever $\{\lambda_n\}$ is a net in $M(\omega_1)$ for which $\lim_{n \rightarrow \infty} \lambda_n * f = \lambda * f$ for all f in $L^1(\omega_1)$ then $\lim_{n \rightarrow \infty} \phi(\lambda_n) * g = \phi(\lambda) * g$ for all g in $L^p(\omega_2)$; that is, ϕ is continuous from the strong operator topology on $M(\omega_1)$ acting on $L^1(\omega_1)$ to the strong operator topology of $M(\omega_2)$ acting on $L^p(\omega_2)$.

The main result of this paper, Theorem (3.1), is that ϕ is always standard for p when $1 < p < \infty$. When $p = 1$, we have previously shown, in joint work with Peter McClure [7, Theorem (2.2), p. 280], that conditions (a), (b), (c) and (d) are equivalent, but when $p = 1$, we have so far only been able to prove that these conditions hold for

a special class of weights [8, Theorem (3.4), p. 284], the regulated weights of Bade and Dales [1].

In §2, we will compare various types of convergence in $L^p(\omega)$ and show that a class of semigroups, which will include all $\mu_t = \phi(\delta_t)$, are strongly continuous when $p > 1$. §3 will be devoted to proving that all ϕ are standard for all $p > 1$. In §4, we observe that, for sequences which do not come from semigroups $\mu_t = \phi(\delta_t)$, the best convergence results require that the weights be regulated, just as when $p = 1$ [7].

2. Types of convergence. Many of the parts of the definition of standard homomorphisms involve comparing various convergence properties for a bounded sequence or net $\{\lambda_n\}$ in $M(\omega)$, just as in the case $p = 1$ (see [10], [7]). For $1 \leq p < \infty$, the dual space of $L^p(\omega)$ is $L^q(1/\omega)$, where q is the conjugate exponent to p , under the usual duality $\langle f, h \rangle = \int_0^\infty f(t)h(t) dt$. Also, when $\omega(t)$ is an algebra weight, $M(\omega)$ is the dual space of $C_0(1/\omega)$ under the analogous duality [10, Theorem 2.2, p. 592]. Here $C_0(1/\omega)$ is the subspace of $L^\infty(1/\omega)$ composed of continuous functions $h(t)$ for which $\lim_{t \rightarrow \infty} h(t)/\omega(t) = 0$. The following is our basic convergence result.

LEMMA (2.1). *Suppose that $\omega(t)$ is an algebra weight and that $\{\lambda_n\}$ is a bounded net in $M(\omega)$. If either*

(i) *There is a $\nu \neq 0$ in $M(\omega)$ for which $\lambda_n * \nu \rightarrow \lambda * \nu$ weak* in $M(\omega) = C_0(1/\omega)^*$*

or

(ii) *There is a $g \neq 0$ in some $L^p(\omega)$ with $1 < p < \infty$, for which $\text{weak} - \lim(\lambda_n * g) = \lambda * g$ in $L^p(\omega)$, then $\lambda_n * \nu \rightarrow \lambda * \nu$ weak* for all ν in $M(\omega)$ and $\lambda_n * g \rightarrow \lambda * g$ weakly in $L^p(\omega)$ for all g in all $L^p(\omega)$ with $1 < p < \infty$.*

The most important ν in $M(\omega)$ is the point mass δ_0 , which is the identity for convolution, so that the assertion of convergence for this ν just says $\lambda_n \rightarrow \lambda$ weak* in $M(\omega)$. We could have considered only sequences instead of bounded nets in the above lemma, since, when restricted to bounded sets, the weak*-topology on $M(\omega)$ and the weak topologies on $L^p(\omega)$ for finite p are metrizable.

Proof of Lemma (2.1). The key to passing between weak* and weak convergence for bounded nets or sequences is the observation

that if f belongs to $L^1(\omega) \cap L^p(\omega) \subseteq M(\omega) \cap L^p(\omega)$ for some $1 < p < \infty$, then

$$(2.2) \quad \lambda_n * f \rightarrow \lambda * f \text{ weak}^* \text{ in } M(\omega) \Leftrightarrow \lambda_n * f \rightarrow \lambda * f \text{ weakly in } L^p(\omega).$$

Formula (2.2) holds because both weak* and weak convergence are equivalent to $\lim \langle \lambda_n * f, h \rangle = \langle \lambda * f, h \rangle$ for all continuous h with compact support in $R^+ = [0, \infty)$, since the continuous functions with compact support are dense in $C_0(1/\omega)$ and in all $L^q(1/\omega)$ for $1 \leq q < \infty$.

Now suppose hypothesis (i) holds; then $\lambda_n * \nu \rightarrow \lambda * \nu$ weak* in $M(\omega)$ for all ν in $M(\omega)$ by [10, Lemma 3.2, p. 595]. Thus formula (2.2) shows that $\lambda_n * f \rightarrow \lambda * f$ weakly for all f in $L^1(\omega) \cap L^p(\omega)$. Since $L^1(\omega) \cap L^p(\omega)$ is dense in $L^p(\omega)$ when $p < \infty$, this proves that the weak limit of $\lambda_n * f$ is $\lambda * f$ for all f in all $L^p(\omega)$ with $1 < p < \infty$.

Now we suppose that hypothesis (ii) holds and verify that $\lambda_n \rightarrow \lambda$ weak* in $M(\omega)$, which is hypothesis (i) for $\nu = \delta_0$. Since $\{\lambda_n\}$ is a bounded net, it follows from weak* compactness that there is a subnet $\{\lambda'_n\}$ which converges weak* to some λ' in $M(\omega)$. To complete the proof we show that we must have $\lambda' = \lambda$. The subsequence $\{\lambda'_n\}$ satisfies hypothesis (i), so $\lambda'_n * g \rightarrow \lambda' * g$ weakly in $L^p(\omega)$. But $\{\lambda'_n * g\}$ is a subnet of the weakly convergent net $\{\lambda_n * g\}$, so that $\lambda * g = \lambda' * g$ and $g \neq 0$. Since it follows from the Titchmarsh convolution theorem [3] that the collection of locally finite measures on R^+ is an integral domain under convolution, we have $\lambda = \lambda'$ as required. This completes the proof. \square

We can now prove that the convolution semigroups we need to consider are strongly continuous on $L^p(\omega)$.

THEOREM (2.3). *Suppose that $\{\mu_t\}$ is a convolution semigroup in $M(\omega)$ with $\|\mu_t\|$ bounded as $t \rightarrow 0^+$. Then $\{\mu_t\}$ is a strongly continuous semigroup on $L^p(\omega)$ for all p with $1 < p < \infty$ if any of the following conditions hold.*

- (i) $\text{weak}^* - \lim_{t \rightarrow 0^+} \mu_t = \delta_0$.
- (ii) *There is some $\nu \neq 0$ in $M(\omega)$ for which $\mu_t * \nu$ is weak*-continuous from the right at some $t \geq 0$.*
- (iii) *There is some $g \neq 0$ in some $L^p(\omega)$ with $1 < p < \infty$ for*

which $\mu_t * g$ is weakly continuous from the right in $L^p(\omega)$ at some $t \geq 0$.

Proof. It follows from Lemma (2.4) that any of conditions (i), (ii) and (iii) implies that μ_t acts as a weakly continuous semigroup on all $L^p(\omega)$ with $1 < p < \infty$. But it is a standard result [12, Theorem 10.6.5, p. 324] that a weakly continuous semigroup is strongly continuous. \square

For $p = 1$, the conditions in the above theorem imply that μ_t is strongly continuous on $L^1(\omega)$ for $t > 0$ and that all $\mu_t * \nu$ are weak*-continuous for $t \geq 0$ [11, Theorem (2.1), p. 160]. For all such semigroups in $M(\omega)$ to be strongly continuous on $L^1(\omega)$ at $t = 0$ is completely equivalent to all continuous homomorphisms from some $L^1(\omega_1)$ to $L^1(\omega)$ being standard for $p = 1$ [11, Theorem 2.9, p. 164].

3. Standard homomorphisms. We are now ready for our major result, verifying that all homomorphisms satisfy the conditions of Definition (1.1) when $p > 1$.

THEOREM (3.1). *If $\phi : L^1(\omega_1) \rightarrow L^1(\omega_2)$ is a continuous nonzero homomorphism, then ϕ is standard for $1 < p < \infty$.*

Proof. Recall that we always extend the homomorphism to a homomorphism $\phi : M(\omega_1) \rightarrow M(\omega_2)$, and that ϕ has the same norm on $L^1(\omega_1)$ and $M(\omega_1)$ [10, Theorem 3.4, p. 596]. We let $\mu_t = \phi(\delta_t)$ and note that since $\|\delta_t\| = \omega_1(t)$ we have

$$(3.2) \quad \|\mu_t\| \leq \|\phi\|\omega_1(t),$$

where the norm $\|\mu_t\|$ is taken in $M(\omega_2)$.

We first prove (a) of Definition (1.1). Choose some f in $L^1(\omega_1)$ with $\phi(f) \neq 0$. Then $\mu_t * \phi(f) = \phi(\delta_t * f)$ is norm continuous, and hence weak*-continuous in $L^1(\omega_2) \subseteq M(\omega_2)$. Formula (3.2) shows that $\|\mu_t\|$ is bounded as $t \rightarrow 0^+$, so it follows from Theorem (2.3) that $\{\mu_t\}$ acts as a strong continuous semigroup on all $L^p(\omega_2)$ with $1 < p < \infty$.

Now use (a) to prove (b) in Definition (1.1). For simplicity we normalize to the case that $\lim_{t \rightarrow \infty} \omega_1(t)^{1/t} < 1$. This normalization is accomplished by replacing $\omega_1(t)$ by some $e^{-rt}\omega_1(t)$ and recalling that the map $f(t) \rightarrow f(t)e^{-rt}$ is an isometric isomorphism from $L^1(e^{-rt}\omega_1(t))$ onto $L^1(\omega_1)$.

By our normalization, $u(t) \equiv 1$ belongs to $L^1(\omega_1)$ and, using formula (3.2), we see that $\lim_{t \rightarrow \infty} \|\mu_t\|^{1/t} < 1$. Thus if we let $-A$ be the generator of the semigroup $\{\mu_t\}$ on $L^p(\omega_2)$, we have that A is a one-one closed operator from a dense subspace of $L^p(\omega_2)$ onto $L^p(\omega_2)$ with

$$(3.3) \quad A^{-1}(g) = \int_0^\infty \mu_t * g \, dt$$

as a Bochner integral in $L^p(\omega_2)$ whenever $g \in L^p(\omega_2)$ [5, pp. 620-622]. (Actually, since $\mu_t * g$ is continuous, the integral is just a vector-valued improper Riemann integral.) The main part of the proof of (b) is proving that if we let $v = \phi(u)$ then

$$(3.4) \quad v * g = \int_0^\infty \mu_t * g \, dt \text{ for all } g \text{ in } L^p(\omega_2).$$

Formula (3.4) will imply that $L^p(\omega_2) * v = \text{Range}(A^{-1}) = \text{Dom}(A)$, which is dense in $L^p(\omega_2)$.

The first step in verifying (3.4) for $p > 1$ is to prove it when $p = 1$. In $L^1(\omega_2)$ we have $\mu_t * g$ strongly measurable, in fact continuous for $t > 0$ [10, Theorem 3.6, p. 599], and $\|\mu_t * g\| \leq \|\phi\| \|g\| \omega_1(t)$ which is integrable by our normalization. Thus the integral in formula (3.4) defines a bounded linear operator on $L^1(\omega_2)$. Since each μ_t is a multiplier of $L^1(\omega_2)$, so is this bounded operator. Hence [10, Theorem (2.2) (E), p. 592] there is a measure λ in $M(\omega_2)$ for which $\lambda * g = \int_0^\infty \mu_t * g \, dt$ for g in $L^1(\omega_2)$. Now choose some $g = \phi(f) \neq 0$ in the range of ϕ . The standard convolution formula says $u * f = \int_0^\infty u(t) \delta_t * f \, dt = \int_0^\infty \delta_t * f \, dt$. Applying ϕ to this formula gives

$$v * g = \phi(u * f) = \int_0^\infty \phi(\delta_t * f) \, dt = \int_0^\infty \mu_t * g \, dt = \lambda * g.$$

Since locally finite measures on R^+ form an integral domain, it follows that $v = \lambda$, so that formula (3.4) holds for $p = 1$.

Now fix $p > 1$. Since $L^1(\omega_2) \cap L^p(\omega_2)$ is dense in $L^p(\omega_2)$ and both the integral and convolution by v are continuous linear operators on g in $L^p(\omega_2)$, it will be enough to verify formula (3.4) for g in $L^1(\omega_2) \cap L^p(\omega_2)$. Suppose that h is a continuous function with compact support, so that h belongs to $(L^p(\omega_2))^* = L^q(1/\omega_2)$ and to

$L^1(\omega_2)^* = L^\infty(1/\omega_2)$. It then follows from formula (3.4) for $p = 1$ that

$$\langle v * g, h \rangle = \left\langle \int_0^\infty \mu_t * g \, dt, h \right\rangle,$$

when the integral is considered as an integral in $L^1(\omega_2)$. But the scalar integral $\int_0^\infty \langle \mu_t * g, h \rangle \, dt$ equals $\langle \int_0^\infty \mu_t * g \, dt, h \rangle$ whether the vector integral is considered in $L^1(\omega_2)$ or $L^p(\omega_2)$. Thus $\langle v * g, h \rangle = \langle \int_0^\infty \mu_t * g \, dt, h \rangle$, when the integral is an $L^p(\omega_2)$ integral, for all h in a dense subspace of the dual space $(L^p(\omega_2))^*$. This proves formula (3.4), and hence that $L^p(\omega_2) * v$ is dense in $L^p(\omega_2)$. Thus we have proved part (b) of Definition (1.1) for the special case where $f = u$.

Now let f be an arbitrary function in $L^1(\omega_1)$ for which $L^1(\omega_1) * f$ is dense. Then there is a sequence $\{h_n\}$ in $L^1(\omega_1)$ for which $f * h_n \rightarrow u$ in $L^1(\omega_1)$, so $\phi(f) * \phi(h_n) \rightarrow v$ in $L^1(\omega_2)$. Thus if g belongs to $L^p(\omega_2)$, then $g * v = \lim_{n \rightarrow \infty} (g * \phi(h_n)) * \phi(f)$ so that the dense subspace $L^p(\omega_2) * v$ is contained in the closure of $L^p(\omega_2) * \phi(f)$. This proves that $L^p(\omega_2) * \phi(f)$ is dense in $L^p(\omega_2)$, as required.

We now use (b) to prove (c) of Definition (1.1). Notice that $L^p(\omega_2)$ is a Banach module over $L^1(\omega_1)$ under the multiplication $f \cdot g = \phi(f) * g$. Let $\{e_n\}$ be a bounded approximate identity for the Banach algebra $L^1(\omega_1)$ (for instance $e_n = n\chi_{[0,1/n]}$). Then (c) will follow from the factorization theorem for modules [2, Theorem 10, p. 61] if we show that $\{e_n\}$ is a module approximate identity.

Choose some f in $L^1(\omega_1)$ for which $L^p(\omega_2) * \phi(f)$ is dense in $L^p(\omega_2)$. Since $\{e_n\}$ is bounded it will be enough to show that $\lim_{n \rightarrow \infty} e_n \cdot (\phi(f) * g) = \phi(f) * g$ for g in $L^p(\omega_2)$. But $e_n \cdot (\phi(f) * g) = \phi(e_n) * \phi(f) * g = \phi(e_n * f) * g \rightarrow \phi(f) * g$, since $\{e_n\}$ is an approximate identity for $L^1(\omega_1)$. This proves (c).

Finally we use (c) to prove (d) in Definition (1.1). Suppose $\{\lambda_n\}$ is a net which converges in the strong operator topology of $M(\omega_1)$ on $L^1(\omega_1)$ to λ . Let $h = \phi(f) * g$ be an arbitrary element of $L^p(\omega_2)$. Then $\phi(\lambda_n) * h = \phi(\lambda_n * f) * g \rightarrow \phi(\lambda * f) * g = \phi(\lambda) * h$, as required. This completes the proof of the theorem. \square

In the same way that we proved formula (3.4) above, we could prove the analogous formula for any $f(t)$ in $L^1(\omega_1)$ in place of $u(t) \equiv 1$. Thus we have:

COROLLARY (3.5). *For all f in $L^1(\omega_1)$ and all g in $L^p(\omega_2)$ where*

$1 \leq p < \infty$, we have $\phi(f) * g = \int_0^\infty f(t) \mu_t * g dt$, where the integral is a Bochner integral in $L^p(\omega_2)$.

In part (c) of Definition (1.1) if $\alpha(h) = \inf(\text{support } h)$, we can require that $\alpha(g) = \alpha(h)$ so that $\alpha(\phi(f)) = 0$. We just consider the functions in $L^p(\omega_2)$ with support in $[\alpha(h), \infty)$ as the $L^1(\omega_1)$ Banach Module.

The proof that (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) in Theorem (3.1) also carries through if $p = 1$. We didn't even need the tricky arguments comparing Bochner integrals in $L^1(\omega_2)$ and $L^p(\omega_2)$. It is also easy to see that (d) \Rightarrow (a) since δ_t is a strongly continuous semigroup, so, when (d) holds, $\mu_t = \phi(\delta_t)$ must also be strongly continuous. Thus our proof of Theorem (3.1) for $p > 1$ also shows that when $p = 1$, the four conditions of Definition (1.1) are equivalent. This gives a slightly simpler proof of the essential parts of our earlier [7, Theorem 2.2, p. 280]. But for $p = 1$, we only know that these conditions hold for a restricted class of weights.

4. Compactness and norm convergence. In our previous studies of the standard homomorphism problem [8] and [7], for $p = 1$, we were able to prove that the weak*-continuous semigroup $\{\mu_t\}$ was strongly continuous by coming up with a condition on the weight $\omega(t)$ which guaranteed that whenever a sequence $\{\lambda_n\}$ converged weak* to λ in $M(\omega)$, then $\lambda_n * f \rightarrow \lambda * f$ in norm in $L^1(\omega)$ for appropriate f . It also turned out [7, Theorem (4.1)] that weak convergence of $\lambda_n * f$ in $L^1(\omega)$ implied norm convergence. Since we have shown that when $p > 1$, the semigroup μ_t is always strongly continuous on $L^p(\omega)$, and since $L^p(\omega)$ is reflexive, so that weak* and weak convergence are the same, one would expect that weak convergence would imply norm convergence of $\lambda_n * f$ in $L^p(\omega)$ for a more general class of weights. Surprisingly, for general sequences or bounded nets $\{\lambda_n\}$, the norm convergence results are essentially the same for $L^p(\omega)$ for all $1 \leq p < \infty$.

The appropriate class of weights are the regulated weights of Bade and Dales [1, Definition 1.3, p. 81]. We say that the algebra weight $\omega(t)$ is *regulated* at $a \geq 0$, if $\lim_{t \rightarrow \infty} \omega(t + b)/\omega(t) = 0$ for all $b > a$. The following two results say that regularity, convergence improvement, and compactness are all equivalent for all $1 \leq p < \infty$. Recall, from Lemma (2.1) above, that the assumption that $\{\lambda_n\}$ converges

weak* to λ is the same as assuming that $\lambda_n * f \rightarrow \lambda * f$ weakly in some (all) $L^p(\omega)$ with $1 < p < \infty$ for some (all) $f \neq 0$ in $L^p(\omega)$. Recall also that, for $f \in L^1_{loc}(R^+)$, we let $\alpha(f) = \inf(\text{support } f)$.

THEOREM (4.1). *If the weight $\omega(t)$ is regulated at a , then for all $1 \leq p < \infty$ and all f in $L^p(\omega)$ with $\alpha(f) \geq a$ we have:*

- (a) *Convolution by f is a compact operator from $M(\omega)$ to $L^p(\omega)$.*
- (b) *If the sequence or bounded net $\{\lambda_n\}$ converges weak* to λ in $M(\omega)$, then $\lambda_n * f \rightarrow \lambda * f$ in norm in $L^p(\omega)$.*

THEOREM (4.2). *If the algebra weight $\omega(t)$ is not regulated at a , then there is a sequence $\{\lambda_n\}$ in $M(\omega)$ with weak*-limit 0 for which, for all $1 \leq p < \infty$ and all f in $L^p(\omega)$ with $\alpha(f) \leq a$, we have:*

- (a) *The sequence $\lambda_n * f$ diverges in norm in $L^p(\omega)$.*
- (b) *Convolution by f is not a compact operator from either $M(\omega)$ or $L^1(\omega)$ to $L^p(\omega)$.*

We will give relatively simple proofs which reduce the results for $p > 1$ to our previous results [8, Theorem (2.3)] [7, pp. 283-284] when $p = 1$, instead of giving a proof for all $1 \leq p < \infty$. For a detailed study of compactness of convolution operators from $L^p(\omega)$ to itself, without the assumption that ω is an algebra weight, see Detre's thesis [4].

Proof of Theorem (4.1). Part (b) is a standard characterization of compactness (cf. [7, Theorem (3.2), p. 284]), so we just need to prove (a). Since $\delta_a * L^p(\omega)$ is dense in $L^p(\omega)_a = \{f \in L^p(\omega) : \alpha(f) \geq a\}$, it is enough to prove (a) for functions $f = \delta_a * g$ with g in $L^p(\omega)$. Since $L^1(\omega)$ has a bounded approximate identity which is also a module approximate identity, it follows from the Cohen factorization theorem that we can write $g = h * k$ with h in $L^1(\omega)$ and k in $L^p(\omega)$. The case $p = 1$ of the theorem [7, Lemma (3.1), p. 283] shows that $(\delta_a * h)$ acts compactly from $M(\omega)$ to $L^1(\omega)$. Also, convolution by k is a bounded operator from $L^1(\omega)$ to $L^p(\omega)$. Hence convolution by $f = (\delta_a * h) * k$ is compact, since it is the composition of a compact operator and a bounded operator. This completes the proof of Theorem (4.1). \square

Proof of Theorem (4.2). First notice that for h in $C_0(1/\omega)$ we have $\langle \delta_s/\omega(s), h \rangle = h(s)/\omega(s)$, which approaches 0 as $s \rightarrow \infty$ by

the definition of $C_0(1/\omega)$. Thus $\delta_s/\omega(s)$, and hence any of its subsequences, approach 0 weak* in $M(\omega)$. Since $\omega(t)$ is not regulated at a , there is a $t_0 > a$ for which $\omega(s + t_0)/\omega(s)$ does not approach 0 as $s \rightarrow \infty$. Hence we can find a sequence $s_n \rightarrow \infty$ with $\omega(s_n + t_0)/\omega(s_n)$ bounded below. We claim that the sequence $\lambda_n = \delta_{s_n}/\omega(s_n)$, which approaches 0 weak* in $M(\omega)$, satisfies the assertion in Theorem (4.2) (a).

Pick some f in some $L^p(\omega)$ with $\alpha(f) \leq a$. Then

$$(\|\lambda_n * f\|_{p,\omega})^p = \int_0^\infty |f(t)|^p (\omega(s_n + t)/\omega(s_n))^p dt.$$

Now the weight $(\omega(t))^p$ is also not regulated and $|f|^p$ belongs to $L^1(\omega(t)^p)$. Also $(\omega(s_n + t)/\omega(s_n))^p$ is bounded away from zero so that, as we showed in our proof of [8, Theorem (2.3)], $\|(\delta(s_n)/\omega(s_n)^p) * |f|^p\|_{1,\omega^p} = \int_0^\infty |f(t)|^p |\omega(s_n + t)/\omega(s_n)|^p dt$ cannot approach 0 as $n \rightarrow \infty$. That is, the $p = 1$ result, for the algebra $L^1(\omega^p)$, which we proved previously, gives the result in Theorem (4.2) (a) for $L^p(\omega)$ for all $1 \leq p < \infty$.

Part (a) shows that convolution by f cannot be a compact operator from $M(\omega)$ to $L^p(\omega)$. We complete the proof by showing that, if convolution by g is compact from $L^1(\omega)$ to $L^p(\omega)$, then it is also compact from $M(\omega)$ to $L^p(\omega)$. The proof we gave for the case $p = 1$, in [7, Lemma (3.1), p. 283] carries through without change for all $1 \leq p < \infty$, since a bounded approximate identity $\{e_n\}$ for $L^1(\omega)$ is also a module approximate identity for $L^p(\omega)$. This completes the proof. \square

The above two theorems show that for arbitrary sequences or nets, as distinct from semigroups, norm convergence in $L^p(\omega)$ for $p > 1$ is no easier to obtain than for $L^1(\omega)$. In contrast, the following small result does give one sense in which $L^p(\omega)$ convergence is easier.

PROPOSITION (4.3). *Suppose that $\{\lambda_n\}$ is a net in $M(\omega)$. If $\lim(\lambda_n * f) = \lambda * f$ in norm in $L^1(\omega)$ for all f in $L^1(\omega)$, then $\{\lambda_n * g\}$ converges in norm to $\lambda * g$ for all g in $L^p(\omega)$ and all $1 < p < \infty$.*

Proposition (4.3) can be viewed as a special case of Theorem (3.1) where the homomorphism is the identity on $L^1(\omega)$. Alternately, and more simply, one can use essentially the same proof of (c) \Rightarrow (d)

in Theorem (3.1) by considering $L^p(\omega)$ as an $L^1(\omega)$ module and applying the Cohen factorization theorem.

REFERENCES

- [1] W.G. Bade and H.G. Dales, *Norms and ideals in radical convolution algebras*, J. Funct. Anal., **41** (1981), 77-109.
- [2] F.F. Bonsall and J. Duncan, *Complete Normed Algebras*, Springer-Verlag, Berlin, 1973.
- [3] H.G. Dales, *Convolution algebras on the real line*, Radical Banach Algebras and Automatic Continuity, Lecture Notes in Math., no. 975, Springer-Verlag, Berlin, 1983.
- [4] P. Detre, *Multipliers of Weighted Lebesgue Spaces*, Ph. D. dissertation, Univ. Calif., Berkeley, 1988.
- [5] N. Dunford and J. Schwartz, *Linear operators, part 1*, Wiley Interscience, New York, 1958.
- [6] F. Ghahramani, *Homomorphisms and derivations on weighted convolution algebras*, J. London Math. Soc., **21** (1980), 149-161.
- [7] F. Ghahramani, S. Grabiner and J.P. McClure, *Standard homomorphisms and regulated weights on weighted convolution algebras*, J. Funct. Anal., **91** (1990), 278-286.
- [8] F. Ghahramani and S. Grabiner, *Standard homomorphisms and convergent sequences in weighted convolution algebras*, Illinois J. Math., no. 3, **3** (1992), 505-527.
- [9] S. Grabiner, *Weighted convolution algebras on the half line*, J. Math. Anal. Appl., **83** (1981), 531-553.
- [10] ———, *Homomorphisms and semigroups in weighted convolution algebras*, Indiana Univ. Math. J., **37** (1988), 589-615.
- [11] ———, *Semigroups and the structure of weighted convolution algebras*, Conference on Automatic Continuity and Banach Algebras, R.J. Loy ed., Proc. Centre Math. Analysis, Australian National University, vol. **21** (1989), 155-169.
- [12] E. Hille and R.S. Phillips, *Functional Analysis and Semi-groups*, Colloquium Publication Series, vol. **31**, Amer. Math. Soc., Providence, Rhode Island, 1957.

Received September 21, 1992 and in revised form March 21, 1993. The first author was supported by NSERC grant OGP0036640. The second author was supported by an NSF grant.

UNIVERSITY OF MANITOBA
WINNIPEG, MANITOBA

CANADA R3T 2N2

AND

POMONA COLLEGE
CLAREMONT, CA 91711-6348

PACIFIC JOURNAL OF MATHEMATICS

Founded by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

EDITORS

Sun-Yung Alice Chang (Managing Editor)

University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

F. Michael Christ
University of California
Los Angeles, CA 90095-1555
christ@math.ucla.edu

Robert Finn
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

Martin Scharlemann
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

Thomas Enright
University of California
San Diego, La Jolla, CA 92093
tenright@ucsd.edu

Vaughan F. R. Jones
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

Gang Tian
Courant Institute
New York University
New York, NY 10012-1100
tiang@taotao.cims.nyu.edu

Nicholas Ercolani
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

Steven Kerckhoff
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

V. S. Varadarajan
University of California
Los Angeles, CA 90095-1555
vsv@math.ucla.edu

SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
STANFORD UNIVERSITY
UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
UNIVERSITY OF UTAH
UNIVERSITY OF WASHINGTON
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$108.00 a year to individual members of supporting institutions.

Subscriptions, orders for back issues published within the last three years, and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,
the American Mathematical Society's TEX macro system.

Copyright © 1995 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 168 No. 1 March 1995

Compact contractible n -manifolds have arc spines ($n \geq 5$)	1
FREDRIC DAVIS ANCEL and CRAIG R. GUILBAULT	
On uniform homeomorphisms of the unit spheres of certain Banach lattices	11
FOUAD CHAATIT	
Applications of subordination chains to starlike mappings in \mathbb{C}^n	33
MARTIN CHUAQUI	
The L^p theory of standard homomorphisms	49
FEREIDOUN GHAHRAMANI and SANDY GRABINER	
Metrics for singular analytic spaces	61
CAROLINE GALT GRANT and PIERRE MILMAN	
Maps on infra-nilmanifolds	157
KYUNG BAI LEE	
Hardy spaces and oscillatory singular integrals. II	167
YIBIAO PAN	
Continuity of convex hull boundaries	183
LINDA JO GOLDWAY KEEN and CAROLINE SERIES	



0030-8730(1995)168:1;1-D