

*Pacific  
Journal of  
Mathematics*

**ITERATED LOOP MODULES AND A FILTRATION FOR  
VERTEX REPRESENTATION OF TOROIDAL LIE ALGEBRAS**

S. ESWARA RAO

**ITERATED LOOP MODULES AND A FILTRATION FOR  
VERTEX REPRESENTATION OF TOROIDAL LIE  
ALGEBRAS**

S. ESWARA RAO

**The purpose of this paper is two fold. The first one is to construct a continuous new family of irreducible (some of them are unitarizable) modules for Toroidal algebras. The second one is to describe the sub-quotients of the (integrable) modules constructed through the use of Vertex operators.**

**Introduction.**

Toroidal algebras  $\tau_{[d]}$  are defined for every  $d \geq 1$  and when  $d = 1$  they are precisely the untwisted affine Lie-algebras. Such an affine algebra  $\mathcal{G}$  can be realized as the universal central extension of the loop algebra  $\dot{\mathcal{G}} \otimes \mathbb{C}[t, t^{-1}]$  where  $\mathcal{G}$  is simple finite dimensional Lie-algebra over  $\mathbb{C}$ . It is well known that  $\mathcal{G}$  is a one dimensional central extension of  $\dot{\mathcal{G}} \otimes \mathbb{C}[t, t^{-1}]$ . The Toroidal algebras  $\tau_{[d]}$  are the universal central extensions of the iterated loop algebra  $\dot{\mathcal{G}} \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$  which, for  $d \geq 2$ , turnout to be infinite central extension. These algebras are interesting because they are related to the Lie-algebra of  $\text{Map}(X, G)$ , the infinite dimensional group of polynomial maps of  $X$  to the complex algebraic group  $G$  where  $X$  is a  $d$ -dimensional torus.

For additional material on recent developments in the theory of Toroidal algebras one may consult [BC], [FM] and [MS].

In [MEY] and [EM] a countable family of modules (also integrable see [EMY]) are constructed for Toroidal algebras on Fock space through the use of Vertex Operators (Theorem 3.4, [EM]). However they are reducible and *not* completely reducible. In §5 we observe that the Fock space is a direct sum of certain  $\underline{h}(\lambda)$ 's. In (5.10) to (5.12) we prove that each  $\underline{h}(\lambda)$  admits a filtration by an increasing sequence of modules such that the successive quotients are all isomorphic to  $V$  defined in §5. In (5.9) we prove that each  $V$  admits a filtration of decreasing sequence of modules such that the successive quotients are all irreducible. In our main Theorem 5.6 we will describe the irreducible modules as certain iterated loop modules twisted by an automorphism of  $\tau_{[d]}$ .

We now describe the contents of the paper.

In §1 we construct an iterated loop algebra of Kac-Moody Lie-algebra  $\mathcal{G}$  and construct a family of completely reducible modules (Theorem (1.8)) using methods similar to [E]. In §2 we prove that some of the above modules are unitarizable (Proposition 2.3). In §3 we specialize these results for  $\mathcal{G}$  finite dimensional (Theorem 3.3) and for  $\mathcal{G}$  an affine Lie-algebra (Theorem 3.6) to get irreducible modules (some of them are unitarizable modules) for  $\tau_{[d]}$ . The modules considered in Theorem 3.6 are the first examples of irreducible modules for  $\tau_{[d]}$  where part of the centre acts non-trivially. It should be mentioned that it is the unitarizable modules (and also integrable modules) which lift to the group.

In §4 we recall the construction of Vertex Operators and the Fock space. We also construct certain automorphisms of  $\tau_{[d]}$  which are necessary in §5.

1.

Let  $d$  and  $k$  be positive integers. Let  $\mathcal{G}$  be a Lie-algebra and let  $\mathcal{G}_k = \oplus \mathcal{G}$  be  $k$  copies of  $\mathcal{G}$ . Let  $A = A_d = \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$  be Laurent polynomial ring in  $d$  variables. Then  $\mathcal{G}_A = \mathcal{G} \otimes A$  is a Lie-algebra with Lie structure  $[X \otimes a, Y \otimes b] = [X, Y] \otimes ab, X, Y \in \mathcal{G} a, b \in A$ .

Let  $\underline{n} = (n_1, n_2, \dots, n_d)$  be a  $d$ -tuple of integers and let

$$t^{\underline{n}} = t_1^{n_1} t_2^{n_2} \dots t_d^{n_d}.$$

For  $1 \leq i \leq k$  let  $\underline{a}_i = (a_i(1), \dots, a_i(d))$  be a  $d$ -tuple of non-zero complex numbers. Let  $a_i^{\underline{n}} = a_i^{n_1}(1) \dots a_i^{n_d}(d)$  be the product. Consider the Lie-algebra homomorphism

$$\begin{aligned} \phi : \mathcal{G}_A &\rightarrow \mathcal{G}_k \\ X \otimes t^{\underline{n}} &\mapsto (X a_1^{\underline{n}}, \dots, X a_k^{\underline{n}}). \end{aligned}$$

It is elementary to check that  $\phi$  is *not* surjective if and only if  $a_i(\ell) = a_j(\ell)$  for some  $i \neq j$  and for all  $\ell$ . First prove it for  $d = 1$  and then using Vandermonde determinant for general  $d$ .

Define derivations  $d_1, d_2, \dots, d_d$  on  $\mathcal{G}_A$  by  $[d_i, X \otimes t^{\underline{n}}] = n_i X \otimes t^{\underline{n}}$  and note  $[d_i, d_j] = 0$ . Let  $D$  be the linear span of  $d_1, d_2, \dots, d_d$  and let  $\tilde{\mathcal{G}}_A = \mathcal{G} \otimes A \oplus D$ .

For any Lie-algebra  $\mathcal{G}$ , let  $U(\mathcal{G})$  denote the universal enveloping algebra. Note that  $A, \mathcal{G}_A$  and  $U(\mathcal{G}_A)$  are obviously  $\mathbb{Z}^d$  graded algebras.

**1.1.** From here onwards we will assume that  $\mathcal{G}$  is a Kac-Moody Lie-algebra. Fix a Cartan subalgebra  $\underline{h}$  of  $\mathcal{G}$ .

Let  $\psi : U(\underline{h}_A) \rightarrow A$  be a  $\mathbb{Z}^d$  graded homomorphism. Let  $A_\psi$  be the image of  $\psi$  which is a  $\mathbb{Z}^d$  graded subalgebra of  $A$ .

$A_\psi$  can be treated as an  $\tilde{\mathfrak{h}}_A$ - module by the  $\psi$  action,

$$h \otimes t^n.t^m = \psi(h \otimes t^n).t^m$$

$$d_i.t^m = m_i t^m.$$

**1.2. Lemma.**  $A_\psi$  is an irreducible  $\tilde{\mathfrak{h}}_A$ - module if and only if homogeneous elements of  $A_\psi$  are invertible in  $A_\psi$ .

*Proof.* Assume the homogeneous elements of  $A_\psi$  are invertible in  $A_\psi$ . To prove irreducibility it is sufficient to prove that given elements  $t^n$  and  $t^m$  there exists  $X$  in  $U(\mathfrak{h}_A)$  such that  $X.t^n = t^m$ . By assumption  $t^{m-n}$  belongs to  $A_\psi$ . Since  $A_\psi$  is the image of  $\psi$  there exists  $X$  in  $U(\mathfrak{h}_A)$  such that  $\psi(X) = t^{m-n}$  and clearly  $X.t^n = \psi(X)t^n = t^m$ .

Now for the converse let  $t^n$  belong to  $A_\psi$ . First note that 1 belongs to  $A_\psi$ . There exists  $X$  in  $U(\mathfrak{h}_A)$  such that  $\psi(X).t^n = 1$  by irreducibility of  $A_\psi$ . Then clearly  $\psi(X) = t^{-n}$  and we are done.  $\square$

**1.3.** The purpose of this section is to construct irreducible modules for  $\tilde{\mathcal{G}}_A$ .

Let  $V(\lambda_1), \dots, V(\lambda_k)$  be irreducible highest weight modules for  $\mathcal{G}$  with highest weights  $\lambda_1, \lambda_2, \dots, \lambda_k$  respectively. Then clearly  $V = \otimes V(\lambda_i)$  is an irreducible module for  $\mathcal{G}_k$ .

**1.4.** Also  $V$  can be treated as an irreducible  $\mathcal{G}_A$ -module via the Lie-algebra homomorphism  $\phi$ . But it is not a  $\mathbb{Z}^d$ - graded module. That is, it cannot be extended to  $\tilde{\mathcal{G}}_A$ .

Consider  $V_A := V \otimes A$  which will be given  $\tilde{\mathcal{G}}_A$  module structure.

$$(1.5) \quad \begin{aligned} X \otimes t^m(v \otimes t^n) &= \varphi(X \otimes t^m)v \otimes t^{m+n} \\ d_i(v \otimes t^n) &= n_i v \otimes t^n \quad (X \in \mathcal{G}, v \in V). \end{aligned}$$

We denote the  $\tilde{\mathcal{G}}_A$  module by  $(V_A, \pi)$ .

Let  $\psi : U(\mathfrak{h}_A) \rightarrow A$  be  $h \otimes t^n \mapsto \sum_i \lambda_i(h) \underline{a}_i^n t^n$  be a  $\mathbb{Z}^d$ - graded homomorphism of algebras.

**1.6.** We believe that the conditions for  $\phi$  to be *surjective* are sufficient to prove that  $A_\psi$  is irreducible, but we could not prove that. Instead we will give a continuous family of examples where  $A_\psi = A$ . In particular for these examples  $A_\psi$  is irreducible.

**1.7.** Assume  $1 \leq \ell \leq d, a_i(\ell)/a_j(\ell)$  is not a  $k$ -th root of unity. Then by Lemma 4.4 of [CP] the following  $\mathbb{Z}$ -graded algebra homomorphism is surjective for all  $\ell$ ,

$$U(\underline{h} \otimes \mathbb{C}[t_\ell, t_\ell^{-1}]) \rightarrow \mathbb{C}[t_\ell, t_\ell^{-1}]$$

$$h \otimes t_\ell^n \mapsto \sum_{j=1}^k \lambda_j(h) a_j^n(\ell) t_\ell^n.$$

Hence  $A_\psi$  contains  $\mathbb{C}[t_\ell, t_\ell^{-1}]$  for all  $\ell$ . So that  $A_\psi = A$ .

*From here on we will assume that  $A_\psi$  is irreducible  $\tilde{\mathcal{H}}_A$ - module.*

We will also note  $\psi$  and the module  $(V_A, \pi)$  depends on the choice of  $\lambda$ 's and  $a_i(\ell)$ 's.

For any  $v \in V$  let  $v(\underline{n}) = v \otimes t^{\underline{n}}$ . We will now prove that  $(V_A, \pi)$  as defined in (1.5) is a completely reducible  $\tilde{\mathcal{G}}_A$ - module.

**1.8. Theorem.** *Let  $G \subseteq \mathbb{Z}^d$  be such that  $\{t^{\underline{m}}, \underline{m} \in G\}$  is a set of coset representatives of  $A/A_\psi$ . Let  $v = v_1 \otimes \dots \otimes v_k$  where each  $v_i$  is a highest weight vector of  $V(\lambda_i)$ . Then*

(1)  $V_A = \bigoplus_{\underline{m} \in G} U(v(\underline{m}))$  as  $\tilde{\mathcal{G}}_A$ - module where  $U(v(\underline{m}))$  is the  $\tilde{\mathcal{G}}_A$  submodule generated by the vector  $v(\underline{m})$ .

(2) Each  $U(v(\underline{m}))$  is an irreducible  $\tilde{\mathcal{G}}_A$ - module.

*In particular  $(V_A, \pi)$  is an irreducible  $\tilde{\mathcal{G}}_A$  -module whenever  $A = A_\psi$ .*

Before we prove the theorem we prove some lemmas.

**1.9. Lemma.** *Any non-zero  $\tilde{\mathcal{G}}_A$ - submodule of  $V_A$  contains  $v(\underline{m})$  for some  $\underline{m}$ .*

*Proof.* Consider the map  $S : V_A \rightarrow V$  defined by  $S(w(\underline{m})) = w$  (extend linearly to  $V_A$ ). Then clearly  $S$  is a surjective  $\mathcal{G}_A$ - module map (it is not a  $\tilde{\mathcal{G}}_A$ -module map).

**Claim.**  $S(W) = V$  for any non-zero  $\tilde{\mathcal{G}}_A$  module  $W$  of  $(V_A, \pi)$ .

Since  $V$  is an irreducible  $\mathcal{G}_A$  -module and  $S(W)$  is a submodule of  $V$ , to see the claim it is sufficient to prove that  $S(W) \neq 0$ . But that is clear. Since  $W$  is a  $\tilde{\mathcal{G}}_A$ -module it therefore contains vector of the form  $w(\underline{m}), w \in V$  and  $S(w(\underline{m})) = w \neq 0$ . This proves the claim.

Now let  $w$  in  $W$  be such that  $S(w) = v$  where  $v$  is the vector defined in the statement of the theorem. Since  $S$  is a  $\underline{h}$ - module map,  $w$  and  $v$  are the same weight.  $\{v(\underline{m}), \underline{m} \in \mathbb{Z}^d\}$  are the only such weight vectors of  $\check{V}_A$  and hence  $w = \sum_i C_i v(\underline{m}^i)$  for some complex numbers  $C_i$  and some  $\underline{m}^i \in \mathbb{Z}^d$ .

But  $W$  is a  $\tilde{\mathcal{G}}_A$  module so is  $(\mathbb{Z}^d$ -graded) and it follows that  $v(\underline{m}^i)$  belongs to  $W$ . □

**1.10. Lemma.** *The following are true.*

- (1)  $v(\underline{m}) \in U(\tilde{\mathcal{G}}_A)v(\underline{n})$  if and only if  $t^{\underline{m}-\underline{n}} \in A_\psi$ .
- (2)  $v(\underline{m}) \in U(\tilde{\mathcal{G}}_A)v(\underline{n})$  if and only if  $t^{\underline{m}-\underline{n}} \in A_\psi$ .
- (3)  $U(\tilde{\mathcal{G}}_A)v(\underline{m}) = U(\tilde{\mathcal{G}}_A)v(\underline{n})$  if and only if  $t^{\underline{m}-\underline{n}} \in A_\psi$ .
- (4)  $N = U(\tilde{\mathcal{G}}_A)v(\underline{m}) \cap U(\tilde{\mathcal{G}}_A)v(\underline{n}) \neq \{0\}$  if and only if  $t^{\underline{m}-\underline{n}} \in A_\psi$ .

*Proof.* (1) Clear. (2) The “if” part follows from (1). For the “only if” part write  $\mathcal{G} = N^- \oplus \mathfrak{h} \oplus N^+$  where  $N^+$  is sum of positive root spaces and  $N^-$  is sum of negative root spaces. Now note that  $U(\tilde{\mathcal{G}}_A)v(\underline{m}) = U(N_A^-)U(\mathfrak{h}_A)v(\underline{m})$  which follows from the fact that  $U(N_A^+)v(\underline{m}) = 0$  and the Poincare-Birkhoff-witt theorem. Now it is easy to see by a weight argument that  $v(\underline{m}) \in U(\tilde{\mathcal{G}}_A)v(\underline{n})$  implies that  $v(\underline{m}) \in U(\mathfrak{h}_A)v(\underline{n})$ . So by (1),(2) is complete. (3) follows from (2). (4) Assume  $N \neq \{0\}$ . By Lemma (1.9) there is a  $\underline{k}$  such that  $v(\underline{k}) \in N$ . Hence by (2)  $t^{\underline{k}-\underline{m}}, t^{\underline{k}-\underline{n}} \in A_\psi$ . But  $t^{\underline{m}-\underline{k}} \in A_\psi$  and therefore  $t^{\underline{m}-\underline{n}} \in A_\psi$ . Converse follows from (3).

*Proof of the Theorem 1.8.* (2) Let  $W$  be a non-zero  $\tilde{\mathcal{G}}_A$  submodule of  $U(\tilde{\mathcal{G}}_A)v(\underline{m})$ . Then  $W$  contains  $v(\underline{n})$  for some  $\underline{n}$  (by Lemma 1.9). So that  $U(\tilde{\mathcal{G}}_A)v(\underline{n}) \subseteq W$ . But by Lemma (1.10),  $t^{\pm(\underline{m}-\underline{n})} \in A_\psi$ . Now by Lemma 1.10 (3) it will follow that  $U(\tilde{\mathcal{G}}_A)v(\underline{n}) = U(\tilde{\mathcal{G}}_A)v(\underline{m})$ . In particular  $W = U(\tilde{\mathcal{G}}_A)v(\underline{m})$ .

(1) Let  $w(\underline{m}) \in V_A, w \in V, \underline{m} \in \mathbb{Z}^d$ . Since  $V$  is irreducible  $\mathcal{G}_A$ - module (see 1.4) there exist  $X \in U(\mathcal{G}_A)$  such that  $\varphi(X)v = w$ . Write  $X = \sum_{\underline{n} \in \mathbb{Z}^d} X_{\underline{n}}$

where  $[d_i, X_{\underline{n}}] = n_i X_{\underline{n}}$ .

Consider  $\sum \pi(X_{\underline{n}})v(-\underline{n} + \underline{m}) = \sum \varphi(X_{\underline{n}})v(\underline{m}) = \varphi(X)v(\underline{m}) = w(\underline{m})$ . Hence we have proved that

$$(1.11) \quad V_A \subseteq \sum_{\underline{m} \in \mathbb{Z}^d} U(\tilde{\mathcal{G}}_A)v(\underline{m}).$$

We also have by Lemma 1.10 (3)

$$(1.12) \quad \sum_{\underline{m} \in \mathbb{Z}^d} U(\tilde{\mathcal{G}}_A)v(\underline{m}) = \sum_{\underline{m} \in G} U(\tilde{\mathcal{G}}_A)v(\underline{m}).$$

In view of (1.11) and (1.12), to see (1) of the Theorem we only have to prove that the sum of RHS in (1.12) is direct.

For that it is sufficient to prove that for all  $\underline{m} \in G$ .

$$(1.13) \quad U(\tilde{\mathcal{G}}_A)v(\underline{m}) \cap \sum_{\underline{n} \in G, \underline{m} \neq \underline{n}} U(\tilde{\mathcal{G}}_A)v(\underline{n}) = \{0\}.$$

Suppose (1.13) does not hold for some  $\underline{m} \in G$ . Then

$$U(\tilde{\mathcal{G}}_A)v(\underline{m}) \subseteq \sum_{\underline{n} \in G, \underline{m} \neq \underline{n}} U(\tilde{\mathcal{G}}_A)v(\underline{n})$$

(since  $U(\tilde{\mathcal{G}}_A)v(\underline{m})$  is irreducible). Since all modules under consideration are  $\underline{h} \oplus D-$  modules  $v(\underline{m})$  has to be linear combinations of  $v(\underline{k})$  where  $v(\underline{k}) \in U(\tilde{\mathcal{G}}_A)v(\underline{n}), \underline{n} \neq \underline{m}$  and  $\underline{n} \in G$ . But this is *not* possible since none of the  $\underline{k}$  can be equal to  $\underline{m}$  by Lemma 1.10 (2) and the choice of  $G$ . This proves that the sum at RHS in (1.12) is direct and that completes the proof of the theorem.

2.

In this section we will prove that modules constructed earlier for  $\tilde{\mathcal{G}}_A$  are unitarizable subject to some conditions on  $\lambda$ 's and  $a_i(\ell)$ 's.

We will start with

**2.1. Definition.** A conjugate-linear anti-involution of a Lie-algebra  $\mathcal{G}'$  is a map  $D : \mathcal{G}' \rightarrow \mathcal{G}'$  such that

$$D(X + Y) = D(X) + D(Y), D(\lambda X) = \bar{\lambda}D(X)$$

$$D[X, Y] = [D(Y), D(X)], D^2 = \text{Id}.$$

for all  $X, Y \in \mathcal{G}'$  and  $\lambda \in C$ .

Such maps are also called forms.

**2.2. Definition.** A  $\mathcal{G}'-$  module  $V$  is said to be unitarizable with respect to conjugate linear anti-involution  $D$  of  $\mathcal{G}'$  if there exists a positive definite hermitian form  $\langle , \rangle$  on  $V$  satisfying,

$$\langle Xv_1, v_2 \rangle = \langle v_1, D(X)v_2 \rangle$$

for all  $v_1, v_2 \in V$  and  $X \in \mathcal{G}'$ .

Let  $w$  be a conjugate-linear anti-involution of  $\mathcal{G}$ . Then  $w_k = \oplus_k w$  is conjugate-linear anti-involution of  $\mathcal{G}_k$ . Define  $\bar{w}$  on  $\tilde{\mathcal{G}}_A$  by  $\bar{w}X(\underline{n}) = w(X)(-\underline{n})$  and  $\bar{w}(d_i) = d_i$  and verify that it extends uniquely to a conjugate-linear anti-involution.

Let  $V(\lambda_i)$  be highest weight unitarizable  $\mathcal{G}-$  module with respect to  $w$ . Then it is a standard fact that  $V = \otimes V(\lambda_i)$  is a unitarizable  $\mathcal{G}_k-$  module with respect to  $w_k$ .

**2.3. Proposition.**  $(V_A, \pi)$  is a unitarizable  $\tilde{\mathcal{G}}_A-$  module with respect to  $\bar{w}$  if  $|a_i(\ell)| = |a_j(\ell)|$  for all  $i, j$  and  $\ell$ .

*Proof.* Define an automorphism  $\mathcal{O} : \mathcal{G}_A \rightarrow \mathcal{G}_A$  by  $\mathcal{O}(X(\underline{n})) = |a_1^{-2n}| X(\underline{n})$ . Then the following diagram commutes.

$$(2.4) \quad \begin{array}{ccc} \mathcal{G}_A & \xrightarrow{\bar{w}} & \mathcal{G}_k \\ \bar{w} \downarrow & & \downarrow w_k \\ \mathcal{G}_A & \xrightarrow{\phi \circ \mathcal{O}} & \mathcal{G}_k \end{array}$$

**2.5.** Hence we have  $\varphi \circ \mathcal{O} \circ \bar{w} = w_k \circ \bar{\varphi}$ . Also we have  $\varphi \circ \mathcal{O} \circ \bar{w}X(\underline{k}) = |a_1^{2k}| \varphi \circ \bar{w}X(\underline{k})$ .

Let  $(,)$  be a positive definite hermitian form on  $V$  satisfying

$$(2.6) \quad (Xv_1, v_2) = (v_1, w_k Xv_2), X \in \mathcal{G}_k, v_1, v_2 \in V.$$

Define a positive definite hermitian form on  $V_A$  by

$$\langle v_1(\underline{m}), v_2(\underline{n}) \rangle = \begin{cases} |a_1^{-2m}| (v_1, v_2) & \text{if } \underline{m} = \underline{n} \\ 0 & \text{otherwise.} \end{cases}$$

The following can be easily seen.

$$(1) \quad \langle \pi d_i v_1(\underline{n}), v_2(\underline{m}) \rangle = \langle v_1(\underline{n}), \pi d_i v_2(\underline{m}) \rangle = \begin{cases} 0 & \text{if } \underline{n} \neq \underline{m} \\ n_i \langle v_1(\underline{n}), v_2(\underline{n}) \rangle & \text{if } \underline{n} = \underline{m} \end{cases}$$

$$(2) \langle \pi X(\underline{k})v_1(\underline{n}), v_2(\underline{m}) \rangle = \langle v_1(\underline{n}), \pi \circ \bar{w}X(\underline{k})v_2(\underline{m}) \rangle = 0 \text{ if } \underline{k} + \underline{n} \neq \underline{m}.$$

Let  $v_1, v_2 \in V, X \in \mathcal{G}, \underline{k}, \underline{n} \in \mathbb{Z}^d$  and  $\underline{m} = \underline{k} + \underline{n}$ .

Consider

$$\begin{aligned} \langle \pi X(\underline{k}).v_1(\underline{n}), v_2(\underline{m}) \rangle &= \langle (\varphi X(\underline{k})v_1)(\underline{k} + \underline{n}), v_2(\underline{m}) \rangle \\ &= |a_1^{-2(\underline{n}+\underline{k})}| (\varphi X(\underline{k})v_1, v_2) \text{ (by def)} \\ &= |a_1^{-2(\underline{n}+\underline{k})}| (v_1, w_k \varphi X(\underline{k})v_2) \text{ (by 2.6)} \\ &= |a_1^{-2(\underline{n}+\underline{k})}| (v_1, \varphi \circ \mathcal{O} \circ \bar{w}X(\underline{k})v_2) \text{ (by (2.5))} \\ &= |a_1^{-2\underline{n}}| (v_1, \varphi \circ \bar{w}X(\underline{k})v_2) \text{ (by (2.5))} \\ &= \langle v_1(\underline{n}), (\varphi \circ \bar{w}X(\underline{k})v_2)(\underline{n}) \rangle \text{ (by def)} \\ &= \langle v_1(\underline{n}), \pi \bar{w}X(\underline{k})v_2(\underline{n} - \underline{k}) \rangle \text{ (by def of } \pi). \end{aligned}$$

It now follows that  $V_A$  is unitarizable as  $\tilde{\mathcal{G}}_A$ - module.

### 3.

We will now use the results of Section 1 and 2 to produce irreducible unitarizable modules for Toroidal algebras.

First recall the construction of a Toroidal algebra  $\tau_{[d]}$  from Section 2 of [MEY].

$\tau_{[d]} = \mathcal{G} \otimes A \oplus \Omega_A / d_A$  where  $\mathcal{G}$  is finite dimensional simple Lie-algebra and  $A$  is a Laurent polynomial ring in  $d$  variables.  $\Omega_d := \Omega_A / d_A$  is central in  $\tau_{[d]}$ .  $\Omega_d$

is a linear space spanned by  $d(t^{\underline{m}})t^{\underline{n}}$  with the relation  $d(t^{\underline{m}})t^{\underline{n}} + d(t^{\underline{n}})t^{\underline{m}} = 0$ . The Lie-algebra structure on  $\tau_{[d]}$  is defined as

$$[X(\underline{m}), Y(\underline{n})] = [X, Y](\underline{m} + \underline{n}) + (X, Y)d(t^{\underline{m}})t^{\underline{n}}$$

where  $(,)$  is a Killing form on  $\mathcal{G}$ .

Let  $d_1, d_2, \dots, d_d$  be derivations on  $\tau_{[d]}$  defined by

$$[d_i, X(\underline{m})] = m_i X(\underline{m}), [d_i, d(t^{\underline{m}})t^{\underline{n}}] = (m_i + n_i)d(t^{\underline{m}})t^{\underline{n}}, [d_i, d_j] = 0.$$

Let  $D$  be the linear span of  $d_1, d_2, \dots, d_d$ . Then  $\tilde{\tau}_{[d]} := \tau_{[d]} \oplus D$  is a Lie-algebra and  $\Omega_d$  is an *abelian ideal*.

**3.1.** Let  $w$  be any conjugate-linear anti-involution of  $\mathcal{G}$  and let  $\bar{w}$  be a conjugate-linear anti-involution of  $\tilde{\mathcal{G}}_A$  as defined in Section 2. Extend  $\bar{w}$  to  $\tilde{\tau}_{[d]}$  by

$$\bar{w}.d(t^{\underline{m}})t^{\underline{n}} = d(t^{-\underline{m}})t^{-\underline{n}}.$$

We also note that  $\tilde{\tau}_{[d]}/\Omega_d \cong \tilde{\mathcal{G}}_A$ . Let  $S$  denote the quotient map. So that any  $\tilde{\mathcal{G}}_A$ - module can be treated as  $\tilde{\tau}_{[d]}$  module via  $S$ .

Now let  $(V(\lambda_i), \tau_i)$  be the highest weight module for  $\mathcal{G}$ . Let  $V = \otimes_i V(\lambda_i)$  and  $V_A = V \otimes A$ . Then  $(V_A, \pi)$  is a  $\tilde{\tau}_{[d]}$ - module in the following way.

$$(3.2) \pi X(\underline{m}).(v_1 \otimes \dots \otimes v_k)(\underline{n}) = \sum a_i^{\underline{m}}(v_1 \otimes \dots \otimes \tau_i(X)v_i \otimes \dots \otimes v_k)(\underline{n} + \underline{m})$$

$$\begin{aligned} \pi d_i(v_1 \otimes \dots \otimes v_k)(\underline{n}) &= n_i(v_1 \otimes \dots \otimes v_k)(\underline{n}) \\ \pi \Omega_d.V_A &= 0. \end{aligned}$$

This is precisely the definition given in Section 1.

As earlier let  $\psi$  be the  $\mathbb{Z}^d$ - graded algebra homomorphism from  $U(\mathfrak{h}_A) \rightarrow A$  given by  $h(\underline{m}) \mapsto \sum \lambda_i(h)a_i^{\underline{m}}t^{\underline{m}}$ . Denote the image by  $A_\psi$ .

**3.3. Theorem.** (1)  $(\dot{V}_A, \pi)$  is a completely reducible  $\tau_{[d]}$ - module if  $A_\psi$  is irreducible as  $\mathfrak{h} \otimes A$ - module.

(2)  $(V_A, \pi)$  is irreducible as  $\tilde{\tau}_{[d]}$ -module if  $a_i(\ell)/a_j(\ell), i \neq j$ , is not a  $k$ -th root of unity.

(3) If  $V(\lambda_i)$  is unitarizable highest weight module for  $\mathcal{G}$  with respect to the form  $w$  and  $|a_i(\ell)| = |a_j(\ell)|$  for all  $i, j$  and  $\ell$ , then  $(V_A, \pi)$  is unitarizable  $\tilde{\tau}_{[d]}$ - module with respect to  $\bar{w}$ .

*Proof.* (1) and (2) follows from Theorem (1.8) by taking Kac-Moody Lie-algebra  $\mathcal{G}$  to be the finite dimensional simple Lie-algebra  $\mathcal{G}$ .

(3) Follows from Proposition (2.3). □

The special case where  $d = 1$  and  $w$  is a compact form of  $\mathcal{G}$  is due to [CP]. The case  $d = 1$  and  $w$  any form, including twisted affine Lie-algebras is due to [E].

**3.4.** Let  $\mathcal{G}$  be finite dimensional simple Lie-algebra and let

$$\bar{\mathcal{G}} = \mathcal{G} \otimes \mathbb{C}[t_{d+1}, t_{d+1}^{-1}] \oplus \mathbb{C}c$$

be the corresponding nontwisted affine Lie-algebra. Let  $\tilde{\mathcal{G}} = \bar{\mathcal{G}} \oplus \mathbb{C}d$  where  $d$  is a derivation. Consider the Lie-algebra  $\bar{\mathcal{G}}_A$  and observe that the center  $\Omega^1$  of  $\bar{\mathcal{G}}_A$  is spanned by  $\{c(\underline{m}), \underline{m} \in \mathbb{Z}^d\}$ . Let  $A^1 = A_{d+1}$  be the Laurent polynomials ring in  $d + 1$  variables. Clearly  $\bar{\mathcal{G}}_A/\Omega^1 \simeq \mathcal{G}_{A^1}$ . Let  $\phi$  denote the quotient map from  $\bar{\mathcal{G}}_A \rightarrow \mathcal{G}_{A^1}$  which is clearly a central extension. Since  $\tau_{[d+1]}$  is the universal central extension of  $\mathcal{G}_{A^1}$  (see Proposition 2.2 of [MEY]), there exists a homomorphism  $\varphi^1 : \tau_{[d+1]} \rightarrow \bar{\mathcal{G}}_A$  such that  $\ker \varphi^1$  is central. It is not difficult to write down the homomorphism (see (5.4)) and then one can see that the  $\ker \varphi^1$  consists of homogeneous elements with respect to the  $\mathbb{Z}^{d+1}$ -grading. Hence we can extend the homomorphism  $\varphi^1 : \tilde{\tau}_{[d+1]} \rightarrow \tilde{\mathcal{G}}_A$  by sending  $d_i \mapsto d_i (1 \leq i \leq d)$  and  $d_{d+1} \mapsto d$ . Also any  $\tilde{\mathcal{G}}_A$ -module can be treated as  $\tilde{\tau}_{[d+1]}$  via  $\varphi^1$ .

**3.5.** Now let  $(V(\lambda_i), \tau_i)$  be an highest weight module for  $\tilde{\mathcal{G}}$  and let  $V = \otimes_i V(\lambda_i)$  and  $V_A = V \otimes A$ . Then  $(V_A, \pi^1)$  is a  $\tilde{\tau}_{[d+1]}$ -module in the following way.

$$\begin{aligned} \pi^1 X(\underline{m})(v_1 \otimes \cdots \otimes v_k)(\underline{n}) &= \sum a_i^{\underline{m}}(v_1 \otimes \cdots \otimes \tau_i(X)v_i \otimes \cdots \otimes v_k)(\underline{n} + \underline{m}) \\ \pi^1 d_i(v_1 \otimes \cdots \otimes v_k)(\underline{n}) &= n_i(v_1 \otimes \cdots \otimes v_k)(\underline{n}) \quad (1 \leq i \leq d) \\ \pi^1 d(v_1 \otimes \cdots \otimes v_k)(\underline{n}) &= \sum_i (v_1 \otimes \cdots \otimes \tau_i(d)v_i \otimes \cdots \otimes v_k)(\underline{n}) \\ \pi^1 \ker \varphi^1 V_A &= 0 \end{aligned}$$

for all  $X \in \bar{\mathcal{G}}, v_i \in V$  and  $\underline{m}, \underline{n} \in \mathbb{Z}^d$ .

This definition is precisely the one given at (1.5) by taking the Kac-Moody Lie-algebra  $\mathcal{G}$  to be the affine Lie-algebra  $\bar{\mathcal{G}}$ .

Let  $\underline{h}$  be a Cartan subalgebra of  $\bar{\mathcal{G}}$  and let  $\psi$  be  $\mathbb{Z}^d$ -graded algebra homomorphism from  $U(\underline{h}_A) \rightarrow A$  given by  $h(\underline{m}) \mapsto \sum \lambda_i(h) a_i^{\underline{m}} t^{\underline{m}}$  and denote the image by  $A_\psi$ .

**3.6. Theorem.** (1)  $(V_A, \pi)$  is a completely reducible  $\tilde{\tau}_{[d+1]}$  module if  $A_\psi$  is an irreducible  $\underline{h}_A$ -module.

(2)  $(V_A, \pi)$  is irreducible as a  $\tilde{\tau}_{[d+1]}$ -module if  $a_i(\ell)/a_j(\ell), i \neq j$  is not a  $k$ th root of unity.

(3) If  $V(\lambda_i)$  is a unitarizable highest weight module for  $\bar{\mathcal{G}}$  with respect to the form  $w$  and  $|a_i(\ell)| = |a_j(\ell)|$  for all  $i, j$  and  $\ell$ , then  $(V_A, \pi^1)$  is unitarizable  $\tilde{\tau}_{[d+1]}$ -module with respect to  $\bar{w}$ .

*Proof.* (1) and (2) follows from Theorem (1.8) by taking the Kac-Moody Lie-algebra  $\mathcal{G}$  to be the affine Lie-algebra  $\bar{\mathcal{G}}$ .

(3) Follows from Proposition (2.3).

**3.7. Remark.** (1) The modules considered in Theorem (3.3) and (3.6) for Toroidal algebras are different. In the first case the full centre acts trivially where as in the second case a part of the centre acts non trivially.

(2) These are the first known irreducible module for Toroidal Lie-algebras where part of the center acts non trivially. The faithful modules constructed in [MEY] and [EM] through the use of vertex operators are all reducible. We will prove towards the end of the paper that the modules constructed in [MEY] and [EM] admits a filtration such that the successive irreducible quotient modules are isomorphic to the one considered here upto an automorphism of the Toroidal Lie-algebra.

4.

In this section we briefly review the construction of Fock space and the vertex operators that act on it. The theory is due to Frenkel-Kac [FK]. For further details one may also consult [FLM], [GO] and [MEY]. We will closely follow the notation from [EM]. We will also construct some automorphisms of Toroidal Lie-algebras.

**4.1.** Let  $Q$  be root lattice of the type  $ADE$  and let  $\alpha_1, \alpha_2, \dots, \alpha_\ell$  be a  $\mathbb{Z}$ -basis. Let  $A = (a_{ij})$  be the cartan matrix and  $a_{ij} = (\alpha_i | \alpha_j)$ . Let  $\Gamma$  be a free  $\mathbb{Z}$ - module on generators  $\alpha_1, \alpha_2, \dots, \alpha_\ell, \delta_1, \delta_2, \dots, \delta_{n-1}, d_1, d_2, \dots, d_{n-1}$ . Let  $(. | .)$  be a  $\mathbb{Z}$ -valued symmetric bilinear form such that  $(\delta_i | \delta_j) = (d_i | d_j) = (\alpha_i | \delta_j) = (\alpha_i | d_j) = 0, (\delta_i | d_j) = \delta_{i,j}$  and  $(\alpha_i | \alpha_j) = a_{ij}$ . Let  $Q$  be the sublattice generated by  $\alpha_1, \alpha_2, \dots, \alpha_\ell, \delta_1, \dots, \delta_{n-1}$ . Here  $n$  is any positive integer.

Let  $\underline{t} = \mathbb{C} \otimes_{\mathbb{Z}} \Gamma, \underline{h} = \mathbb{C} \otimes_{\mathbb{Z}} Q$  and  $\underline{c} = \mathbb{C} \otimes_{\mathbb{Z}} Q$ . We will define a Heisenberg algebra structure  $\hat{\underline{b}} = \oplus_{k \in \mathbb{Z}} \underline{t}(k) \oplus \mathbb{C}c$  where each  $\underline{t}(k)$  is an isomorphic copy of  $\underline{t}$  and the isomorphism is by  $\alpha \mapsto \alpha(k)$ . The Lie-algebra structure is defined by

$$[\alpha(k), \beta(m)] = k(\alpha | \beta)\delta_{k+m,0}c.$$

Let  $\hat{\underline{a}} = \oplus_{k \in \mathbb{Z}} \underline{h}(k) \oplus \mathbb{C}c$ . Define  $\underline{b} = \oplus_{k \neq 0} \underline{t}(k) \oplus \mathbb{C}c \subseteq \hat{\underline{b}}$  and  $\underline{b}_{\pm} = \oplus_{k > 0} \underline{t}(k)$ .

Similarly define  $\underline{a}, \underline{a}_{\pm}$  by replacing  $\underline{t}$  by  $\underline{h}$  and  $\underline{a}$  by replacing  $\underline{t}$  by  $\underline{h}$ .

The Fock space representation of  $\underline{b}$  is the symmetric algebra  $S(\underline{b}-)$  of  $\underline{b}-$  together with an action of  $\underline{b}$  on  $S(\underline{b}-)$  defined by the following:

$c$  acts as 1,  $a(-m)$  acts as multiplication by  $a(-m), m > 0$   
 $a(m)$  acts as the unique derivation on  $S(\underline{b}-)$  for which  $b(-n) \rightarrow \delta_{m,-n}m(a | b).$  ( $a, b \in \underline{t}, m, n > 0$ ).

$S(\underline{b}-)$  affords an irreducible representation of  $\underline{b}$ . However  $S(\underline{a}-)$  does not afford an irreducible representation of  $\underline{a}$  since the form is degenerate on  $\underline{h}$ .

A vector  $\delta$  in  $Q$  is called a null root if  $\delta = n_1\delta_1 + n_2\delta_2 + \dots + n_{n-1}\delta_{n-1}$  for some integers  $n_i$ . Note that  $(\delta | \delta) = 0$ .

Following [EMY] we let  $\epsilon : Q \times Q \rightarrow \{\pm 1\}$  be a bi- multiplicative map satisfying

- (1)  $\epsilon(\alpha, \alpha) = (-1)^{(\alpha|\alpha)}$
- (2)  $\epsilon(\alpha, \beta)\epsilon(\beta, \alpha) = (-1)^{(\alpha|\beta)}$
- (3)  $\epsilon(\alpha, \delta) = 1$ ,

where  $\alpha, \beta \in Q$  and  $\delta$  a null root. Extend  $\epsilon$  to a bimultiplicative map of  $Q \times \Gamma \rightarrow \{\pm 1\}$  and form the vector space  $\mathbb{C}[\Gamma]$ , with  $\mathbb{C}$  basis  $\{e^\nu, \nu \in \Gamma\}$ . Then  $C[\Gamma]$  contains the space  $\mathbb{C}[Q]$  similarly defined. Following Borchers [B] make  $\mathbb{C}[\Gamma]$  a  $\mathbb{C}[Q]$  module by defining

$$e^\alpha \cdot e^\nu = \epsilon(\alpha, \nu)e^{\alpha+\nu} (\alpha \in Q, \nu \in \Gamma).$$

Let  $M \subseteq S(\underline{b}-)$  be any  $\underline{a}-$  submodule (with respect to the Fock space action). We define  $V(\Gamma, M) = \mathbb{C}[\Gamma] \otimes_{\mathbb{C}} M$ . Of particular interest in the sequel will be  $V(\Gamma, S(\underline{a}-))$  and  $V(\Gamma, S(\underline{b}-))$  which we simply denote by  $V(\Gamma)$  and  $V(\Gamma, \underline{b})$  respectively. We extend the action of  $\underline{a}$  on  $M$  to  $\hat{\underline{a}}$  on  $V(\Gamma, M)$  by

$$\begin{aligned} a(m)(e^\gamma \otimes u) &= e^\gamma \otimes a(m)u, m \neq 0 \\ a(0)e^\gamma \otimes u &= (a | \gamma)e^\gamma \otimes u. \end{aligned}$$

**Vertex operators:** Let  $z$  be a complex valued variable and let  $\alpha \in Q$ . Define

$$T_{\pm}(\alpha, z) = - \sum_{\substack{n \\ n > 0}} \frac{1}{n} \alpha(n) z^{-n}.$$

Then the vertex operator for  $\alpha$  in  $Q$  is defined as  $X(\alpha, z) = z^{\frac{(\alpha|\alpha)}{2}} \exp T(\alpha, z)$  where  $\exp T(\alpha, z) = \exp T_-(\alpha, z)e^{\alpha}z^{\alpha(0)} \exp T_+(\alpha, z)$  and the operators  $z^{\alpha(0)}$  is defined by

$$z^{\alpha(0)}e^\gamma \otimes u = z^{(\alpha|\gamma)}e^\gamma \otimes u.$$

$X(\alpha, z)$  can be formally expanded in powers of  $z$  to give

$$X(\alpha, z) = \sum_{n \in \mathbb{Z}} X_n(\alpha) z^{-n}$$

and the moments  $X_n(\alpha) = \frac{1}{2\pi i} \int X(\alpha, z) z^n \frac{dz}{z}$  are operators on  $V(\Gamma, M)$ .

Let  $\Delta$  be a root system and let  $X_\alpha, \alpha \in \Delta, h \in \underline{h}$  be a Chevalley basis. Let  $\underline{r} = (r_1, r_2, \dots, r_{n-1}) \in \mathbb{Z}^{n-1}$  and  $\bar{r} = (\underline{r}, r_n) \in \mathbb{Z}^n$ . Let  $\delta_{\underline{r}} = r_1\delta_1 + \dots + r_{n-1}\delta_{n-1}$ .

We will now recall the main Theorem 3.14 of [EM].

**4.2. Theorem.** *The Lie-algebra generated by operators  $X_m(\alpha + \delta)$ , ( $\alpha \in \Delta, m \in \mathbb{Z}$  and  $\delta$  a null root) on  $V(\Gamma, \underline{b})$  is isomorphic to the Toroidal Lie-algebra  $\tau_{[n]}$ . The isomorphism is given by*

$$\begin{aligned} X_{r_n}(\alpha + \delta_r) &\mapsto X_\alpha \otimes t^{\bar{r}}, \alpha \in \Delta \\ T_{r_n}^h(\delta_r) &\mapsto h \otimes t^{\bar{r}} \\ T_{r_n}^{\delta_i}(\delta_r) &\mapsto t^{\bar{r}} t_i^{-1} dt_i \quad (1 \leq i \leq n-1) \\ X_{r_n}(\delta_r) &\mapsto t^{\bar{r}} t_n^{-1} dt_n \end{aligned}$$

where  $t^{\bar{r}} = t_1^{r_1} t_2^{r_2} \dots t_n^{r_n}$  and

$$T_m^g(\alpha) = \frac{1}{2\pi i} \int : g(z) X(\alpha, z) : \frac{dz}{z} z^m.$$

**4.3.** Here we construct certain automorphisms  $\tau_{[d]}$  which are needed in our main result in section 5. Let  $SL(d, \mathbb{Z})$  be the group of matrices of order  $d \times d$  with integral entries and determinate 1.

Let  $A = (a_{ji})_{\substack{1 \leq i \leq d \\ 1 \leq j \leq d}}$  be an element of  $SL(d, \mathbb{Z})$ . Let  $\underline{r} = (r_1, r_2, \dots, r_d)$  and  $\underline{s} = (s_1, s_2, \dots, s_d)$  be such that  $r_i, s_i \in \mathbb{Z}$ . Let  $e_i = (0, \dots, 1, \dots, 0)$  where 1 is at the  $i$ th place. Let  $A\underline{r}^T = \underline{m}^T, A\underline{s}^T = \underline{n}^T$  and  $Ae_i^T = a(i)^T$  where  $\underline{m} = (m_1, m_2, \dots, m_d), \underline{n} = (n_1, n_2, \dots, n_d)$  and  $a(i) = (a_{1i}, a_{2i}, \dots, a_{di})$  and  $T$  denotes the transpose. We now define an automorphism of  $\tau_{[d]}$  again denoted by  $A$ .

$$\begin{aligned} A(X(\underline{r})) &= X(\underline{m}) \\ A(d(t^{\underline{r}})t^{\underline{s}}) &= d(t^{\underline{m}})t^{\underline{n}}. \end{aligned}$$

It is straightforward to check that  $A$  is an automorphism of  $\tau_{[d]}$ . We will now extend it to  $\tilde{\tau}_{[d]}$ . Let  $(d_1^1, d_2^1, \dots, d_d^1)^T = A^{-1}(d_1, d_2, \dots, d_d)^T$ . Define  $A(d_i) = d_i^1$  and check that it defines an automorphisms of  $\tilde{\tau}_{[d]}$ .

**Remark.**  $A$  does not quite preserve the natural  $\mathbb{Z}^d$ -gradation of  $\tau_{[d]}$ .

### 5.

The purpose of this section is to describe the subquotients of  $V(\Gamma) = \mathbb{C}[\Gamma] \otimes S(\underline{a}-)$ . We will follow the notation of §5 of [MEY].

Let  $\lambda \in \Gamma$  and define  $V(\lambda) := e^{\lambda+Q} \otimes S(\underline{a}-)$ . Note that  $V(\Gamma) = \bigoplus_{\lambda \in \Gamma/Q} V(\lambda)$ . Each  $V(\lambda)$  is a cyclic  $\tau_{[n]}$ -module with generator  $e^\lambda \otimes 1$ . Towards the end of the section we will exhibit a decreasing filtration of modules and describe its successive irreducible quotients.

Fix  $\lambda \in \Gamma$  and let  $W(\lambda) := W = e^{\lambda+Q} \otimes S(\underline{a}-)$ . This is a module for the non-twisted affine Lie-algebra  $\mathcal{G}_{aff} = \mathcal{G} \otimes \mathbb{C}[t_n, t_n^{-1}] \oplus \mathbb{C}c$  generated by the

operators  $X_n(\alpha), \alpha \in \Delta$ . But we know by Frenkel-Kac theory [FK] that this is an irreducible level 1 representation of  $\mathcal{G}_{aff}$ . Let  $V = V(\lambda)$ .

The following Lemma is trivial to see.

**5.1. Lemma.**

$$V = \bigoplus_{n_1, n_2, \dots, n_k > 0, k \geq 0} \delta_{i_1}(-n_1) \cdots \delta_{i_k}(-n_k) e^{\lambda+Q} \otimes S(\underline{a} -).$$

Let  $M = \bigoplus_{n_1, n_2, \dots, n_k > 0, k > 0} \delta_{i_1}(-n_1) \cdots \delta_{i_k}(-n_k) e^{\lambda+Q} \otimes S(\underline{a} -)$ .

By Lemma (5.1) it is easy to verify that  $M$  is a  $\tau_{[n]}$ - module.

We will recall the following Lemma (5.2) of [MEY], which though stated for the  $n = 2$  case there, is true for any  $n$ .

**5.2. Lemma .** *Let  $\lambda \in \Gamma$ , let  $\delta$  be a null root, let  $N = (\lambda \mid \delta)$  and let  $m \in \mathbb{Z}$ . Then*

$$X_m(\delta)e^\lambda \otimes 1 = \begin{cases} \epsilon(\delta, \lambda)e^{\lambda+\delta} \otimes S_{-m-N}(\delta), m + n < 0 \\ \epsilon(\delta, \lambda)e^{\lambda+\delta} \otimes 1, m + n = 0 \\ 0, m + n > 0 \end{cases}$$

where the operators  $S_p(\delta)$  are defined by  $\exp T_-(\delta, z) = \sum_{p=0}^{\infty} S_p(\delta)z^p$ .

**5.3.** Set  $\tau_i = X_{-N_i}(\delta_i)$  where  $N_i = (\lambda \mid \delta_i)$ . We note that  $\tau_i^k = X_{-kN_i}(k\delta_i)$ . Let  $\tau^{\underline{m}} = \tau_1^{m_1} \tau_2^{m_2} \cdots \tau_{n-1}^{m_{n-1}}$  and let

$$W \otimes t^{\underline{m}} := \tau^{\underline{m}}W = e^{\lambda + \sum m_i \delta_i + Q} \otimes S(\underline{a} -).$$

**Lemma.**  $V/M \cong W \otimes A_{[n-1]}$  as Vector spaces.

Our aim is to describe  $V/M$  as a  $\tilde{\tau}_{[n]}$ - module.

**5.4.** Note that  $\mathcal{G}_{aff} \otimes A_{[n-1]} = \dot{\mathcal{G}} \otimes A_{[n]} \oplus \sum_{\underline{m} \in \mathbb{Z}^{n-1}} \mathbb{C}c \otimes t^{\underline{m}}$ .

See Section (3.4) where we considered such a Lie-algebra.

Define a Lie-algebra homomorphism  $\varphi^1 : \tau_{[n]} \rightarrow \dot{\mathcal{G}}_{aff} \otimes A_{[n-1]}$  by

1)  $\varphi^1$  is Id. on  $\dot{\mathcal{G}} \otimes A_{[n]}$

$$2) \varphi^1(t^{\underline{m}}t_i^{-1}dt_i) = \begin{cases} 0 \text{ if } m_n \neq 0 \\ 0 \text{ if } m_n = 0, 1 \leq i \leq n - 1 \\ c \otimes t^{\underline{m}} \text{ if } m_n = 0 \text{ and } i = n. \end{cases}$$

**5.5.** Let  $(\bar{V}, \pi)$  be an irreducible highest weight module for  $\mathcal{G}_{aff}$ . Then  $\bar{V} \otimes A_{[n-1]}$  admits an action by  $\mathcal{G}_{aff} \otimes A_{[n-1]}$  in the following way,

$$X(\underline{m})v(\underline{n}) = Xv(\underline{m} + \underline{n}), (X \in \mathcal{G}_{aff}, v \in \bar{V}).$$

Now we extend the module action to one of  $\tau_{[n]}$  via the surjective homomorphism  $\varphi^1$ . That is

$$X.v(\underline{m}) = \varphi^1(X).v(\underline{m}).$$

It is easy to see that  $\bar{V} \otimes A_{[n-1]}$  admits a natural  $\mathbb{Z}^{n-1}$ - gradation and is graded irreducible as a  $\tau_{[n]}$ - module. In other words  $\tilde{\tau}_{[n]}$  irreducible (one can apply Theorem 3.6(2) taking  $d + 1 = n$  and  $k = 1$ ).

We can twist the module action of  $\tilde{\tau}_{[n]}$  by an automorphism  $A$  considered in (4.3), so that  $(\bar{V} \otimes A_{[n-1]}, Ao\varphi^1o\pi)$  is a  $\tilde{\tau}_{[n]}$  irreducible module.

With the notations in (5.3) we have the following result.

**5.6. Theorem.**  $V/M$  is isomorphic to  $(W \otimes A_{[n-1]}, Ao\varphi^1o\pi)$  as  $\tau_{[n]}$ - module where  $A = (b_{ij})$  and  $b_{ii} = 1, b_{in} = (\lambda \mid \delta_i), 1 \leq i \leq n - 1, b_{ij} = 0$  otherwise.

We need some lemmas before we give a proof of the Theorem.

**5.7. Lemma.** The following equality holds as operators on  $V/M$ .

- (1)  $\delta_{\underline{m}}(n) = \begin{cases} 0 & \text{if } n \neq 0 \\ (\lambda \mid \delta_{\underline{m}}) & \text{if } n = 0 \end{cases}$
- (2)  $X_n(\delta_{\underline{m}}) = \begin{cases} 0 & \text{if } n + (\lambda \mid \delta_{\underline{m}}) \neq 0 \\ \tau^{\underline{m}} & \text{if } n + (\lambda \mid \delta_{\underline{m}}) = 0 \end{cases}$
- (3)  $X_n(\alpha + \delta_{\underline{m}}) = X_{-(\lambda \mid \delta_{\underline{m}})}(\delta_{\underline{m}})X_{n+(\lambda \mid \delta_{\underline{m}})}(\alpha) = X_{n+(\lambda \mid \delta_{\underline{m}})}(\alpha)\tau^{\underline{m}}, \alpha \in \dot{\Delta}$ .
- (4)  $T_n^h(\delta_{\underline{m}}) = h(n + (\lambda \mid \delta_{\underline{m}}))X_{-(\lambda \mid \delta_{\underline{m}})}(\delta_{\underline{m}}) = h(n + (\lambda \mid \delta_{\underline{m}}))\tau^{\underline{m}}, h \in \dot{h}$ .

*Proof.* First observe that  $\delta_{\underline{m}}(n)$  and  $X_n(\delta_{\underline{m}})$  are central operators on  $V$  and hence are determined on the generator  $e^\lambda \otimes 1$ .

1. By definition  $\delta_{\underline{m}}(n)e^\lambda \otimes 1 = 0$  for  $n > 0$  and  $\delta_{\underline{m}}(0)e^\lambda \otimes 1 = (\lambda \mid \delta_{\underline{m}})e^\lambda \otimes 1$ . Hence  $\delta_{\underline{m}}(n)$  on  $V/M$  is zero for  $n > 0$  and  $(\lambda \mid \delta_{\underline{m}})$  for  $n = 0$ . For  $n < 0, \delta_{\underline{m}}(n)e^\lambda \otimes 1 = e^\lambda \otimes \delta_{\underline{m}}(n) \in M$  and hence  $\delta_{\underline{m}}(n)$  is zero on  $V/M$ .
2. The case  $n + (\lambda \mid \delta_{\underline{m}}) > 0$  follows from the Lemma (5.2). Also for the case  $n + (\lambda \mid \delta_{\underline{m}}) < 0$ , note that

$$X_n(\delta_{\underline{m}})e^\lambda \otimes 1 \in M \text{ (by Lemma 5.2).}$$

Hence  $X_n(\delta_{\underline{m}})$  is zero on  $V/M$ . Now let  $n + (\lambda \mid \delta_{\underline{m}}) = 0$ . First we will note that it will follow from the definition of vertex operators that

$X(\alpha + \beta, z) = X(\alpha, z)X(\beta, z)$  whenever  $(\alpha | \beta) = 0$ . Hence we have the following:

$$\begin{aligned} X_n(\delta_{\underline{m}}) &= \sum_{k_1+k_2+\dots+k_{n-1}=n} X_{k_1}(m_1\delta_1) \cdots X_{k_{n-1}}(m_{n-1}\delta_{n-1}) \\ &= \sum_{\substack{k_1+k_2+\dots+k_{n-1}=n, \\ k_i+m_i(\lambda | \delta_i) \leq 0}} X_{k_1}(m_1\delta_1) \cdots X_{k_{n-1}}(m_{n-1}\delta_{n-1}) \text{ (by Lemma 5.2)}. \end{aligned}$$

But  $0 = n + (\lambda | \delta_{\underline{m}}) = \sum_i (k_i + m_i(\lambda | \delta_i)) \leq 0$ . It follows that  $k_i + m_i(\lambda | \delta_i) = 0$  for all  $i$ . Now by Lemma 5.2 (2) we have  $X_{k_i}(m_i(\lambda | \delta_i)) = \tau_i^{m_i}$ . Hence  $X_n(\delta_{\underline{m}}) = \tau^{\underline{m}}$ .

3. Since  $(\alpha | \delta_{\underline{m}}) = 0, \alpha \in \Delta$  we have

$$\begin{aligned} X_n(\alpha + \delta_{\underline{m}}) &= \sum_k X_{m-k}(\alpha)X_k(\delta_{\underline{m}}) \\ &= X_{n+(\lambda|\delta_{\underline{m}})}(\alpha) X_{-(\lambda|\delta_{\underline{m}})}(\delta_{\underline{m}}) \text{ on } V/M \text{ (by(2))} \\ &= X_{n+(\lambda|\delta_{\underline{m}})}(\alpha) \tau^{\underline{m}} \text{ (by (2))}. \end{aligned}$$

4. The proof is similar to (3). We only note that

$$T_n^h(\delta_{\underline{m}}) = \sum_k h(n - k)X_n(\delta_{\underline{m}}) \text{ (by def).}$$

□

*Proof of the Theorem 5.6.* We will first describe the  $\tau_{[n]}$ - module  $(W \otimes A_{[n-1]}, A\circ\varphi^1\circ\pi)$ .

$$X(\bar{r})v(\underline{k}) = (\pi X \otimes t_n^{r_n+(\lambda|\delta_{\underline{r}})}v)(\underline{r} + \underline{k}).$$

$$d(t^{\bar{r}})t^{\bar{s}}v(\underline{k}) = \begin{cases} 0 & \text{if } r_n + s_n + (\lambda | \delta_{\underline{r}+\underline{s}}) \neq 0 \\ (r_n + (\lambda | \delta_{\underline{r}}))v(\underline{k} + \underline{r} + \underline{s}) & \text{if } r_n + s_n + (\lambda | \delta_{\underline{r}+\underline{s}}) = 0 \end{cases}$$

This follows from definition of  $A, \varphi^1$  and  $\pi$ .

We will now compute the action of  $\tau_{[n]}$  on  $V/M$  using Lemma (5.7) and then verify it to be exactly as above. That will complete the proof.

$$\begin{aligned} \text{(a)} \quad X_\alpha(\bar{r}).w(\underline{k}) &= X_{r_n}(\alpha + \delta_{\underline{r}})w(\underline{k}) \text{ (by Theorem 4.1)} \\ &= X_{r_n+(\lambda|\delta_{\underline{r}})}(\alpha)w(\underline{r} + \underline{k}) \text{ (by Lemma 5.7 and Notation at 5.3)} \\ &= \pi(X_\alpha \otimes t_n^{r_n+(\lambda|\delta_{\underline{r}})})w(\underline{r} + \underline{k}) \text{ (by [FK])}, \end{aligned}$$

$$\begin{aligned}
 \text{(b)} \quad h(\bar{r}) \cdot w(\underline{k}) &= T_{r_n}^h(\alpha + \delta_r)w(\underline{k}) \text{ (by Theorem 4.1)} \\
 &= h(r_n + (\lambda \mid \delta_r))w(\underline{r} + \underline{k}) \text{ (by Lemma 5.7 and Notation 5.3)} \\
 &= \pi(h \otimes t^{r_n + (\lambda \mid \delta_r)})w(\underline{r} + \underline{k}) \text{ (by [FK])},
 \end{aligned}$$

$$\begin{aligned}
 \text{(c)} \quad d(t^{\bar{r}}) \cdot t^{\bar{s}}w(\underline{k}) &= (T_{r_n+s_n}^{\delta_r}(\delta_{\underline{r}+\underline{s}}) + r_n X_{r_n+s_n}(\delta_{\underline{r}+\underline{s}}))w(\underline{k}) \text{ (by Theorem 4.1)} \\
 &= \delta_r(r_n + s_n + (\lambda \mid \delta_{\underline{r}+\underline{s}}))w(\underline{k} + \underline{r} + \underline{s}) \\
 &\quad + r_n X_{r_n+s_n}(\delta_{\underline{r}+\underline{s}})w(\underline{k} + \underline{r} + \underline{s}) \text{ (by Lemma 5.7 and Notation at 5.3)} \\
 &= \begin{cases} ((\lambda \mid \delta_r) + r_n)w(\underline{k} + \underline{r} + \underline{s}) & \text{if } r_n + s_n + (\lambda \mid \delta_{\underline{r}+\underline{s}}) = 0 \\ 0 & \text{otherwise} \end{cases}
 \end{aligned}$$

(follows from Lemma 5.7 (1) and 5.7 (2)). □

**5.8. Remark.** Note that the derivations  $d_1, d_2, \dots, d_{n-1}$  act on  $V$  as  $d_i(0) \cdot e^\lambda \otimes a = (\lambda \mid d_i)e^\lambda \otimes a$  and remain the same on the quotient  $V/M$ . Also note that  $A^{-1}d_i = d_i^1 = d_i(1 \leq i \leq n-1)$  (see 4.3). Hence the isomorphism in the theorem preserves the natural  $\mathbb{Z}^{n-1}$  gradation.

**5.9. Remark.** Let  $V_N = \oplus_{k \geq N} \delta_{i_1}(-n_1) \cdots \delta_{i_k}(-n_k)e^{\lambda+Q} \otimes S(\underline{a}-)$  which is a  $\tau_{[n]}$  submodule of  $V$ . Further  $V = V_o \supseteq V_1 \supseteq V_2 \supseteq \dots$  is decreasing sequence of  $\tau_{[n]}$ -modules. Consider  $V_N/V_{N+1} = \oplus \delta_{i_1}(-n_1) \cdots \delta_{i_N}(-n_N)e^{\lambda+Q} \otimes S(\underline{a}-)$  as  $\tau_{[n]}$ -module. Then for a fixed  $i_1, i_2, \dots, i_N$  and  $n_1, n_2, \dots, n_N$ ,  $F = \delta_{i_1}(-n_1) \cdots \delta_{i_N}(-n_N)e^{\lambda+Q} \otimes S(\underline{a}-)$  is a  $\tau_{[n]}$  submodule of  $V_N/V_{N+1}$ . It is also easy to see that  $F \cong V/M$  as  $\tau_{[n]}$ -module. In other words there exists a filtration of  $\tau_{[n]}$  submodules of  $V$  such that the successive quotients are  $\mathbb{Z}^{n-1}$ -graded irreducible and isomorphic to  $V/M$ .

**5.10.** Recall  $d_1, d_2, \dots, d_{n-1}$  from Section 4. Let  $N \geq 0$  and let

$$W_N = \oplus_{0 \leq k \leq N} d_{i_1}(-n_1) \cdots d_{i_k}(-n_k)V.$$

Clearly  $V = W_o \subseteq W_1 \subseteq W_2 \subseteq \dots$ . We will first prove that  $W_N$  is a  $\tau_{[n]}$ -module. Since  $X_n(\alpha)$  generate  $\tau_{[n]}$  as a Lie-algebra (see Theorem 3.14 of [EM]) it is sufficient to prove the

**5.11. Lemma.** For  $\alpha, m, k$  and for all  $v$  in  $V$

$$X_m(\alpha)d_{i_1}(-n_1) \cdots d_{i_k}(-n_k)v \in W_k.$$

*Proof.* By induction on  $k$ . Clearly this is true for  $k = 0$ . For  $k = 1$ ,

$$X_m(\alpha)d_{i_1}(-n_1)v = d_{i_1}(-n_1)X_m(\alpha)v + (\alpha \mid d_{i_1})X_{m-n_1}(\alpha)v$$

(by 3.8 (1) of [EM]). Clearly  $d_{i_1}(-n_1)X_n(\alpha)v \in W_1$  and  $X_{m-n_1}(\alpha) \in W_o \subseteq W_1$ . Hence we are done.

We will now assume the Lemma for all  $\ell, 1 \leq \ell \leq k$  and prove it for  $\ell = k + 1$ .

Consider

$$\begin{aligned} & X_m(\alpha)d_{i_1}(-n_1) \cdots d_{i_{k+1}}(-n_{k+1})v \\ &= \sum_{1 \leq j \leq k+1} T_j + d_{i_1}(-n_1) \cdots d_{i_{k+1}}(-n_{k+1})X_m(\alpha)v \end{aligned}$$

where  $T_j = d_{i_1}(-n_1) \cdots [X_m(\alpha), d_{i_j}(-n_j)] \cdots d_{i_{k+1}}(-n_{k+1})v$ . For  $1 \leq j \leq k + 1$ , we have  $k + 1 - j \leq k$  and by induction hypothesis we have

$$(\alpha \mid d_j)X_{m-n_j}(\alpha)d_{i_{j+1}}(-n_{j+1}) \cdots d_{i_{k+1}}(-n_{k+1})v \in W_{k+1-j}.$$

Hence  $T_j \in W_k \subseteq W_{k+1}$ . (See 3.8 (1) of [EM]). This completes the proof of the Lemma. □

From the above we also have

$$(5.12) \quad X_m(\alpha)d_{i_1}(-n_1) \cdots d_{i_{k+1}}(-n_{k+1})v = d_{i_1}(-n_1) \cdots d_{i_{k+1}}(-n_{k+1})X_m(\alpha)v \text{ in } W_{k+1}/W_k.$$

Further  $W_{k+1}/W_k = \oplus d_{i_1}(-n_1) \cdots d_{i_{k+1}}(-n_{k+1})V$  and from (5.12) each  $d_{i_1}(-n_1) \cdots d_{i_{k+1}}(-n_{k+1})V$  is a submodule of  $W_{k+1}/W_k$  isomorphic to  $V$ .

Put  $\cup_{i \geq 0} W_i = \underline{b}(\lambda)$  and remember each  $W_i$  depends on  $V$  and  $V$  in turn depends on  $\lambda$ .

**5.13. Remark.** (1) The full Fock space  $V(\Gamma, \underline{b}) = \oplus_{\lambda \in \Gamma/Q} \underline{b}(\lambda)$  as  $\tau_{[n]}$ -module.

(2) Each  $\underline{b}(\lambda)$  admits a filtration by an increasing sequence of modules whose successive quotients are isomorphic to  $V$  (see above).

(3) Each  $V$  above admits a filtration by a decreasing sequence of modules such that the successive quotients are irreducible (see Remark 5.9).

### 6. Acknowledgements.

This paper is dedicated for those courageous people who believed in me and helped me in my most testing year 1984-85.

### References

- [B] R. Borcherds, *Vertex algebras, Kac-Moody algebras and the Monster*, Proc. Natl. Acad. Sci. USA, **83** (1986), 3068-3071.
- [BC] S. Berman and B. Cox, *Enveloping algebras and representations of toroidal Lie-algebras*, Pacific J. Math., **165** (1994), 239-267.

- [CP] V. Chari and A. Pressley, *New Unitary Representations of Loop Groups*, Math. Ann., **275** (1986), 87-104.
- [E] S. Eswara Rao, *On Representations of Loop Algebras*, Communications in Algebra, **21(6)** (1993), 2131-2153.
- [EM] S. Eswara Rao and R.V. Moody, *Vertex Representations for  $N$ -Toroidal Lie-algebras and a generalization of the Virasoro algebra*, Commun. Math. Phys., **159** (1994), 239-264.
- [EMY] S. Eswara Rao, R.V. Moody and T. Yokonuma, *Lie algebras and Weyl groups arising from Vertex operator representations*, Nova J. of Algebra and Geometry, **1** (1992), 15-58.
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Academic Press, Boston, 1989.
- [FM] M. Fabbri and R.V. Moody, *Irreducible representations of Virasoro-toroidal Lie algebra*, **159** (1994), 1-13.
- [FK] I. Frenkel and V.G. Kac, *Basic representations of Affine Lie-algebras and dual resonance models*, Invent. Math., **62** (1980), 23-66.
- [GO] P. Goddard and D. Olive, *Algebras, lattices and strings*, Vertex operators in Mathematics and Physics, Publ. Math. Sci. Res. Inst., **3**, Springer-Verlag, 1984.
- [KMPS] S. Kass, R.V. Moody, J. Patera and R. Slansky, *Representations of Affine Algebras and Branching Rules*, University of California Press, Berkeley, 1990.
- [MEY] R.V. Moody, S. Eswara Rao and T. Yokonuma, *Toroidal Lie-algebras and Vertex representations*, Geom. Ded., **35** (1990), 283-307.
- [MS] R.V. Moody and Z. Shi, *Toroidal Weyl groups*, Nova J. Algebra and Geometry, **I** (1992), 317-337.

Received May 20, 1993 and revised February 16, 1994.

TATA INSTITUTE OF FUNDAMENTAL RESEARCH  
HOMI BHABHA ROAD  
BOMBAY - 5  
INDIA  
*E-mail address:* Senapati@tifrvax.bitnet

## PACIFIC JOURNAL OF MATHEMATICS

Founded by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

### EDITORS

Sun-Yung Alice Chang (Managing Editor)  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

F. Michael Christ  
University of California  
Los Angeles, CA 90095-1555  
christ@math.ucla.edu

Robert Finn  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

Martin Scharlemann  
University of California  
Santa Barbara, CA 93106  
mgscharl@math.ucsb.edu

Thomas Enright  
University of California  
San Diego, La Jolla, CA 92093  
tenright@ucsd.edu

Vaughan F. R. Jones  
University of California  
Berkeley, CA 94720  
vfr@math.berkeley.edu

Gang Tian  
Courant Institute  
New York University  
New York, NY 10012-1100  
tiang@taotao.cims.nyu.edu

Nicholas Ercolani  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

Steven Kerckhoff  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

V. S. Varadarajan  
University of California  
Los Angeles, CA 90095-1555  
vsv@math.ucla.edu

### SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
STANFORD UNIVERSITY  
UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
UNIVERSITY OF CALIFORNIA  
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA  
UNIVERSITY OF NEVADA, RENO  
UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
UNIVERSITY OF UTAH  
UNIVERSITY OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$108.00 a year to individual members of supporting institutions.

Subscriptions, orders for back issues published within the last three years, and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,  
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,  
the American Mathematical Society's TEX macro system.  
Copyright © 1995 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 171 No. 2 December 1995

---

On $H^p$ -solutions of the Bezout equation	297
ERIC AMAR, JOAQUIM BRUNA FLORIS and ARTUR NICOLAU	
Amenable correspondences and approximation properties for von Neumann algebras	309
CLAIRE ANANTHARAMAN-DELAROCHE	
On moduli of instanton bundles on $\mathbb{P}^{2n+1}$	343
VINCENZO ANCONA and GIORGIO MARIA OTTAVIANI	
Minimal surfaces with catenoid ends	353
JORGEN BERGLUND and WAYNE ROSSMAN	
Permutation model for semi-circular systems and quantum random walks	373
PHILIPPE BIANE	
The Neumann problem on Lipschitz domains in Hardy spaces of order less than one	389
RUSSELL M. BROWN	
Matching theorems for twisted orbital integrals	409
REBECCA A. HERB	
Uniform algebras generated by holomorphic and pluriharmonic functions on strictly pseudoconvex domains	429
ALEXANDER IZZO	
Quantum Weyl algebras and deformations of $U(g)$	437
NAIHUAN JING and JAMES ZHANG	
Calcul du nombre de classes des corps de nombres	455
STÉPHANE LOUBOUTIN	
On geometric properties of harmonic $Lip_1$ -capacity	469
PERTTI MATTILA and P. V. PARAMONOV	
Reproducing kernels and composition series for spaces of vector-valued holomorphic functions	493
BENT ØRSTED and GENKAI ZHANG	
Iterated loop modules and a filtration for vertex representation of toroidal Lie algebras	511
S. ESWARA RAO	
The intrinsic mountain pass	529
MARTIN SCHECHTER	
A Frobenius problem on the knot space	545
RON G. WANG	
On complete metrics of nonnegative curvature on 2-plane bundles	569
DAVID YANG	
Correction to: "Free Banach-Lie algebras, couniversal Banach-Lie groups, and more"	585
VLADIMIR G. PESTOV	
Correction to: "Asymptotic radial symmetry for solutions of $\Delta u + e^u = 0$ in a punctured disc"	589
KAI SENG (KAISING) CHOU (TSO) and TOM YAU-HENG WAN	



0030-8730(1995)171:2;1-G