

*Pacific
Journal of
Mathematics*

THE INTRINSIC MOUNTAIN PASS

MARTIN SCHECHTER

Volume 171 No. 2

December 1995

THE INTRINSIC MOUNTAIN PASS

MARTIN SCHECHTER

We show how the mountain pass and saddle point theorems can be formulated with out the use of “auxiliary” sets. Moreover, we show that results can still be obtained when some basic hypotheses of these theorems are not satisfied. We then apply our results to semilinear problems for partial differential equations.

1. Introduction.

In the mountain pass and saddle point theorems one is concerned with a C^1 functional G on a Banach space E . One wishes to find a solution of $G'(u) = 0$ or at least a sequence $\{u_k\} \subset E$ such that

$$(1.1) \quad G(u_k) \rightarrow c, \quad G'(u_k) \rightarrow 0$$

for some $c \in R$. A general procedure was formulated in Brezis-Nirenberg [BN] as follows. One finds a compact metric space K and selects a closed subset K^* of K such that $K^* \neq \emptyset, K^* \neq K$. One then picks a map $p^* \in C(K^*, E)$ and defines

$$A = \{p \in C(K, E) : p = p^* \text{ on } K^*\}$$

$$(1.2) \quad a = \inf_{p \in A} \max_{\xi \in K} G(p(\xi)).$$

Brezis-Nirenberg assume

(A) For each $p \in A$, $\max_{\xi \in K} G(p(\xi))$ is attained at a point in $K \setminus K^*$. They then prove that there is a sequence satisfying

$$(1.3) \quad G(u_k) \rightarrow a, \quad G'(u_k) \rightarrow 0.$$

In reference to the procedure one can ask three questions

1. Are the sets K, K^* essential to the method, or can they be eliminated?
2. How can one verify (A)?
3. What can be said if (A) fails to hold?

The purpose of the present paper is to address these questions. Concerning the first, we show that indeed a quantity corresponding to (1.2) can be introduced which is “intrinsic” in nature and does not depend on the “auxiliary” sets K, K^* . For this purpose we use a definition given in [ST]. We define

$$(1.4) \quad a = \inf_{\Gamma \in \Phi} \sup_{0 \leq s \leq 1, u \in A} G(\Gamma(s)u)$$

where A is a given subset of E and Φ is a family of maps in $C(E \times [0, 1], E)$ with certain properties (for a precise definition cf. Section 2). The quantity (1.4) depends only on G and the set A . It is not required that $A = p^*(K^*)$ for some K^* and $p^* \in C(K, E)$. In dealing with (1.4) we replace hypothesis (A) with

(B) For each $\Gamma \in \Phi$,

$$\max_{0 \leq s \leq 1, u \in A} G(\Gamma(s)u)$$

is attained at a point not in A .

We then show that hypothesis (B) implies the existence of a sequence satisfying (1.3) (actually, we use a hypothesis weaker than (B)). From this it follows that

$$(1.5) \quad a_0 := \sup_A G < a < \infty$$

implies the existence of a sequence satisfying (1.3).

Concerning the second question, we show that hypothesis (B) holds if and only if there is a subset B of E such that A links B in the sense of [ST] and

$$(1.6) \quad a_0 := \sup_A G \leq b_0 := \inf_B G.$$

The definition of linking given in [ST] differs from that usually found. Essentially, it says that A and B link if they cannot be “slipped” apart without intersection. In our opinion, this definition is more in keeping with the concept of linking (for a formal definition cf. Section 2). The results of the present paper show that the only practical way of applying hypothesis (B) is to find a subset B of E such that A links B and (1.6) holds (the same is true for hypothesis (A)).

Concerning the third question, we note that the only case not covered is when there is a set B such that A links B and

$$(1.7) \quad -\infty < b_0 < a_0 = a < \infty.$$

Surprisingly, something can be accomplished in this case as well. We have

Theorem 1.1. *Assume (1.7) and let*

$$B' = \{v \in B : G(v) < a_0\}.$$

Let α, T be positive numbers such that

$$(1.8) \quad a_0 - b_0 < \alpha T, \quad T < d' = d(A, B').$$

Then for each $\delta > 0$ sufficiently small there is a $u \in E$ satisfying

$$(1.9) \quad b_0 - \delta \leq G(u) \leq a_0 + \delta, \quad \|G'(u)\| \leq \alpha$$

and either

$$(1.10) \quad d(u, B') < T \text{ or } d(u, B \setminus B') < \delta/\alpha.$$

Corollary 1.2. *Let $\{B_n\}$ be a sequence of subsets of E such that A links B_n for each n and $d(A, B'_n) \rightarrow \infty$, where*

$$B'_n = \{v \in B_n : G(v) < a_0\}.$$

Assume that $a < \infty$ and that

$$(1.11) \quad \inf_{B_n} G \geq b_0 > -\infty.$$

Then there is a sequence $\{u_k\}$ satisfying (1.1) with $b_0 \leq c \leq a_0$.

In essence, Corollary 1.2 says that if A links B_n for each n and the part B'_n of B_n on which G is $< a_0$ moves out to infinity and (1.11) holds, then the mountain pass methods still apply. Special cases of this theorem were given in [Sc1-4, Si]. We present an application in Section 4.

2. A Generalized Mountain Pass.

Before stating our main theorems we recall the definition of linking sets given in [ST]. Let E be a Banach space and Let Φ be the set of all continuous maps $\Gamma(t)$ from $E \times [0, 1]$ to E such that

- (a) $\Gamma(0) = I$
- (b) there is an $x_0 \in E$ such that $\Gamma(1)x = x_0$ for each $x \in E$
- (c) $\Gamma(t)x \rightarrow x_0$ as $t \rightarrow 1$ uniformly on bounded subsets of E
- (d) for each $t \in [0, 1]$, $\Gamma(t)$ is a homeomorphism of E onto itself and Γ^{-1} is continuous on $E \times [0, 1]$.

Definition. A subset A of E links a subset B of E if $A \cap B = \phi$ and for each $\Gamma \in \Phi$ there is a $t \in (0, 1]$ such that $\Gamma(t)A \cap B \neq \phi$.

Let A, B be subsets of a Banach space E such that A links B , and let G be a C^1 functional on E . Define

$$(2.1) \quad a := \inf_{\Gamma \in \Phi} \sup_{0 \leq s \leq 1, u \in A} G(\Gamma(s)u), b_0 = \inf_B G.$$

Since A links B , we have

$$(2.2) \quad b_0 \leq a.$$

Assume that

$$(2.3) \quad d(A, B) > 0.$$

Let

$$(2.4) \quad B' := \{v \in B : G(v) < a\}.$$

We note that

$$(2.6) \quad B' = \phi \text{ iff } b_0 = a$$

(If D is any set, we write $d(D, \phi) = \infty$. Thus $d' = \infty$ when $B' = \phi$.) Let α, T be any positive numbers satisfying

$$(2.7) \quad a - b_0 < \alpha T, T < d'.$$

We have

Theorem 2.1. *Assume in addition that*

$$(2.8) \quad -\infty < b_0, a < \infty.$$

Then for every $\delta > 0$ sufficiently small there is a $u \in E$ such that

$$(2.9) \quad b_0 - \delta \leq G(u) \leq a + \delta, \quad \|G'(u)\| \leq \alpha$$

and either

$$(2.10) \quad d(u, B') < T$$

or

$$(2.11) \quad d(u, B \setminus B') < \delta/\alpha.$$

Corollary 2.2. *If $b_0 = a$, then there is a sequence $\{u_k\} \subset E$ such that*

$$(2.12) \quad G(u_k) \rightarrow a, G'(u_k) \rightarrow 0, d(u_k, B) \rightarrow 0.$$

Corollary 2.3. *Let $\{A_n\}, \{B_n\}$ be sequences of subsets of E , and define*

$$(2.13) \quad a_n = \inf_{\Gamma \in \Phi} \sup_{0 \leq s \leq 1, u \in A_n} G(\Gamma(s)u), b_{0n} = \inf_{B_n} G$$

$$(2.14) \quad a = \liminf a_n, b_0 = \limsup b_{0n}$$

$$(2.15) \quad B'_n = \{v \in B_n : G(v) < a_n\}$$

$$(2.16) \quad d'_n = d(A_n, B'_n).$$

Assume that A_n links B_n , $d(A_n, B_n) > 0$ and that

$$(2.17) \quad -\infty < b_0 \leq a < \infty$$

$$(2.18) \quad d'_n \rightarrow \infty.$$

Then there is a sequence $\{u_k\} \subset E$ such that

$$(2.19) \quad G(u_k) \rightarrow c, b_0 \leq c \leq a, G'(u_k) \rightarrow 0.$$

Similarly we have

Theorem 2.4. *Assume that $A, B \subset E$, B links A , and $G \in C^1(E, R)$. Define*

$$(2.20) \quad a_0 := \sup_A G, b := \sup_{\Gamma \in \Phi} \inf_{0 \leq s \leq 1, v \in B} G(\Gamma(s)v)$$

$$(2.21) \quad A' = \{u \in A : G(u) > b\}, d'' = d(A', B)$$

and let α, T be any positive constants satisfying

$$(2.22) \quad a_0 - b < \alpha T, T < d''.$$

Assume also that

$$(2.23) \quad -\infty < b, a_0 < \infty$$

and that (2.3) holds. Then for every $\delta > 0$ sufficiently small there is a $u \in E$ satisfying

$$(2.24) \quad b - \delta \leq G(u) \leq a_0 + \delta, \|G'(u)\| \leq \alpha$$

and either

$$(2.25) \quad d(u, A') < T$$

or

$$(2.26) \quad d(u, A \setminus A') < \delta/\alpha.$$

Corollary 2.5. Let $\{A_n\}, \{B_n\}$ be sequences of subsets of E , and define

$$(2.27) \quad a_{on} = \sup_{A_n} G, b_n = \sup_{\Gamma \in \Phi} \inf_{0 \leq s \leq 1, v \in B_n} G(\Gamma(s)v)$$

$$(2.28) \quad a_0 = \liminf a_{on}, b = \limsup b_n.$$

$$(2.29) \quad A'_n = \{u \in A_n : G(u) > b_n\}, d''_n = d(A'_n, B_n).$$

Assume that B_n links A_n , $d(A_n, B_n) > 0$ and that

$$(2.30) \quad -\infty < b \leq a_0 < \infty, d''_n \rightarrow \infty.$$

Then there is a sequence $\{u_k\} \subset E$ such that

$$(2.31) \quad G(u_k) \rightarrow c, b \leq c \leq a_0, G'(u_k) \rightarrow 0.$$

Theorem 2.6. If $a_0 \leq b_0$ and $a < \infty$, then there is a sequence $\{u_k\} \subset E$ such that

$$(2.32) \quad G(u_k) \rightarrow a, G'(u_k) \rightarrow 0.$$

We now show that we can remove the reference to the set B . We have

Theorem 2.7. Assume that $a < \infty$ and that for each $\Gamma \in \Phi$ the set

$$(2.33) \quad g_\Gamma := \{v = \Gamma(s)u : s \in (0, 1], u \in A, v \notin A, G(v) \geq a_0\}$$

is not empty. Then there is a sequence satisfying (2.32).

Corollary 2.8. *If $a < \infty$ and $a_0 \neq a$, then there is a sequence satisfying (2.32).*

Corollary 2.9. *If $a < \infty$ and for each $\Gamma \in \Phi$*

$$(2.34) \quad \max_{0 \leq s \leq 1, u \in A} G(\Gamma(s)u)$$

is attained at a point not in A , then there is a sequence satisfying (2.32).

We can summarize by

Theorem 2.10. *Assume that A links B and that they satisfy (2.8). Then one of the following holds*

- (i) $a_0 \neq a$
- (ii) $b_0 = a_0 = a$
- (iii) $b_0 < a_0 = a$.

In case (i) there is a sequence satisfying (2.32). In case (ii) there is a sequence satisfying (2.12). In case (iii) the conclusions of Theorem (2.1) hold.

Another consequence of Theorem 2.1 is

Theorem 2.11. *Let M, N be complementary subspaces of a Banach space E with one of them being finite dimensional. Let G be a C^1 functional on E , and define*

$$(2.35) \quad m_0 := \sup_{v \in N} \inf_{w \in M} G(v + w), m_1 := \inf_{w \in M} \sup_{v \in N} G(v + w).$$

Assume that

$$(2.36) \quad -\infty < m_0 \text{ and } m_1 < \infty.$$

Then there is a sequence $\{u_k\} \subset E$ such that

$$(2.37) \quad G(u_k) \rightarrow c, m_0 \leq c \leq m_1, G'(u_k) \rightarrow 0.$$

Corollary 2.2 was proved in [ST] as well as Theorem 2.6. Theorem 2.11 generalizes theorems of [Sc1, Si]. The results of this section will be proved in Section 3.

Finally, we note the following

Theorem 2.12. *There is a $B \subset E$ such that A links B and $a_0 \leq b_0$ if and only if g_Γ defined by (2.33) is not empty for each $\Gamma \in \Phi$.*

Remarks. There are basic differences between the approach of [BN] and that of [ST] and the present paper. The sets K, K^* must be compact; otherwise their arguments fail. In [ST] the set A need not be compact (or even closed). Also, both K and K^* must be chosen initially. In [ST] only the set A is chosen. The sets (2.33) vary with Γ and are determined automatically. For our linking, the set B is only required to intersect the set (2.33) for each Γ . It is not required to intersect $p(K)$ for each continuous map p (a much larger set). There are no counterparts of Theorem 2.1, Corollary 2.3, Theorem 2.4, Corollary 2.5 in the [BN] theory.

Assumption (2.3) is not required in Theorems 2.6, 2.7 and Corollary 2.9.

3. The Flow.

In this section we give the proofs of the results of Section 2.

Proof of Theorem 2.1. Assume that the theorem is false. Then there are α, T satisfying (2.7) and a $\delta > 0$ such that

$$(3.1) \quad \|G'(u)\| > \alpha$$

when

$$u \in Q := \{u \in E : b_0 - 3\delta \leq G(u) \leq a + 3\delta \text{ and either } d(u, B') \leq T + 3\delta \text{ or } d(u, B \setminus B') \leq 5\delta/\alpha\}.$$

Let

$$\begin{aligned} Q_0 &= \{u \in E : b_0 - 2\delta \leq G(u) \leq a + 2\delta \\ &\quad \text{and either } d(u, B') \leq T + 2\delta \text{ or } d(u, B \setminus B') \leq 4\delta/\alpha\} \\ Q_1 &= \{u \in E : b_0 - \delta \leq G(u) \leq a + \delta \text{ and either } d(u, B') \\ &\quad \leq T + \delta \text{ or } d(u, B \setminus B') \leq 3\delta/\alpha\}. \end{aligned}$$

Define

$$Q_2 = E \setminus Q_0, \eta(u) = d(u, Q_2)/[d(u, Q_1) + d(u, Q_2)].$$

Note that $\eta = 1$ on $Q_1, \eta = 0$ on \bar{Q}_2 and $0 < \eta < 1$ otherwise. Let $d_1 = d(B \setminus B', A)$ and reduce δ , if necessary, so that

$$(3.3) \quad 2\delta < \alpha d_1, \delta < \alpha T - (a_0 - b).$$

Let θ satisfy

$$(3.4) \quad \frac{2}{3} < \theta < 1, 2\delta < \theta\alpha d_1, \delta < \theta\alpha T - (a_0 - b).$$

One can show that there is a locally Lipschitz continuous map $Y(u) : \hat{E} \rightarrow \hat{E} = \{u \in E : G'(u) \neq 0\}$ satisfying

$$(3.5) \quad \|Y(u)\| \leq 1, (G'(u), Y(u)) \geq \theta \|G'(u)\|, u \in \hat{E}$$

(cf, e.g., [BN]). Let $\sigma(t)$ be the flow generated by the vector field $\eta(u)Y(u)$ which is I for $t = 0$. Then

$$(3.6) \quad \begin{aligned} \|\sigma(t)u - u\| &= \left\| \int_0^t \eta(\sigma(\tau)u)Y(\sigma(\tau)u)d\tau \right\| \\ &\leq \Psi(t, u) := \int_0^t \eta(\sigma(\tau)u)d\tau, u \in E, \end{aligned}$$

$$(3.7) \quad G(\sigma(t)u) - G(u) = \int_0^t \eta(\sigma(\tau)u)(G'(\sigma(\tau)u), Y(\sigma(\tau)u))d\tau \geq \theta\alpha\psi(t, u), u \in E.$$

Thus

$$(3.8) \quad \|u - v\| \leq \|u - \sigma(t)v\| + \Psi(t, v).$$

If $u \in A$ and $v \in B'$, this implies

$$(3.9) \quad \|u - \sigma(t)v\| \geq d' - T, 0 \leq t \leq T, u \in A, v \in B'.$$

On the other hand, if $v \in B \setminus B'$, then $G(v) \geq a$ and $G(\sigma(t)v) \leq \max[G(v), a + 2\delta]$ by the definition of η . Hence by (3.7)

$$(3.10) \quad \theta\alpha\Phi(t, v) \leq 2\delta, t \geq 0, v \in B \setminus B'.$$

Consequently, by (3.8) - (3.10) we have

$$(3.11) \quad \|u - \sigma(t)v\| \geq \min[d' - T, d_1 - (2\delta/\theta\alpha)] > 0, 0 \leq t \leq T, u \in A, v \in B.$$

Let $B_1 = \sigma(T)B$. I claim that A links B_1 . By (3.11), $A \cap \sigma(t)B = \emptyset$ for $0 \leq t \leq T$. Let $\Gamma \in \Phi$, and put $\Gamma_1(s) = \sigma(2sT)^{-1}$ for $0 \leq s \leq \frac{1}{2}$, $\Gamma_1(s) = \sigma(T)^{-1}\Gamma(2s - 1)$ for $\frac{1}{2} < s \leq 1$. Then $\Gamma_1 \in \Phi$. Since A links B , there is an $s_1 \in [0, 1]$ such that $\Gamma_1(s_1)A \cap B \neq \emptyset$. But $\sigma(2s_1T)^{-1}A \cap B = \emptyset$ for $0 \leq s \leq \frac{1}{2}$ by (3.11). Thus $\frac{1}{2} < s_1 \leq 1$, and $\sigma(T)^{-1}\Gamma(2s_1 - 1)A \cap B \neq \emptyset$, or equivalently,

$\Gamma(2s_1 - 1)A \cap B_1 \neq \phi$. Thus A links B_1 . Suppose there is a $t_1 \leq T$ such that $\sigma(t_1)v \notin Q_1$. Then either

$$(3.12) \quad G(\sigma(t_1)v) > a + \delta$$

or

$$(3.13) \quad d(\sigma(t_1)v, B') > T + \delta \text{ and } d(\sigma(t_1)v, B \setminus B') > 3\delta/\alpha.$$

But if $v \in B'$, then $d(\sigma(t_1)v, B') \leq t_1 \leq T$ by (3.6), and if $v \in B \setminus B'$, then $d(\sigma(t_1)v, B \setminus B') \leq \Phi(t_1, v) \leq 2\delta/\theta\alpha < 3\delta/\alpha$ by (3.6) and (3.10). Thus (3.13) is false if $v \in B$. Hence (3.12) must hold. This implies

$$(3.14) \quad G(\sigma(T)v) > a + \delta.$$

On the other hand, if $\sigma(t)v \in Q_1$ for $0 \leq t \leq T$, then (3.7) gives

$$G(\sigma(T)v) \geq b_0 + \theta\alpha T > a + \delta.$$

Hence (3.14) holds for all $v \in B$. Thus means that

$$(3.15) \quad \inf_{B_1} G \geq a + \delta.$$

But by the definition of a , there is a $\Gamma \in \Phi$ such that

$$(3.16) \quad \sup_{0 \leq s \leq 1, u \in A} G(\Gamma(s)u) < a + (\delta/2)$$

and since A links B_1 , there is an $s \in [0, 1]$ such that $\Gamma(s)A \cap B_1 \neq \phi$, and consequently (3.16) contradicts (3.15). Thus (3.1) cannot hold for u satisfying (3.2), and the theorem is proved. \square

Proof of Corollary 2.2. In this case $B' = \phi$ and $d' = \infty$. For each n we take $T_n = 1, \delta_n = 1/n^2, \alpha_n = 1/n$ and apply Theorem 2.1. \square

Proof of Corollary 2.3. For each n we take $\delta_n = 1/n, T_n = d'_n/2, \alpha_n = [(a_n - b_{on})/T_n] + \delta_n$. Then $\alpha_n \rightarrow 0$ and there is a u_n such that

$$b_{on} - \delta_n \leq G(u_n) \leq a_n + \delta_n, \|G'(u_n)\| \leq \alpha_n.$$

We take a renamed subsequence such that $a_n \rightarrow a, b_{on} \rightarrow b_0$ and $G(u_n) \rightarrow c$. \square

Proof of Theorem 2.4. We interchange A and B and replace G by $-G$ in Theorem 2.1. Then b_0 becomes $-a_0$ and a becomes $-b$. B' becomes A' . We then apply Theorem 2.1. \square

The proof of Corollary 2.5 is similar to that of Corollary 2.3.

Proof of Theorem 2.7. Let

$$(3.17) \quad B = \bigcup_{\Gamma \in \Phi} g_{\Gamma}.$$

Then $A \cap B = \phi$ and for each $\Gamma \in \Phi$ there is a $v \in B$, an $s \in (0, 1]$ and a $u \in A$ such that $\Gamma(s)u = v$. Hence

$$\Gamma(s)A \cap B \neq \phi.$$

Hence A links B . Since $G(v) \geq a_0$ for each $v \in B$, we have $a_0 \leq b_0$. We can now apply Theorem 2.6 to conclude that a sequence satisfying (2.32) exists. \square

Proof of Corollary 2.8. If $a_0 < a$, then for each $\Gamma \in \Phi$ there is a $u \in A$ and an $s \in [0, 1]$ such that $G(\Gamma(s)u) > a_0$. Clearly $v = \Gamma(s)u \notin A$. Thus $g_{\Gamma} \neq \phi$. We can now apply Theorem 2.7. \square

Proof of Corollary 2.9. If the maximum (2.34) is attained at a point outside A , then this point is in g_{Γ} . Hence g_{Γ} is not empty for all $\Gamma \in \Phi$. Apply Theorem 2.7. \square

Proof of Theorem 2.11. Assume, for definiteness, that $\dim N < \infty$. Let $\{\epsilon_k\}$ be a sequence tending to 0. For each k there are $v_k \in N, w_k \in M$ such that

$$(3.18) \quad \inf_{M_k} G > m_0 - \epsilon_k, \sup_{N_k} G < m_1 + \epsilon_k$$

where $M_k = v_k \oplus M$ and $N_k = w_k \oplus N$. Let

$$A_k = \{v + w_k : v \in N, \|v - v_k\| = k\}, B_k = M_k.$$

Note that $A_k \subset N_k$ and $A_k \cap B_k = \phi$. It is readily shown that A_k links B_k for each k (cf., Proposition 1.2 of [ST]). By (3.18) we have

$$b_{ok} \geq m_0 - \epsilon_k, a_{ok} \leq m_1 + \epsilon_k.$$

Moreover, for each k ,

$$\Gamma_k(s)u = s(v_k + w_k) + (1 - s)u$$

is in Φ . Consequently, definition (2.13) gives

$$a_k \leq \sup_{N_k} G < m_1 + \epsilon_k.$$

Moreover, by (2.6)

$$d'_k = d(A_k, B'_k) \geq d(A_k, B_k) = k \rightarrow \infty.$$

If a and b_0 are given by (2.14), we see that $m_0 \leq b_0$ and $a \leq m_1$. We can now apply Corollary 2.3 to conclude that a sequence satisfying (2.37) exists. The proof for the case $\dim M < \infty$ is similar. \square

Proof of Theorem 2.12. The “if” part was already proved in the proof of Theorem 2.7. Conversely, suppose A links B and $a_0 \leq b_0$. If $g_\Gamma = \phi$ for some $\Gamma \in \Phi$, then $\Gamma(s)A \cap B = \phi$ for each $s \in [0, 1]$. This says that A does not link B , a contradiction. \square

Remark. The proof of Theorem 2.6 was not given because it was proved in [ST]. However, it is a corollary of Theorem 2.1. To see this, note that if $a_0 \leq b_0 = a$, then $B' = \phi, d' = \infty$, and we can take $\alpha = \delta, T = 1$. If $a_0 \leq b_0 < a$, let B_1 be given by (3.17). Then A links B_1 and $a_0 \leq b_1 \equiv \inf_{B_1} G = a$. We can now apply Theorem 2.1 again with B replaced by B_1 .

4. An Application.

In this section we show how the theorems of Section 2 can be applied. Let Ω be a smooth, bounded domain in R^n , and let A be a selfadjoint operator on $L^2(\Omega)$ with discrete spectrum $0 < \lambda_0 < \lambda_1 < \dots < \lambda_j < \dots$. We assume that $C_0^\infty(\Omega) \subset D := D(A^{1/2}) \subset H^m(\Omega)$ for some $m > 0$ (m need not be an integer). Let q be a number satisfying

$$(4.1) \quad 2 \leq q < 2m/(n - 2m), 2 \leq q < \infty, 2m < n;$$

and let $f(x, t)$ be a Caratheodory function on $\Omega \times R$. Assume

$$(I) \quad |f(x, t)| \leq V_0(x)^q |t|^{q-1} + V_0(x)V_1(x)$$

where $V_0 \in L^q(\Omega), V_1 \in L^{q'}(\Omega)$ and multiplication by V_0 is a compact operator from D to $L^q(\Omega)$.

(II) For some $\ell > 0$ the function

$$F(x, t) := \int_0^t f(x, s)ds$$

satisfies

$$(4.2) \quad \lambda_{\ell-1}t^2 - W_0(x) \leq 2F(x, t) \leq \nu t^2 + V(x)p|t|^p + W_1(x)$$

and

$$(4.3) \quad \lambda_\ell t^2 - W_2(x) \leq 2 F(x, t)$$

where $\nu < \lambda_\ell, p > 2$,

$$(4.4) \quad B_j := \int_{\Omega} W_j(x) dx < \infty, j = 0, 1, 2$$

and

$$(4.5) \quad \|Vu\|_p^p \leq C \|A^{1/2}u\|^p, u \in D.$$

(III) For some $\mu > 2$ the function

$$H_\mu(x, t) := \mu F(x, t) - tf(x, t)$$

satisfies

$$(4.6) \quad H_\mu(x, t) \leq V_3(x)^2 t^2 \sigma(t) + W_3(x)$$

where multiplication by V_3 is bounded from D to $L^2(\Omega)$, $W_3 \in L^1(\Omega)$ and $\sigma(t)$ is a continuous function such that $\sigma(t) \rightarrow 0$ as $|t| \rightarrow \infty$.

$$(IV) \quad B_0 + B_1 < (1 - \frac{2}{p})(1 - \frac{\nu}{\lambda_\ell})^{p/p-2} (\frac{2}{pC})^{2/p-2}.$$

We have

Theorem 4.1. *Under hypotheses (I) - (IV) there is a solution of*

$$(4.7) \quad Au = f(x, u), \quad u \in D.$$

Proof. Under hypothesis (I) the functional

$$(4.8) \quad G(u) = a(u) - 2 \int_{\Omega} F(x, u) dx$$

has a continuous Frechet derivative on D given by

$$(4.9) \quad (G'(u), v) = 2a(u, v) - 2(f(\cdot, u), v)$$

where

$$(4.10) \quad a(u, v) = (Au, v), a(u) = a(u, u), u, v \in D.$$

Let N be the subspace spanned by the eigenfunctions corresponding to the eigenvalues $\lambda_0, \dots, \lambda_{\ell-1}$, and let $M = N^\perp \cap D$, the orthogonal complement of N on D . On M we have

$$G(w) \geq a(w) - \nu \|w\|^2 - \|Vw\|_p^p - B_1 \geq \left(1 - \frac{\nu}{\lambda_\ell}\right) \|w\|_D^2 - C \|w\|_D^p - B_1$$

by (4.2) and (4.5). If we take $\|w\|_D = \delta$ with $\delta^{p-2} = 2 \left(1 - \frac{\nu}{\lambda_\ell}\right) / p$, we have

$$(4.11) \quad G(w) \geq \left(1 - \frac{2}{p}\right) \left(1 - \frac{\nu}{\lambda_\ell}\right) \delta^2 - B_1 \geq \beta - B_1, \|w\|_D = \delta$$

where β is the expression on the right hand side in hypothesis (IV). On the other hand, we have by (4.2)

$$(4.12) \quad G(v) \leq a(v) - \lambda_{\ell-1} \|v\|^2 + B_0 \leq B_0, v \in N.$$

Let w_0 be an eigenfunction corresponding to the eigenvalue λ_ℓ with unit norm, and let N_1 denote the subspace spanned by N and w_0 . By (4.3) we have

$$(4.13) \quad G(v) \leq a(v) - \lambda_\ell \|v\|^2 + B_2 \leq B_2, v \in N_1.$$

Let

$$\begin{aligned} B &= \{w \in M : \|w\|_D = \delta\}, B_n = B \\ A_{1n} &= \{v \in N : \|v\|_D \leq n\} \\ A_{2n} &= \{u = v + sw_0 : v \in N, s \geq 0, \|u\|_D = n\}. \end{aligned}$$

One can show that $A_n = A_{1n} \cup A_{2n}$ links $B_n = B$ for each n (cf., [ST, Proposition 1.2]). By (4.11)

$$(4.14) \quad b_{on} = b_0 \geq \beta - B_1, b \geq -B_1.$$

If $b_0 \neq b$, then we can find a sequence in D satisfying

$$(4.15) \quad G(u_k) \rightarrow b, G'(u_k) \rightarrow 0$$

by Corollary 2.8. If $b_0 = b$, then we have

$$A'_n = \{u \in A_n : G(u) > b\} \subset A_{2n}$$

since

$$G(v) \leq B_0 < \beta - B_1, v \in A_{1n}$$

by (4.12) and hypothesis (IV). Hence

$$d''_n = d(A'_n, B) \geq d(A_{2n}, B) \geq n - \delta \rightarrow \infty.$$

By (4.13), $a_{on} \leq B_2$ for each n . Hence we may apply Corollary 2.5 to conclude that there is a sequence in D such that

$$(4.16) \quad G(u_k) \rightarrow c, \beta - B_1 \leq c \leq B_2, G'(u_k) \rightarrow 0.$$

In either case we claim that the sequence is bounded in D . For suppose $\rho_k = \|u_k\|_D \rightarrow \infty$. Let $\tilde{u}_k = u_k/\rho_k$. Then by (4.15) or (4.16) we have

$$\begin{aligned} \|u_k\|_D^2 - 2 \int_{\Omega} F(x, u_k) dx &= o(1) \\ \|u_k\|_D^2 - \int_{\Omega} f(x, u_k) u_k dx &= o(\rho_k). \end{aligned}$$

Thus

$$(4.17) \quad (\mu - 2)\rho_k^2 - 2 \int_{\Omega} H(x, u_k) dx = o(\rho_k).$$

Now

$$(4.18) \quad \int \rho_k^{-2} H(x, u_k) dx \leq \int_{\Omega} V_3(x)^2 \tilde{u}_k^2 \sigma(\rho_k \tilde{u}_k) dx + \rho_k^{-2} \int_{\Omega} W_3(x) dx.$$

There is a renamed subsequence for which $\tilde{u}_k(x)$ converges a.e. By hypothesis $V_3 \tilde{u}_k$ is bounded in $L^2(\Omega)$ as well as $V_3 \tilde{u}_k \sigma(u_k)$. Moreover, $V_3 \tilde{u}_k \sigma(u_k) \rightarrow 0$ a.e. Hence the right hand side of (4.18) converges to 0. But this together with (4.17) implies that $\mu \leq 2$, contrary to assumption. Hence the sequence $\{u_k\}$ is bounded. It now follows by standard arguments that $\{u_k\}$ has a subsequence which converges in D and that the limit satisfies $G'(u) = 0$ (cf., e.g., [Ra2]). It now follows from (4.9) that the limit is a solution of (4.7). \square

References

- [BBF] P. Bartolo, V. Benci and D. Fortunato, *Abstract critical point theorems and applications to some nonlinear problems with "strong" resonance at infinity*, Nonlinear Anal. TMA, **7** (1983), 981-1012.
- [BN] H. Brezis and L. Nirenberg, *Remarks on finding critical points*, Comm. Pure Appl. Math., **44** (1991), 939-964.
- [E] I. Ekeland, *On the variational principle*, J. Math. Anal. Appl., **47** (1974), 324-353.
- [Ra2] P.H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, Cont. Board of Math. Sci. Reg. Cont. Ser. in Math., No. 65, Amer. Math. Soc., 1986.
- [Sc1] M. Schechter, *New saddle point theorems*, Proceedings of an International Symposium on Generalized Functions and their Applications, Varanasi, India, December 23-26, 1991.
- [Sc2] M. Schechter, *A generalization of the saddle point method with applications*, Ann. Polonici Math., **57** (1992), 269-281.
- [Sc3] M. Schechter, *Critical points over splitting subspaces*, Nonlinearity, **6** (1993), 417-427.

- [Sc4] M. Schechter, *Splitting subspaces and critical points*, *Applicable Analysis*, **49** (1993), 33-48.
- [Si] E.A. de B.e. Silva, *Linking theorems and applications to semilinear elliptic problems at resonance*, *Nonlinear Analysis TMA*, **16** (1991), 455-477.
- [ST] M. Schechter and K. Tintarev, *Pairs of critical points produced by linking subsets with applications to semilinear elliptic problems*, *Bull. Soc. Math. Belg.*, **44** (1992), 249-261.

Received April 5, 1993 and revised January 19, 1994. Research supported in part by an NSF grant.

UNIVERSITY OF CALIFORNIA
IRVINE, CA 92717

E-mail: mschecht@math.uci.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

EDITORS

Sun-Yung Alice Chang (Managing Editor)
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

F. Michael Christ
University of California
Los Angeles, CA 90095-1555
christ@math.ucla.edu

Robert Finn
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

Martin Scharlemann
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

Thomas Enright
University of California
San Diego, La Jolla, CA 92093
tenright@ucsd.edu

Vaughan F. R. Jones
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

Gang Tian
Courant Institute
New York University
New York, NY 10012-1100
tiang@taotao.cims.nyu.edu

Nicholas Ercolani
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

Steven Kerckhoff
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

V. S. Varadarajan
University of California
Los Angeles, CA 90095-1555
vsv@math.ucla.edu

SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
STANFORD UNIVERSITY
UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
UNIVERSITY OF UTAH
UNIVERSITY OF WASHINGTON
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$108.00 a year to individual members of supporting institutions.

Subscriptions, orders for back issues published within the last three years, and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,
the American Mathematical Society's TEX macro system.
Copyright © 1995 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 171 No. 2 December 1995

On H^p -solutions of the Bezout equation	297
ERIC AMAR, JOAQUIM BRUNA FLORIS and ARTUR NICOLAU	
Amenable correspondences and approximation properties for von Neumann algebras	309
CLAIRE ANANTHARAMAN-DELAROCHE	
On moduli of instanton bundles on \mathbb{P}^{2n+1}	343
VINCENZO ANCONA and GIORGIO MARIA OTTAVIANI	
Minimal surfaces with catenoid ends	353
JORGEN BERGLUND and WAYNE ROSSMAN	
Permutation model for semi-circular systems and quantum random walks	373
PHILIPPE BIANE	
The Neumann problem on Lipschitz domains in Hardy spaces of order less than one	389
RUSSELL M. BROWN	
Matching theorems for twisted orbital integrals	409
REBECCA A. HERB	
Uniform algebras generated by holomorphic and pluriharmonic functions on strictly pseudoconvex domains	429
ALEXANDER IZZO	
Quantum Weyl algebras and deformations of $U(g)$	437
NAIHUAN JING and JAMES ZHANG	
Calcul du nombre de classes des corps de nombres	455
STÉPHANE LOUBOUTIN	
On geometric properties of harmonic Lip_1 -capacity	469
PERTTI MATTILA and P. V. PARAMONOV	
Reproducing kernels and composition series for spaces of vector-valued holomorphic functions	493
BENT ØRSTED and GENKAI ZHANG	
Iterated loop modules and a filtration for vertex representation of toroidal Lie algebras	511
S. ESWARA RAO	
The intrinsic mountain pass	529
MARTIN SCHECHTER	
A Frobenius problem on the knot space	545
RON G. WANG	
On complete metrics of nonnegative curvature on 2-plane bundles	569
DAVID YANG	
Correction to: "Free Banach-Lie algebras, couniversal Banach-Lie groups, and more"	585
VLADIMIR G. PESTOV	
Correction to: "Asymptotic radial symmetry for solutions of $\Delta u + e^u = 0$ in a punctured disc"	589
KAI SENG (KAISING) CHOU (TSO) and TOM YAU-HENG WAN	



0030-8730(1995)171:2;1-G