

*Pacific
Journal of
Mathematics*

**SINGULAR MODULI SPACES OF STABLE VECTOR BUNDLES
ON P^3**

ROSA M. MIRÓ-ROIG

Volume 172 No. 2

February 1996

SINGULAR MODULI SPACES OF STABLE VECTOR BUNDLES ON \mathbf{P}^3

ROSA M. MIRÓ-ROIG

The goal of this paper is to give an example of singular moduli space of rank 3 stable vector bundles on \mathbf{P}^3 .

Introduction.

In 1977/78, M. Maruyama proved the existence of a moduli scheme $M_{\mathbf{P}^n}(r; c_1, \dots, c_{\min(r,n)})$ parametrizing isomorphic classes of rank r stable vector bundles on \mathbf{P}^n with given Chern classes $c_1, \dots, c_{\min(r,n)}$ (cf. [M1, M2]). The goal of this note is to give, to the best of my knowledge, the first example of singular moduli space of stable vector bundles on \mathbf{P}^3 . It has been motivated by a recent work of Ancona and Ottaviani where they show that the moduli space $MI_{\mathbf{P}^5}(k)$ of stable instanton bundles on \mathbf{P}^5 with quantum number $k=3$ or 4 is singular. Moreover they claim that $MI_{\mathbf{P}^5}(3)$ and $MI_{\mathbf{P}^5}(4)$ are the first examples of singular moduli spaces of stable vector bundles on projective spaces (cf. [AO]). Ancona-Ottaviani's result together with the well known fact that $M_{\mathbf{P}^2}(r; c_1, c_2)$ is a smooth quasi-projective variety of dimension $2rc_2 - (r-1)c_1^2 + 1 - r^2$ gives rise the following question:

Is there any example of singular moduli space of stable vector bundles on \mathbf{P}^3 ?

As I pointed out before my aim is to give an affirmative answer to this question (cf. Theorem 2.10).

1. Preliminaries.

In this section we recall some well known results needed later on.

1.1. Let $H(18, 39)$ be the open subscheme of $Hilb_{\mathbf{P}^3}$ parametrizing smooth connected curves $C \subset \mathbf{P}^3$ of degree 18 and genus 39. (See [EF] for a precise description of $H(18, 39)$.) Let $H_1 \subset H(18, 39)$ be the 72-dimensional irreducible, generically smooth component whose general point parametrizes an arithmetically Cohen-Macaulay curve $X \subset \mathbf{P}^3$ having a locally free resolution of the following type:

$$(1) \quad 0 \rightarrow \mathcal{O}(-7)^4 \rightarrow \mathcal{O}(-6)^4 \oplus \mathcal{O}(-4) \rightarrow I_X \rightarrow 0.$$

Let $H_2 \subset H(18, 39)$ be the 72-dimensional irreducible, generically smooth component whose general point parametrizes an arithmetically Cohen-Macaulay curve $Y \subset \mathbf{P}^3$ having a locally free resolution of the following type:

$$(2) \quad 0 \rightarrow \mathcal{O}(-6)^2 \oplus \mathcal{O}(-8) \rightarrow \mathcal{O}(-5)^4 \rightarrow I_Y \rightarrow 0.$$

It is well known that there exists an irreducible subset $H = H_1 \cap H_2 \subset H(18, 39)$ of dimension 71 whose general point parametrizes an arithmetically Buchsbaum curve $C \subset \mathbf{P}^3$ having a locally free resolution of the following type:

$$(3) \quad 0 \rightarrow \mathcal{O}(-8) \rightarrow \mathcal{O}(-7)^4 \oplus \mathcal{O}(-8) \rightarrow \mathcal{O}(-6)^4 \oplus \mathcal{O}(-4) \rightarrow I_C \rightarrow 0.$$

1.2. Remark. For all curve $Z \in H_1 \cup H_2$, $\omega_Z(2)$ is globally generated. From now on, for all curve $Z \in H_1 \cup H_2$, we set $\alpha := \dim H^0(\omega_Z(2))$ ($=74$; by Riemann-Roch's Theorem).

1.3. Fact. Let $0 \rightarrow E \rightarrow F \rightarrow G \rightarrow 0$ be an exact sequence of vector bundles. Then, we have the following exact sequence involving alternating and symmetric powers:

$$0 \rightarrow S^q E \rightarrow S^{q-1} E \otimes F \rightarrow \dots \rightarrow E \otimes \Lambda^{q-1} F \rightarrow \Lambda^q F \rightarrow \Lambda^q G \rightarrow 0.$$

1.4. Hoppe's criterion for the stability of a vector bundle. Let X be a projective manifold with $\text{Pic}(X) \cong \mathbf{Z}$ and let E be a vector bundle on X . If $H^0(X, (\Lambda^q E)_{\text{norm}}) = 0$ for $1 \leq q \leq \text{rk}(E) - 1$, then E is stable. As usual, given a vector bundle E on X , we denote by E_{norm} the twist of E whose first Chern class c_1 verifies $-\text{rk}(E) + 1 \leq c_1 \leq 0$.

2. Main results.

2.1. Let us call \mathcal{L}_1 the irreducible family of sheaves E on \mathbf{P}^3 constructed as an extension:

$$\sigma = (\sigma_1, \dots, \sigma_\alpha) : \quad 0 \rightarrow \mathcal{O}^\alpha \rightarrow E(1) \rightarrow I_X(2) \rightarrow 0$$

where $X \in H_1$ and $\sigma_1, \dots, \sigma_\alpha \in H^0(\omega_X(2)) \cong \text{Ext}^1(I_X(2), \mathcal{O})$ are general global sections which generate the sheaf $\omega_X(2)$ everywhere.

It is easy to see that E is a vector bundle on \mathbf{P}^3 of rank $\alpha + 1$.

2.2. Let us call \mathcal{L}_2 the irreducible family of sheaves F on \mathbf{P}^3 constructed as an extension:

$$\lambda = (\lambda_1, \dots, \lambda_\alpha) : \quad 0 \rightarrow \mathcal{O}^\alpha \rightarrow F(1) \rightarrow I_Y(2) \rightarrow 0$$

where $Y \in H_2$ and $\lambda_1, \dots, \lambda_\alpha \in H^0(\omega_Y(2)) \cong \text{Ext}^1(I_Y(2), \mathcal{O})$ are general global sections which generate the sheaf $\omega_Y(2)$ everywhere.

Again it is easy to see that F is a vector bundle on \mathbf{P}^3 of rank $\alpha + 1$.

2.3. And let $\mathcal{L} \subset \mathcal{L}_1 \cap \mathcal{L}_2$ be the irreducible family of sheaves G on \mathbf{P}^3 constructed as an extension:

$$\mu = (\mu_1, \dots, \mu_\alpha) : \quad 0 \rightarrow \mathcal{O}^\alpha \rightarrow G(1) \rightarrow I_C(2) \rightarrow 0$$

where $C \in H \subset H_1 \cap H_2$ and $\mu_1, \dots, \mu_\alpha \in H^0(\omega_C(2)) \cong Ext^1(I_C(2), \mathcal{O})$ are general global sections which generate the sheaf $\omega_Z(2)$ everywhere.

Again it is easy to see that G is a vector bundle on \mathbf{P}^3 of rank $\alpha + 1$.

Proposition 2.4.

- (1) *A general vector bundle $E \in \mathcal{L}_1$ has a locally free resolution of the following type:*

$$0 \rightarrow \mathcal{O}(-5)^4 \rightarrow \mathcal{O}(-4)^4 \oplus \mathcal{O}(-2) \oplus \mathcal{O}^\alpha \rightarrow E(1) \rightarrow 0.$$

- (2) *A general vector bundle $F \in \mathcal{L}_2$ has a locally free resolution of the following type:*

$$0 \rightarrow \mathcal{O}(-6) \oplus \mathcal{O}(-4)^2 \rightarrow \mathcal{O}(-3)^4 \oplus \mathcal{O}^\alpha \rightarrow F(1) \rightarrow 0.$$

- (3) *A general vector bundle $G \in \mathcal{L}$ has a locally free resolution of the following type:*

$$0 \rightarrow \mathcal{O}(-6) \rightarrow \mathcal{O}(-6) \oplus \mathcal{O}(-5)^4 \rightarrow \mathcal{O}(-4)^4 \oplus \mathcal{O}(-2) \oplus \mathcal{O}^\alpha \rightarrow G(1) \rightarrow 0.$$

Proof. (1) From the exact sequence:

$$0 \rightarrow \mathcal{O}^\alpha \rightarrow E(1) \rightarrow I_X(2) \rightarrow 0$$

and the locally free resolution of $I_X(2)$ (See 1.1):

$$0 \rightarrow \mathcal{O}(-5)^4 \rightarrow \mathcal{O}(-4)^4 \oplus \mathcal{O}(-2) \rightarrow I_X(2) \rightarrow 0$$

we get the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{O}(-5)^4 & \xlongequal{\quad} & \mathcal{O}(-5)^4 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \mathcal{O}^\alpha & \longrightarrow & \mathcal{O}^\alpha \oplus \mathcal{O}(-2) \oplus \mathcal{O}(-4)^4 & \longrightarrow & \mathcal{O}(-2) \oplus \mathcal{O}(-4)^4 \longrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathcal{O}^\alpha & \longrightarrow & E(1) & \longrightarrow & I_X(2) \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

which gives what we want.

(2) and (3) Analogous. □

Corollary 2.5. *Given a vector bundle $E \in \mathcal{L}_1 \cup \mathcal{L}_2$, $E(t)$ is globally generated for all $t \geq 5$.*

2.6. Let \mathcal{F}_1 be the irreducible family of rank 3 vector bundles P on \mathbf{P}^3 defined as the cokernel:

$$0 \rightarrow \mathcal{O}(-5)^{\alpha-2} \xrightarrow{s_1, \dots, s_2} E \rightarrow P \rightarrow 0$$

where $E \in \mathcal{L}_1$ and $s_i \in H^0(E(5))$ are general global sections of $E(5)$.

2.7. Let \mathcal{F}_2 be the irreducible family of rank 3 vector bundles Q on \mathbf{P}^3 defined as the cokernel:

$$0 \rightarrow \mathcal{O}(-5)^{\alpha-2} \xrightarrow{f_1, \dots, f_2} F \rightarrow Q \rightarrow 0$$

where $F \in \mathcal{L}_2$ and $f_i \in H^0(F(5))$ are general global sections of $F(5)$.

2.8. Let $\mathcal{F} \subset \mathcal{L}_1 \cap \mathcal{L}_2$ be the irreducible family of rank 3 vector bundles R on \mathbf{P}^3 defined as the cokernel:

$$0 \rightarrow \mathcal{O}(-5)^{\alpha-2} \xrightarrow{g_1, \dots, g_2} G \rightarrow R \rightarrow 0$$

where $G \in \mathcal{L}$ and $g_i \in H^0(G(5))$ are general global sections of $G(5)$.

Proposition 2.9.

- (1) *A general vector bundle P of \mathcal{F}_1 is a rank 3 stable vector bundle on \mathbf{P}^3 with Chern classes (287, 42065, 4195775).*
- (2) *A general vector bundle Q of \mathcal{F}_2 is a rank 3 stable vector bundle on \mathbf{P}^3 with Chern classes (287, 42065, 4195775).*
- (3) *A general vector bundle R of \mathcal{F} is a rank 3 stable vector bundle on \mathbf{P}^3 with Chern classes (287, 42065, 4195775).*

Proof. (1) Using the exact sequence:

$$(*) \quad 0 \rightarrow \mathcal{O}(-5)^{\alpha-2} \rightarrow E \rightarrow P \rightarrow 0$$

and the locally free resolution of E given in Proposition 2.4(1) we get:

$$c_t(P) = (1 - 3t)(1 - t)^{74} / ((1 - 6t)^4(1 - 5t)^{68}).$$

Hence $c_1(P) = 287$, $c_2(P) = 42065$ and $c_3(P) = 4195775$.

Let us see that P is stable. Using Hoppe's criterion we need to prove that $H^0(P)_{\text{norm}} = H^0(\Lambda^2 P)_{\text{norm}} = 0$. Since $c_1(P) > 0$ and $c_1(\Lambda^2 P) > 0$, we have $(P)_{\text{norm}} = P(\lambda)$ and $(\Lambda^2 P)_{\text{norm}} = (\Lambda^2 P)(\rho)$ for some $\rho, \lambda \leq -1$. So it suffices to prove that $H^0(P)(-1) = H^0(\Lambda^2 P)(-1) = 0$.

Using the exact sequence (*) and the locally free resolution of E given in Proposition 2.4(1) we easily get that $H^0 E(-1) = H^0 P(-1) = 0$. Again using the exact sequence (*) and taking wedge powers we get the exact sequence

$$0 \rightarrow S^2 \mathcal{O}(-5)^{\alpha-2} \rightarrow \mathcal{O}(-5)^{\alpha-2} \otimes E \rightarrow \Lambda^2 E \rightarrow \Lambda^2 P \rightarrow 0$$

cutting in short exact sequences we get $H^0(\Lambda^2 P)(-1) = H^0(\Lambda^2 E)(-1) = 0$ where the last equality follows from the locally free resolution of E given in Proposition 2.4(1) taking wedge powers and cutting in short exact sequences.

(2) and (3) are analogous. \square

Theorem 2.10. *The moduli space $M_{\mathbf{P}^3}(3; -1, 14609, 1917791)$ is singular.*

Proof. We have constructed two irreducible families \mathcal{F}_1 and \mathcal{F}_2 of rank 3 stable vector bundles on \mathbf{P}^3 with Chern classes (287,42065,4195775) which meets along an irreducible family \mathcal{F} . Hence in order to see that $M := M_{\mathbf{P}^3}(-1, 14609, 1917791) \cong M_{\mathbf{P}^3}(-287, 42065, 4195775)$ is singular it is enough to prove that \mathcal{F}_1 and \mathcal{F}_2 belongs to two different components of M . Using proposition 2.9 and 2.4 we get:

(1) If P is a general vector bundle of \mathcal{F}_1 then:

$$\begin{aligned} H_*^1 P &= H^3 P(3) = 0 \\ h^0 P(3) &= 1 + 10\alpha, \quad h^2 P(3) = 0. \end{aligned}$$

(2) If Q is a general vector bundle of \mathcal{F}_2 then:

$$\begin{aligned} H_*^1 Q &= H^3 Q(3) = 0 \\ h^0 Q(3) &= 10\alpha, \quad h^2 Q(3) = 1. \end{aligned}$$

Therefore, by semicontinuity \mathcal{F}_1 and \mathcal{F}_2 are contained in different components of M which gives what we want. \square

References

- [AO] V. Ancona and G. Ottaviani, *On moduli of instanton bundles on \mathbf{P}^{2n+1}* , Pacific J. Math., to appear.
- [EF] Ph. Ellia and M. Fiorentini, *Défaut de postulation et singularités du schéma de Hilbert*, Ann. Univ. Ferrara, **30** (1984), 185-198.

- [H] H. Hoppe, *Generischer spaltungstypum un zweite Chernklasse stabiler Vektorraum-bündel vom rang 4 auf \mathbf{P}^4* , Math. Z., **187** (1984), 345-360.
- [M1] M. Maruyama, *Moduli of stable sheaves, I*, J. Math. Kyoto Univ., **17** (1977), 91-126.
- [M2] ———, *Moduli of stable sheaves, II*, J. Math. Kyoto Univ., **18** (1978), 557-614.

Received September 13, 1993. The author was partially supported by DGICYT PB94-0850.

DEPARTMENT ALGEBRA Y GEOMETRÍA
UNIVERSIDAD DE BARCELONA, 08007 BARCELONA, SPAIN
E-mail address: miro@cerber.ub.es

Rosa M. Miró-Roig , Singular moduli spaces of stable vector bundles on \mathbf{P}^3	477
Hitoshi Moriyoshi and Toshikazu Natsume , The Godbillon-Vey Cyclic Cocycle and Longitudinal Dirac Operators	483
J.C. Naranjo , The positive dimensional fibres of the Prym map	223
Artur Nicolau and Arne Stray , Nevanlinna's coefficients and Douglas algebras	541
K.K. Park , Entropy of a skew product with a Z^2 -action	227
María Cristina Pereyra , Sobolev spaces on Lipschitz curves	553
T. Sano , Commuting co-commuting squares and finite dimensional Kac algebras	243
H.B. Thompson , Second order ordinary differential equations with fully nonlinear two point boundary conditions	255
H.B. Thompson , Second order ordinary differential equations with fully nonlinear two point boundary conditions II	279
F. Xu , The flat part of non-flat orbifolds	299
Hidenobu Yoshida , A type of uniqueness for the Dirichlet problem on a half-space with continuous data	591

PACIFIC JOURNAL OF MATHEMATICS

Volume 172 No. 2 February 1996

On the failure cycles for the quadratic normality of a projective variety	307
EDOARDO BALLICO	
On the minimal free resolution of general embeddings of curves	315
EDOARDO BALLICO	
On normality of the closure of a generic torus orbit in G/P	321
ROMUALD DABROWSKI	
Paragroupe d'Adrian Ocneanu et algèbre de Kac	331
MARIE-CLAUDE DAVID	
Irreducibility and dimension theorems for families of height 3 Gorenstein algebras	365
SUSAN J. DIESEL	
On the cohomology of the Lie algebra L_2	399
ALICE FIALOWSKI	
Generic differentiability of convex functions on the dual of a Banach space	413
JOHN R. GILES, P. S. KENDEROV, WARREN BRIAN MOORS and S. D. SCIFFER	
Moon hypersurfaces and some related existence results of capillary hypersurfaces without gravity and of rotational symmetry	433
FEI-TSEN LIANG	
Stable relations. II. Corona semiprojectivity and dimension-drop C^* -algebras	461
TERRY ATHERTON LORING	
Singular moduli spaces of stable vector bundles on \mathbf{P}^3	477
ROSA M. MIRÓ-ROIG	
The Godbillon-Vey cyclic cocycle and longitudinal Dirac operators	483
HITOSHI MORIYOSHI and TOSHIKAZU NATSUME	
Nevanlinna's coefficients and Douglas algebras	541
ARTUR NICOLAU and ARNE STRAY	
Sobolev spaces on Lipschitz curves	553
MARÍA CRISTINA PEREYRA	
A type of uniqueness for the Dirichlet problem on a half-space with continuous data	591
HIDENOBU YOSHIDA	



0030-8730(1996)172:2;1-E