

*Pacific
Journal of
Mathematics*

ON QUADRATIC RECIPROCITY OVER FUNCTION FIELDS

KATHY DONOVAN MERRILL AND LYNNE WALLING

Volume 173 No. 1

March 1996

ON QUADRATIC RECIPROCITY OVER FUNCTION FIELDS

KATHY D. MERRILL AND LYNNE H. WALLING

A proof of quadratic reciprocity over function fields is given using the inversion formula of the theta function.

Over the years, many authors have produced proofs of the law of quadratic reciprocity. In 1857, Dedekind [2] stated that quadratic reciprocity holds over function fields; this was later proved by Artin [1]. One of the simplest proofs over the rational numbers relies on the functional equation of the classical theta function (see, for example, [3]); this technique was later generalized by Hecke [4] to number fields. In this note we use an analogous technique to give a simple and direct proof of quadratic reciprocity over rational function fields. We thank David Grant for suggesting this application of Theorem 2.3 of [6].

The reader is referred to [5] for a more complete discussion of the history of the Law of Quadratic Reciprocity.

Let $\mathbf{F} = \mathbf{F}_p$ be a finite field with p elements; for the sake of clarity we assume p is an odd prime. Let T be an indeterminate, and set $\mathbf{A} = \mathbf{F}[T]$. Then for $\alpha, \beta \in \mathbf{A}$ with α irreducible, let

$$\left(\frac{\beta}{\alpha}\right) = \begin{cases} 1 & \text{if } \beta \text{ is a (nonzero) quadratic residue modulo } \alpha, \\ -1 & \text{if } \beta \text{ is a (nonzero) quadratic nonresidue modulo } \alpha, \\ 0 & \text{if } \alpha \text{ divides } \beta. \end{cases}$$

We will show that for $\alpha, \beta \in \mathbf{A}$ distinct monic irreducible polynomials,

$$\left(\frac{\beta}{\alpha}\right) = \begin{cases} \left(\frac{-1}{p}\right) \left(\frac{\alpha}{\beta}\right) & \text{if } \deg \alpha, \deg \beta \text{ are both odd,} \\ \left(\frac{\alpha}{\beta}\right) & \text{otherwise.} \end{cases}$$

We require the following definitions.

Let $\mathbf{K} = \mathbf{F}(T)$; let \mathbf{K}_∞ denote the completion of \mathbf{K} with respect to the “infinite” valuation $|\cdot|_\infty$ given by $|\alpha/\beta|_\infty = p^{\deg \alpha - \deg \beta}$ where $\alpha, \beta \in \mathbf{A}$. (We adopt the convention that $\deg 0 = -\infty$, and hence $|0|_\infty = 0$.) One easily sees that $\mathbf{K}_\infty = \mathbf{F}\left(\left(\frac{1}{T}\right)\right)$, formal Laurent series in $\frac{1}{T}$; for $x \in \mathbf{K}_\infty$, we write $x = \sum_{j=-\infty}^n x_j T^j$. The “unit ball” or “ring of integers” in \mathbf{K}_∞ is

$\mathcal{O}_\infty = \{x \in \mathbf{K}_\infty : |x|_\infty \leq 1\} = \mathbf{F} \left[\left[\frac{1}{T} \right] \right]$, formal Taylor series in $\frac{1}{T}$. Set $G = PSL_2(\mathbf{K}_\infty)$; then the maximal compact subgroup of G (with respect to the standard topology induced on G by $|\cdot|_\infty$) is $PSL_2(\mathcal{O}_\infty)$. Thus we set

$$\mathbf{H} = PSL_2(\mathbf{K}_\infty) / PSL_2(\mathcal{O}_\infty).$$

We can view $PSL_2(*)$ as a subgroup of $PGL_2(*)$; so we consider a matrix of $PSL_2(*)$ equivalent to every nonzero scalar multiple of the matrix. Then as shown in [6],

$$\left\{ \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} : y = T^{2m}, m \in \mathbf{Z}, x \in T^{2m+1} \mathbf{A} \right\}$$

is a complete set of representatives for \mathbf{H} . For each $z \equiv \begin{pmatrix} y & x \\ 0 & 1 \end{pmatrix} \in \mathbf{H}$, set

$$\theta(z) = \sum_{\delta \in \mathbf{A}} \chi_{\mathcal{O}_\infty}((T\delta)^2 y) e\{(T\delta)^2 x\}$$

where $e\{\gamma\} = e\left\{\sum_{j \geq N} \gamma_j T^j\right\} = \exp(2\pi i \gamma_1 / p)$ and $\chi_{\mathcal{O}_\infty}$ is the characteristic function for \mathcal{O}_∞ .

As in the classical setting, we will connect this theta series to quadratic reciprocity through Gauss sums. Accordingly, for $\alpha, \beta \in \mathbf{A}$ with α irreducible and α not dividing β , define the Gauss sum $G_\alpha(\beta)$ to be $G_\alpha(\beta) = \sum_{\delta \in \mathbf{A}/\alpha\mathbf{A}} e\{\beta\delta^2 T^2 / \alpha\}$.

Lemma 1. For $\alpha, \beta \in \mathbf{A}$ with α irreducible and $\alpha \nmid \beta$, $\left(\frac{\beta}{\alpha}\right) = \frac{G_\alpha(\beta)}{G_\alpha(1)}$.

Proof. We have

$$G_\alpha(\beta) = \sum_{\delta \in \mathbf{A}/\alpha\mathbf{A}} \left(1 + \left(\frac{\delta}{\alpha}\right)\right) e\{\beta\delta T^2 / \alpha\} = \sum_{\delta \in \mathbf{A}} \left(\frac{\delta}{\alpha}\right) e\{\beta\delta T^2 / \alpha\}$$

and for $\beta' \in \mathbf{A}$ such that $\beta\beta' \equiv 1 \pmod{\alpha}$

$$= \sum_{\delta \in \mathbf{A}/\alpha\mathbf{A}} \left(\frac{\delta\beta'}{\alpha}\right) e\{\beta\delta\beta' T^2 / \alpha\} = \left(\frac{\beta'}{\alpha}\right) G_\alpha(1) = \left(\frac{\beta}{\alpha}\right) G_\alpha(1).$$

□

Lemma 2. For α, β relatively prime irreducible polynomials, $G_\alpha(\beta)G_\beta(\alpha) = G_{\alpha\beta}(1)$.

Proof. Notice that the map $(\delta + \alpha\beta\mathbf{A}, \gamma + \alpha\beta\mathbf{A}) \mapsto \delta + \gamma + \alpha\beta\mathbf{A}$ is an injective homomorphism from $(\beta\mathbf{A}/\alpha\beta\mathbf{A}) \times (\alpha\mathbf{A}/\alpha\beta\mathbf{A})$ into $\mathbf{A}/\alpha\beta\mathbf{A}$; since the cardinalities of the domain and the codomain are finite and equal, the map is an

isomorphism. Also notice that for $\delta \in \beta\mathbf{A}$ and $\gamma \in \alpha\mathbf{A}$, $e\{(\delta + \gamma)^2 T^2 / \alpha\beta\} = e\{\delta^2 T^2 / \alpha\beta\} e\{\gamma^2 T^2 / \alpha\beta\}$. Thus

$$G_{\alpha\beta}(1) = \sum_{\delta \in \mathbf{A}/\alpha\mathbf{A}} e\{(\beta\delta)^2 T^2 / \alpha\beta\} \sum_{\gamma \in \mathbf{A}/\beta\mathbf{A}} e\{(\alpha\gamma)^2 T^2 / \alpha\beta\} = G_{\alpha}(\beta)G_{\beta}(\alpha).$$

□

Combining these two lemmata, we have that for α, β relatively prime irreducible polynomials, $\left(\frac{\alpha}{\beta}\right) \left(\frac{\beta}{\alpha}\right) = \frac{G_{\alpha\beta}(1)}{G_{\alpha}(1)G_{\beta}(1)}$. Thus for formulate the law of Quadratic Reciprocity, we need only evaluate $G_{\gamma}(1)$ for $\alpha \in \mathbf{A}$. This is the content of our final lemma.

Lemma 3. For any $\gamma \in \mathbf{A}$, $G_{\gamma}(1) = p^{\frac{d}{2}} \left(\frac{\gamma_d}{p}\right)^d \sqrt{\left(\frac{-1}{p}\right)^d}$ where $d = \deg \gamma$ and γ_d denotes the coefficient of T^d in γ .

Proof. First notice that by the Euclidean Algorithm, $\{\delta \in \mathbf{A} : \deg \delta < d\}$ is a complete set of representatives for $\mathbf{A}/\gamma\mathbf{A}$. Thus

$$G_{\gamma}(1) = \sum_{\delta \in \mathbf{A}} \chi_{\mathcal{O}_{\infty}}((T\delta)^2 T^{-2d}) e\{(T\delta)^2 / \gamma\}.$$

Letting $z = \begin{pmatrix} T^{-2d} & 1 \\ 0 & 1 \end{pmatrix}$, we see that $G_{\gamma}(1) = \theta(z)$ where $\theta(z)$ is as in [6]. By

the Inversion Formula, we have $\theta(z) = p^{\frac{d}{2}} \left(\frac{\gamma_d}{p}\right)^d \sqrt{\left(\frac{-1}{p}\right)^d} \theta\left(-\frac{1}{z}\right)$ where $-\frac{1}{z} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} z \equiv \begin{pmatrix} 1 & -\gamma \\ 0 & 1 \end{pmatrix}$. Since the only $\delta \in \mathbf{A}$ satisfying $\chi_{\mathcal{O}_{\infty}}((T\delta)^2) = 1$ is $\delta = 0$, $\theta\left(-\frac{1}{z}\right) = 1$. □

These Lemmata easily imply the following

Theorem. Let α, β be relatively prime irreducible polynomials of degrees d and d' respectively. Then

$$\left(\frac{\alpha}{\beta}\right) = \epsilon \left(\frac{\alpha_d}{p}\right)^{d'} \left(\frac{\beta_{d'}}{p}\right)^d \left(\frac{\beta}{\alpha}\right)$$

where

$$\epsilon = \begin{cases} \left(\frac{-1}{p}\right) & \text{if } d, d' \text{ are both odd,} \\ 1 & \text{otherwise.} \end{cases}$$

In particular, when α and β are distinct monic irreducible polynomials,

$$\left(\frac{\alpha}{\beta}\right) = \begin{cases} \left(\frac{-1}{p}\right) \left(\frac{\beta}{\alpha}\right) & \text{if } d, d' \text{ are both odd,} \\ \left(\frac{\beta}{\alpha}\right) & \text{otherwise.} \end{cases}$$

References

- [1] E. Artin, *Quadratische Körper im Gebiete der höheren Kongruenzen*, Math. Zeit., **19** (1924), 153-246.
- [2] R. Dedekind, *Abriss einer Theorie der höheren Congruenzen in Bezug auf einer reellen Primzahl-Modulus*, J. reine und angew. Math., **54** (1857), 1-26.
- [3] H. Dym and H.P. McKean, *Fourier Series and Integrals*, Academic Press, New York, 1972.
- [4] E. Hecke, *Lectures on the Theory of Algebraic Numbers*, Springer-Verlag, New York-Heidelberg-Berlin, 1981.
- [5] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, New York-Berlin-Heidelberg-London-Paris-Tokyo-Hong Kong, 1990.
- [6] K.D. Merrill and L.H. Walling, *Sums of squares over function fields*, Duke Math. J., **71**(3) (1993), 665-684.

Received September 27, 1993 and revised March 25, 1994. The second author was partially supported by NSF grant DMS 9103303.

COLORADO COLLEGE
 COLORADO SPRINGS, CO 80903
E-mail address: merrill@cc.colorado.edu

AND

UNIVERSITY OF COLORADO
 BOULDER, CO 80309-0426
E-mail address: walling@euclid.colorado.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. Beckenbach (1906-1982)

F. Wolf (1904-1989)

EDITORS

Sun-Yung A. Chang (Managing Editor)
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

F. Michael Christ
University of California
Los Angeles, CA 90095-1555
christ@math.ucla.edu

Nicholas Ercolani
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

Robert Finn
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

Steven Kerckhoff
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

Martin Scharlemann
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

Gang Tian
Massachusetts Institute of Technology
Cambridge, MA 02139
tian@math.mit.edu

V. S. Varadarajan
University of California
Los Angeles, CA 90095-1555
vsv@math.ucla.edu

Dan Voiculescu
University of California
Berkeley, CA 94720
dvv@math.berkeley.edu

SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
STANFORD UNIVERSITY
UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
UNIVERSITY OF UTAH
UNIVERSITY OF WASHINGTON
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The table of contents and the abstracts of the papers in the current issue, as well as other information about the Pacific Journal of Mathematics, may be found on the Internet at <http://www.msri.org>.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$245.00 a year (10 issues). Special rate: \$123.00 a year to individual members of supporting institutions. Subscriptions, back issues published within the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143 Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,
the American Mathematical Society's TEX macro system.
Copyright © 1995 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 173 No. 1 March 1996

Isometric immersions of H_1^n into H_1^{n+1}	1
KINETSU ABE	
Rotationally symmetric hypersurfaces with prescribed mean curvature	29
MARIE-FRANÇOISE BIDAUT-VÉRON	
The covers of a Noetherian module	69
JIAN-JUN CHUAI	
On the odd primary cohomology of higher projective planes	77
MARK FOSKEY and MICHAEL DAVID SLACK	
Unit indices of some imaginary composite quadratic fields. II	93
MIKIHITO HIRABAYASHI	
Mixed automorphic vector bundles on Shimura varieties	105
MIN HO LEE	
Trace ideal criteria for Toeplitz and Hankel operators on the weighted Bergman spaces with exponential type weights	127
PENG LIN and RICHARD ROCHBERG	
On quadratic reciprocity over function fields	147
KATHY DONOVAN MERRILL and LYNNE WALLING	
(A_2) -conditions and Carleson inequalities in Bergman spaces	151
TAKAHIKO NAKAZI and MASAHIRO YAMADA	
A note on a paper of E. Boasso and A. Larotonda: "A spectral theory for solvable Lie algebras of operators"	173
C. OTT	
Tensor products with anisotropic principal series representations of free groups	181
CARLO PENSAVALLE and TIM STEGER	
On Ricci deformation of a Riemannian metric on manifold with boundary	203
YING SHEN	
The Weyl quantization of Poisson $SU(2)$	223
ALBERT JEU-LIANG SHEU	
Weyl's law for $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$	241
ERIC GEORGE STADE and DOROTHY IRENE WALLACE (ANDREOLI)	
Minimal hyperspheres in two-point homogeneous spaces	263
PER TOMTER	
Subalgebras of little Lipschitz algebras	283
NIKOLAI ISAAC WEAVER	



0030-8730(1996)173:1;1-E