

*Pacific  
Journal of  
Mathematics*

WEYL'S LAW FOR  $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$

ERIC GEORGE STADE AND DOROTHY IRENE WALLACE (ANDREOLI)

## WEYL'S LAW FOR $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$

ERIC STADE AND D.I. WALLACE

**In this paper we derive a Weyl's law, or asymptotic description of the distribution of eigenvalues of the Laplacian, on  $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$ . Our main tool in this derivation is the Selberg trace formula for the space. Our Weyl's law, which refines the present theory for the space in question, is also seen to coincide with known results in the case where  $SL(3, \mathbb{Z})$  is replaced by a cocompact discrete subgroup.**

### §1. Introduction; statement of results.

The distribution of eigenvalues of the Laplacian for a general bounded domain  $B$  in  $\mathbb{R}^n$  was first studied systematically by H. Weyl [We]. There he gave a precise asymptotic formula, of the form  $N_B(x) \sim \kappa_0 x^{n/2}$  ( $\kappa_0$  a constant), for the number  $N_B(x)$  of eigenvalues less than  $x$  of this Laplacian. It is now standard to refer to such a result as a "Weyl's law" for the domain  $B$ .

Since then, many authors have considered generalizations of Weyl's law to other kinds of Riemannian manifolds. Particular attention, in large part because of their relevance to the theory of automorphic forms, has been given to the spaces  $\Gamma \backslash SL(2, \mathbb{R}) / SO(2, \mathbb{R})$ , where  $\Gamma$  is a cofinite (but, usually, not cocompact) discrete subgroup of  $SL(2, \mathbb{R})$ . Results concerning eigenvalues of the Laplacian on such spaces, and on products of such spaces, may be found in [Ef, Hej1, Hej2, Hu1], and [Hu-Te] (among other works).

In the present article, we wish to develop a Weyl's law in the higher-rank setting of  $\Gamma \backslash H^3$ , where  $\Gamma = SL(3, \mathbb{Z})$  and  $H^3 = SL(3, \mathbb{R}) / SO(3, \mathbb{R})$ . Actually, what we shall count here is the number  $N(x)$  of linearly independent *cuspidal forms* on  $\Gamma \backslash H^3$  whose eigenvalue for the Laplacian is less than  $x$ . By "cuspidal form" we mean a non-constant, square-integrable, *joint eigenfunction* (on  $\Gamma \backslash H^3$ ) of the algebra  $D$  of all  $SL(3, \mathbb{R})$ -invariant differential operators on  $H^3$ . (In higher-rank settings, the definition of "cuspidal form" also requires the vanishing of certain "cuspidal integrals" of the eigenfunction. However, in the current situation, such a definition of cuspidal form coincides with the one we have given.) The cuspidal forms are central to harmonic analysis on  $L^2(\Gamma \backslash H^3)$ , and to the corresponding theory of automorphic forms on this space. (Technically, our count will also include the constant functions on

$\Gamma \backslash H^3$ , which have eigenvalue 0 under the Laplacian. However, note that the result of Theorem 1 below is unaltered by the addition or exclusion of a subspace of dimension one.)

The algebra  $D$  is known [Hel, p. 432] to have two generators. We shall denote these by  $\Delta_1$  and  $\Delta_2$  respectively, where  $\Delta_1$  is the Laplacian on our space and  $\Delta_2$  is a third-order differential operator. (See §2 below for details.) Because  $D$  comprises substantially more than the Laplacian alone, it may seem that counting joint eigenfunctions of this algebra is a problem somewhat different than that of obtaining a Weyl's law for  $\Gamma \backslash H^3$ . However the problems are in fact the same, according to the following observations:  $\Delta_1$  and  $\Delta_2$  commute, and any eigenspace of  $\Delta_1$  (in  $L^2(\Gamma \backslash H^3)$ ) is finite-dimensional. (The first statement is a consequence of [Hel, p. 432]; the second is a general fact regarding Laplacian operators that also follows, in the present setting, from Theorem 1 below.) So any  $\Delta_1$ -eigenspace has a basis consisting of finitely many eigenvectors for  $\Delta_2$ —such an eigenvector is, by the above, either a cusp form or a constant. Therefore,  $N(x)$  in fact equals the dimension of the subspace of  $L^2(\Gamma \backslash H^3)$  spanned by functions that are eigenvectors of  $\Delta_1$ , and have  $\Delta_1$ -eigenvalue  $< x$ . (The above argument was pointed out to us by David Farmer.)

To state our Weyl's law, let us denote by  $\text{vol}(\Gamma \backslash H^3)$  the “hyperbolic volume” (computed according to the  $SL(3, \mathbb{R})$ -invariant volume element given in §2) of this space. We then have

**Theorem 1.** *Let all notation be as above. Then*

$$N(x) \sim \frac{\text{vol}(\Gamma \backslash H^3)}{(4\pi)^{5/2} \Gamma(7/2)} x^{5/2}$$

as  $x \rightarrow \infty$ .

**Remark.** The asymptotic result of our theorem is identical to the one appearing in Weyl's law for bounded domains in  $\mathbb{R}^5$ , and for compact Riemannian 5-manifolds (see Weyl [We] and Chavel [Ch]). Moreover, our theorem sharpens results of Donnelly [Do], and of Huntley [Hu2] (who consider more general situations). Specifically, Donnelly obtains the value  $\text{vol}(\Gamma \backslash H^3)/((4\pi)^{5/2} \Gamma(7/2))$  as an upper bound for  $\overline{\lim} x^{-5/2} N(x)$ ; according to our theorem, the  $\overline{\lim}$  is actually a limit, and is equal to the stated upper bound. Huntley does show that  $x^{-5/2} N(x)$  has a finite limit, but does not determine this limit explicitly.

We further observe that Huntley's count includes certain functions that are square-integrable, but *not* cusp forms. Specifically, for the eigenfunctions of  $\Delta_1$  contributing to his Weyl's law, the relevant cuspidal integrals need not vanish identically, but only outside a specified compact subset of  $\Gamma \backslash H^3$ .

(Such functions are not generally eigenfunctions in the true sense, or even in the sense of distributions, but rather in a certain  $L^2$  sense described in [Hu2].) As our Weyl's law counts only the cusp forms, it is technically somewhat different than his.

We now turn to the proof of Theorem 1. Our principal tool in this proof will be the Selberg trace formula—as realized by Wallace [Wa]—for nice functions  $g$  on the space  $SO(3, \mathbb{R}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$ . The details of this trace formula are given below; the basic idea behind its application is as follows. The left-hand, or “spectral,” side of the trace formula may be interpreted, for appropriate choice of  $g$ , as the Laplace transform  $L_N(T)$  of  $N(x)$ . By studying the various “orbital integrals” on the right-hand side of the trace formula, one may obtain an asymptotic equation, as  $T \rightarrow 0^+$ , for  $L_N(T)$ . Then, by applying a Tauberian theorem [Wi] for the Laplace transform, information on  $L_N(T)$  may be translated into the desired formula for  $N(x)$  itself.

The arguments just described will be fleshed out in §3 below, following a brief discussion, in §2, of harmonic analysis on  $H^3$ . In §4, an inversion formula for a certain integral transform, known as the “Helgason transform”  $\widehat{g}$  (see §2), of functions  $g$  will be proved. This formula will be required in what follows, since the trace formula relates expressions involving  $g$  to others involving  $\widehat{g}$ , and moreover the particular  $g$  of interest to us will in fact be defined as a function with a given  $\widehat{g}$ . In §§5–7 we will carry out the computation of  $L_N(T)$ , by considering the orbital terms on the right-hand side of the trace formula. In the last section, we will combine this computation with the relevant Tauberian theorem to obtain the result embodied by Theorem 1.

The authors would like to thank David Grant, Jonathan Huntley, and Thomas Shemanske for many helpful conversations.

## §2. Harmonic analysis on $SL(3, \mathbb{R}) / SO(3, \mathbb{R})$ .

In this section, we recall some basic ideas concerning coordinate systems and differential operators on  $H^3 = SL(3, \mathbb{R}) / SO(3, \mathbb{R})$ .

First, using the Iwasawa decomposition for  $SL(3, \mathbb{R})$ , we may identify  $H^3$  with the space  $AN$ , where  $A$  consists of diagonal matrices

$$a = \text{diag}(a_1, a_2, (a_1 a_2)^{-1}) \quad (a_i > 0)$$

and  $N$  consists of upper triangular unipotents

$$n = \begin{pmatrix} 1 & x_2 & x_3 \\ 0 & 1 & x_1 \\ 0 & 0 & 1 \end{pmatrix}$$

( $x_i \in \mathbb{R}$ ). In these coordinates, the  $SL(3, \mathbb{R})$ -invariant measure on  $H^3$  is given by

$$(2.1) \quad dn \, da = 54 \, dx_1 \, dx_2 \, dx_3 \frac{da_1 \, da_2}{a_1 \, a_2}.$$

The factor of 54 is included to provide a measure equal to the one specified in [Ch] and [Do].

As mentioned in §1, the algebra  $D$  of  $SL(3, \mathbb{R})$ -invariant differential operators on  $H^3$  has two generators: the Laplacian  $\Delta_1$ —which, properly normalized, is a *positive* operator (cf. [Bo])—and a third-order operator  $\Delta_2$ . The former may be computed according to a formula in Chavel [Ch]. On the other hand, both operators have been determined explicitly by Bump [Bu] (using a fundamentally different coordinatization of  $H^3$ ). In our coordinates, we have

**Proposition 2.1.**

$$\begin{aligned} \Delta_1 &= -\frac{1}{3}(H_1^2 - H_1 + H_2^2 - H_2 - H_1H_2 + X_1^2 + X_2^2 + Z_0^2); \\ \Delta_2 &= -H_1^2H_2 + H_1H_2^2 + H_1^2 - H_2^2 - H_1 + H_2 \\ &\quad - X_1^2H_2 + X_2^2H_1 - Z_0^2H_1 + Z_0^2H_2 + Z_0X_2X_1 + Z_0X_1X_2, \end{aligned}$$

where

$$\begin{aligned} H_1 &= \frac{a_1}{3} \frac{\partial}{\partial a_1} + \frac{a_2}{3} \frac{\partial}{\partial a_2} - x_2 \frac{\partial}{\partial x_2} - x_3 \frac{\partial}{\partial x_3}; \\ H_2 &= \frac{2a_1}{3} \frac{\partial}{\partial a_1} - \frac{a_2}{3} \frac{\partial}{\partial a_2} - x_1 \frac{\partial}{\partial x_1} - x_2 \frac{\partial}{\partial x_2}; \\ X_1 &= x_1 \frac{\partial}{\partial x_2} + \frac{\partial}{\partial x_3}; \quad X_2 = \frac{\partial}{\partial x_1}; \quad Z_0 = \frac{\partial}{\partial x_2}. \end{aligned}$$

*Proof.* These are exactly equations (2.31) and (2.36) in Bump [Bu], except that we have multiplied by  $-1/3$  his definition of  $\Delta_1$ : this normalization yields the same Laplacian as appears in [Ch] and [Do]. We must restate Bump's differential operators  $H_1, H_2$ , etc. in terms of our coordinates on  $H^3$ , but this is straightforward: Bump gives us these operators as matrices in the Lie algebra of  $SL(3, \mathbb{R})$ , where such a matrix  $X$  defines a differential operator in the usual way:

$$Xf(z) = \left. \frac{d}{dt} f(ze^{tX}) \right|_{t=0}$$

( $f \in C^\infty(SL(3, \mathbb{R}))$ ,  $z \in H^3$ ). In particular, we have

$$H_1 = \text{diag}\left(\frac{1}{3}, \frac{1}{3}, -\frac{2}{3}\right); \quad H_2 = \text{diag}\left(\frac{2}{3}, -\frac{1}{3}, -\frac{1}{3}\right);$$

$$X_1 = E_{2,3}; \quad X_2 = E_{1,2}; \quad Z_0 = E_{1,3},$$

where  $E_{i,j}$  denotes the matrix with a one in the  $i, j$  slot and zeroes elsewhere. We merely choose  $z$  to have the  $AN$ -coordinates given above, and our proposition follows after a few brief calculations. □

We now wish to consider integral transforms on  $H^3$ . (See, for example, [Te2] §4.2.) Let  $K = SO(3, \mathbb{R})$ . Suppose  $g$  is a bi- $K$ -invariant function on  $SL(3, \mathbb{R})$ : that is, we may consider  $g$  as a function on  $H^3$  satisfying  $g(kz) = g(z)$  for all  $k \in K$ ,  $z \in H^3$ . Then the Helgason transform  $\widehat{g}$  of  $g$  is defined, for  $s, t \in \mathbb{C}$ , by

$$(2.2) \quad \widehat{g}(s, t) = \int_{AN} g(an) a_1^{3s+t} a_2^{3s-t} dn da.$$

(Our definition of the Helgason transform certainly makes sense for  $g$  infinitely differentiable and of compact support; however it is pointed out in [Te2] that this transform extends to an isometry on all of  $L^2(K \backslash G / K)$ .) We remark that the function  $H_{s,t}(an) = a_1^{3s+t} a_2^{3s-t}$  is an eigenfunction of  $D$ , for any  $s, t$ ; this follows, for example, from Proposition 2.1. We write  $\Delta_1 H_{s,t} = \lambda H_{s,t}$  and  $\Delta_2 H_{s,t} = \mu H_{s,t}$ ; then it is readily computed that

$$(2.3) \quad \lambda = s(1-s) + \frac{1}{3}t(1-t); \quad \mu = (1-2s)(s+t-1)(s-t).$$

(The seemingly unusual normalization of the exponents in the definition of  $H_{s,t}$  is justified by the simple form that this provides for the inversion formula of §4. Moreover, it may be seen that  $H_{s,t}$  has the same eigenvalues under  $D$  as does the Eisenstein series  $E(z, s, t)$ , as defined in [Wa]; therefore we may use results from the latter work without significant modification.)

A related transform of  $g$  is the Harish-Chandra transform  $\widetilde{g}$ , defined for a diagonal matrix  $a$  as above by

$$(2.4) \quad \widetilde{g}(a) = \int_N g(an) dn.$$

Note that Mellin inversion applied to (2.2) gives us

$$(2.5) \quad \widetilde{g}(a) = \frac{1}{(2\pi i)^2} \int_{\text{Re}(v)=c_2} \int_{\text{Re}(u)=c_1} \widehat{g}\left(\frac{u+v}{6}, \frac{u-v}{2}\right) a_1^{-u} a_2^{-v} du dv,$$

for appropriate  $c_1, c_2$ .

To state the required version of the trace formula for  $SL(3, \mathbb{Z}) \backslash H^3$ , we will need a convenient way of indexing eigenvalues of cusp forms on this space. We do this in a way consistent with the notation previously introduced for  $H_{s,t}$ . Specifically, we let  $\{\varphi_n\}_{n=1}^\infty$  denote a maximal orthonormal set of such cusp forms, arranged in order of increasing eigenvalue for the Laplacian, and let  $\varphi_0$  denote the constant function 1. Writing  $\Delta_1 \varphi_n = \lambda_n \varphi_n$  for all  $n$ , we then have

$$0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots .$$

We will also write  $\mu_n$  for the eigenvalue of  $\varphi_n$  under  $\Delta_2$ . Finally, we introduce “eigenparameters”  $s_n, t_n$  in a way analogous to equation (2.3): namely, we write

$$(2.6) \quad \lambda_n = s_n(1 - s_n) + \frac{1}{3}t_n(1 - t_n); \quad \mu_n = (1 - 2s_n)(s_n + t_n - 1)(s_n - t_n).$$

It is not hard to show that, for arbitrary  $\lambda_n, \mu_n$ , these equations may be solved for  $s_n, t_n$ .

We now turn to our discussion of the trace formula for  $SL(3, \mathbb{Z}) \backslash H^3$ , and of its application to Weyl’s law.

### §3. The trace formula and Weyl’s law on $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$ .

It will be instructive to recall the notion of a trace formula in a quite general setting. To this end, we begin with a noncompact real semisimple Lie group  $G$ , a maximal compact subgroup  $K$ , and a cofinite discrete subgroup  $\Gamma$ . Then  $G/K$  is a Riemannian symmetric space on which  $\Gamma$  acts. Let  $g(x) : G \rightarrow \mathbb{C}$  be a  $K$ -bi-invariant function. One constructs the kernel of an integral operator on  $L^2(\Gamma \backslash G/K)$  by setting

$$F(z, w) = \sum_{\gamma \in \Gamma} g(z^{-1} \gamma w).$$

The Selberg trace formula, introduced by Selberg in the 1950’s (cf., for example, [Se1, Se2]), equates two expressions for the trace of this operator on the discrete part of the spectrum of  $D$ . As above in the setting of  $SL(3, \mathbb{R}) / SO(3, \mathbb{R})$ ,  $D$  denotes the algebra of  $G$ -invariant differential operators on  $G/K$ , or on  $\Gamma \backslash G/K$ .

The two sides of the trace formula take the following forms. On the one hand (the left-hand side of the trace formula), one may use Mercer’s Theorem for Hilbert–Schmidt operators to write this trace as a sum over eigenvalues of  $D$ . The quantity being summed is the “Helgason” or “Selberg”

transform  $\widehat{g}$  of  $g$ :  $\widehat{g}$  is an integral of  $g$  against an eigenfunction  $\varphi$  of  $D$  on  $G/K$ . (The arguments of  $\widehat{g}$  will be the eigenvalues of  $\varphi$  under generators for  $D$ , or certain parameters indexing these eigenvalues, as we have seen above in the specific case of interest to us.) On the other hand (the right-hand side of the trace formula), one can break up the sum defining  $F(z, w)$  according to the conjugacy class of  $\gamma$ , and use this decomposition to break up the integral defining the trace of  $F$ . Each of the resulting integrals is called an “orbital integral” for  $g$ . (Care must be exercised with the contribution from the “parabolic” orbital integrals; it is only because the “continuous” part of the kernel  $F$  has been subtracted that this contribution is finite.) The various orbital integrals may themselves be expressed as integrals and sums of the Helgason transform, and the related Harish–Chandra transform, of  $g$ .

Let us now restrict our attention to the situation considered in §1: namely, to the case  $G = SL(3, \mathbb{R})$ ;  $K = SO(3, \mathbb{R})$ ;  $\Gamma = SL(3, \mathbb{Z})$ . In this setting, the above generalities regarding a trace formula have been made explicit by Wallace [Wa]. These results may be summarized by

**Proposition 3.1.** *Let  $g \in C_c^\infty(SO(3, \mathbb{R}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R}))$  (however, see the remarks below); let all notation be as in §§ 1 and 2 above. Then*

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{g}(s_n, t_n) &= \text{vol}(\Gamma \backslash H^3) g(I) \\ &+ \sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) |\varepsilon_1^2 \varepsilon_2|}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} \widetilde{g}(\text{diag}(|\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|)) \\ &+ \sum_{(r, \theta)} \frac{|\ln r_0| \text{Cl}(\mathbb{Z}[r])}{|1 - 2r^{-3} \cos \theta + r^{-6}|} \int_{\text{Re}(s)=1/2} \int_{\text{Re}(t)=1/2} \widehat{g}(s, t) \frac{r^{1-s} e^{-2\theta \text{Im}(t)}}{1 + e^{-2\pi \text{Im}(t)}} ds dt \\ &+ K_1 \widehat{g}\left(\frac{1}{2}, \frac{1}{2}\right) + K_2 \widehat{g}\left(\frac{1}{3}, 1\right) + K_3 \widehat{g}\left(\frac{2}{3}, 1\right) \end{aligned}$$

where:

- a.  $I$  denotes the identity coset (the first term on the right-hand side of Proposition 3.1 is called the “identity” or “central” term).
- b. The sum in  $(\varepsilon_1, \varepsilon_2, \varepsilon_3)$  runs over all distinct real triples with  $|\varepsilon_1| > |\varepsilon_2| > |\varepsilon_3|$  (no  $\varepsilon_i = 1$ ), such that there is some (by definition, hyperbolic) matrix in  $SL(3, \mathbb{Z})$  with eigenvalues  $\varepsilon_1, \varepsilon_2, \varepsilon_3$ . Also,  $\text{Reg}$  denotes the regulator of the order, and  $\text{Cl}$  the narrow class number. (The second term on the right-hand side above is called the “hyperbolic” term.)
- c. The sum in  $(r, \theta)$  runs over all distinct pairs with  $r > 0$  ( $r \neq 1$ ) and  $0 < \theta \leq \pi$ , such that there is some (by definition, loxodromic) matrix  $g$  in  $SL(3, \mathbb{Z})$  with eigenvalues  $r^{-2}, re^{i\theta}, re^{-i\theta}$ . Again,  $\text{Cl}$  denotes the narrow class

number; also,  $r_0$  is such that the centralizer of  $g$  (in  $SL(3, \mathbb{Z})$ ), which is infinite cyclic, has generator conjugate (in  $SL(3, \mathbb{C})$ ) to  $\text{diag}(r_0^{-2}, r_0 e^{i\theta_0}, r_0 e^{-i\theta_0})$ . (The third term on the right-hand side above is called the “loxodromic” term.)

*d.*  $K_1, K_2, K_3$  are universal constants. (The sum of the last three terms on the right-hand side above is called the “parabolic” term.)

**Remarks on Proposition 3.1.** We note that the version of the Selberg trace formula that we have stated above is slightly different than that given in [Wa, §8]. Namely, in the latter work both the hyperbolic and loxodromic terms are broken down into sums over powers of “primitive” conjugacy classes. The forms given above for the hyperbolic and loxodromic terms will be more convenient for our purposes. (See §§ 6 and 7, below.)

We observe further that the hyperbolic term above is, as in [Wa], actually a sum over all distinct sets of eigenvalues of hyperbolic matrices. The stipulation  $|\varepsilon_1| > |\varepsilon_2| > |\varepsilon_3|$  precisely assures that each such set is counted only once. Similarly, the loxodromic term is a sum over all sets of eigenvalues of loxodromic matrices; in this case the requirements  $r > 0$  and  $0 < \theta \leq \pi$  assure that no set of eigenvalues occurs twice.

Finally, we make the crucial observation that the trace formula may be extended to include a broader class of “test function”  $g$  than that stipulated in the above proposition. In particular, it is straightforward to show, using techniques analogous to those employed in [He1, Chapter 1] and [He2, Chapter 8], that Proposition 3.1 remains valid for smooth functions  $g \in L^2(K \backslash G / K)$ , such that  $\widehat{g}(s, t)$  decays exponentially as  $\text{Im}(s)$  or  $\text{Im}(t)$  approaches  $\pm\infty$  (and the corresponding real part remains bounded). We will apply Proposition 3.1 to such a function  $g$ , as described immediately below (the class of allowable  $g$  may be made broader still, with a small amount of additional work, but we will not require such a generalization).

The application of the above trace formula to Weyl’s law will proceed as follows. First, for  $T > 0$ , we will let  $g_T$  be a left  $K$ -invariant function on  $H^3$  such that

$$(3.1) \quad \widehat{g}_T(s, t) = e^{-\lambda T} = e^{-T[s(1-s)+t(1-t)/3]}.$$

Such a function  $g_T$  must exist for the following reason: we have seen that  $s(1-s) + t(1-t)/3$  is the eigenvalue of  $\Delta_1$  belonging to the eigenfunction  $H_{s,t}(z) = a_1^{3s+t} a_2^{3s-t}$ . Such an eigenvalue is invariant under the action of the Weyl group on  $(s, t)$ , as described in [Te2], Chapter 4 or [Bu], Chapter 2 (in the respective coordinate systems). But it follows from [Te2], Theorem 1, p. 88 (after restriction to the determinant-one surface  $SP_3$ ), that the Helgason transform is an isometry from square-integrable,  $K$ -invariant functions

on  $H^3$  onto the space of functions of  $(s, t)$  that are square-integrable with respect to the spectral measure (cf. §4) and are invariant under this action.

We will then apply Proposition 3.1 to our function  $\widehat{g}_T$ : on the left-hand side we get

$$L_N(T) \equiv \sum_{n=0}^{\infty} e^{-\lambda_n T}.$$

In particular  $L_N(T)$  is a Laplace, or Laplace–Stieltjes, transform of  $N(x)$ . By estimating each term occurring on the right-hand side of the trace formula, we may obtain asymptotic information (as  $T \rightarrow 0^+$ ) regarding  $L_N(T)$ . Theorem 1 will then follow, by the Tauberian theorem mentioned in §1.

We now proceed with our derivation of the inversion formula for the Helgason transform.

#### §4. Inversion of the Helgason transform on $H^3 = SL(3, \mathbb{R})/SO(3, \mathbb{R})$ .

In this section, we will prove

**Proposition 4.1.** *Let  $g$  be a smooth, square-integrable function on  $H^3$  that is  $K = SO(3, \mathbb{R})$ -invariant, i.e.  $g(kz) = g(z)$  for all  $k \in K$ . Then*

$$g(I) = \frac{1}{2^4 3^3 \pi^3} \int_{r_1, r_2 \in \mathbb{R}} \widehat{g}\left(\frac{1}{2} + ir_1, \frac{1}{2} + ir_2\right) \cdot r_2 (r_2^2 - 9r_1^2) \tanh \pi r_2 \tanh \pi \frac{r_2 + 3r_1}{2} \tanh \pi \frac{r_2 - 3r_1}{2} dr_1 dr_2.$$

*Proof.* Our starting point for the proof of Proposition 4.1 will be the “Helgason inversion formula” that appears in [Te2] (Theorem 1, p. 88). This formula pertains to functions on the space

$$\mathcal{P}_n = \{\text{positive definite, symmetric real } n \times n \text{ matrices}\},$$

which is a  $GL(n, \mathbb{R})$ -space via the action  $(Y, g) \rightarrow Y[g] = {}^t g Y g$  ( $Y \in \mathcal{P}_n, g \in GL(n, \mathbb{R})$ ). We will be concerned with the case  $n = 3$ . By first identifying integrals on  $AN$  with integrals on the determinant-one surface  $SP_3$  of  $\mathcal{P}_3$ , and then restricting the inversion formula in [Te2] to  $SP_3$ , we will deduce from Helgason inversion the required inversion formula for  $H^3$ .  $\square$

We begin with

**Lemma 4.1.** *Let  $f$  be integrable on  $SP_3$ . Then, for a suitably normalized  $SL(3, \mathbb{R})$ -invariant measure  $dW$ ,*

$$\int_{W \in \mathcal{SP}_3} f(W) dW = \frac{2}{27} \int_{AN} f(I[(an)^{-1}]) dn da.$$

*Proof of Lemma.* We begin with formula (1.39), Chapter 4 in [Te2] for integrable functions on  $\mathcal{P}^3$ :

$$\int_{Y \in \mathcal{P}_3} F(Y) dY = 8 \int_{A'N} F \left( I \left[ \left( \begin{pmatrix} a_1 & & \\ & a_2 & \\ & & a_3 \end{pmatrix} \begin{pmatrix} 1 & x_2 & x_3 \\ & 1 & x_1 \\ & & 1 \end{pmatrix} \right)^{-1} \right] \right) dn' da'$$

where  $dY$  is  $GL(3, \mathbb{R})$ -invariant measure on  $\mathcal{P}_3$ ,  $A' = \{\text{diag}(a_1, a_2, a_3) \mid a_i > 0\}$ ,  $dn' = dx_1 dx_2 dx_3$ , and  $da' = da_1 da_2 da_3 / a_1 a_2 a_3$ . We choose the  $SL(3, \mathbb{R})$ -invariant measure  $dW$  on  $\mathcal{SP}_3$  so that  $dY = v^{-1} dv dW$ , where  $v = \det Y$  and  $Y = v^{1/3} W$ . (That such a choice of measures is possible is pointed out in [Te2], §4.1.) Also let  $k$  be a function on  $\mathbb{R}^+$  whose integral with respect to the Haar measure  $dv/v$  exists and is nonzero; put  $F(Y) = k(v)f(W)$ . Then

$$\int_{v>0} k(v) \frac{dv}{v} \int_{W \in \mathcal{SP}_3} f(W) dW = 8 \int_{A'N} k((a_1 a_2 a_3)^{-2}) \cdot f \left( I \left[ \left( \begin{pmatrix} (a_1^2/a_2 a_3)^{1/3} & & \\ & (a_2^2/a_1 a_3)^{1/3} & \\ & & (a_3^2/a_1 a_2)^{1/3} \end{pmatrix} \begin{pmatrix} 1 & x_2 & x_3 \\ & 1 & x_1 \\ & & 1 \end{pmatrix} \right)^{-1} \right] \right) dn' da'.$$

We substitute  $v = (a_1 a_2 a_3)^{-2}$ ;  $u_1 = (a_1^2/a_2 a_3)^{1/3}$ ;  $u_2 = (a_2^2/a_1 a_3)^{1/3}$  into the integral on the right. It is seen that, under this substitution,

$$da' \rightarrow \frac{1}{2} \frac{dv du_1 du_2}{v u_1 u_2}.$$

So we get

$$\int_{v>0} k(v) \frac{dv}{v} \int_{W \in \mathcal{SP}_3} f(W) dW = 4 \int_{v>0} k(v) \frac{dv}{v} \cdot \int_{u_1, u_2 > 0} \int_N f \left( I \left[ \left( \begin{pmatrix} u_1 & & \\ & u_2 & \\ & & (u_1 u_2)^{-1} \end{pmatrix} \begin{pmatrix} 1 & x_2 & x_3 \\ & 1 & x_1 \\ & & 1 \end{pmatrix} \right)^{-1} \right] \right) dn' \frac{du_1 du_2}{u_1 u_2}$$

whence, upon replacing  $u_i$  by  $a_i$  and recalling the factor of 54 present in (2.1), the lemma. □

Next, using Lemma 4.1, we wish to write the Helgason transform  $\hat{g}$  in terms of an analogous transform for functions on  $\mathcal{SP}_3$ . We begin by writing,

for any  $Y \in \mathcal{P}_3$  and  $s = (s_1, s_2, s_3) \in \mathbb{C}^3$ ,  $p_s(Y) = \prod_{i=1}^3 |Y_i|^{s_i}$  where  $Y_i$  denotes the top left-hand  $i \times i$  corner of  $Y$ . Note that  $p_s(W)$  is independent of  $s_3$  for  $W \in SP_3$ ; we will write  $p_{(s_1, s_2)}(W)$  for  $p_s(W)$ . We have:

**Lemma 4.2.** *Let  $g$  be a function on  $H^3$  whose Helgason transform (2.2) converges absolutely; let  $f$  be the function on  $SP_3$  defined by  $f(I[(an)^{-1}]) = g(an)$ . Then*

$$\frac{2}{27} \widehat{g}(s, t) = f^* \left( -\bar{t}, \frac{t - 3s}{2} \right)$$

where, by definition,

$$f^*(s_1, s_2) = \int_{W \in SP_3} f(W) \overline{p_{(s_1, s_2)}(W)} dW.$$

*Proof of Lemma.* This follows immediately from Lemma 4.1, upon noting that

$$H_{s,t}(z) \equiv a_1^{3s+t} a_2^{3s-t} = p_{(-t, (t-3s)/2)}(I[(an)^{-1}]).$$

□

In our next lemma, we obtain an inversion formula on  $SP_3$ .

**Lemma 4.3.** *If  $f$  is square-integrable and  $K$ -invariant on  $SP_3$ , i.e.  $f(W[k]) = f(W)$  for all  $k \in K, W \in SP_3$ , then*

$$f(I) = 2\pi i \omega_3 \int_{\text{Re}(s_i) = -1/2} f^*(s_1, s_2) |c_3(s)|^{-2} ds_1 ds_2$$

where the “spectral measure”  $\omega_3 |c_3(s)|^{-2}$  is given in [Te2] Theorem 1, p. 88 (or equations (4.1), below).

*Proof of Lemma.* Let  $F$  be square-integrable and  $O(3, \mathbb{R})$ -invariant on  $\mathcal{P}_3$ . The Helgason transform  $\widehat{F}(s)$  of  $F$ , for  $s \in \mathbb{C}^3$  as above, may be defined by

$$\widehat{F}(s) = \int_{v>0} v^{\bar{r}} \int_{W \in SP_3} F(v^{1/3}W) \overline{p_s(W)} dW \frac{dv}{v}$$

where  $r = (1/3) \sum_{j=1}^3 js_j$  (cf. [Te2], p. 94). We now take  $F(Y) = k(v)f(W)$ , where  $k$  is a suitable function on  $\mathbb{R}^+$  that does not vanish at  $v = 1$ . (Note that  $f$   $K$ -invariant  $\Rightarrow F$   $O(3, \mathbb{R})$ -invariant.) For simplicity, let us in fact take  $k(v) = e^{-v}$ . Then

$$\widehat{F}(s) = \int_{v>0} e^{-v} v^{\bar{r}} \frac{dv}{v} \int_{W \in SP_3} f(W) \overline{p_s(W)} dW = \Gamma(\bar{r}) f^*(s_1, s_2)$$

(cf. Lemma 4.2 above). We recall the Helgason inversion formula for  $\mathcal{P}_3$  (cf. [Te2], Theorem 1, p. 88):

$$F(Y) = \omega_3 \int_{\substack{\operatorname{Re}(s_1)=\operatorname{Re}(s_2)=-1/2 \\ \operatorname{Re}(s_3)=1/2}} \widehat{F}(s) h_s(Y) |c_3(s)|^{-2} ds$$

where  $h_s$  is a “spherical function” obtained by averaging the power function  $p_s$  over the orthogonal group. For our purposes, it suffices to know that  $h_s(I) = 1$ . Also, one notes from the definition that  $c_3(s)$  is independent of  $s_3$ . So the above inversion formula gives us

$$F(I) = k(1)f(I) = \omega_3 \int_{\operatorname{Re}(s_1)=\operatorname{Re}(s_2)=-1/2} f^*(s_1, s_2) |c_3(s)|^{-2} \cdot \left[ \int_{\operatorname{Re}(s_3)=1/2} \Gamma\left(\frac{1}{s_3} + \frac{2}{3}\frac{1}{s_2} + \frac{1}{3}\frac{1}{s_1}\right) ds_3 \right] ds_1 ds_2.$$

But  $k(1) = 1/e$ , which is  $(2\pi i)^{-1}$  times the value of the integral in  $s_3$  (we are merely invoking Mellin inversion to evaluate this integral). So

$$f(I) = 2\pi i \omega_3 \int_{\operatorname{Re}(s_1)=\operatorname{Re}(s_2)=-1/2} f^*(s_1, s_2) |c_3(s)|^{-2} ds_1 ds_2,$$

as was to be proved. □

We now put the above lemmas together to deduce our inversion formula on  $H^3$ . Let  $g$  be square-integrable and  $K$ -invariant on  $H^3$ ; let  $f$  be the function on  $\mathcal{S}P_3$  with  $g(an) = f(I[(an)^{-1}])$ . Noting that  $f$  is  $K$ -invariant on  $\mathcal{S}P_3$ , we have by Lemma 4.3

$$g(I) = f(I) = 2\pi i \omega_3 \int_{\operatorname{Re}(s_i)=-1/2} f^*(s_1, s_2) |c_3(s)|^{-2} ds_1 ds_2.$$

We substitute  $s_i \rightarrow -\bar{s}_i$ , to get

$$g(I) = 2\pi i \omega_3 \int_{\operatorname{Re}(s_i)=1/2} f^*(-\bar{s}_1, -\bar{s}_2) |c_3(-\bar{s}_1, -\bar{s}_2)|^{-2} ds_1 ds_2;$$

following this with the substitution

$$s = \frac{s_1 + 2s_2}{3}; \quad t = s_1,$$

we get

$$\begin{aligned} g(I) &= 3\pi i \omega_3 \int_{\operatorname{Re}(s)=\operatorname{Re}(t)=1/2} f^*\left(-\bar{t}, \frac{\bar{t}-3\bar{s}}{2}\right) \left| c_3\left(-\bar{t}, \frac{\bar{t}-3\bar{s}}{2}\right) \right|^{-2} ds dt \\ &= \frac{2}{9} \pi i \omega_3 \int_{\operatorname{Re}(s)=\operatorname{Re}(t)=1/2} \widehat{g}(s, t) \left| c_3\left(-\bar{t}, \frac{\bar{t}-3\bar{s}}{2}\right) \right|^{-2} ds dt \end{aligned}$$

by Lemma 4.2. Let us finally substitute  $s = \frac{1}{2} + ir_1$ ,  $t = \frac{1}{2} + ir_2$ . We find from [Te2], p. 88 (and equation (3.23), p. 144 in [Te1]), that

$$(4.1a) \quad \left| c_3 \left( -\bar{t}, \frac{\overline{t-3s}}{2} \right) \right|^{-2} \\ = \pi r_2 (r_2^2 - 9r_1^2) \tanh \pi r_2 \tanh \pi \frac{r_2 + 3r_1}{2} \tanh \pi \frac{r_2 - 3r_1}{2}.$$

Since our substitution takes  $ds dt$  to  $-dr_1 dr_2$ , and as

$$(4.1b) \quad \omega_3 = \prod_{j=1}^3 \frac{\Gamma(j/2)}{j(2\pi i)\pi^{j/2}} = \frac{\Gamma(1/2)\Gamma(1)\Gamma(3/2)}{6(2\pi i)^3\pi^3} = \frac{i}{96\pi^5}$$

(cf. [Te2], p. 88, again), we get

$$g(I) = \frac{1}{2^4 3^3 \pi^3} \int_{r_1, r_2 \in \mathbb{R}} \widehat{g} \left( \frac{1}{2} + ir_1, \frac{1}{2} + ir_2 \right) \\ \cdot r_2 (r_2^2 - 9r_1^2) \tanh \pi r_2 \tanh \pi \frac{r_2 + 3r_1}{2} \tanh \pi \frac{r_2 - 3r_1}{2} dr_1 dr_2.$$

This proves Proposition 4.1.

### §5. The identity term.

The first term that contributes to the right-hand side of the trace formula, for a given test function  $g$ , is the identity term

$$\text{vol}(\Gamma \backslash H^3) g(I)$$

(cf. Proposition 3.1). For  $g = g_T$  having the Helgason transform given in (3.1), we will prove

**Proposition 5.1.**

$$g_T(I) = \frac{T^{-5/2}}{(4\pi)^{5/2}} + O(T^{-3/2})$$

as  $T \rightarrow 0^+$ .

*Proof.* Recalling Proposition 4.1, we have

$$g_T(I) = \frac{1}{2^4 3^3 \pi^3} \int_{r_1, r_2 \in \mathbb{R}} \widehat{g}_T \left( \frac{1}{2} + ir_1, \frac{1}{2} + ir_2 \right) r_2 (r_2^2 - 9r_1^2) \\ \cdot \tanh \pi r_2 \tanh \pi \frac{r_2 + 3r_1}{2} \tanh \pi \frac{r_2 - 3r_1}{2} dr_1 dr_2 \\ = \frac{1}{2^4 3^3 \pi^3} \int_{r_1, r_2 \in \mathbb{R}} e^{-T(1+3r_1^2+r_2^2)/3} r_2 (r_2^2 - 9r_1^2) \\ \tanh \pi r_2 \tanh \pi \frac{r_2 + 3r_1}{2} \tanh \pi \frac{r_2 - 3r_1}{2} dr_1 dr_2.$$

To simplify this expression, we use the trig identity

$$\tanh(a + b) \tanh(a) \tanh(b) = \tanh(a) + \tanh(b) - \tanh(a + b)$$

to obtain

$$g_T(I) = \frac{1}{2^4 3^3 \pi^3} \int_{r_1, r_2 \in \mathbb{R}} e^{-T(1+3r_1^2+r_2^2)/3} r_2 (r_2^2 - 9r_1^2) \cdot \left( \tanh \pi \frac{r_2 + 3r_1}{2} + \tanh \pi \frac{r_2 - 3r_1}{2} - \tanh \pi r_2 \right) dr_1 dr_2.$$

Into the first of the resulting three integrals we substitute  $p = (-r_1 + r_2)/2, q = (r_2 + 3r_1)/2$ ; into the second we substitute  $p = (r_1 + r_2)/2, q = (r_2 - 3r_1)/2$ ; and into the third,  $p = r_1, q = r_2$ . Each of these substitutions has Jacobian equal to one, and moreover the first and second merely negate the quantity

$$e^{-T(1+3r_1^2+r_2^2)/3} r_2 (r_2^2 - 9r_1^2)$$

(and give the new names  $p, q$  to  $r_1, r_2$  respectively). This latter phenomenon is a consequence of a general invariance of Helgason transforms, cf. the discussion following equation (3.1). At any rate, our substitutions yield

(5.1)

$$\begin{aligned} g_T(I) &= -\frac{1}{2^4 3^2 \pi^3} \int_{p, q \in \mathbb{R}} e^{-T(1+3p^2+q^2)/3} q (q^2 - 9p^2) \tanh \pi q dp dq \\ &= \frac{1}{2^2 3^2 \pi^3} \int_0^\infty \int_0^\infty e^{-T(1+3p^2+q^2)/3} q (9p^2 - q^2) \tanh \pi q dp dq. \end{aligned}$$

Now

$$(5.2) \quad \tanh \pi q = \frac{e^{\pi q} - e^{-\pi q}}{e^{\pi q} + e^{-\pi q}} = 1 - 2 \frac{e^{-2\pi q}}{1 + e^{-2\pi q}} = 1 + O(e^{-2\pi q})$$

for  $q \geq 0$ . Since

$$\begin{aligned} &\int_0^\infty \int_0^\infty e^{-T(1+3p^2+q^2)/3} q (9p^2 - q^2) e^{-2\pi q} dp dq \\ &= O \left( \int_0^\infty \int_0^\infty e^{-Tp^2} q (9p^2 + q^2) e^{-2\pi q} dp dq \right) \\ &= O \left( \int_0^\infty e^{-Tp^2} p^2 dp + \int_0^\infty e^{-Tp^2} dp \right) \\ &= O(T^{-3/2} + T^{-1/2}) = O(T^{-3/2}) \end{aligned}$$

as  $T \rightarrow 0^+$ , we have from (5.1) and (5.2):

$$\begin{aligned}
 (5.3) \quad g_T(I) &= \frac{1}{2^2 3^2 \pi^3} \int_0^\infty \int_0^\infty e^{-T(1+3p^2+q^2)/3} q(9p^2 - q^2) dp dq + O(T^{-3/2}) \\
 &= \frac{1}{2^4 3^2 \pi^3} \int_0^\infty \int_0^\infty e^{-T(1+3p+q)/3} p^{1/2} q(9p - q) \frac{dp dq}{pq} + O(T^{-3/2})
 \end{aligned}$$

as  $T \rightarrow 0^+$ .

Making the substitution

$$p = \frac{xy}{y+1}; \quad q = \frac{3x}{y+1},$$

we get

$$g_T(I) = \frac{e^{-T/3}}{2^4 \pi^3} \int_0^\infty \int_0^\infty e^{-Tx} x^{5/2} y^{1/2} \frac{(3y-1)}{(y+1)^{5/2}} \frac{dx dy}{xy} + O(T^{-3/2}).$$

The integral in  $y$  may be evaluated via the standard beta function formula

$$\int_0^\infty y^a (y+1)^{-(a+b)} \frac{dy}{y} = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$$

( $\text{Re}(a), \text{Re}(b) > 0$ ), whereby

$$\begin{aligned}
 (5.4) \quad g_T(I) &= \frac{e^{-T/3}}{2^4 \pi^3} \left( 3 \frac{\Gamma(3/2)\Gamma(1)}{\Gamma(5/2)} - \frac{\Gamma(1/2)\Gamma(2)}{\Gamma(5/2)} \right) \\
 &\quad \cdot \int_0^\infty e^{-Tx} x^{5/2} \frac{dx}{x} + O(T^{-3/2}) \\
 &= \frac{e^{-T/3}}{2^4 \pi^3} \left( 3 \frac{\sqrt{\pi}/2}{\Gamma(5/2)} - \frac{\sqrt{\pi}}{\Gamma(5/2)} \right) \frac{\Gamma(5/2)}{T^{5/2}} + O(T^{-3/2}) \\
 &= \frac{T^{-5/2} e^{-T/3}}{(4\pi)^{5/2}} + O(T^{-3/2}).
 \end{aligned}$$

Since  $T^{-5/2} e^{-T/3} = T^{-5/2} + O(T^{-3/2})$  as  $T \rightarrow 0^+$ , Proposition 5.1 clearly follows. □

### §6. The hyperbolic term.

We now wish to consider the contribution from hyperbolic conjugacy classes to the orbital side of the trace formula. In particular we wish to prove, for  $g = g_T$  as above (and all notation as in §3):

**Proposition 6.1.**

$$\sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) |\varepsilon_1^2 \varepsilon_2|}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} \widetilde{g}_T(\text{diag}(|\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|)) = O(T^{-1})$$

as  $T \rightarrow 0^+$ .

*Proof.* Let us write, for brevity,  $E_\varepsilon = \text{diag}(|\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|)$ . By equation (2.5),

$$\begin{aligned} \widetilde{g}_T(E_\varepsilon) &= \frac{1}{(2\pi i)^2} \int_{\text{Re}(v)=c_2} \int_{\text{Re}(u)=c_1} \widehat{g}_T\left(\frac{u+v}{6}, \frac{u-v}{2}\right) |\varepsilon_1|^{-u} |\varepsilon_2|^{-v} du dv \\ &= \frac{6}{(2\pi i)^2} \int_{\text{Re}(t)=(c_1-c_2)/2} \int_{\text{Re}(s)=(c_1+c_2)/6} \widehat{g}_T(s, t) |\varepsilon_1|^{-(3s+t)} |\varepsilon_2|^{-(3s-t)} ds dt \end{aligned}$$

for suitable  $c_1, c_2$ . Now equation (3.1) for  $\widehat{g}_T$  clearly allows us to take  $c_1 = 2, c_2 = 1$ . Then

$$\begin{aligned} \widetilde{g}_T(E_\varepsilon) &= \frac{6}{(2\pi)^2} \int_{\mathbb{R}^2} \widehat{g}_T\left(\frac{1}{2} + ir_1, \frac{1}{2} + ir_2\right) \\ &\quad \cdot |\varepsilon_1|^{-(2+i(3r_1+r_2))} |\varepsilon_2|^{-(1+i(3r_1-r_2))} dr_1 dr_2 \\ &= \frac{6 |\varepsilon_1^{-2} \varepsilon_2^{-1}| e^{-T/3}}{(2\pi)^2} \int_{\mathbb{R}} e^{-Tr_1^2} |\varepsilon_1^3 \varepsilon_2^3|^{-ir_1} dr_1 \int_{\mathbb{R}} e^{-Tr_2^2/3} |\varepsilon_1/\varepsilon_2|^{-ir_2} dr_2. \end{aligned}$$

Using the fact that

$$\int_{\mathbb{R}} e^{-\alpha r^2} \varepsilon^{-ir} dr = \sqrt{\frac{\pi}{\alpha}} e^{-(\log^2 \varepsilon)/4\alpha}$$

for  $\varepsilon, \alpha > 0$  (this identity amounts to the statement that  $f(x) = e^{-\pi x^2}$  is invariant under the Fourier transform), we find that

$$\begin{aligned} \widetilde{g}_T(E_\varepsilon) &= \frac{6 |\varepsilon_1^{-2} \varepsilon_2^{-1}| e^{-T/3}}{(2\pi)^2} \left( \sqrt{\frac{\pi}{T}} e^{-(\log^2 |\varepsilon_1^3 \varepsilon_2^3|)/4T} \right) \left( \sqrt{\frac{3\pi}{T}} e^{-3(\log^2 |\varepsilon_1/\varepsilon_2|)/4T} \right) \\ &= \frac{3\sqrt{3} |\varepsilon_1^{-2} \varepsilon_2^{-1}| e^{-T/3}}{2\pi T} e^{-(9 \log^2 |\varepsilon_1| + 3 \log^2 |\varepsilon_1 \varepsilon_2^2|)/4T} \\ &= O(|\varepsilon_1^{-2} \varepsilon_2^{-1}| T^{-1} e^{-(9 \log^2 |\varepsilon_1|)/4T}). \end{aligned}$$

The implied constant is absolute and holds for all  $T > 0$ .

Thus we find that

$$\begin{aligned} (6.1) \quad \sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) |\varepsilon_1^2 \varepsilon_2|}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} \widetilde{g}_T(\text{diag}(|\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|)) \\ = O\left(T^{-1} \sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) e^{-(9 \log^2 |\varepsilon_1|)/4T}}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|}\right). \end{aligned}$$

To estimate the sum on the right-hand side of (6.1), we need the following

**Lemma 6.1.** *For some positive constant  $b$ ,*

$$\frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1])}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} = O(\varepsilon_1^b).$$

*Proof of Lemma.* Let  $d = [(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)]^2$  be the absolute value of the discriminant of  $\mathbb{Z}[\varepsilon_1]$ . It follows from [Bo-Sh], Lemma 3, p. 121, as well as Lemmas 2 and 3, p. 127, that

$$\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) = O(d^{b_1})$$

for some positive number  $b_1$ . Therefore

$$\begin{aligned} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1])}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} \\ = O(|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|^{2b_1-1}) = O(\varepsilon_1^b), \end{aligned}$$

for  $b = 6b_1 - 3$  (we have used the fact that  $|\varepsilon_1| > |\varepsilon_2| > |\varepsilon_3|$ ). Thus our lemma is proved.  $\square$

Now let

$$\pi(x) = \left\{ \begin{array}{l} \text{distinct sets } (\varepsilon_1, \varepsilon_2, \varepsilon_3) \text{ of eigenvalues} \\ \text{of hyperbolic matrices : } |\varepsilon_3| < |\varepsilon_2| < |\varepsilon_1| \leq x \end{array} \right\}.$$

Then, since  $\pi(x) = 0$  for  $x \leq 1$ , equation (6.1) becomes

$$\begin{aligned} \sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) |\varepsilon_1^2 \varepsilon_2|}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} \widetilde{g}_T(\text{diag}(|\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|)) \\ = O\left(T^{-1} \sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \varepsilon_1^b e^{-(9 \log^2 |\varepsilon_1|)/4T}\right) \\ = O\left(T^{-1} \int_1^\infty x^b e^{-(9 \log^2 x)/4T} d\pi(x)\right). \end{aligned}$$

But  $e^{-(9 \log^2 x)/4T} = O(x^{-b-4})$  for  $x > 1$  and for  $T$  bounded above, so

$$\begin{aligned} (6.2) \quad \sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) |\varepsilon_1^2 \varepsilon_2|}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} \widetilde{g}_T(\text{diag}(|\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|)) \\ = O\left(T^{-1} \int_1^\infty x^{-4} d\pi(x)\right), \end{aligned}$$

for  $T$  bounded above.

To complete our proof of Proposition 6.1, we need only show the above integral converges. We do this via

**Lemma 6.2.**  $\pi(x) = O(x^3)$  for  $x \geq 1$ .

*Proof.* Consider the map

$$(\varepsilon_1, \varepsilon_2, \varepsilon_3) \rightarrow \lambda^3 - (\varepsilon_1 + \varepsilon_2 + \varepsilon_3)\lambda^2 + (\varepsilon_1\varepsilon_2 + \varepsilon_2\varepsilon_3 + \varepsilon_1\varepsilon_3)\lambda - 1.$$

This map is an injection from the set  $\{(\varepsilon_1, \varepsilon_2, \varepsilon_3)\}$  figuring in the definition of  $\pi(x)$ , into the set of cubic monic polynomials  $\lambda^3 - v\lambda^2 + w\lambda - 1 \in \mathbb{Z}[x]$  with

$$|v| \leq |\varepsilon_1| + |\varepsilon_2| + |\varepsilon_3| < 3x; \quad |w| \leq |\varepsilon_1\varepsilon_2| + |\varepsilon_2\varepsilon_3| + |\varepsilon_1\varepsilon_3| < 3x^2.$$

This set of polynomials clearly has cardinality  $O(x^3)$ , whence the lemma. □

We now integrate the right-hand side of equation (6.2) by parts. By Lemma 6.2, and since  $\pi(1) = 0$ , we find

$$\begin{aligned} \sum_{(\varepsilon_1, \varepsilon_2, \varepsilon_3)} \frac{\text{Reg}(\mathbb{Z}[\varepsilon_1]) \text{Cl}(\mathbb{Z}[\varepsilon_1]) |\varepsilon_1^2 \varepsilon_2|}{|(\varepsilon_1 - \varepsilon_2)(\varepsilon_1 - \varepsilon_3)(\varepsilon_2 - \varepsilon_3)|} \widetilde{g}_T(\text{diag}(|\varepsilon_1|, |\varepsilon_2|, |\varepsilon_3|)) \\ = O\left(T^{-1} \int_1^\infty x^{-5} \pi(x) dx\right) = O\left(T^{-1} \int_1^\infty x^{-2} dx\right) = O(T^{-1}) \end{aligned}$$

as  $T \rightarrow 0^+$ . Thus Proposition 6.1 is proved. □

### §7. The loxodromic and parabolic terms.

In this section we complete our study of the orbital side of the trace formula. The calculations in this section will be brief, as the arguments regarding the loxodromic term are quite similar to those of the previous section, and the parabolic terms may be evaluated readily.

We begin with the loxodromic term

$$\sum_{(r, \theta)} \frac{|\ln r_0| \text{Cl}(\mathbb{Z}[r])}{|1 - 2r^{-3} \cos \theta + r^{-6}|} \int_{\text{Re}(s)=1/2} \int_{\text{Re}(t)=1/2} \widehat{g}_T(s, t) \frac{r^{1-s} e^{-2\theta \text{Im}(t)}}{1 + e^{-2\pi \text{Im}(t)}} ds dt$$

(cf. Proposition 3.1). By pairing each summand arising from a conjugacy class  $\{g\}$  in the sum with the one arising from  $\{g^{-1}\}$  (and noting that  $r$  is a

unit in  $\mathbb{Z}[r]$ , we may write this term as

$$\sum_{\substack{(r,\theta) \\ r>1}} \frac{|\ln r_0| \operatorname{Cl}(\mathbb{Z}[r]) (1+r^{-6})}{|1-2r^{-3}\cos\theta+r^{-6}|} \cdot \int_{\operatorname{Re}(s)=1/2} \int_{\operatorname{Re}(t)=1/2} \widehat{g}_T(s,t) \frac{(r^{1-s}+r^{s-1})e^{-2\theta\operatorname{Im}(t)}}{1+e^{-2\pi\operatorname{Im}(t)}} ds dt.$$

The integral in  $t$  above is readily shown to be  $O(T^{-1/2})$ . The integral in  $s$  may be evaluated by techniques very similar to those employed in §6. We may thus show that

$$\int_{\operatorname{Re}(s)=1/2} \int_{\operatorname{Re}(t)=1/2} \widehat{g}_T(s,t) \frac{(r^{1-s}+r^{s-1})e^{-2\theta\operatorname{Im}(t)}}{1+e^{-2\pi\operatorname{Im}(t)}} ds dt = O(T^{-1}r^{1/2}e^{-(\log^2 r)/4T}).$$

Next, as in Lemma 6.1, and using the fact that  $|\ln r_0| \leq |\ln r|$ , we may show that  $|\ln r_0| \operatorname{Cl}(\mathbb{Z}[r]) = O(r^c)$  for some  $c > 0$ . We therefore find that, for  $T$  bounded above,

$$\begin{aligned} &\sum_{(r,\theta)} \frac{|\ln r_0| \operatorname{Cl}(\mathbb{Z}[r])}{|1-2r^{-3}\cos\theta+r^{-6}|} \int_{\operatorname{Re}(s)=1/2} \int_{\operatorname{Re}(t)=1/2} \widehat{g}_T(s,t) \frac{r^{1-s}e^{-2\theta\operatorname{Im}(t)}}{1+e^{-2\pi\operatorname{Im}(t)}} ds dt \\ &= O\left(T^{-1} \sum_{\substack{(r,\theta) \\ r>1}} r^{c+1/2} e^{-(\log^2 r)/4T}\right) \\ &= O\left(T^{-1} \sum_{\substack{(r,\theta) \\ r>1}} r^{-4}\right) = O\left(T^{-1} \int_1^\infty x^{-4} d\sigma(x)\right) \end{aligned}$$

where

$$\sigma(x) = \left\{ \begin{array}{l} \text{distinct pairs } (r, \theta) \text{ corresponding} \\ \text{to loxodromic matrices : } 1 < r \leq x, 0 < \theta \leq \pi \end{array} \right\}.$$

To complete our study of the loxodromic term, we need to consider the quantity  $\sigma(x)$ . But minor modifications applied to Lemma 6.2 tell us that  $\sigma(x) = O(x^3)$ , whence

(7.1)

$$\begin{aligned} &\sum_{(r,\theta)} \frac{|\ln r_0| \operatorname{Cl}(\mathbb{Z}[r])}{|1-2r^{-3}\cos\theta+r^{-6}|} \cdot \int_{\operatorname{Re}(s)=1/2} \int_{\operatorname{Re}(t)=1/2} \widehat{g}_T(s,t) \frac{r^{1-s}e^{-2\theta\operatorname{Im}(t)}}{1+e^{-2\pi\operatorname{Im}(t)}} ds dt \\ &= O\left(T^{-1} \int_1^\infty x^{-2} dx\right) = O(T^{-1}), \end{aligned}$$

as  $T \rightarrow 0^+$ .

We now need only evaluate the parabolic terms in Proposition 3.1. It follows immediately from equation (3.1) that

$$(7.2) \quad K_1 \widehat{g}\left(\frac{1}{2}, \frac{1}{2}\right) + K_2 \widehat{g}\left(\frac{1}{3}, 1\right) + K_3 \widehat{g}\left(\frac{2}{3}, 1\right) \\ = K_1 e^{-T/3} + (K_2 + K_3) e^{-2T/9},$$

and we are done.

### §8. Proof of Theorem 1.

In this section, we put together previous results to obtain an asymptotic expression for  $L_N(T)$ . A Tauberian theorem is finally used to derive from this the desired formula for  $N(x)$ .

We begin by combining Proposition 3.1, for  $g = g_T$  given by (3.1), with Propositions 5.1, 6.1; equations (7.1) and (7.2). We then have

$$(8.1) \quad L_N(T) \equiv \sum_{n=0}^{\infty} e^{-\lambda_n T} = \frac{\text{vol}(\Gamma \backslash H^3)}{(4\pi)^{5/2}} T^{-5/2} + O(T^{-3/2} + T^{-1} + 1) \\ \sim \frac{\text{vol}(\Gamma \backslash H^3)}{(4\pi)^{5/2}} T^{-5/2}$$

as  $T \rightarrow 0^+$ .

We now note that

$$L_N(T) = \int_0^{\infty} e^{-xT} dN(x),$$

the Laplace transform of  $N(x)$ . Then a standard Tauberian theorem (see Widder, [Wi]) yields

$$N(x) \sim \frac{\text{vol}(\Gamma \backslash H^3)}{(4\pi)^{5/2} \Gamma(7/2)} x^{5/2},$$

as  $x \rightarrow \infty$ . This completes the proof of Theorem 1.

### References

- [Bo] A. Borel, *Cohomology of arithmetic groups*, Proc. Internat. Cong. Math., I (1974), 435–442.
- [Bo–Sh] Z.I. Borevich and I.R. Shafarevich, *Number Theory*, Academic Press, 1966
- [Bu] D. Bump, *Automorphic Forms on  $GL(3, \mathbb{R})$* , Springer Lecture Notes in Mathematics #1083, 1984.

- [Ch] I. Chavel, *Eigenvalues in Riemannian Geometry*, Academic Press, 1984.
- [Do] H. Donnelly, *On the cuspidal spectrum for finite volume symmetric spaces*, J. Diff. Geom., **17** (1982), 239–253.
- [Ef] I. Efrat, *Selberg trace formulas, rigidity, and Weyl's law*, Ph. D. Thesis, New York University, 1981.
- [Hej1] D. Hejhal, *The Selberg Trace Formula for  $PSL(2, \mathbb{R})$  I*, Springer Lecture Notes in Mathematics, #548, 1976.
- [Hej2] ———, *The Selberg Trace Formula for  $PSL(2, \mathbb{R})$  II*, Springer Lecture Notes in Mathematics, #1001, 1983.
- [Hel] S. Helgason, *Differential Geometry and Symmetric Spaces*, Academic Press, 1962.
- [Hu1] J. Huntley, *Spectral multiplicity on products of hyperbolic spaces*, Proc. Amer. Math. Soc., **111** (1991), 1–12.
- [Hu2] ———, *Spectral multiplicity for  $GL(n, \mathbb{R})$* , Trans. Amer. Math. Soc., **332** (2) (1991), 875–888.
- [Hu–Te] J. Huntley and D. Tepper, *A local Weyl's law, the angular distribution and multiplicity of cusp forms on product spaces*, Trans. Amer. Math. Soc., **330** (1) (1992), 97–110.
- [Se1] A. Selberg, *Lectures on the Trace Formula*, University of Göttingen, 1954.
- [Se2] ———, *Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series*, J. Indian Math. Soc., **20** (1956), 47–87.
- [Te1] A. Terras, *Harmonic Analysis on Symmetric Spaces, I*, Springer–Verlag, 1985.
- [Te2] ———, *Harmonic Analysis on Symmetric Spaces, II*, Springer–Verlag, 1988.
- [Wa] D. Wallace, *The Selberg trace formula for  $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$* , Trans. Amer. Math. Soc., **345** (1) (1994), 1–36.
- [We] H. Weyl, *Über die asymptotische Verteilung der Eigenwerte*, Nachr. Akad. Wiss. Göttingen Math.– Phys. Kl., II, (1911), 110–117.
- [Wi] D.V. Widder, *The Laplace Transform*, Princeton University Press, 1941.

Received August 17, 1993 and revised March 31, 1994.

UNIVERSITY OF COLORADO  
BOULDER CO 80309

*E-mail address:* stade@euclid.colorado.edu

AND

DARTMOUTH COLLEGE  
HANOVER NH 03755



## PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. Beckenbach (1906-1982)

F. Wolf (1904-1989)

### EDITORS

Sun-Yung A. Chang (Managing Editor)  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

F. Michael Christ  
University of California  
Los Angeles, CA 90095-1555  
christ@math.ucla.edu

Nicholas Ercolani  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

Robert Finn  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

Steven Kerckhoff  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

Martin Scharlemann  
University of California  
Santa Barbara, CA 93106  
mgscharl@math.ucsb.edu

Gang Tian  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
tian@math.mit.edu

V. S. Varadarajan  
University of California  
Los Angeles, CA 90095-1555  
vsv@math.ucla.edu

Dan Voiculescu  
University of California  
Berkeley, CA 94720  
dvv@math.berkeley.edu

### SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
STANFORD UNIVERSITY  
UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
UNIVERSITY OF CALIFORNIA  
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA  
UNIVERSITY OF NEVADA, RENO  
UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
UNIVERSITY OF UTAH  
UNIVERSITY OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

---

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The table of contents and the abstracts of the papers in the current issue, as well as other information about the Pacific Journal of Mathematics, may be found on the Internet at <http://www.msri.org>.

---

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$245.00 a year (10 issues). Special rate: \$123.00 a year to individual members of supporting institutions. Subscriptions, back issues published within the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143 Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,  
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,  
the American Mathematical Society's TEX macro system.  
Copyright © 1995 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 173 No. 1 March 1996

---

Isometric immersions of $H_1^n$ into $H_1^{n+1}$	1
KINETSU ABE	
Rotationally symmetric hypersurfaces with prescribed mean curvature	29
MARIE-FRANÇOISE BIDAUT-VÉRON	
The covers of a Noetherian module	69
JIAN-JUN CHUAI	
On the odd primary cohomology of higher projective planes	77
MARK FOSKEY and MICHAEL DAVID SLACK	
Unit indices of some imaginary composite quadratic fields. II	93
MIKIHITO HIRABAYASHI	
Mixed automorphic vector bundles on Shimura varieties	105
MIN HO LEE	
Trace ideal criteria for Toeplitz and Hankel operators on the weighted Bergman spaces with exponential type weights	127
PENG LIN and RICHARD ROCHBERG	
On quadratic reciprocity over function fields	147
KATHY DONOVAN MERRILL and LYNNE WALLING	
$(A_2)$ -conditions and Carleson inequalities in Bergman spaces	151
TAKAHIKO NAKAZI and MASAHIRO YAMADA	
A note on a paper of E. Boasso and A. Larotonda: "A spectral theory for solvable Lie algebras of operators"	173
C. OTT	
Tensor products with anisotropic principal series representations of free groups	181
CARLO PENSAVALLE and TIM STEGER	
On Ricci deformation of a Riemannian metric on manifold with boundary	203
YING SHEN	
The Weyl quantization of Poisson $SU(2)$	223
ALBERT JEU-LIANG SHEU	
Weyl's law for $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$	241
ERIC GEORGE STADE and DOROTHY IRENE WALLACE (ANDREOLI)	
Minimal hyperspheres in two-point homogeneous spaces	263
PER TOMTER	
Subalgebras of little Lipschitz algebras	283
NIKOLAI ISAAC WEAVER	



0030-8730(1996)173:1;1-E