

*Pacific
Journal of
Mathematics*

**A MEAN VALUE INEQUALITY WITH APPLICATIONS TO
BERGMAN SPACE OPERATORS**

PATRICK ROBERT AHERN AND ZELJKO CUCKOVIC

Volume 173 No. 2

April 1996

A MEAN VALUE INEQUALITY WITH APPLICATIONS TO BERGMAN SPACE OPERATORS

PATRICK AHERN AND ŽELJKO ČUČKOVIĆ

If u is integrable over the unit disc and $u = Tu$, where T is the Berezin operator then it is known that u must be harmonic. In this paper we give examples to show that the condition $Tu \geq u$ does not imply that u is subharmonic, but we are able to show that the condition $Tu \geq u$ does imply that u must be “almost” subharmonic near the boundary in an appropriate sense. We give two versions of this “almost” subharmonicity, a “pointwise” version and a “weak-star” version. We give applications of these results to hyponormal Toeplitz operators on the Bergman space.

Introduction.

Let D be the open unit disc in the complex plane. We let $H^\infty(D)$ denote the space of bounded holomorphic functions in D and let $B(D)$ denote the Bergman space on D ; the set of holomorphic functions f on D such that

$$\int_D |f(z)|^2 dA(z) < \infty$$

where dA denotes planar Lebesgue measure on D . $B(D)$ is a closed subspace of the Hilbert space $L^2(dA)$ and so there is an orthogonal projection $P : L^2(dA) \rightarrow B(D)$. If $\varphi \in L^\infty(dA)$ we define the Toeplitz operator $T_\varphi : B(D) \rightarrow B(D)$ by $T_\varphi f = P(\varphi f)$. For each $z \in D$ we have the kernel function $k_z(\zeta) = \frac{1}{\pi(1 - \bar{z}\zeta)^2}$. For each $f \in B(D)$ we have $f(z) = \langle f, k_z \rangle$ where $\langle f, g \rangle$ denotes the inner product in $L^2(dA)$. We use the usual notation of $\|f\|_2^2 = \langle f, f \rangle$ for $f \in L^2(dA)$. Note that $\|k_z\|_2^2 = \langle k_z, k_z \rangle = k_z(z) = \frac{1}{\pi(1 - |z|^2)^2}$. For each $z \in D$ we have the biholomorphic involution $\varphi_z : D \rightarrow D$ given by $\varphi_z(\zeta) = \frac{z - \zeta}{1 - \bar{z}\zeta}$. With these involutions we can define the Berezin transform Tu of any $u \in L^1(dA)$, by

$$Tu(z) = \frac{1}{\pi} \int_D u \circ \varphi_z dA.$$

Equivalently, after a change of variables, we have

$$Tu(z) = \frac{(1 - |z|^2)^2}{\pi} \int_D \frac{u(\zeta)}{|1 - \bar{\zeta}z|^4} dA(\zeta).$$

Finally, if A is a bounded operator on a Hilbert space X , with norm $\|x\|$, we say A is hyponormal if $A^*A \geq AA^*$, or in other words, if

$$\|Ax\| \geq \|A^*x\| \text{ for all } x \in X.$$

It is a simple matter to check that if u is harmonic in D , i.e., $\Delta u(z) = \frac{\partial^2}{\partial z \partial \bar{z}} u(z) \equiv 0$, and $u \in L^1(dA)$, then $Tu(z) = u(z)$ for all $z \in D$. In [1], the converse was established, i.e., if $Tu = u$ in D then u must be harmonic. Now if u is subharmonic and in $L^1(dA)$ then it follows easily that $Tu \geq u$ in D . We start Section 1 by showing the converse of this statement to be false, i.e., we show that there exists u (indeed a large class of such u) so that $Tu \geq u$ in D but u is not subharmonic. However in Theorem 2 we show that the condition $Tu \geq u$ in D implies some sort of vestigial subharmonicity near the boundary. We show, under a rather mild integrability condition on Δu , that if $Tu \geq u$ in D then $\overline{\lim}_{z \rightarrow \zeta} \Delta u(z) \geq 0$ for all $\zeta \in \partial D$. Actually Theorem 2 gives a more precise “local” theorem. The main tool in the proof is a formula that represents $Tu - u$ as an integral of Δu times a positive kernel. This is the content of Theorem 1.

Our second result of this type says that if $Tu \geq u$ in D and if the measures $\Delta u(re^{i\theta})d\theta$ have a weak-star limit as $r \rightarrow 1$ on some interval I , then that limit is a positive measure on I . This is Theorem 3.

In the second section we give two applications of the results of the first section. In [2] H. Sadraoui showed that if $f, g \in H^\infty(D)$ and if $T_{f+\bar{g}}$ is hyponormal and if we assume that f', g' both lie in the Hardy class H^2 , then $|f'(e^{i\theta})| \geq |g'(e^{i\theta})|$ a.e. on the unit circle. Our first result says that if $f, g \in H^\infty(D)$ and $T_{f+\bar{g}}$ is hyponormal, then $\overline{\lim}_{z \rightarrow e^{i\theta}} (|f'(z)| - |g'(z)|) \geq 0$ for all $e^{i\theta}$. Our second result says that if, in addition, there is an arc I on the circle such that $f' \in H^2(I)$, (this is defined precisely in Section 2), then g' has the same property and $|f'(e^{i\theta})| \geq |g'(e^{i\theta})|$ a.e. on I . This last result can be viewed as a local version of Sadraoui’s result and it contains his theorem as a special case.

Section 1.

We begin with an example of a function u such that $Tu \geq u$ in D but u is not subharmonic. Note that $\Lambda(a) = \int_D |\varphi_a| \frac{dA}{\pi}$ is continuous and $\Lambda(0) = 2/3$ so there exists $\delta > 0$ such that $\Lambda(a) > \frac{1}{2}$ if $|a| < \delta$. Now let u be any

strictly convex function that is continuous and integrable on $[0, 1)$ such that $u(0) = u(\alpha) = 0$ for some $0 < \alpha < \frac{1}{2}$. Then we have $u(r) < 0$ for $0 < r < \alpha$ and u has a minimum at a unique point β , $0 < \beta < \alpha$. We further assume that $\beta < \delta$. We regard u as a radial function on D . We claim any such u satisfies $Tu \geq u$. First suppose $|a| \leq \beta$ then $u(a) = u(|a|) < 0$. On the other hand

$$\int |\varphi_a| \frac{dA}{\pi} \geq \frac{1}{2} > \alpha$$

so

$$0 < u \left(\int |\varphi_a| \frac{dA}{\pi} \right) \leq \int u \circ \varphi_a \frac{dA}{\pi},$$

the latter inequality is Jensen's. Hence

$$u(a) \leq \int u \circ \varphi_a \frac{dA}{\pi}$$

in this case.

If $|a| > \beta$ we have $a = \int \varphi_a \frac{dA}{\pi}$ and hence $|a| \leq \int |\varphi_a| \frac{dA}{\pi}$ and therefore

$$u(a) \leq u \left(\int |\varphi_a| \frac{dA}{\pi} \right),$$

because u is strictly increasing on $(\beta, 1)$. Another application of Jensen's inequality proves that $u(a) \leq \int u \circ \varphi_a \frac{dA}{\pi}$ in this case. Clearly u is not subharmonic since $u(0) = 0$ and

$$\frac{1}{2\pi} \int_0^{2\pi} u(re^{i\theta})d\theta = u(r) < 0$$

if $0 < r < \alpha$.

Suppose $u \in C^2(D)$ and $0 < r < 1$, then starting from one of Green's identities we obtain the familiar formula

$$(1) \quad u(0) = \frac{1}{2\pi} \int_0^{2\pi} u(re^{i\theta})d\theta + \frac{2}{\pi} \int_{|\zeta| \leq r} \Delta u(\zeta) \log \frac{|\zeta|}{r} dA(\zeta),$$

which we may rewrite as

$$(2) \quad \frac{1}{2\pi} \int_0^{2\pi} u(re^{i\theta})d\theta - u(0) = \frac{2}{\pi} \int_{|\zeta| \leq r} \Delta u(\zeta) \log \frac{r}{|\zeta|} dA(\zeta).$$

Next we multiply both sides of (2) by $2r$ and integrate on r from 0 to 1. We obtain

$$(3) \quad (Tu)(0) - u(0) = \int_{|\zeta| < 1} \Delta u(\zeta) K(\zeta) dA(\zeta),$$

where

$$(4) \quad K(\zeta) = \frac{4}{\pi} \int_{|\zeta|}^1 r \log \frac{r}{|\zeta|} dr = \frac{1}{\pi} \left[\log \frac{1}{|\zeta|^2} - (1 - |\zeta|^2) \right].$$

So far this is a purely formal calculation. To see what conditions are required on u , we look at the kernel K . We let $f(x) = \log \frac{1}{x} - (1 - x)$, then an application of Taylor's formula with remainder shows that

$$(5) \quad f(x) = \frac{1}{2t^2}(x - 1)^2 \quad \text{where } 0 < x < t < 1.$$

From this we see that $f(x) \geq 0$, $0 < x < 1$ and

$$(6) \quad f(x) \geq \frac{1}{2}(1 - x)^2 \text{ for } 0 < x < 1, \text{ and } f(x) \leq 2(1 - x)^2 \text{ for } \frac{1}{2} < x < 1.$$

So (3) holds if $u \in C^2(D)$ and if

$$\int_{|\zeta| < 1} |u(\zeta)| dA(\zeta) < \infty \quad \text{and} \quad \int_{|\zeta| < 1} |\Delta u(\zeta)|(1 - |\zeta|^2)^2 dA(\zeta) < \infty.$$

Now we wish to apply (3) not to u but to $u \circ \varphi_z$. This yields

$$Tu(z) - u(z) = \int_{|\zeta| < 1} \Delta(u \circ \varphi_z)(\zeta) K(\zeta) dA(\zeta).$$

Recalling that $\Delta(u \circ \varphi_z)(\zeta) = (\Delta u)(\varphi_z(\zeta)) |\varphi'_z(\zeta)|^2$ and making the change of variables $\omega = \varphi_z(\zeta)$ we arrive at the following

Theorem 1. *Suppose that $u \in C^2(D)$ and that*

$$\int_{|\zeta| < 1} |u(\zeta)| dA(\zeta) < \infty$$

and

$$\int_{|\zeta| < 1} |\Delta u(\zeta)|(1 - |\zeta|^2)^2 dA(\zeta) < \infty.$$

Then

$$Tu(z) - u(z) = \int_{|\zeta| < 1} \Delta u(\zeta) K(z, \zeta) dA(\zeta)$$

where

$$K(z, \zeta) = \frac{1}{\pi} \left[\log \frac{1}{|\varphi_z(\zeta)|^2} - (1 - |\varphi_z(\zeta)|^2) \right].$$

Moreover the kernel K satisfies:

$$(7) \quad K(z, \zeta) \geq \frac{1}{2\pi} \left[\frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - \bar{z}\zeta|^2} \right]^2 \quad \text{for } z, \zeta \in D$$

and

$$(8) \quad K(z, \zeta) \leq \frac{2}{\pi} \left[\frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - \bar{z}\zeta|^2} \right]^2 \quad \text{if} \\ \frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - \bar{z}\zeta|^2} < \frac{1}{2}.$$

Proof. Everything has been proved except (7) and (8) but they follow from (6) and the well-known identity

$$1 - |\varphi_z(\zeta)|^2 = \frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - \bar{z}\zeta|^2}.$$

□

The following well-known estimate is proved by a straightforward calculation that we omit.

Lemma 1. *There exists a constant $C_0 > 0$ such that*

$$\int_{|\zeta| < 1} \frac{(1 - |\zeta|^2)^2}{|1 - \bar{z}\zeta|^4} dA(\zeta) \geq C_0 \log \frac{1}{1 - |z|}.$$

Theorem 2. *Suppose that $u \in C^2(D)$,*

$$\int_{|\zeta| < 1} |u(\zeta)| dA(\zeta) < \infty, \\ \int_{|\zeta| < 1} |\Delta u(\zeta)|(1 - |\zeta|^2)^2 dA(\zeta) < \infty,$$

and that $\overline{\lim}_{z \rightarrow \zeta_0} \Delta u(z) < 0$ for some $\zeta_0 \in \partial D$. Then there exists $\delta > 0$ such that $Tu(z) < u(z)$ for all $z \in D$ such that $|z - \zeta_0| < \delta$.

Proof. For convenience we assume that $\zeta_0 = 1$. By assumption there exists $a > 0$ and $\epsilon > 0$ such that if $z \in D$ and $|z - 1| < \epsilon$, then $\Delta u(z) \leq -a$. If $D(1, \epsilon)$ denotes the set of points in D with $|z - 1| < \epsilon$ and $D(1, \epsilon)'$ the complement of $D(1, \epsilon)$ in D , then we have

$$\int_D \Delta u(\zeta) K(z, \zeta) dA(\zeta) = \int_{D(1, \epsilon)} \Delta u(\zeta) K(z, \zeta) dA(\zeta) \\ + \int_{D(1, \epsilon)'} \Delta u(\zeta) K(z, \zeta) dA(\zeta).$$

We deal with the second integral: if $|z - 1| < \epsilon/2$ and $\zeta \in D(1, \epsilon)'$, then $|1 - \bar{\zeta}z|$ is bounded away from 0 and hence

$$\frac{(1 - |z|^2)(1 - |\zeta|^2)}{|1 - \bar{\zeta}z|^2} \leq C(1 - |z|^2) < 1/2$$

if $1 - |z|^2$ is sufficiently small, and hence by (8) we have $K(z, \zeta) \leq C(1 - |z|^2)^2(1 - |\zeta|^2)^2$, so

$$\left| \int_{D(1, \epsilon)'} \Delta u(\zeta) K(z, \zeta) dA(\zeta) \right| \leq C(1 - |z|^2)^2 \int_{D(1, \epsilon)'} |\Delta u(\zeta)| (1 - |\zeta|^2)^2 dA(\zeta).$$

Note that this is $O((1 - |z|^2)^2)$. Next

$$\begin{aligned} \int_{D(1, \epsilon)} \Delta u(\zeta) K(z, \zeta) dA(\zeta) &\leq -a \int_{D(1, \epsilon)} K(z, \zeta) dA(\zeta) \\ &= -a \int_D K(z, \zeta) dA(\zeta) + a \int_{D(1, \epsilon)'} K(z, \zeta) dA(\zeta) \\ &\leq -\frac{a}{2\pi} \int_D \frac{(1 - |z|^2)^2(1 - |\zeta|^2)^2}{|1 - \bar{\zeta}z|^4} dA(\zeta) \\ &\quad + \frac{2a}{\pi} \int_{D(1, \epsilon)'} \frac{(1 - |z|^2)^2(1 - |\zeta|^2)^2}{|1 - \bar{\zeta}z|^4} dA(\zeta) \\ &\leq -C_0 a (1 - |z|^2)^2 \log \frac{1}{1 - |z|} + O((1 - |z|^2)^2). \end{aligned}$$

Here we have used (7) and (8) again as well as Lemma 1. Combining these estimates we have, for $|1 - z| < \epsilon/2$,

$$Tu(z) - u(z) \leq -C_0 a (1 - |z|^2)^2 \log \frac{1}{1 - |z|} + O((1 - |z|^2)^2),$$

which becomes negative as z approaches 1. □

The next lemma shows that the inequality $Tu \geq u$ is preserved under certain convolutions.

Lemma 2. *Suppose $u \in L^1(D)$ and $Tu \geq u$ in D . Suppose $w \geq 0$ is a bounded measurable function on the circle. Define, for $z \in D$,*

$$(9) \quad U(z) = \frac{1}{2\pi} \int_0^{2\pi} u(ze^{-it}) w(e^{it}) dt.$$

Then $U \in L^1(D)$ and

$$TU \geq U \quad \text{in } D.$$

Proof. Note that if $z = re^{i\theta}$, then

$$\begin{aligned} U(re^{i\theta}) &= \frac{1}{2\pi} \int_0^{2\pi} u(re^{i(\theta-t)}) w(e^{it}) dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} u(re^{it}) w(e^{i(\theta-t)}) dt. \end{aligned}$$

By hypothesis,

$$u(re^{it}) \leq \frac{(1-r^2)^2}{\pi} \int_0^1 \rho \int_0^{2\pi} \frac{u(\rho e^{i(t-s)})}{|1-r\rho e^{is}|^4} ds d\rho.$$

Since $w \geq 0$ we can multiply both sides of this inequality by $w(e^{i(\theta-t)})$ and integrate on t . After interchanging the order of integration we get

$$\begin{aligned} U(re^{i\theta}) &= \frac{1}{2\pi} \int_0^{2\pi} u(re^{it}) w(e^{i(\theta-t)}) dt \\ &\leq \frac{(1-r^2)^2}{\pi} \int_0^1 \rho \int_0^{2\pi} \frac{1}{|1-r\rho e^{is}|^4} \int_0^{2\pi} u(\rho e^{i(t-s)}) w(e^{i(\theta-t)}) \frac{dt}{2\pi} ds d\rho \\ &= \frac{(1-r^2)^2}{\pi} \int_0^1 \rho \int_0^{2\pi} \frac{1}{|1-r\rho e^{is}|^4} \int_0^{2\pi} u(\rho e^{it}) w(e^{i(\theta-t-s)}) \frac{dt}{2\pi} ds d\rho \\ &= \frac{(1-r^2)^2}{\pi} \int_0^1 \rho \int_0^{2\pi} \frac{1}{|1-r\rho e^{is}|^4} U(\rho e^{i(\theta-s)}) ds d\rho \\ &= (TU)(re^{i\theta}). \end{aligned}$$

□

The next theorem says that if $Tu \geq u$ and Δu has a weak* limit on some interval, that limit is non-negative.

Theorem 3. *Suppose that $u \in C^2(D) \cap L^1(D)$, and that $\int_D |\Delta u(\zeta)|(1 - |\zeta|^2)^2 dA(\zeta) < \infty$. Suppose further that $Tu \geq u$ in D and that there is a closed arc I on the boundary of the unit circle and a finite Borel measure μ on I such that for all continuous functions φ on I we have*

$$\lim_{r \rightarrow 1} \int_I \Delta u(re^{i\theta}) \varphi(e^{i\theta}) \frac{d\theta}{2\pi} = \int_I \varphi d\mu,$$

then $\mu \geq 0$ on $\overset{\circ}{I}$, the interior of I .

Proof. Let $w(e^{-it})$ be a continuous non-negative function with compact support in $\overset{\circ}{I}$, let

$$U(z) = \frac{1}{2\pi} \int_0^{2\pi} u(ze^{-it}) w(e^{it}) dt.$$

From Lemma 2 we know that $TU \geq U$ in D . Since the Laplacian commutes with rotations it follows from (9) that

$$(10) \quad \Delta U(z) = \frac{1}{2\pi} \int_0^{2\pi} (\Delta u)(ze^{-it})w(e^{it})dt,$$

and hence that

$$\int_D |\Delta U(\zeta)|(1 - |\zeta|^2)^2 dA(\zeta) < \infty.$$

It follows from Theorem 2 that there exists $r_k \rightarrow 1$ and $\theta_k \rightarrow 0$ such that $\lim_{k \rightarrow \infty} \Delta U(r_k e^{i\theta_k}) \geq 0$. Now it follows from (10) that

$$\Delta U(r_k e^{i\theta_k}) = \frac{1}{2\pi} \int_0^{2\pi} \Delta u(r_k e^{it})w(e^{i(\theta_k-t)}) dt.$$

Notice that for all k sufficiently large $w(e^{i(\theta_k-t)})$ will have its support in $\overset{\circ}{I}$. We have

$$\begin{aligned} & \int_I w(e^{-it})d\mu(t) - \Delta U(r_k e^{i\theta_k}) \\ &= \int_I w(e^{-it})d\mu(t) - \int_I \Delta u(r_k e^{it})w(e^{-it})\frac{dt}{2\pi} \\ & \quad + \int_I \Delta u(r_k e^{it}) \left[w(e^{-it}) - w(e^{i(\theta_k-t)}) \right] \frac{dt}{2\pi}. \end{aligned}$$

The first difference above goes to 0 as $r_k \rightarrow 1$ by hypothesis. The second difference is bounded in modulus by

$$\left(\sup_k \int_I |\Delta u(r_k e^{it})|\frac{dt}{2\pi} \right) \left(\sup_t |w(e^{-it}) - w(e^{i(\theta_k-t)})| \right).$$

The first factor is bounded, by the principle of uniform boundedness and the second goes to zero as $k \rightarrow \infty$ by the uniform continuity of w . We have shown that $\int_I w(e^{-it})d\mu(t) \geq 0$ for all non-negative $w(e^{-it})$ continuous with compact support in $\overset{\circ}{I}$; the result follows. □

Section 2.

Now suppose that f and g are holomorphic in D and $f + \bar{g} = \varphi$ is bounded. We wish to calculate $\|T_\varphi F\|_2^2$ for $F \in H^\infty(D)$

$$T_\varphi F = P(f + \bar{g})F = fF + P(\bar{g}F),$$

so

$$\begin{aligned} \|T_\varphi F\|_2^2 &= \langle fF + P(\bar{g}F), fF + P(\bar{g}F) \rangle \\ &= \|fF\|_2^2 + \|P\bar{g}F\|_2^2 + \langle P\bar{g}F, fF \rangle + \langle fF, P\bar{g}F \rangle \\ &= \|fF\|_2^2 + \|P\bar{g}F\|_2^2 + \langle \bar{f}\bar{g}F, F \rangle + \langle fgF, F \rangle, \end{aligned}$$

since P is self-adjoint.

By interchanging the roles of f and g we see that

$$\|T_{\bar{\varphi}}F\|_2^2 = \|gF\|_2^2 + \langle \bar{f}\bar{g}F, F \rangle + \langle fgF, F \rangle + \|P\bar{f}F\|_2^2.$$

Hence T_φ is hyponormal if and only if

$$(9) \quad \|fF\|_2^2 + \|P\bar{g}F\|_2^2 \geq \|gF\|_2^2 + \|P\bar{f}F\|_2^2$$

for all $F \in H^\infty(D)$.

In particular (9) holds if $F = k_z$ for some $z \in D$. Now it is immediate that $\bar{g}k_z - \overline{g(z)}k_z \perp B(D)$ for any $g \in H^\infty(D)$ and hence that $P(\bar{g}k_z) = \overline{g(z)}k_z$.

Theorem 4. *Suppose that f and g are holomorphic in D , that $f + \bar{g} = \varphi$ is bounded in D and that T_φ is hyponormal, then $Tu \geq u$ in D where $u(z) = |f(z)|^2 - |g(z)|^2$.*

Proof. By the above discussion, if we let $F = k_z$ in (9) we get

$$(10) \quad \|fk_z\|_2^2 + |g(z)|^2\|k_z\|_2^2 \geq \|gk_z\|_2^2 + |f(z)|^2\|k_z\|_2^2.$$

Since $\|k_z\|_2^2 = \frac{1}{\pi(1 - |z|^2)^2}$, a minor rearrangement of (10) proves the theorem. □

Corollary. *Suppose that f and g are holomorphic in D , that $f + \bar{g} = \varphi$ is bounded in D and that T_φ is hyponormal, then $\overline{\lim}_{z \rightarrow \zeta} (|f'(z)|^2 - |g'(z)|^2) \geq 0$ for every $\zeta \in \partial D$. In particular, if f' and g' are continuous at $\zeta \in \partial D$, then $|f'(\zeta)| \geq |g'(\zeta)|$.*

Proof. The proof follows from the theorem and the simple observation that $\Delta|f|^2 = |f'|^2$ for any holomorphic f . □

Suppose that f is holomorphic in an open set of the form

$$\{re^{i\theta} : r_0 < r < 1 \text{ and } e^{i\theta} \in I\}$$

where I is some open arc on the boundary of the unit circle. We say that $f \in H^2(I)$ if

- (i) f has polynomial growth i.e., there exists $A > 0$ such that $f(re^{i\theta}) = O((1 - r)^{-A})$ for all $e^{i\theta} \in I$.
- (ii) There exists $r_k \rightarrow 1$ such that

$$\int_I |f(r_k e^{i\theta})|^2 d\theta \leq C < \infty, \quad \text{all } k.$$

The next lemma is standard. Since we know of no convenient references we indicate the proof.

Lemma 3. *Suppose $f \in H^2(I)$, then there exists $F \in L^2(I)$ such that $\lim_{r \rightarrow 1} f(re^{i\theta}) = F(e^{i\theta})$ a.e. on I and for every compact subinterval $J \subset I$*

$$\lim_{r \rightarrow 1} \int_J |f(re^{i\theta}) - F(e^{i\theta})|^2 d\theta = 0.$$

In particular, $\overline{\lim}_{r \rightarrow 1} \int_J |f(re^{i\theta})|^2 d\theta < \infty$.

Proof. Pick a compact interval L such that $J \subseteq \overset{\circ}{L} \subseteq L \subseteq I$. Let $e^{i\theta_1}, e^{i\theta_2}$ be the end points of L and choose N such that

$$\lim_{r \rightarrow 1} [(re^{i\theta} - e^{i\theta_1})(re^{i\theta} - e^{i\theta_2})]^N f(re^{i\theta}) = 0$$

if $\theta = \theta_1$ or θ_2 . This is possible by i). Let $g(z) = [(z - e^{i\theta_1})(z - e^{i\theta_2})]^N f(z)$. Let $r_0 < r_1 < 1$ and $\Delta_k = \{re^{i\theta} : r_1 \leq r \leq r_k, e^{i\theta} \in L\}$. Let $\partial\Delta_k = \Gamma_k \cup L_k$ where $L_k = \{r_k e^{i\theta} : e^{i\theta} \in L\}$. If $z \in \overset{\circ}{\Delta}_k$ we have

$$g(z) = \frac{1}{2\pi i} \int_{\Gamma_k} \frac{g(\zeta)}{\zeta - z} d\zeta + \frac{1}{2\pi i} \int_{L_k} \frac{g(\zeta)}{\zeta - z} d\zeta.$$

If we let $k \rightarrow \infty$ we get $g(z) = g_1(z) + g_2(z)$ where $g_1(z)$ is holomorphic on $\overset{\circ}{L}$ and $g_2(z)$ is the Cauchy integral of an L^2 function on the circle. It follows that the conclusions of the lemma hold for g and hence for f . □

Theorem 5. *Suppose that f and g are holomorphic in D , that $f + \bar{g} = \varphi$ is bounded in D and that T_φ is hyponormal. Suppose further that there is an open interval I such that $f' \in H^2(I)$. Then for any open subinterval $J \subseteq \bar{J} \subseteq I$ $g' \in H^2(J)$ and $|f'(e^{i\theta})| \geq |g'(e^{i\theta})|$ almost everywhere on I .*

Proof. Let $w(e^{-it})$ be a continuous function with compact support in I such that $0 \leq w \leq 1$ and $w(e^{-it}) \equiv 1$ on a neighborhood of \bar{J} , combining Theorems 2 and 3 with Lemma 2 we have the existence of $r_k \rightarrow 1$ and $\theta_k \rightarrow 0$

such that

$$\lim_{k \rightarrow \infty} \int_0^{2\pi} (|f'(r_k e^{it})|^2 - |g'(r_k e^{it})|^2) w(e^{i(\theta_k - t)}) dt \geq 0.$$

Let L be compact interval so that $\bar{J} \subseteq \overset{\circ}{L} \subseteq L \subseteq I$. As before, for large k , $w(e^{i(\theta_k - t)})$ has support in L and hence,

$$\begin{aligned} & \int_0^{2\pi} |f'(r_k e^{it})|^2 w(e^{i(\theta_k - t)}) dt \\ & \leq \int_L |f'(r_k e^{it})|^2 dt \leq C < \infty, \quad \text{by Lemma 3.} \end{aligned}$$

Also, for large k , $w(e^{i(\theta_k - t)}) \equiv 1$ on J from which it follows that

$$\underline{\lim}_{k \rightarrow \infty} \int_J |g'(r_k e^{it})|^2 dt \leq C < \infty.$$

Now since $g \in H^\infty(D)$, g' has polynomial growth and hence $g' \in H^2(J)$. It now follows that the measures $(|f'(r e^{i\theta})|^2 - |g'(r e^{i\theta})|^2) \frac{d\theta}{2\pi}$ have a weak * limit as $r \rightarrow 1$, $e^{i\theta} \in J$, and that this limit is $(|f'(e^{i\theta})|^2 - |g'(e^{i\theta})|^2) \frac{d\theta}{2\pi}$. It follows that $|f'(e^{i\theta})| \geq |g'(e^{i\theta})|$ a.e. on J , and hence on I since $J \subseteq \bar{J} \subseteq I$, was arbitrary. \square

References

- [1] P. Ahern, M. Flores and W. Rudin, *An invariant volume-mean-value property*, Journal of Functional Analysis, vol. 111, 2 (1993), 380–397.
- [2] H. Sadraoui, *Hyponormal Toeplitz operators on the Bergman space*, preprint.

Received October 12, 1993.

UNIVERSITY OF WISCONSIN-MADISON
MADISON, WI 53706
E-mail address: ahern@math.wisc.edu

AND

UNIVERSITY OF TOLEDO
TOLEDO, OH 43606
E-mail address: zcuckovi@math.utoledo.edu

Peng Lin and Richard Rochberg , Trace ideal criteria for Toeplitz and Hankel operators on the weighted Bergman spaces with exponential type weights	127
Donald E. Marshall and Arne Stray , Interpolating Blaschke products ..	491
Kathy D. Merrill and Lynne H. Walling , On quadratic reciprocity over function fields	147
Takahiko Nakazi and Masahiro Yamada , (A_2) -conditions and Carleson inequalities in Bergman spaces	151
C. Ott , A note on a paper of E. Boasso and A. Larotonda	173
Victor Patrangenaru , Classifying 3 and 4 dimensional homogeneous Riemannian manifolds by Cartan triples	511
Carlo Pensavalle and Tim Steger , Tensor products with anisotropic principal series representations of free groups	181
Ying Shen , On Ricci deformation of a Riemannian metric on manifold with boundary	203
Albert Jeu-Liang Sheu , The Weyl quantization of Poisson $SU(2)$	223
Alexandra Shlapentokh , Polynomials with a given discriminant over fields of algebraic functions of positive characteristic	533
Eric Stade and D.I. Wallace , Weyl's law for $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$	241
Christopher W. Stark , Resolutions modeled on ternary trees	557
Per Tomter , Minimal hyperspheres in two-point homogeneous spaces	263
Jun Tomiyama , Topological Full groups and structure of normalizers in transformation group C^* -algebras	571
Nik Weaver , Subalgebras of little Lipschitz algebras	283

PACIFIC JOURNAL OF MATHEMATICS

Volume 173 No. 2 April 1996

A mean value inequality with applications to Bergman space operators	295
PATRICK ROBERT AHERN and ZELJKO CUCKOVIC	
H^p -estimates of holomorphic division formulas	307
MATS ANDERSSON and HASSE CARLSSON	
Group structure and maximal division for cubic recursions with a double root	337
CHRISTIAN JEAN-CLAUDE BALLOT	
The Weil representation and Gauss sums	357
ANTONIA WILSON BLUHER	
Duality for the quantum $E(2)$ group	375
ALFONS VAN DAELE and S. L. WORONOWICZ	
Cohomology complex projective space with degree one codimension-two fixed submanifolds	387
KARL HEINZ DOVERMANN and ROBERT D. LITTLE	
On the mapping intersection problem	403
ALEXANDER DRANISHNIKOV	
From the L^1 norms of the complex heat kernels to a Hörmander multiplier theorem for sub-Laplacians on nilpotent Lie groups	413
XUAN THINH DUONG	
Isoperimetric inequalities for automorphism groups of free groups	425
ALLEN E. HATCHER and KAREN VOGTMANN	
Approximation by normal elements with finite spectra in C^* -algebras of real rank zero	443
HUAXIN LIN	
Interpolating Blaschke products	491
DONALD EDDY MARSHALL and ARNE STRAY	
Interpolating Blaschke products generate H^∞	501
JOHN BRADY GARNETT and ARTUR NICOLAU	
Classifying 3- and 4-dimensional homogeneous Riemannian manifolds by Cartan triples	511
VICTOR PATRANGENARU	
Polynomials with a given discriminant over fields of algebraic functions of positive characteristic	533
ALEXANDRA SHLAPENTOKH	
Resolutions modeled on ternary trees	557
CHRISTOPHER W. STARK	
Topological full groups and structure of normalizers in transformation group C^* -algebras	571
JUN TOMIYAMA	



0030-8730(1996)173:2;1-D