

*Pacific
Journal of
Mathematics*

**EXPLICIT SOLUTIONS FOR THE CORONA PROBLEM WITH
LIPSCHITZ DATA IN THE POLYDISC**

STEVEN KRANTZ AND SONG-YING LI

Volume 174 No. 2

June 1996

EXPLICIT SOLUTIONS FOR THE CORONA PROBLEM WITH LIPSCHITZ DATA IN THE POLYDISC

STEVEN G. KRANTZ AND SONG-YING LI

This paper contains considerations of various versions of the classical corona problem on domains in complex n -dimensional space. Although we do not solve the H^∞ corona problem, we do obtain positive results in other topologies. We also provide explicit constructions for solutions.

1. Introduction.

Let Δ be the unit disc in the complex plane, and let Δ^n be the unit polydisc in \mathbb{C}^n . We let $\mathcal{H}(\Delta^n)$ denote the space of all holomorphic functions on the polydisc, and $\mathcal{H}^p(\Delta^n)$ the holomorphic Hardy space on Δ^n (see [Rud]). For each $0 < \alpha < \infty$, we let $\Lambda_\alpha(\Delta^n)$ denote the holomorphic Zygmund spaces over Δ^n (see [KR2]). Suppose that $f_1, \dots, f_m \in \mathcal{H}^\infty(\Delta^n)$ are such that

$$(1.1) \quad 0 < \delta^2 \leq \sum_{j=1}^m |f_j(z)|^2 \leq 1, \quad z \in \Delta^n.$$

In case $n = 1$, L. Carleson [C] solved the Corona problem and proved that there exist $g_j \in \mathcal{H}^\infty(\Delta)$ such that

$$\sum_{j=1}^m f_j(z)g_j(z) \equiv 1, \quad \|g_j\|_{\mathcal{H}^\infty(\Delta)} \leq C(m, \delta).$$

The question of whether the Corona problem can be solved in several complex variables has attracted much attention (for example, see [Am], [An], [AC], [Ch], [FS1, 2], [HS], [KL], [Li], [Lin], [S], and [V1, V2], etc.). On a strongly pseudoconvex domain, there have been attempts to generalize the method of Hörmander [H] and of Wolff [KO] to higher dimensions. This entails solving a problem of the form $\bar{\partial}u = \mu$, with μ a Carleson measure. One seeks a bounded solution u . Such a bounded solution *does not always exist* when the dimension exceeds 1 (see [V1]). However it should be noted that the result of [V1] does *not* imply that the Corona problem fails in several variables—only that the $\bar{\partial}$ technique with *that particular definition of Carleson measure* fails.

The point of the present paper is to obtain favorable results for Lipschitz solutions of the Corona problem with the corona data being Lipschitz — using iteration of one variable techniques. We shall construct a Λ_α solution of the Corona problem in one variable that allows us to treat a vector-valued problem, thus allowing induction on the number of variables. We carry out this plan by constructing an explicit formula for an $\Lambda_\alpha(\Delta^n)$ solution of the Corona problem with Corona data $f_j \in \Lambda_\alpha(\Delta^n)$.

We now give a formal statement of our theorem. The construction of our solution will be given in Section 2. The proof of the theorem is completed in Section 3.

Theorem 1.1. *Let $f_1, \dots, f_m \in \Lambda_\alpha(\Delta^n)$ ($0 < \alpha < \infty$) satisfy inequality (1.1) and*

$$(1.2) \quad \sum_{j=1}^m \|f_j\|_{\Lambda_\alpha(\Delta^n)} \leq 1.$$

Then there are functions $g_1, \dots, g_m \in \Lambda_\alpha(\Delta^n)$ such that

$$(1.3) \quad \sum_{j=1}^m f_j(z)g_j(z) \equiv 1, \quad \text{and} \quad \sum_{j=1}^m \|g_j\|_{\Lambda_\alpha(\Delta^n)} \leq C(n, m)\alpha^{1-2n}\delta^{-3n}.$$

The last estimate, in terms of α^{1-2n} and a negative power of δ , gives an indication of how the problem blows up as $\alpha \rightarrow 0^+$.

We refer the reader to [KR2] for careful definitions and discussion of the Lipschitz spaces Λ_α .

The first author thanks Peter W. Jones for a helpful communication.

2. Construction of the Solution.

In this section we shall construct an explicit solution of the Corona problem in Δ^n without yet proving any regularity properties. Let $f_1, \dots, f_m \in \mathcal{H}^\infty(\Delta^n)$ satisfy (1.1). Without loss of generality, we may use a normal families argument and reduce the proof of Theorem 1.1 to the case of $f_j \in \mathcal{H}^\infty(\overline{\Delta^n})$. We define

$$\phi_j(z) = \bar{f}_j(z) \left[\sum_{k=1}^m |f_k(z)|^2 \right]^{-1}, \quad j = 1, \dots, m.$$

For $\lambda, \eta \in \Delta$, we let

$$(2.1.1) \quad K[g](\lambda) = \int_{\Delta} K(\lambda, \eta)g(\eta) dA(\eta),$$

where

$$(2.1.2) \quad K(\lambda, \eta) = \frac{1}{2\pi i} \frac{1 - |\eta|^2}{(1 - \bar{\eta}\lambda)(\eta - \lambda)}, \quad dA(\eta) = d\eta \wedge d\bar{\eta}.$$

Notice that K is essentially the Poisson-Szegö kernel for the disc. It is known that $\bar{\partial}_\lambda K[g] = g$ in the sense of distributions. In fact

$$\frac{1 - |\eta|^2}{(1 - \bar{\eta}\lambda)(\eta - \lambda)} = \frac{1}{\eta - \lambda} - \frac{\bar{\eta}}{1 - \lambda\bar{\eta}}.$$

The first kernel on the right is well known (see [KR1]) to be a solution operator for $\bar{\partial}$, up to a constant factor; and the second one is holomorphic in $\lambda \in \Delta$. By standard arguments (integration by parts), one can show that this integral operator K maps $C^\infty(\bar{\Delta})$ to $C^\infty(\bar{\Delta})$.

For convenience, when $1 \leq j \leq n$ and $g \in L^2(\Delta^n)$, we let

$$(2.2) \quad K_j[g](z_j) = K[g(z_1, \dots, z_{j-1}, \cdot, z_{j+1}, \dots, z_n)](z_j).$$

The integral acts only on the variable z_j . We set

$$(2.3) \quad u_{jk}^1(z) = K_1[\phi_k \bar{\partial}_1 \phi_j](z_1),$$

$j, k = 1, \dots, n$, where

$$\bar{\partial}_j g \equiv \frac{\partial g}{\partial \bar{z}_j}, \quad g \in C^1(\Delta^n).$$

Also set

$$(2.4) \quad g_j^0 = \phi_j, \quad g_j^1 = \phi_j - \sum_{k=1}^m f_k(u_{jk}^1 - u_{kj}^1) \in C^\infty(\bar{\Delta}^n).$$

It is obvious that $\bar{\partial}_i g_j^1(\cdot, z')$, $g_j^1(\cdot, z') \in \mathcal{H}(\Delta)$ for each fixed $z' \in \Delta^{n-1}$ and $1 \leq i \leq n$.

Suppose that $g_j^k \in C^\infty(\bar{\Delta}^n)$ is already defined so that g_j^k and $\bar{\partial}_i g_j^k$ are holomorphic in z_1, \dots, z_k for all $1 \leq i \leq n$. Inductively, we set

$$(2.5) \quad g_j^{k+1}(z) = g_j^k(z) - \sum_{\ell=1}^m f_\ell(z)(u_{j\ell}^{k+1}(z) - u_{\ell j}^{k+1}(z))$$

where

$$(2.6) \quad u_{j\ell}^{k+1}(z) = K_{k+1} \left[g_\ell^k(z^{k+1}) \bar{\partial}_{k+1} g_j^k \right] (z_{k+1}).$$

Then our definitions imply that $g_j^{k+1}(z) \in C^\infty(\overline{\Delta^n})$. Moreover, since the inductive hypothesis implies that $g_j^k(z)$ and $g_\ell^k \cdot \bar{\partial}_{k+1} g_j^k$ are holomorphic in z_1, \dots, z_k then

$$\bar{\partial}_i(u_{j\ell}^{k+1}) = 0, \quad \text{and} \quad \bar{\partial}_i g_j^k = 0, \quad i = 1, \dots, k.$$

Therefore

$$\bar{\partial}_i g_j^{k+1} = 0, \quad i = 1, \dots, k + 1.$$

This implies that g_j^{k+1} (also $\bar{\partial}_i g_j^{k+1}$) are holomorphic in z_1, \dots, z_{k+1} for all $1 \leq i \leq n$. Finally, notice that

$$\sum_{j=1}^m f_j g_j^{k+1} \equiv 1$$

for all $k = 0, \dots, n - 1$. Therefore the functions g_1^n, \dots, g_m^n form a solution to the Corona problem with data f_1, \dots, f_m . In order to prove Theorem 1.1, it suffices now to prove the following result:

Theorem 2.1. *Let $f_j, \dots, f_m \in \Lambda_\alpha(\Delta^n)$ satisfy (1.1) and (1.2). Then*

$$(2.7) \quad \|g_j^n\|_{\Lambda_\alpha(\Delta^n)} \leq C(n, m) \alpha^{1-2n} \delta^{-3n}$$

for all $0 < \alpha < \infty$. Notice that the same solution set $\{g_j^n\}$ suffices for all α .

We shall consider the proof of Theorem 2.1 in the next section.

3. The Proof of Theorem 2.1.

In this section, we shall complete the proof of Theorem 2.1. Let us start with the following well-known simple lemma.

Lemma 3.1. *Let $g \in C^k(\Delta^n), \alpha > 0$. Suppose that $[\alpha] + 1 \leq k$, and*

$$|D^k g(z)| \leq C \text{dist}(z, \partial\Delta^n)^{\alpha-k}.$$

Then $g \in \Lambda_\alpha(\Delta^n)$. Moreover,

$$\|g\|_{\Lambda_\alpha(\Delta^n)} \leq Cn\alpha^{-1}.$$

Here D^k denotes any derivatives of g of order k , and $[\alpha] + 1$ denotes the least integer which is greater than α .

The proof of the above lemma and of more general results can be found in [KR2].

Corollary 3.2. *Let $0 < \alpha < \infty$, and let $f \in \mathcal{H}(\Delta^n)$. Then $f \in \Lambda_\alpha(\Delta^n)$ if and only if*

$$|\partial_j \partial^\beta f(z)| \leq C(1 - |z_j|^2)^{\alpha-1},$$

for all $|\beta| = \beta_1 + \dots + \beta_n \leq [\alpha]$, $j = 1, \dots, n$.

For simplicity, we shall prove Theorem 2.1 only for the case $0 < \alpha < 1$ and $n = 2$. For the case $\alpha \geq 1$ and $n > 2$, the proof may be done similarly, but it is much more tedious. For this special purpose, we shall prove the following lemma:

Lemma 3.3. *Let $0 < \alpha < 1$ and let $f, g \in C^\infty(\Delta^n)$ satisfy*

$$(3.1) \quad |D_j g(z)| + |D_j f(z)| \leq C_1(1 - |z_j|^2)^{\alpha-1}, \quad j = 1, 2, \dots, n.$$

Then

$$(3.2) \quad |D_j K_j[f \bar{\partial}_j g](z)| \leq CC_1 \alpha^{-2} (1 - |z_j|^2)^{\alpha-1}.$$

If we also assume that

$$(3.3) \quad |D_k P_j[g](z)| \leq C_1(1 - |z_k|^2)^{\alpha-1}, \quad k \neq j,$$

then

$$(3.4) \quad |D_k K_j[f \bar{\partial}_j g](z)| \leq CC_1 \alpha^{-2} (1 - |z_k|^2)^{\alpha-1}$$

where $D_k = \partial_k$ or $\bar{\partial}_k$, $k = 1, \dots, n$, and P_j denotes the Bergman projection from $L^2(\Delta)$ onto the Bergman space $A^2(\Delta)$ when we restrict attention to functions of z_j .

Proof. By symmetry, it suffices to treat the cases $j = 1$ and $k = 1$, or 2 . Let us prove (3.2) first. Since $\bar{\partial}_1 K_1[f \bar{\partial}_1 g] = f \bar{\partial}_1 g$, it suffices to prove (3.2) for $D^1 = \partial_1$. For this case, without loss of generality, we may assume that f, g are function only of z_1 , in other words, we assume $n = 1$. First of all,

$$\begin{aligned} \partial_1 K_1[f \bar{\partial}_1 g](z) &= \frac{1}{2\pi i} \partial_1 \int_{\Delta} \frac{(1 - |\eta|^2) f(\eta) \bar{\partial}_1 g(\eta)}{(\eta - z_1)(1 - z_1 \bar{\eta})} d\eta \wedge d\bar{\eta} \\ &= \frac{1}{2\pi i} \partial_1 \int_{\Delta} \frac{(1 - |\eta|^2)(f(\eta) - f(z_1)) \bar{\partial}_1 g(\eta)}{(\eta - z_1)(1 - z_1 \bar{\eta})} d\eta \wedge d\bar{\eta} \\ &\quad + \frac{1}{2\pi i} \partial_1 \left[f(z_1) \int_{\Delta} \frac{(1 - |\eta|^2) \bar{\partial}_1 g(\eta)}{(\eta - z_1)(1 - z_1 \bar{\eta})} d\eta \wedge d\bar{\eta} \right] \\ &= I_1(z) + I_2(z). \end{aligned}$$

Now

$$\begin{aligned}
 -\partial_{\bar{\eta}} \frac{1 - |\eta|^2}{(\eta - z_1)(1 - z_1\bar{\eta})} &= \frac{\eta}{(\eta - z_1)(1 - z_1\bar{\eta})} - \frac{z_1(1 - |\eta|^2)}{(\eta - z_1)(1 - z_1\bar{\eta})^2} \\
 (3.5) \qquad \qquad \qquad &= \frac{1}{(1 - z_1\bar{\eta})^2}.
 \end{aligned}$$

Integrating by parts, we then have

$$I_2(z) = \partial_1[f(z)g(z)] + \partial_1 \left[f(z) \frac{1}{2\pi i} \int_{\Delta} \frac{g(\eta)}{(1 - z_1\bar{\eta})^2} d\eta \wedge d\bar{\eta} \right].$$

It is easy to see that $|I_2(z)| \leq C(1 - |z_1|^2)^{\alpha-1}$ since the assumption (3.1) and

$$\begin{aligned}
 \left| \partial_1 \left[\frac{1}{2\pi i} \int_{\Delta} \frac{g(\eta)}{(1 - z_1\bar{\eta})^2} d\eta \wedge d\bar{\eta} \right] \right| &= \left| \frac{1}{2\pi i} \int_{\Delta} \frac{2\bar{\eta}g(\eta)}{(1 - z_1\bar{\eta})^3} d\eta \wedge d\bar{\eta} \right| \\
 &\leq \left| g(z) + \frac{1}{2\pi i} \int_{\Delta} \frac{\bar{\eta}(g(\eta) - g(z_1))}{(1 - z_1\bar{\eta})^3} d\eta \wedge d\bar{\eta} \right| \\
 &\leq \|g(z)\|_{\infty} + \int_{\Delta} C_1\alpha^{-1} \frac{|z_1 - \eta|^{\alpha}}{|1 - z_1\bar{\eta}|^3} |dA(\eta)| \\
 &\leq CC_1\alpha^{-1}(1 - |z_1|^2)^{\alpha-1}.
 \end{aligned}$$

Now we consider $I_1(z)$. It is clear that

$$\begin{aligned}
 2\pi i I_1(z) &= \partial_1 \int_{\Delta} \frac{(1 - |\eta|^2)(f(\eta) - f(z_1))\bar{\partial}_1 g(\eta)}{(\eta - z_1)(1 - z_1\bar{\eta})} d\eta \wedge d\bar{\eta} \\
 &= -\partial_1 f(z) \int_{\Delta} \frac{(1 - |\eta|^2)\bar{\partial}_1 g(\eta)}{(1 - z_1\bar{\eta})(\eta - z_1)} d\eta \wedge d\bar{\eta} \\
 &\quad + \int_{\Delta} \frac{(1 - |\eta|^2)(f(\eta) - f(z_1))\bar{\partial}_1 g(\eta)}{(1 - z_1\bar{\eta})(\eta - z_1)^2} d\eta \wedge d\bar{\eta} \\
 &\quad + \int_{\Delta} \frac{(1 - |\eta|^2)\bar{\eta}(f(\eta) - f(z_1))\bar{\partial}_1 g(\eta)}{(1 - z_1\bar{\eta})^2(\eta - z_1)} d\eta \wedge d\bar{\eta} \\
 &= I_{11}(z) + I_{12}(z) + I_{13}(z).
 \end{aligned}$$

Since

$$(1 - |\eta|^2)|\bar{\partial}_1 g| \leq C(1 - |\eta|^2)^{\alpha},$$

we see that by assumption (3.1) and simple calculation

$$|I_{11}(z)| \leq C\alpha^{-1}|\partial_1 f(z)| \leq C\alpha^{-1}(1 - |z_1|^2)^{\alpha-1}.$$

By assumption (3.1), we have

$$|f(\eta) - f(z_1)| \leq CC_1\alpha^{-1}|\eta - z_1|^{\alpha},$$

we have

$$\begin{aligned} & \left| \int_{\Delta} \frac{(1 - |\eta|^2)(f(\eta) - f(z_1))\bar{\partial}_1 g(\eta)}{(1 - z_1\bar{\eta})(\eta - z_1)^2} d\eta \wedge d\bar{\eta} \right| \\ & \leq CC_1\alpha^{-1} \int_{\Delta} |\bar{\partial}_1 g(\eta)| |\eta - z_1|^{-2+\alpha} |dA(\eta)| \\ & \leq CC_1^2\alpha^{-1} \int_{\Delta} (1 - |\eta|^2)^{-1+\alpha} |1 - z_1\bar{\eta}|^{-2+\alpha} |dA(\eta)| \\ & \leq CC_1^2\alpha^{-2}(1 - |z_1|^2)^{2\alpha-1}. \end{aligned}$$

Therefore

$$|I_{12}(z)| \leq C\alpha^{-2}(1 - |z_1|^2)^{\alpha-1}.$$

Similarly, we have $|I_{13}(z)| \leq C\alpha^{-2}(1 - |z_1|^2)^{\alpha-1}$. Hence $|I_1(z)| \leq C\alpha^{-2}(1 - |z_1|^2)^{\alpha-1}$. Therefore, combining the above estimates, the proof of (3.2) is complete.

Next we prove that (3.4) holds for $D^1 = \partial_2$. Notice that

$$\partial_2 K_1[f\bar{\partial}_1 g] = K_1[\partial_2(f\bar{\partial}_1 g)] = K_1[\partial_2 f \bar{\partial}_1 g] + K_1[f\bar{\partial}_1 \partial_2 g]$$

and

$$K_1[f\bar{\partial}_1 \partial_2 g] = \frac{1}{2\pi i} \int_{\Delta} \frac{f \partial_2 g}{(1 - z_1\bar{\eta})^2} dA(\eta) + f(z) \partial_2 g(z) - K_1[\bar{\partial}_1 f \partial_2 g].$$

We shall estimate all these terms. First

$$\begin{aligned} & |K_1[\bar{\partial}_1 f \partial_2 g](z)| \\ & \leq C(1 - |z_2|^2)^{\alpha-1} \int_{\Delta} \frac{C(1 - |\eta|^2)^{-1+\alpha}}{|\eta - z_1|} dA(\eta) \\ & \leq C\alpha^{-1}(1 - |z_2|^2)^{-1+\alpha}. \end{aligned}$$

Similarly

$$|K_1[\bar{\partial}_1 g \partial_2 f](z)| \leq C(1 - |z_2|^2)^{\alpha-1}.$$

It is easy to see that

$$|f(z)| |\partial_2 g(z)| \leq CC_1(1 - |z_2|^2)^{\alpha-1}.$$

Moreover, we have

$$\begin{aligned} & \left| \frac{1}{2\pi i} \int_{\Delta} \frac{f(\eta, z') \partial_2 g}{(1 - z_1\bar{\eta})^2} dA(\eta) \right| \\ & = \left| \frac{1}{2\pi i} \int_{\Delta} \frac{(f(\eta, z') - f(z)) \partial_2 g(\eta, z')}{(1 - z_1\bar{\eta})^2} dA(\eta) + f(z) \partial_2 P_1[g] \right| \\ & \leq \int_{\Delta} C_1 |1 - z_1\bar{\eta}|^{\alpha-2} |\partial_2 g| |dA(\eta)| + C_1(1 - |z_2|^2)^{\alpha-1} \\ & \leq C\alpha^{-1} C_1(1 - |z_2|^2)^{\alpha-1}. \end{aligned}$$

Therefore (3.4) holds for $D_k = \partial_2$. Similarly, we have (3.4) holds for $D_k = \bar{\partial}_2$. Therefore, the proof of Lemma 3.3 is complete. \square

Combining Lemmas 3.1 and 3.3, to complete the proof of Theorem 2.1, it suffices to prove the following lemma:

Lemma 3.4. *If $f_j \in \Lambda_\alpha(\Delta^n)$ satisfies (1.1) and (1.2). Then*

$$|D_i P_j [g_\ell^k](z)| \leq C C_1 (1 - |z_i|^2)^{\alpha-1},$$

for all $i, j = 1, \dots, n, i \neq j, 0 \leq k \leq n - 1, \text{ and } 1 \leq \ell \leq m.$

For convenience, we shall prove Lemma 3.4 for $n = 2$. [The case $n > 2$ is similar, but more tedious.] To achieve this special goal, we first prove:

Lemma 3.5. *Suppose that $f_j \in \Lambda_\alpha(\Delta^n)$ satisfies (1.1) and (1.2). Then Lemma 3.4 holds when $k = 0$, i.e. when $g_j^0 = \phi_j$.*

Proof. By symmetry, it suffices to prove Lemma 3.5 with $j = 1$ and $i = 2$. Moreover, we need only consider the case $D_2 = \bar{\partial}_2$ since $D_2 = \partial_2$ is similar. With the notation $|f(z)|^2 = \sum_{k=1}^m |f_k(z)|^2$, we have

$$\begin{aligned} & |D_2 P_1 [\phi_j](z)| \\ & \leq \left| \frac{1}{2\pi} \int_{\Delta} \frac{\bar{\partial}_2 \phi_j(\eta, z')}{(1 - z_1 \bar{\eta})^2} dA(\eta) \right| \\ & \leq \left| \frac{1}{2\pi} \int_{\Delta} \frac{\bar{\partial}_2 \bar{f}_j |f(\eta, z')|^2 - \bar{f}_j(\eta, z') \sum_{k=1}^m f_k(\eta, z') \bar{\partial}_2 \bar{f}_k(\eta, z')}{|f(\eta, z')|^4 (1 - z_1 \bar{\eta})^2} dA(\eta) \right| \\ & \leq \left| \frac{1}{2\pi} \int_{\Delta} \frac{\bar{\partial}_2 \bar{f}_j}{|f(\eta, z')|^2 (1 - z_1 \bar{\eta})^2} dA(\eta) \right| \\ & \quad + \left| \frac{1}{2\pi} \int_{\Delta} \frac{\bar{f}_j(\eta, z') \sum_{k=1}^m f_k(\eta, z') \bar{\partial}_2 \bar{f}_k(\eta, z')}{|f(\eta, z')|^4 (1 - z_1 \bar{\eta})^2} dA(\eta) \right| \\ & = \frac{1}{2\pi} |J_1(z) + J_2(z)|. \end{aligned}$$

Now we let

$$h(z) = 1/|f(z)|^2, \quad h_{jk}(z) = \bar{f}_j(z) f_k(z) / |f(z)|^4.$$

Then

$$\begin{aligned} |J_1(z)| & \leq \left| \int_{\Delta} \frac{(h(\eta, z') - h(z)) \bar{\partial}_2 \bar{f}_j}{(1 - z_1 \bar{\eta})^2} dA(\eta) \right| + \left| 2\pi i h(z) \bar{\partial}_2 \bar{f}_j(0, z') \right| \\ & \leq C(m, n) \alpha^{-1} \delta^{-3} C_1 (1 - |z_2|^2)^{\alpha-1}, \end{aligned}$$

and

$$|J_2(z)| = \left| \sum_{k=1}^m \int_{\Delta} \frac{(h_{jk}(\eta, z') - h_{jk}(z))\bar{\partial}_2 \bar{f}_k}{(1 - z_1 \bar{\eta})^2} dA(\eta) + 2\pi h_{jk}(z)\bar{\partial}_2 \bar{f}_k(0, z') \right| \leq C(m, n)\delta^{-3}\alpha^{-1}C_1(1 - |z_2|^2)^{\alpha-1}.$$

This completes the proof of Lemma 3.5. □

Corollary 3.6. *Let $f_j \in \Lambda_{\alpha}(\Delta^n)$ satisfy (1.1) and (1.2). Then*

$$|D_i g_j^1(z)| \leq C(m, n)\alpha^{-2}\delta^{-3}(1 - |z_i|^2)^{\alpha-1}$$

for all $0 < \alpha < 1$, $1 \leq i \leq n$ and $1 \leq j \leq m$.

Proof. This follows from Lemmas 3.1, 3.2, 3.3 and 3.5.

To prove Lemma 3.4 for the case $n = 2$, we need only to prove:

$$(3.6) \quad |D_k P_j[g_\ell^1](z)| \leq C\alpha^{-3}\delta^{-5}(1 - |z_k|^2)^{\alpha-1}$$

for all $k = 1$ with $j = 2$ and $1 \leq \ell \leq m$.

Notice that

$$\begin{aligned} &|D_k P_2[g_\ell^1](z)| \\ &= \left| \frac{1}{2\pi} D_k \int_{\Delta} \frac{g_\ell^1(z_1, \eta, z'')}{(1 - z_2 \bar{\eta})^2} dA(\eta) \right| \\ &= \left| \frac{1}{2\pi} D_k \int_{\Delta} \frac{\phi_\ell(z_1, \eta, z'') - \sum_{j=1}^m f_j(z_1, \eta, z'')(u_{\ell j}^1 - u_{j \ell}^1)(z_1, \eta, z'')}{(1 - z_2 \bar{\eta})^2} dA(\eta) \right| \\ &\leq \left| D_k P_2[\phi_\ell](z) - \frac{1}{2\pi i} D_k \int_{\Delta} \frac{\sum_{j=1}^m f_j(z_1, \eta, z'')(u_{\ell j}^1 - u_{j \ell}^1)(z_1, \eta, z'')}{(1 - z_2 \bar{\eta})^2} dA(\eta) \right|. \end{aligned}$$

By Lemma 3.5, we have

$$|D_k P_2[\phi_\ell](z)| \leq C\alpha^{-3}\delta^{-4}(1 - |z_1|^2)^{\alpha-1}.$$

Thus, combining this with $f_j \in \Lambda_{\alpha}(\Delta^n)$, the estimation of $D_k P_2[g_\ell^1]$ can be reduced to prove:

$$(3.7) \quad |D_k P_2[u_{j \ell}^1](z)| \leq C(n, m)\alpha^{-3}\delta^{-4}(1 - |z_1|^2)^{\alpha-1}$$

for $k = 1$. In order to prove (3.7) for $k = 1$, we need the following lemma.

Lemma 3.7. *Let $f_j \in \Lambda_{\alpha}(\Delta^n)$ satisfy (1.1) and (1.2). Then*

$$(3.8) \quad (1 - |z_1|^2)|\partial_1 \phi_j(z_1, z_2, z'') - \partial_1 \phi_j(z_1, w_2, z'')| \leq C\delta^{-3}|z_2 - w_2|^\alpha.$$

Proof. Since

$$\begin{aligned}
 & (1 - |z_1|^2) |\partial_1 f_k(z_1, z_2, z'') - \partial_1 f_k(z_1, w_2, z'')| \\
 &= \left| (1 - |z_1|^2) \partial_1 \int_T \frac{f_k(\eta, z_2, z'') - f_k(\eta, w_2, z'')}{1 - \bar{\eta}z_1} d\sigma(\eta) \right| \\
 &\leq C |z_2 - w_2|^\alpha (1 - |z_1|^2) \int_T \frac{1}{|1 - z_1\eta|^2} d\sigma(\eta) \\
 &\leq C |z_2 - w_2|^\alpha.
 \end{aligned}$$

Now

$$\partial_1 \phi_j(z) = \bar{f}_j(z) \left(\sum_{k=1}^m |f_k(z)|^2 \right)^{-2} \sum_{k=1}^m \bar{f}_k(z) \partial_1 f_k(z).$$

Thus

$$\begin{aligned}
 & (1 - |z_1|^2) |\partial_1 \phi_j(z_1, z_2, z'') - \partial_1 \phi_j(z_1, w_2, z'')| \\
 &\leq C \delta^{-2} \sum_{k=1}^m (1 - |z_1|^2) |\partial_1 f_k(z_1, z_2, z'') - \partial_1 f_k(z_1, w_2, z'')| \\
 &\quad + (1 - |z_1|^2) \sum_{k=1}^m |\partial_1 f_k| \left| \sum_{k=1}^m \left(\bar{f}_j \bar{f}_k \left(\sum_{k=1}^m |f_k|^2 \right)^{-2} \right) (z_1, z_2, z'') \right. \\
 &\quad \left. - \left(\bar{f}_j \bar{f}_k \left(\sum_{k=1}^m |f_k|^2 \right)^{-2} \right) (z_1, w_2, z'') \right| \\
 &\leq C \delta^{-3} |z - w|^\alpha.
 \end{aligned}$$

This completes the proof of Lemma 3.7. \square

Now we are ready to prove (3.7) for $k = 1$. Since g_j^1 is holomorphic in z_1 , it suffices to prove (3.7) for $D_1 = \partial_1$. Observe that

$$\begin{aligned}
 & \left| D_1 P_2[u_j^1](z) \right| \\
 &= \left| \partial_1 P_2[K_1[\phi_\ell \bar{\partial}_1 \phi_j]](z) \right| \\
 &\leq \left| \partial_1 P_2 \left[K_1 \left[(\phi_\ell(\eta, z') - \phi_\ell(z)) \bar{\partial}_1 \phi_j(\eta, z') \right] \right] \right| + \left| \partial_1 P_2 \left[\phi_\ell K_1[\bar{\partial}_1 \phi_j] \right] (z) \right| \\
 &= I_3(z) + I_4(z).
 \end{aligned}$$

We consider $I_3(z)$ first. Now

$$\begin{aligned} I_3(z) &= \left| \partial_1 P_2 \left[K_1 \left[(\phi_\ell(\eta, z') - \phi_\ell(z)) \bar{\partial}_1 \phi_j(\eta, z') \right] \right] \right| \\ &= \left| P_2 \left[\partial_1 K_1 \left[(\phi_\ell(\eta, z') - \phi_\ell(z)) \bar{\partial}_1 \phi_j(\eta, z') \right] \right] \right| \\ &\leq \left| P_2 \left[\int_\Delta \frac{(1 - |\eta|^2)(\phi_\ell(\eta, z') - \phi_\ell(z)) \bar{\partial}_1 \phi_j(\eta, z')}{(\eta - z)^2(1 - z_1 \bar{\eta})} \right] (z) \right| \\ &\quad + \left| P_2 \left[\int_\Delta \frac{(1 - |\eta|^2) \bar{\eta}(\phi_\ell(\eta, z') - \phi_\ell(z)) \bar{\partial}_1 \phi_j(\eta, z')}{(\eta - z)(1 - z_1 \bar{\eta})^2} \right] (z) \right| \\ &\quad + \left| P_2 \left[\int_\Delta \frac{(1 - |\eta|^2) \partial_1(\phi_\ell(z)) \bar{\partial}_1 \phi_j(\eta, z')}{(\eta - z)(1 - z_1 \bar{\eta})} \right] (z) \right| \\ &= I_{31}(z) + I_{32}(z) + I_{33}(z). \end{aligned}$$

With the notation $B(z_1, \eta) = (1 - |\eta|^2)(\eta - z_1)^{-2}(1 - z_1 \bar{\eta})^{-1}$ and an application of Lemma 3.7, we have

$$\begin{aligned} &|I_{31}(z)| \\ &\leq \left| \int_\Delta B(z_1, \eta) \int_\Delta \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(z_1, \lambda, z'')) \bar{\partial}_1 \phi_j(\eta, \lambda, z'')}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\ &\leq \left| \int_\Delta B(z_1, \eta) \right. \\ &\quad \left. \int_\Delta \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(z_1, \lambda, z'')) [\bar{\partial}_1 \phi_j(\eta, \lambda, z'') - \bar{\partial}_1 \phi_j(\eta, z')]}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\ &\quad + \left| \int_\Delta B(z_1, \eta) \bar{\partial}_1 \phi_j(\eta, z') \int_\Delta \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(z_1, \lambda, z''))}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\ &\leq C \delta^{-4} \int_\Delta |\eta - z_1|^{-2+\alpha} |1 - z_1 \bar{\eta}|^{-1} \int_\Delta |1 - z_2 \lambda|^{-2+\alpha} dA(\lambda) dA(\eta) \\ &\quad + \left| \int_\Delta B(z_1, \eta) \bar{\partial}_1 \phi_j(\eta, z') \int_\Delta \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(z_1, \lambda, z''))}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\ &\leq C \alpha^{-2} \delta^{-4} (1 - |z_1|^2)^{\alpha-1} + I_{311}. \end{aligned}$$

Notice that:

$$\begin{aligned} &\int_\Delta \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(z_1, \lambda, z''))}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) \\ &= \int_\Delta \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(\eta, z_2, z'')) + \phi_\ell(z) - \phi_\ell(z_1, \lambda, z'')}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) \\ &\quad + 2\pi i [\phi_\ell(\eta, z') - \phi_\ell(z)]. \end{aligned}$$

By using integration by parts and Lemmas 3.5 and 3.7, we have

$$\begin{aligned}
 & |I_{311}(z)| \\
 & \leq \left| \int_{\Delta} \frac{\bar{\eta} \bar{\partial}_1 \phi_j(\eta, z')}{(\eta - z_1)(1 - z_1 \bar{\eta})} \int_{\Delta} \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(z_1, \lambda, z''))}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\
 & \quad + \left| \int_{\Delta} \frac{(1 - |\eta|^2) \partial_1 \bar{\partial}_1 \phi_j(\eta, z'')}{(\eta - z_1)(1 - z_1 \bar{\eta})} \int_{\Delta} \frac{(\phi_\ell(\eta, \lambda, z'') - \phi_\ell(z_1, \lambda, z''))}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\
 & \quad + \left| \int_{\Delta} \frac{(1 - |\eta|^2) \bar{\partial}_1 \phi_j(\eta, z'')}{(\eta - z_1)(1 - z_1 \bar{\eta})} \int_{\Delta} \frac{\partial_1 \phi_\ell(\eta, \lambda, z'')}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\
 & \leq C\delta^{-4} \int_{\Delta} \frac{(1 - |\eta|^2)^{\alpha-1}}{|1 - z_1 \bar{\eta}|^2} \int_{\Delta} \frac{|\lambda - z_2|^\alpha}{|1 - z_2 \bar{\lambda}|^2} dA(\eta) dA(\lambda) \\
 & \quad + C\delta^{-5} \int_{\Delta} \frac{(1 - |\eta|^2)^{2\alpha-1}}{|1 - z_1 \bar{\eta}|^2} \int_{\Delta} \frac{|\lambda - z_2|^\alpha}{|1 - z_2 \bar{\lambda}|^2} dA(\lambda) dA(\eta) \\
 & \quad + C\delta^{-2} \int_{\Delta} \frac{(1 - |\eta|^2)^\alpha}{|1 - z_1 \bar{\eta}|^2} \left| \int_{\Delta} \frac{\partial_1 \phi_j(\eta, \lambda, z'')}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\
 & \leq C\alpha^{-2} \delta^{-4} (1 - |z_1|^2)^{\alpha-1} + C\delta^{-5} \int_{\Delta} \frac{(1 - |\eta|^2)^\alpha}{|1 - z_1 \bar{\eta}|^2} |\partial_1 P_2[\phi_j](\eta, z')| dA(\eta) \\
 & \leq C\alpha^{-2} \delta^{-4} (1 - |z_1|^2)^{\alpha-1} + C\delta^{-5} \int_{\Delta} \frac{(1 - |\eta|^2)^{2\alpha-1}}{|1 - z_1 \bar{\eta}|^2} dA(\eta) \\
 & \leq C\alpha^{-2} \delta^{-4} (1 - |z_1|^2)^{\alpha-1}.
 \end{aligned}$$

Since the estimation of $I_{32}(z)$ is similar to and easier than that of $I_{31}(z)$, we therefore have

$$I_{31}(z) + I_{32}(z) \leq C\alpha^{-2} \delta^{-5} (1 - |z_1|^2)^{\alpha-1}.$$

Now we consider $I_{33}(z)$. By Lemmas 3.5 and 3.7 again, we have

$$\begin{aligned}
 I_{33}(z) & \leq \left| \int_{\Delta} \frac{(1 - |\eta|^2)}{(\eta - z_1)(1 - z_1 \bar{\eta})} \int_{\Delta} \frac{\partial_1 \phi_\ell \bar{\partial}_1 \phi_j(\eta, \lambda, z'')}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\
 & \leq \left| \int_{\Delta} \frac{(1 - |\eta|^2)}{(\eta - z_1)(1 - z_1 \bar{\eta})} \right. \\
 & \quad \left. \int_{\Delta} \frac{(\partial_1 \phi_\ell(\eta, \lambda, z'') - \partial_1 \phi_\ell(\eta, z_2, z'')) \bar{\partial}_1 \phi_j(\eta, \lambda, z'')}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right| \\
 & \quad + \left| \int_{\Delta} \frac{(1 - |\eta|^2)}{(\eta - z_1)(1 - z_1 \bar{\eta})} \partial_1 \phi_\ell(\eta, z') \int_{\Delta} \frac{\bar{\partial}_1 \phi_j(\eta, \lambda, z'')}{(1 - z_2 \bar{\lambda})^2} dA(\lambda) dA(\eta) \right|
 \end{aligned}$$

$$\begin{aligned} &\leq C\alpha^{-1}\delta^{-4} \int_{\Delta} \frac{1}{|1 - z_1\bar{\eta}|^2} (1 - |\eta|^2)^{\alpha-1} dA(\eta) \\ &\quad + C\delta^{-4} \int_{\Delta} \frac{1 - |\eta|^2}{|1 - z_1\bar{\eta}|^2} (1 - |\eta|^2)^{2\alpha-2} dA(\eta) \\ &\leq C\delta^{-4}\alpha^{-2}(1 - |z_1|^2)^{\alpha-1}. \end{aligned}$$

Therefore

$$|I_3(z)| \leq C\alpha^{-2}\delta^{-4}(1 - |z_1|^2)^{\alpha-1}.$$

Now we estimate $I_4(z)$. Observe that

$$\begin{aligned} I_4(z) &= \left| \partial_1 P_2 \left[\phi_\ell K_1 \left[\bar{\partial}_1 \phi_j \right] \right] \right| \\ &\leq \left| \partial_1 \int_{\Delta} \frac{\phi_\ell(z_1, \lambda, z'') K_1 \left[\bar{\partial}_1 \phi_j \right] (z_1, \lambda, z'')}{(1 - z_2\bar{\lambda})^2} dA(\lambda) \right| \\ &\leq \left| \int_{\Delta} \frac{\partial_1 \phi_\ell(z_1, \lambda, z'') K_1 \left[\bar{\partial}_1 \phi_j \right] (z_1, \lambda, z'')}{(1 - z_2\bar{\lambda})^2} dA(\lambda) \right| \\ &\quad + \left| \int_{\Delta} \frac{\phi_\ell(z_1, \lambda, z'') \partial_1 K_1 \left[\bar{\partial}_1 \phi_j \right] (z_1, \lambda, z'')}{(1 - z_2\bar{\lambda})^2} dA(\lambda) \right| \\ &= I_{41}(z) + I_{42}(z). \end{aligned}$$

By Lemmas 3.3 and 3.5, we have

$$\left| \partial_1 K_1 \left[\bar{\partial}_1 \phi_k \right] \right| \leq C\alpha^{-1}\delta^{-3}(1 - |z_1|^2)^{\alpha-1}.$$

Thus

$$\begin{aligned} I_{42}(z) &\leq \int_{\Delta} \frac{|\phi_\ell(z_1, \lambda, z'') - \phi_\ell(z)| \left| \partial_1 K_1 \left[\bar{\partial}_1 \phi_k \right] (z_1, \lambda, z'') \right|}{|1 - z_2\bar{\lambda}|^2} |dA(\lambda)| \\ &\quad + \left| \phi_\ell(z) \int_{\Delta} \frac{\partial_1 K_1 \left[\bar{\partial}_1 \phi_k \right] (z_1, \lambda, z'')}{(1 - z_2\bar{\lambda})^2} dA(\lambda) \right| \\ &\leq C\delta^{-5}\alpha^{-1}(1 - |z_1|^2)^{\alpha-1} \int_{\Delta} \frac{|\lambda - z_2|^\alpha}{|1 - z_2\bar{\lambda}|^2} |dA(\lambda)| + I_{421}(z) \\ &\leq C\delta^{-5}\alpha^{-2}(1 - |z_1|^2)^{\alpha-1} + I_{421}(z). \end{aligned}$$

Since

$$\begin{aligned} I_{421}(z) &\leq \left| \partial_1 P_2 \left[K_1 \left[\bar{\partial}_1 \phi_k \right] \right] \right| \\ &= \left| \partial_1 K_1 \left[P_2 \left[\bar{\partial}_1 \phi_k \right] \right] \right| \\ &= \left| \partial_1 K_1 \left[\bar{\partial}_1 P_2 \left[\phi_k \right] \right] \right|. \end{aligned}$$

By Lemma 3.5, we have

$$|\bar{\partial}_1 P_2 [\phi_k] (z)| \leq C\alpha^{-1}(1 - |z_1|^2)^{\alpha-1}.$$

Moreover, we have

$$\begin{aligned} |D_2 P_2 [\phi_k] (z)| &\leq |\phi_k| + \left| \int_{\Delta} \frac{\phi_k(z_1, \xi, z'') - \phi(z)}{(1 - z_2 \bar{\xi})^3} dA(\xi) \right| \\ &\leq C\delta^{-1} + C\delta^{-2} \int_{\Delta} \frac{|\xi - z_2|}{|1 - z_2 \bar{\xi}|^3} |dA(\xi)| \\ &\leq C\delta^{-2}\alpha^{-1}(1 - |z_2|^2)^{-1+\alpha}. \end{aligned}$$

Thus (3.1) is satisfied for $f = 1$ and $g = P_2 [\phi_k]$. Now we apply Lemma 3.3, we have

$$I_{421} \leq \left| \partial_1 K_1 [\bar{\partial}_1 P_2 [\phi_k]] \right| \leq C\alpha^{-2}\delta^{-2}(1 - |z_1|^2)^{\alpha-1}.$$

Therefore, we have

$$I_{42}(z) \leq C\alpha^{-2}\delta^{-5}(1 - |z_1|^2)^{\alpha-1}.$$

Next we estimate $I_{41}(z)$. By Lemma 3.5, we have

$$\begin{aligned} I_{41}(z) &\leq |P_2 [\partial_1 \phi_\ell \phi_j]| + C \left| P_2 \left[\partial_1 \phi_j \int_{\Delta} \left[\frac{2\phi_k(\eta, z')}{(1 - z_1 \bar{\eta})} + \frac{\phi_k(\eta, z')}{(1 - z_1 \bar{\eta})^2} \right] dA(\eta) \right] \right| \\ &\leq C\alpha^{-2}\delta^{-4}(1 - |z_1|^2)^{\alpha-1} + C \left| P_2 \left[\partial_1 \phi_j \int_{\Delta} \frac{\phi_k(\eta, z')}{(1 - z_1 \bar{\eta})^2} dA(\eta) \right] \right| \\ &\leq C\alpha^{-2}\delta^{-4}(1 - |z_1|^2)^{\alpha-1} + C |P_2 [\partial_1 \phi_j P_1 [\phi_k]]| \\ &\leq C\alpha^{-2}\delta^{-4}(1 - |z_1|^2)^{\alpha-1} + C |P_1 [\phi_k] (z) P_2 [\partial_1 \phi_j]| \\ &\quad + C |P_2 [\partial_1 \phi_j (P_1 [\phi_k]) - P_1 [\phi_k] (z_1, z_2, z'')]| \\ &\leq C\alpha^{-2}\delta^{-4}(1 - |z_1|^2)^{\alpha-1} \\ &\quad + C\alpha^{-1}\delta^{-2}(1 - |z_1|^2)^{\alpha-1} \\ &\quad \times \int_{\Delta} \frac{|P_1 [\phi_k] (z_1, \lambda, z'') - P_1 [\phi_k] (z_1, z_2, z'')|}{|1 - z_2 \bar{\lambda}|^2} |dA(\lambda)| \\ &\leq C\alpha^{-2}\delta^{-4}(1 - |z_1|^2)^{\alpha-1} \end{aligned}$$

Therefore, $I_4(z) \leq C\alpha^{-2}\delta^{-4}(1 - |z_1|^2)^{\alpha-1}$. Combining all of these estimates, the proof of (3.7) for the case $k = 1$ is complete. Therefore, the proof of Lemma 3.4 for the case $n = 2$ is complete. □

As a consequence of Lemmas 3.1, 3.2, 3.3, 3.6 and (3.7) , we have that the proof of Theorem 2.1 for the case $n = 2$ and $0 < \alpha < 1$ is complete. The other cases can be done similarly, but the details are tedious.

We note in closing that the solution to the Corona problem presented in this paper is essentially linear in nature. It is well known that solutions to the original \mathcal{H}^∞ Corona problem are perforce non-linear in nature.

References

- [Am] E. Amar, *On the corona problem*, J. of Geometric Analysis, **4** (1991), 291–305.
- [An] M. Anderson, *The \mathcal{H}^2 corona problem and $\bar{\partial}_b$ in weakly pseudoconvex domains*, Tran. A.M.S., to appear.
- [AC] M. Anderson and H. Carlsson, *\mathcal{H}^p -estimates of holomorphic division formulas*, preprint.
- [C] L. Carleson, *Interpolation by bounded analytic functions and the corona problem*, Ann. of Math., **76** (1962), 547–559.
- [Ch] S-Y. A. Chang, *Two remarks of \mathcal{H}^1 and BMO on the bidisc*, Conference on harmonic analysis in honor of A. Zygmund, Vol II, 373–393.
- [FS1] J. Fornaess and N. Sibony, *Counterexamples to the Corona and $\bar{\partial}$ Problems*, Proceedings of a Special Year at the Mittag-Leffler Institute, Princeton Lecture Notes, Princeton University Press, 1993.
- [FS2] ———, *Smooth pseudoconvex domain for which the Corona and L^p estimates for $\bar{\partial}$ fail*, preprint.
- [FES] C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math., **129** (1972), 137–193.
- [G] J. B. Garnett, *Bounded analytic functions*, Academic Press, New York, 1981.
- [H] L. Hörmander, *Estimates for (Pluri-)subharmonic functions*, Math. Scand., **20** (1967), 65–78.
- [HS] M. Hakim and N. Sibony, *Spectre de $A(\bar{\Omega})$ pour les domaines faiblement pseudoconvexes*, J. Functional Analysis, **37** (1980), 127–135.
- [J] P. Jones, *L^∞ estimates for $\bar{\partial}$ problems in a half-plane*, Acta Math., **150** (1983), 137–152.
- [Ko] P. Koosis, *Lectures on H^p Spaces*, Cambridge Univ. Press, Cambridge, UK, 1980.
- [KR1] S.G. Krantz, *Function Theory of Several Complex Variables*, 2nd. Ed., Wadsworth, 1992.
- [KR2] ———, *Lipschitz spaces, smoothness of functions, and approximation theory*, Expositiones Math., **3** (1983), 193–260.
- [KL] S.G. Krantz and S-Y. Li, *Some remarks on the Corona problem on strongly pseudoconvex Domains in \mathbb{C}^n* , Illinois J. of Math., **39** (1995), 323–349.
- [Li] S-Y. Li, *Corona Problem of Several Complex Variables*, Contemporary Mathematics, **137** (1992), 307–328.
- [Lin] K.C. Lin, *\mathcal{H}^p solutions for the corona problem on the polydisc in \mathbb{C}^n* , Bull A.M.S., **110** (1986), 69–84.
- [Rud] W. Rudin, *Function Theory in Polydiscs*, Benjamin, New York, 1969.
- [S] N. Sibony, *Problème de la Couronne pour des domaines pseudoconvexes à bord lisse*, Annals of Math., **126** (1987), 675–682.

- [V1] N.Th. Varopoulos, *BMO functions and the $\bar{\partial}$ -equation*, Pacific J. of Math., **71** (1977), 221–272.
- [V2] ———, *Probabilistic approach to some problems in complex analysis*, Bull. S. C. Math. Paris, **105** (1981), 181–224.

Received February 1, 1994 and revised August 30, 1994. The first author was supported in part by a grant from the National Science Foundation.

WASHINGTON UNIVERSITY
ST. LOUIS, MO 63130
E-mail address: sk@math.wustl.edu
songying@math.wustl.edu

| | |
|--|-----|
| Andreas Seeger , Endpoint inequalities for Bochner-Riesz multipliers in the plane | 543 |
| Ted Stanford , Braid commutators and Vassiliev invariants | 269 |
| Xiangsheng Xu , On the Cauchy problem for a singular parabolic equation | 277 |
| Xingwang Xu , On the existence of extremal metrics | 555 |
| Rugang Ye , Constant mean curvature foliation: singularity structure and curvature estimate | 569 |

PACIFIC JOURNAL OF MATHEMATICS

Volume 174 No. 2 June 1996

| | |
|--|-----|
| Quantum affine algebras and affine Hecke algebras | 295 |
| VYJAYANTHI CHARI and ANDREW PRESSLEY | |
| On the zero sets of bounded holomorphic functions in the bidisc | 327 |
| PHILIPPE CHARPENTIER and JOAQUIM ORTEGA-CERDÀ | |
| Bloch constants in one and several variables | 347 |
| IAN GRAHAM and DROR VAROLIN | |
| Characters of the centralizer algebras of mixed tensor representations of $GL(r, \mathbb{C})$ and the quantum group ${}^{\mathcal{O}}U_q(gl(r, \mathbb{C}))$ | 359 |
| TOM HALVERSON | |
| Derivations of C^* -algebras and almost Hermitian representations on Π_k -spaces | 411 |
| EDWARD KISSIN, ALEKSEI I. LOGINOV and VICTOR S. SHULMAN | |
| Twisted Alexander polynomial and Reidemeister torsion | 431 |
| TERUAKI KITANO | |
| Explicit solutions for the corona problem with Lipschitz data in the polydisc | 443 |
| STEVEN KRANTZ and SONG-YING LI | |
| Prepolar deformations and a new Lê-Iomdine formula | 459 |
| DAVID MASSEY | |
| KK -groups of twisted crossed products by groups acting on trees | 471 |
| KEVIN PAUL MCCLANAHAN | |
| Coherent states, holomorphic extensions, and highest weight representations | 497 |
| KARL-HERMANN NEEB | |
| Endpoint inequalities for Bochner-Riesz multipliers in the plane | 543 |
| ANDREAS SEEGER | |
| On the existence of extremal metrics | 555 |
| XINGWANG XU | |
| Constant mean curvature foliation: singularity structure and curvature estimate | 569 |
| RUGANG YE | |



0030-8730(1996)174:2;1-C