

*Pacific  
Journal of  
Mathematics*

**TRIANGLE SUBGROUPS OF HYPERBOLIC TETRAHEDRAL  
GROUPS**

COLIN MACLACHLAN

## TRIANGLE SUBGROUPS OF HYPERBOLIC TETRAHEDRAL GROUPS

C. MACLACHLAN

It is known that there are nine compact tetrahedra in three-dimensional hyperbolic space for which the group of isometries generated by reflections in the faces is discrete. The subgroup of orientation-perserving isometries in such a group will be called a tetrahedral Kleinian group. Exactly eight such  $\Gamma$  are arithmetic and also a complete list of the finitely many arithmetic Fuchsian triangle groups  $G$  is known. In this paper, we determine for which pairs of groups  $(G, \Gamma)$  as above, with one possible exception, one can embed  $G$  into  $\Gamma$ . We find that there are many such pairs, contrasting with the single pair  $(G, \Gamma)$  which is known to arise when, instead of arithmeticity, the condition that  $G$  be realised as the subgroup of elements of  $\Gamma$  which centralise a reflection in one of the faces of the associated tetrahedron, is imposed.

### 1. Introduction.

There are nine compact tetrahedra in hyperbolic 3-space whose dihedral angles are submultiples of  $\pi$  so that the group generated by reflections in the faces of the tetrahedron is a discrete subgroup of the isometry group,  $\text{Isom}(\mathbf{H}^3)$ . The centraliser of a reflection in one of the faces of the tetrahedron is then a discrete subgroup of the isometry group of the 2-dimensional hyperbolic plane on which that face lies. Thus if we restrict to orientation-preserving subgroups in both the ambient group and the subgroup, these tetrahedral Kleinian groups will contain Fuchsian subgroups. For only one of the tetrahedra and one of the faces of that tetrahedron, is the Fuchsian subgroup a triangle group [1]. Eight of the tetrahedral groups are arithmetic [20], and in these cases, these tetrahedral groups contain infinitely many commensurability classes of Fuchsian subgroups [12]. Here, using arithmetic techniques, it will be shown that these tetrahedral groups contain Fuchsian triangle groups in addition to the one "visible" on a face described above. For all except one of these arithmetic triangle groups complete information is obtained.

These tetrahedral groups arise in connection with various extremal problems in hyperbolic 3-manifolds and orbifolds. For example, one of them

is the arithmetic orbifold of minimal volume [3] and conjecturally the 3-dimensional orbifold of minimal volume [5]. Others arise in the investigation of hyperbolic orbifolds with elliptic elements whose axes are as close as possible [6, 7]. Indeed it was in investigating one such case, that the question addressed in this paper came to light and I am grateful to Gaven Martin for discussions on this.

## 2. Tetrahedral Groups.

The group of orientation-preserving isometries of hyperbolic 3-space  $\mathbf{H}^3$ , is isomorphic to  $\mathrm{PSL}(2, \mathbf{C})$  via the Poincaré extension. Then 3-dimensional hyperbolic orbifolds (respectively manifolds) arise as the quotient of  $\mathbf{H}^3$  by Kleinian groups which are discrete (discrete and torsion-free) subgroups  $\Gamma$  of  $\mathrm{PSL}(2, \mathbf{C})$ . Here we will only be concerned with the cases where  $\Gamma$  is cocompact i.e. the quotient space  $\mathbf{H}^3/\Gamma$  is compact.

Let  $T$  denote a compact tetrahedron in  $\mathbf{H}^3$  with vertices  $A, B, C, D$  such that the dihedral angles are submultiples of  $\pi$ . If the dihedral angles along the edges  $AB, BC, CA, CD, DA, BD$  are  $\pi/l_1, \pi/l_2, \pi/l_3, \pi/m_1, \pi/m_2, \pi/m_3$  respectively, we denote the tetrahedron  $[l_1, l_2, l_3; m_1, m_2, m_3]$ . There are nine such tetrahedra [8]. If  $T_i$  denotes the tetrahedron let  $\Gamma_i$  denote the subgroup of index two which consists of orientation-preserving isometries in the group generated by reflections in the faces of  $T_i$ . These are the tetrahedral groups and with the notation just described are listed below.

$T_1[2, 2, 3; 3, 5, 2]$	$T_2[2, 2, 3; 2, 5, 3]$	$T_3[2, 2, 4; 2, 3, 5]$
$T_4[2, 2, 5; 2, 3, 5]$	$T_5[2, 3, 3; 2, 3, 4]$	$T_6[2, 3, 4; 2, 3, 4]$
$T_7[2, 3, 3; 2, 3, 5]$	$T_8[2, 3, 4; 2, 3, 5]$	$T_9[2, 3, 5; 2, 3, 5]$

Table 1

Recall that  $\Gamma_i$  then has presentation

$$\Gamma_i = \{a, b, c \mid a^{l_1} = b^{l_2} = c^{l_3} = (bc)^{m_1} = (ca)^{m_2} = (ab)^{m_3} = 1\}.$$

Each face of the tetrahedron is a triangle on a hyperbolic plane in  $\mathbf{H}^3$ . If  $\rho$  denotes the reflection in that plane, let  $C_{\Gamma_i}(\rho)$  denote the centraliser of  $\rho$  in  $\Gamma_i$  so that  $C_{\Gamma_i}(\rho)$  will be a group of isometries of that hyperbolic plane. The subgroup  $C_{\Gamma_i}^+(\rho)$  of elements which preserve the orientation of this plane is thus a Fuchsian group. A fundamental region for  $C_{\Gamma_i}^+(\rho)$  on that hyperbolic plane can be obtained. This applies more generally to any discrete subgroup generated by reflections in the faces of a polyhedron and the structure of the Fuchsian groups so obtained in the cases of the nine tetrahedral groups has been determined (for all this see [1]).

For  $T_3$ , let  $\rho$  be the reflection in the face  $ABC$ . Then  $C_{\Gamma_3}(\rho)$  has, as fundamental region, the triangular face  $ABC$  whose face angles are  $\pi/4, \pi/5, \pi/2$  respectively. Thus  $\Gamma_3$  contains the  $(2, 4, 5)$  triangle group. Now the  $(2, 4, 5)$  triangle group is well known to contain the triangle subgroups  $(2, 5, 5), (4, 4, 5)$  as subgroups of indices 2 and 6 respectively. Thus  $\Gamma_3$  also contains these. However, this is the only tetrahedron  $T_i$  such that a subgroup of the form  $C_{\Gamma_i}^+(\rho)$  is a triangle group [1].

It should be noted that one can construct triangular prisms in  $\mathbf{H}^3$  whose dihedral angles are all submultiples of  $\pi$ , with one triangular face meeting its neighbours orthogonally and having face angles  $\pi/2, \pi/3, \pi/q$  for any  $q \geq 7$  [4, 13]. It follows that these Kleinian pentahedral groups contain the Fuchsian triangle groups  $(2, 3, q)$ .

### 3. Arithmetic Groups.

Let  $k$  be a number field with one complex place and let  $A$  be a quaternion algebra over  $k$  which is ramified at the real places. If  $\sigma : A \rightarrow M(2, \mathbf{C})$  is a representation of  $A$  and  $\mathcal{O}$  an order in  $A$ , then  $P\sigma(\mathcal{O}^1)$  is a discrete subgroup of  $\text{PSL}(2, \mathbf{C})$  of finite covolume, where  $\mathcal{O}^1$  is the group of elements in  $\mathcal{O}$  of norm 1. An *arithmetic Kleinian group* is any Kleinian group commensurable with some such  $P\sigma(\mathcal{O}^1)$ . In fact if  $\Gamma$  is an arithmetic Kleinian group, then  $\Gamma^{(2)} \subset P\sigma(\mathcal{O}^1)$  for some order  $\mathcal{O}$  where

$$\Gamma^{(2)} = \langle \gamma^2 \mid \gamma \in \Gamma \rangle.$$

Arithmetic Fuchsian groups are defined in a similar way with in this case  $k$  being a totally real number field and  $A$  ramified at all real places except one. For information on arithmetic Fuchsian and Kleinian groups see [19, 2].

For any finite volume Kleinian group  $\Gamma$  then

$$k\Gamma = \mathbf{Q}(\text{tr } \delta \mid \delta \in \Gamma^{(2)})$$

is a finite extension of the rationals and an invariant of the commensurability class of  $\Gamma$  [14], called the invariant field of  $\Gamma$ . Furthermore

$$A\Gamma = \left\{ \sum a_i \delta_i \mid a_i \in k\Gamma, \delta_i \in \Gamma^{(2)} \right\}$$

is a quaternion algebra over  $k\Gamma$  [11]. In the cases where  $\Gamma$  is arithmetic,  $k\Gamma, A\Gamma$  define the arithmetic structure [9]. This last statement is also true for Fuchsian groups [16].

For a circle or straight line  $\mathcal{C}$  in  $\mathbf{C}$ , and a Kleinian group  $\Gamma$ , define

$$\text{Stab}(\mathcal{C}, \Gamma) = \{ \gamma \in \Gamma \mid \gamma(\mathcal{C}) = \mathcal{C} \text{ and } \gamma \text{ preserves the components of } \mathbf{C} \setminus \mathcal{C} \}$$

Then  $\text{Stab}(\mathcal{C}, \Gamma)$  will be a maximal non-elementary Fuchsian subgroup if its limit set on  $\mathcal{C}$  consists of more than two points.

In the cases where  $\Gamma$  is arithmetic, these groups  $\text{Stab}(\mathcal{C}, \Gamma)$  will be arithmetic Fuchsian groups  $F$  [9]. Furthermore, if  $\Gamma$  is determined by the quaternion algebra  $A$  over the number field  $k$ , then  $F$  will be determined by a quaternion algebra  $B$  over the totally real number field  $l = k \cap \mathbf{R}$  where  $[l : k] = 2$  and  $A \cong B \otimes_l k$  [9]. This then forces the primes that ramify in  $A$  to have a certain form and also implies a relationship between these primes and the primes of  $l$  which are ramified in  $B$  [12]. For the applications, this information is gathered in the following theorem.

**Theorem 3.1.** *Let  $\Gamma$  be an arithmetic Kleinian group whose quaternion algebra  $A$  is defined over the number field  $k$ . Let  $F$  be a maximal non-elementary Fuchsian subgroup of  $\Gamma$ , which will then be arithmetic with quaternion algebra  $B$  defined over some totally real number field  $l$ . In addition*

- (1)  $k \cap \mathbf{R} = l$  and  $[k : l] = 2$ .
- (2)  $B \otimes_l k \cong A$ .
- (3)  $A$  has finite ramification at a (possibly empty) set of pairs of prime ideals  $\mathcal{Q}_i, \mathcal{Q}'_i (i = 1, 2, \dots, n)$  of  $k$  such that  $\mathcal{Q}_i \cap R_l = \mathcal{Q}'_i \cap R_l = \mathcal{P}_i$ .
- (4)  $B$  has finite ramification at the ideals  $\mathcal{P}_i (i = 1, 2, \dots, n)$  together with a (possibly empty) set of prime ideals  $\mathcal{P}$  of  $l$  such that  $\mathcal{P}$  is either inert or ramified in the extension  $k | l$ .

The above theorem gives us information on which triangle groups are candidates to be subgroups of a given tetrahedral group. We now give results aimed at showing the existence of triangle subgroups of tetrahedral groups.

In the above circumstances,  $B \otimes_l k \cong A$  and so there is an embedding  $j : B \rightarrow A$ . Now if  $\mathcal{L}$  is an order in  $B$ , then  $\mathcal{L} \otimes_{R_l} R_k$  is an order in  $B \otimes_l k$ . If  $\mathcal{L}$  is maximal in  $B$ ,  $\mathcal{L} \otimes_{R_l} R_k$  is not in general maximal in  $B \otimes_l k$ . Indeed, if  $d(\mathcal{L})$  is the discriminant of  $\mathcal{L}$  then  $d(\mathcal{L} \otimes_{R_l} R_k) = d(\mathcal{L})R_k$  [15]. In all cases, we obtain an embedding  $j : \mathcal{L} \rightarrow \mathcal{O}$  where  $\mathcal{O}$  is a maximal order in  $A$ . Thus via  $j$  we obtain embeddings of Fuchsian groups into arithmetic Kleinian groups.

**Theorem 3.2.** *If  $B \otimes_l k \cong A$ , thus yielding  $j : B \rightarrow A$  as in the above theorem, and  $\mathcal{L}$  is a maximal order in  $B$ , then  $j(\mathcal{L}^1) \subset \mathcal{O}^1$  for some maximal order in  $A$ .*

The type number of  $A$  is the number of conjugacy classes of maximal orders in  $A$  and can be determined from the arithmetic data of  $A$  [19]. For

a maximal order  $\mathcal{O}$ , define

$$N\mathcal{O} = \{x \in A^* \mid x^{-1}\mathcal{O}x = \mathcal{O}\}.$$

Then if  $\sigma$  is a representation  $\sigma : A \rightarrow M(2, \mathbf{C})$ ,  $P\sigma(\mathcal{O}^1) \subset P\sigma(N\mathcal{O})$  and  $P\sigma(N\mathcal{O})$  is a maximal Kleinian group [2]. The same applies in the case of  $B$  and Fuchsian groups.

#### 4. Arithmetic tetrahedral and triangle groups.

For all  $T_i$ ,  $i \neq 8$ , the groups  $\Gamma_i$  are arithmetic [20], and the defining fields  $k_i$  and quaternion algebras  $A_i$  have been determined [10]. This information is recorded in Table 2 below together with some additional information readily deduced from the details in [10]. Here  $\Delta_k$  denotes the discriminant of the field  $k$  and  $Ram_f$ , the set of finite primes at which the algebra  $A$  is ramified.

Group	$\Delta_k$	$Ram_f$	$[P\sigma(N\mathcal{O}) : P\sigma(\mathcal{O}^1)]$	$P\sigma(\mathcal{O}^1)$	Type Number
$\Gamma_1$	-400	$\phi$	2	$\Gamma_1$	1
$\Gamma_2$	-275	$\phi$	2	$\Gamma_2$	1
$\Gamma_3$	-400	$\phi$	2	$\Gamma_3^{(2)}$	1
$\Gamma_4$	-475	$\phi$	2	$\Gamma_4$	1
$\Gamma_5$	-448	$\phi$	2	$\Gamma_5$	1
$\Gamma_6$	-7	$\mathcal{Q}_2, \mathcal{Q}'_2$	8	$\Gamma_6^{(2)}$	1
$\Gamma_7$	-775	$\phi$	2	$\Gamma_7$	1
$\Gamma_9$	-1375	$\phi$	4	$\Gamma_9$	2

Table 2

Note that since only elements of orders 2,3,4,5 can occur in these tetrahedral groups, we need only consider the triangle groups in the table below. These are gathered together in commensurability classes and adjoined at the end of each class is the maximal triangle group in which these lie.

- a) (2, 4, 5), (2, 5, 5), (4, 4, 5); (2, 4, 5)
- b) (3, 3, 4), (4, 4, 4); (2, 3, 8)
- c) (3, 3, 5)(5, 5, 5) : (2, 3, 10)
- d) (3, 4, 4); (2, 4, 6)
- e) (3, 5, 5); (2, 5, 6)
- f) (4, 5, 5); (2, 5, 8)
- g) (3, 4, 5); (3, 4, 5)

Table 3

Precisely which triangle groups are arithmetic has been determined [17]. Recalling that arithmetic Fuchsian groups are commensurable if and only

if their quaternion algebras are isomorphic, the commensurability classes of these triangle groups were obtained by classifying their quaternion algebras [18]. From that information we note that the group  $(3, 4, 5)$  is not arithmetic and also that the defining field of the group  $(4, 5, 5)$  is  $Q(\sqrt{5}, \sqrt{2})$ . Thus by Theorem 3.1, neither of these can be subgroups of any of the arithmetic tetrahedral groups. For the remainder the detailed information is given below. In this Table, we let  $F_\alpha$ ,  $\alpha = a, b, c, d, e$  denote the maximal triangle group in the class and label the corresponding field  $k_\alpha$  and defining algebra  $B_\alpha$ .

	$F_\alpha$	$F_\alpha^{(2)}$	$k_\alpha$	Finite Ramification
<i>a</i>	$(2, 4, 5)$	$(2, 5, 5)$	$Q(\sqrt{5})$	$\mathcal{P}_2$
<i>b</i>	$(2, 3, 8)$	$(3, 3, 4)$	$Q(\sqrt{2})$	$\mathcal{P}_2$
<i>c</i>	$(2, 3, 10)$	$(3, 3, 5)$	$Q(\sqrt{5})$	$\mathcal{P}_5$
<i>d</i>	$(2, 4, 6)$	$(2, 2, 3, 3)$	$Q$	$\mathcal{P}_2, \mathcal{P}_3$
<i>e</i>	$(2, 5, 6)$	$(3, 5, 5)$	$Q(\sqrt{5})$	$\mathcal{P}_3$

Table 4

It should be noted in each of these cases that  $F_\alpha^{(2)} = P\mu(\mathcal{L}^1)$  where  $\mu$  is a representation of the quaternion algebra  $B_\alpha$  and  $\mathcal{L}$  is a maximal order of  $B_\alpha$ .

## 5. Triangle Subgroups of Tetrahedral Groups.

In this section, the arithmetic tetrahedral groups  $\Gamma_i$   $i \neq 8$  are examined in turn to decide which triangle groups they contain.

Consider first the information obtained in Section 2, noting that  $\Gamma_1 = \Gamma_3^{(2)}$  is a subgroup of index 2 in  $\Gamma_3$ . For the tetrahedral groups, this is the only pair that are commensurable [10]. From Section 2, it follows that  $F_a \subset \Gamma_3$  and so  $F_a^{(2)} \subset \Gamma_3^{(2)}$ . Thus  $(2, 5, 5)$  is a subgroup of  $\Gamma_1$  but, since  $\Gamma_1$  has no elements of order 4, the other two triangle groups in the commensurability class are not.

From Table 2,  $A_1 (= A_3)$  has type number 1 so that for a maximal order  $\mathcal{O}$  in  $A_1$  we can assume, up to conjugacy, and again using Table 2, that  $\Gamma_3^{(2)} = P\sigma(\mathcal{O}^1)$ . So  $\Gamma_3$  and  $P\sigma(N\mathcal{O})$  lie in the normaliser of  $P\sigma(\mathcal{O}^1)$  which is discrete of finite covolume. Since  $P\sigma(N\mathcal{O})$  is a maximal Kleinian group, it follows that  $\Gamma_3 = P\sigma(N\mathcal{O})$ . Now  $k_1 \cap \mathbf{R} = \mathbf{Q}(\sqrt{5})$  so we consider  $F_c, F_e$  by Theorem 3.1. Using Kummer's theorem applied to the extension  $k_1 \mid \mathbf{Q}(\sqrt{5})$ , both  $\mathcal{P}_3, \mathcal{P}_5$  are inert in this extension. Then by the comments following Table 4, and the fact the  $A_1$  has type number one,

$$F_c^{(2)} = P\sigma(j\mathcal{L}^1) \subset P\sigma(\mathcal{O}^1) = \Gamma_1$$

by Theorem 3.2. In exactly the same way  $F_e^{(2)} \subset \Gamma_1$ .

The cases  $\Gamma_i, i = 2, 4, 5, 7$  follow in a very similar way using the data from Tables 2 and 4. The results are given in Theorem 5.1 below.

The group  $\Gamma_6$  will now be considered. Note that  $A_6$  is ramified at  $\mathcal{Q}_2, \mathcal{Q}'_2$  and that  $\mathcal{P}_3$  is inert in  $\mathbf{Q}(\sqrt{-7}) \mid \mathbf{Q}$ . Thus as before  $F_d^{(2)}$  embeds in  $\Gamma_6^{(2)}$ . Additional information on  $A_6$  is required. Recall from [2] that, in this case, the norm mapping defines an epimorphism  $n : N\mathcal{O} \rightarrow R_f^*$  which induces an isomorphism

$$(1) \quad \frac{P\sigma(N\mathcal{O})}{P\sigma(\mathcal{O}^1)} \cong \frac{R_f^*}{R_f^{*2}}$$

where  $R_f^*$  is the group of units in the ring consisting of those elements of  $k_6$  which are integral at all finite places not in the ramification set of  $A_6$ . The quotient group  $Q$  in (1) is isomorphic to  $\mathbf{Z}_2 \oplus \mathbf{Z}_2 \oplus \mathbf{Z}_2$  and is generated by the images of  $x_1 = (1 + \sqrt{-7})/2, x_2 = (1 - \sqrt{-7})/2, x_3 = -1$ . Now  $\Gamma_6$  is a subgroup of  $P\sigma(N\mathcal{O})$  of index 4, containing  $P\sigma(\mathcal{O}^1)$  and so corresponds to one of the subgroups of order 2 in  $Q$ . Now  $\Gamma_6$  contains elements of order 4. Since  $\sqrt{2} \notin k_6$ , there are no elements of order 4 in  $P\sigma(\mathcal{O}^1)$ . Let  $\alpha \in N(\mathcal{O})$  be such that  $P\sigma(\alpha)$  has order 4. Let  $n, t$  denote the norm and trace of  $\alpha$ . Then  $n = xy^2$ , where  $x$  is a product of the  $x_i$ s and  $y \in R_f^*$ . Since  $P\sigma(\alpha)$  has order 4,  $t^2 = 2n$ . Thus  $2x \in k_6^{*2}$  and so  $x = x_1x_2 = 2$ . Thus the subgroup of order 2 in  $Q$  generated by the image of  $x_1x_2$  gives the unique subgroup of  $P\sigma(N\mathcal{O})$  containing  $P\sigma(\mathcal{O}^1)$  which contains elements of order 4. This unique subgroup must be  $\Gamma_6$ . Furthermore, if we choose in  $A_6$  where

$$A_6 = \left( \frac{-1, -1}{\mathbf{Q}(\sqrt{-7})} \right)$$

the maximal order  $\mathcal{O} = R_{k_6}[1, i, j, 1/2(1 + i + j + ij)]$  then  $\alpha = 1 + i$  lies in  $N\mathcal{O}$  and  $n(\alpha) = 2$ . Thus  $\Gamma_6 = \langle P\sigma(\mathcal{O}^1), P\sigma(\alpha) \rangle$ . Furthermore, if we choose  $\beta = i + j + ij$  so that  $n(\beta) = 3$  then  $\Omega = \mathcal{O} \cap \beta\mathcal{O}\beta^{-1}$  is an Eichler order of level  $\mathcal{Q}_3$  in  $A_6$ . Up to conjugacy there is only one class of Eichler orders of this level in  $A_6$  [19] and one can check directly that  $\alpha \in N\Omega$ .

Now consider the group  $F_d$ . Recall the  $F_d^{(2)} = P\sigma(j(\mathcal{L}^1))$  has signature  $(2, 2, 3, 3)$ . Also  $F_d = P\sigma(j(N\mathcal{L}))$  and the group  $(3, 4, 4)$  is its unique subgroup of index 2 which contains elements of order 4. The order  $\mathcal{L} \otimes_{\mathbf{Z}} R_{k_6}$  has discriminant  $\mathcal{Q}_2\mathcal{Q}'_2\mathcal{Q}_3$ , which since it is square-free, is an Eichler order of level  $\mathcal{Q}_3$ . Thus we can assume that  $\mathcal{L} \otimes_{\mathbf{Z}} R_{k_6} = \Omega$  and hence the image of one of the elements of order 4 in the  $(3, 4, 4)$  group will coincide, after conjugacy, with  $P\sigma(\alpha)$ . Thus  $(3, 4, 4)$  is a subgroup of  $\Gamma_6$ .

Finally, consider the group  $\Gamma_9$ . From Table 2, note that the type number of  $A_9$  is 2 and that  $\Gamma_9 = P\sigma(\mathcal{O}_1)$  for some maximal order  $\mathcal{O}_1$  in  $A_9$ . But in this case there is a maximal order  $\mathcal{O}_2$  in  $A_9$  which is not conjugate to  $\mathcal{O}_1$  in



$A_9$ . Now  $P\sigma(\mathcal{O}_2^1)$  cannot be isomorphic to  $\Gamma_9$ , otherwise by Mostow rigidity,  $P\sigma(\mathcal{O}_2^1)$  would be conjugate to  $P\sigma(\mathcal{O}_1^1)$  in  $\text{Isom}(\mathbf{H}^3)$ . But that implies that  $\mathcal{O}_1^1$  and  $\mathcal{O}_2^1$  are conjugate in  $A_9^*$ . Let  $t\mathcal{O}_1^1t^{-1} = \mathcal{O}_2^1$  and set  $\mathcal{M} = t\mathcal{O}_1t^{-1}$ , so that  $\mathcal{M}$  is a maximal order. Let  $\mathcal{E} = \mathcal{M} \cap \mathcal{O}_2$  so that  $\mathcal{E}^1 = \mathcal{O}_2^1$ . If  $\mathcal{M} \neq \mathcal{O}_2$ , then  $\mathcal{E}$  is an Eichler order, and for some prime  $\mathcal{P} \in k_9$ ,  $[((\mathcal{O}_2)_{\mathcal{P}})^1 : (\mathcal{E}_{\mathcal{P}})^1] > 1$ . Since the index  $[\mathcal{O}_2^1 : \mathcal{E}^1]$  is a product of local indices, this contradiction shows that  $\mathcal{M} = \mathcal{O}_2$ .

Let  $P\sigma(\mathcal{O}_2^1) = \Gamma'_9$ . Now  $k_9 = \mathbf{Q}(\sqrt{5})(\theta)$  where  $\theta$  satisfies  $x^2 - x + (9 + 4\sqrt{5}) = 0$  and is such that  $\{1, \theta\}$  is a relative integral basis of  $k_9 \mid \mathbf{Q}(\sqrt{5})$  [10]. Now investigating the candidates  $F_a, F_c, F_e$  we find that  $\mathcal{P}_2$  splits in the extension,  $\mathcal{P}_5$  ramifies and  $\mathcal{P}_3$  is inert. Thus we obtain that  $F_c^{(2)}, F_e^{(2)}$  are contained in either  $\Gamma_9$  or  $\Gamma'_9$ . In fact, by choosing a basis, one can prove directly that if  $\mathcal{L}$  is a maximal order in  $B_c$  then the order  $\mathcal{L} \otimes_{R_{\mathbf{Q}(\sqrt{5})}} R_{k_9}$ , whose discriminant is  $\mathcal{Q}_5^2$ , lies inside an  $\Omega$  whose discriminant is  $\mathcal{Q}_5$ . But then  $\Omega$  is an Eichler order of level  $\mathcal{Q}_5$  and there is only one class of Eichler orders of level  $\mathcal{Q}_5$  in  $A_9$ , since  $\mathcal{Q}_5$  is generated by the element  $1/2(\sqrt{5} - \sqrt{5 - 4\sqrt{5}})$  which is positive at one real ramified place and negative at the other [19]. It follows that the groups  $(3, 3, 5), (5, 5, 5)$  lie in both  $\Gamma_9, \Gamma'_9$ . A similar argument does not work for  $\mathcal{Q}_3$  as it is generated by 3 and thus we have a less precise statement.

**Theorem 5.1.** *The tetrahedral groups  $\Gamma_i, i = 1, 2 \dots, 7$  contain precisely the triangle groups given in Table 5 below. The group  $\Gamma_9$  contains the groups  $(3, 3, 5), (5, 5, 5)$  and  $\Gamma_9$  or  $\Gamma'_9$  contains  $(3, 5, 5)$ .*

Tetrahedral Group	Triangle Subgroups
$\Gamma_1$	$(2, 5, 5), (3, 3, 5), (5, 5, 5), (3, 5, 5)$
$\Gamma_2$	$(2, 5, 5), (3, 3, 5), (5, 5, 5)$
$\Gamma_3$	$(2, 4, 5), (2, 5, 5), (4, 4, 5), (3, 3, 5), (5, 5, 5), (3, 5, 5)$
$\Gamma_4$	$(2, 5, 5), (3, 5, 5)$
$\Gamma_5$	$(3, 3, 4), (4, 4, 4)$
$\Gamma_6$	$(3, 4, 4)$
$\Gamma_7$	$(3, 3, 5), (5, 5, 5), (3, 5, 5)$

**Table 5.**

**References**

[1] T. Baskan and A.M. Macbeath, *Centralisers of reflections in crystallographic groups*, Math. Proc. Cam. Phil. Soc., **92** (1982), 79-91.

- [2] A. Borel, *Commensurability classes and volumes of hyperbolic 3-manifolds*, Ann. Sc. Norm. Sup. Pisa, **8** (1981), 1-33.
- [3] T. Chinburg and E. Friedman, *The smallest arithmetic hyperbolic three - orbifold*, Invent. Math., **86** (1986), 507-527.
- [4] M.D.E. Conder and G. Martin, *Cusps, triangle group and hyperbolic 3-folds*, J. Austral. Math. Soc., **55** (1993), 149-182.
- [5] F. Gehring and G. Martin, *On the minimal volume hyperbolic 3-orbifold*, Math. Research Letters, **1** (1994), 107-114.
- [6] F. Gehring, G. Martin and A.W. Reid, *Arithmeticity, discreteness and volume*, to appear.
- [7] F. Gehring, C. Maclachlan, G. Martin and A.W. Reid, *Simple elliptic axes in a Kleinian group*, to appear.
- [8] F. Lanner, *On complexes with transitive groups of automorphisms*, Comm. Sem. Math. Univ. Lund, 1950.
- [9] C. Maclachlan and A.W. Reid, *Commensurability classes of arithmetic Kleinian groups and their Fuchsian subgroups*, Math. Proc. Cam. Phil. Soc., **102** (1987), 251-257.
- [10] ———, *The arithmetic structure of tetrahedral groups of hyperbolic isometries*, Mathematika, **36** (1989), 221-240.
- [11] W. Neumann and A.W. Reid, *Arithmetic of hyperbolic manifolds*, Topology '90 Ed. B. Apanasov et al de Gruyter (1992), 273-310.
- [12] A.W. Reid, *Totally geodesic surfaces in hyperbolic 3-manifolds*, Proc. Edin. Math. Soc., **34** (1991), 77-88.
- [13] ———, *Arithmetic Kleinian groups and their Fuchsian subgroups*, Ph. D. Thesis, Aberdeen University, 1987.
- [14] ———, *A note on trace fields of Kleinian groups*, Bull. London Math. Soc., **22** (1990), 349-352.
- [15] I. Reiner, *Maximal Orders*, Academic Press, 1975.
- [16] K. Takeuchi, *A characterisation of arithmetic Fuchsian groups*, J. Math. Soc. Japan, **27** (1975), 600-612.
- [17] ———, *Arithmetic triangle groups*, J. Math. Soc. Japan, **29** (1977), 91-106.
- [18] ———, *Commensurability classes of arithmetic triangle groups*, J. Fac. Sci. Univ. Tokyo, Sect. 1A, Math., **24** (1977), 201-212.
- [19] M.-F. Vignéras, *Arithmétique des algèbres de quaternions*, Lecture notes in Maths, **800**, Springer-Verlag, 1980.
- [20] E.B. Vinberg, *Discrete groups generated by reflections in Lobachevskii space*, Math. Sb., **114** (1967), 429-444.

Received July 13, 1994 and revised June 8, 1995.

ABERDEEN UNIVERSITY  
ABERDEEN, SCOTLAND



## PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by

E. F. Beckenbach (1906-1982)

F. Wolf (1904-1989)

### EDITORS

Sun-Yung A. Chang (Managing Editor)  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

F. Michael Christ  
University of California  
Los Angeles, CA 90095-1555  
christ@math.ucla.edu

Nicholas Ercolani  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

Robert Finn  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

Steven Kerckhoff  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

Martin Scharlemann  
University of California  
Santa Barbara, CA 93106  
mgscharl@math.ucsb.edu

Gang Tian  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
tian@math.mit.edu

V. S. Varadarajan  
University of California  
Los Angeles, CA 90095-1555  
vsv@math.ucla.edu

Dan Voiculescu  
University of California  
Berkeley, CA 94720  
dvv@math.berkeley.edu

### SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIF. INST. OF TECHNOLOGY  
CHINESE UNIV. OF HONG KONG  
HONG KONG UNIV. OF SCI. & TECH.  
KEIO UNIVERSITY  
MACQUARIE UNIVERSITY  
MATH. SCI. RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.  
PEKING UNIVERSITY  
RITSUMEIKAN UNIVERSITY  
STANFORD UNIVERSITY

TOKYO INSTITUTE OF TECHNOLOGY  
UNIVERSIDAD DE LOS ANDES  
UNIV. OF ARIZONA  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIF., BERKELEY  
UNIV. OF CALIF., DAVIS  
UNIV. OF CALIF., IRVINE  
UNIV. OF CALIF., LOS ANGELES  
UNIV. OF CALIF., RIVERSIDE  
UNIV. OF CALIF., SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF HAWAII  
UNIV. OF MELBOURNE  
UNIV. OF MONTANA  
UNIV. NACIONAL AUTONOMA DE MEXICO  
UNIV. OF NEVADA, RENO  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

---

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The table of contents and the abstracts of the papers in the current issue, as well as other information about the Pacific Journal of Mathematics, may be found on the Internet at <http://www.math.uci.edu/pjm.html>.

---

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$245.00 a year (10 issues). Special rate: \$123.00 a year to individual members of supporting institutions.

Subscriptions, back issues published within the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,  
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,  
the American Mathematical Society's TEX macro system.  
Copyright © 1995 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 176    No. 1    November 1996

---

Moduli spaces of isometric pluriharmonic immersions of Kähler manifolds into indefinite Euclidean spaces	1
HITOSHI FURUHATA	
On a theorem of Koch	15
FARSHID HAJIR	
Degree-one maps onto lens spaces	19
CLAUDE HAYAT-LEGRAND, SHICHENG WANG and HEINER ZIESCHANG	
Unitary representation induced from maximal parabolic subgroups for split $F_4$	33
CHENG CHON HU	
New constructions of models for link invariants	71
FRANÇOIS JAEGER	
Solvability of Dirichlet problems for semilinear elliptic equations on certain domains	117
ZHIREN JIN	
Hadamard-Frankel type theorems for manifolds with partially positive curvature	129
KATSUEI KENMOTSU and CHANGYU XIA	
Boundary behavior of the Bergman curvature in strictly pseudoconvex polyhedral domains	141
KANG-TAE KIM and JIYE YU	
Existence and behavior of the radial limits of a bounded capillary surface at a corner	165
KIRK LANCASTER and DAVID SIEGEL	
Triangle subgroups of hyperbolic tetrahedral groups	195
COLIN MACLACHLAN	
Chern classes of vector bundles on arithmetic varieties	205
TOHRU NAKASHIMA and YUICHIRO TAKEDA	
Haar measure on $E_q(2)$	217
ARUP KUMAR PAL	
Domains of partial attraction in noncommutative probability	235
VITTORINO PATA	
Partitioning products of $\mathcal{P}(\omega)/\text{fin}$	249
OTMAR SPINAS	
Dimensions of nilpotent algebras over fields of prime characteristic	263
CORA M. STACK	
Tensor products of structures with interpolation	267
FRIEDRICH WEHRUNG	
Fourier multipliers for $L_p(\mathbb{R}^n)$ via $q$ -variation	287
QUANHUA XU	