

*Pacific
Journal of
Mathematics*

TVERBERG PARTITIONS AND BORSUK–ULAM
THEOREMS

K.S. SARKARIA

TVERBERG PARTITIONS AND BORSUK–ULAM THEOREMS

K.S. SARKARIA

An N -dimensional real representation E of a finite group G is said to have the “Borsuk–Ulam Property” if any continuous G -map from the $(N + 1)$ -fold join of G (an N -complex equipped with the diagonal G -action) to E has a zero. This happens iff the “Van Kampen characteristic class” of E is nonzero, so using standard computations one can explicitly characterize representations having the B-U property. As an application we obtain the “continuous” Tverberg theorem for all prime powers q , i.e., that some q disjoint faces of a $(q - 1)(d + 1)$ -dimensional simplex must intersect under any continuous map from it into affine d -space. The “classical” Tverberg, which makes the same assertion for all linear maps, but for all q , is explained in our set-up by the fact that any representation E has the analogously defined “linear B-U property” iff it does not contain the trivial representation.

1. Introduction.

This paper is essentially an analysis of a method which I had used in a manuscript [19] circulated in 1988-89. Some of its results have in the meantime been independently obtained by others, and it is possible that the newer methods of [21] might lead to better results. Nevertheless, it does give a complete account of one aspect of “*the method of deleted joins*”: it delineates clearly its power and limitations, as far as the two topics mentioned in the title are concerned, *if one uses only finite groups*, as against [21], where we use a continuous group action.

In 1966 Tverberg [24] showed that any cardinality $(q - 1)(d + 1) + 1$ subset of a real affine d -dimensional space can be partitioned into q disjoint subsets whose convex hulls have a nonempty intersection; a much easier proof is given in [20]. There is a “continuous” analogue which asks more: given any continuous map f from a $(q - 1)(d + 1)$ -simplex into d -space, can one always find q disjoint faces $\sigma_1, \dots, \sigma_q$ of this simplex such that $f(\sigma_1) \cap \dots \cap f(\sigma_q)$ is nonempty? For q prime this was established by Bárány-Shlosman-Szücs [4]. In [18] I gave an easy proof of this result using a deleted \mathbb{Z}/q -join of the N -simplex, $N = (q - 1)(d + 1)$, viz. the $(N + 1)$ -fold join $E_N(\mathbb{Z}/q) = \mathbb{Z}/q \dots \mathbb{Z}/q$.

In [19] I attempted to generalize this argument to all q by using, in addition, the “Van Kampen obstruction” class e .

The importance of this characteristic class $e(\mathbb{E}) \in H^N(G, \widehat{\mathbb{Z}})$, $n = \dim(\mathbb{E})$, which we define in 2.6 for any real representation \mathbb{E} of any finite group G , stems from the fact — see Theorem 1, 2.6.2 — that it is nonzero iff \mathbb{E} has the **Borsuk-Ulam property**, i.e., any continuous G -map $E_N(G) \rightarrow \mathbb{E}$ has a zero. Using the argument of [18], the “continuous” Tverberg holds if one has an order q group G for which $\mathbb{L}^\perp(G)$, the $(d + 1)$ -fold direct sum of the non-trivial part of the regular representation, has this B-U property. Our Theorem 2, 2.6.3 gives a complete characterization of complex \mathbb{Z}/q -representations having the B-U property. In particular, it shows that the representations $\mathbb{L}^\perp(\mathbb{Z}/q)$ all have this property iff q is prime, which gives of course the B-S-S theorem, and shows also that to go beyond one needs to look at finite non-cyclic groups. Amusingly, the original Tverberg theorem also fits neatly into this B-U framework: we check that the argument of [20] or [10] is really just the same, except that one now invokes a *linear analogue 2.4 of the B-U property* which holds for all q . The next Theorem 3, 2.8.1 generalizes the “continuous” Tverberg to all *prime powers* $q = p^k$ and has also been proved independently by Ozaydin [16] and Volovikov [25]. It follows at once from Theorem 4, 2.8.2 which says that a representation of $(\mathbb{Z}/p)^k$ has the B-U property iff it does not contain the trivial representation. Finally in 2.9, we embed the \mathbb{Z}/q -action of $\mathbb{L}^\perp(\mathbb{Z}/q)$ in an action of the symmetric group Σ_q , and show — see Theorem 5, 2.9.3 — that the characteristic class of this Σ_q representation is zero iff q is not a prime power. To go beyond prime powers it thus seems necessary to use continuous group actions.

The exposition below is self-contained except that we refer to the literature for standard facts regarding Chern classes of finite group actions. For more background material see also Mark de Longueville’s notes [13] of a seminar based on this paper.

2. Borsuk-Ulam representations.

The main character of our story is a real N -dimensional group representation \mathbb{E} which does not contain the trivial representation, mostly $\mathbb{E} = \mathbb{L}^\perp$ (defined in 2.2 below) which has dimension $N = (q - 1)(d + 1)$.

2.1. . By the q -th *deleted join* [17] $K * \dots * K$ of a simplicial complex K one understands the subcomplex of its q -fold join $K \cdot \dots \cdot K$ consisting of all simplices $(\sigma_1, \dots, \sigma_q)$ with $\sigma_i \cap \sigma_j = \emptyset \forall i \neq j$. Mostly $K = [N] =$ all faces of the N -simplex $\{e_1, \dots, e_{N+1}\}$. Let Q be a cardinality q set. Denoting the q copies of each e_α by ge_α , $g \in Q$, $[N] \cdot \dots \cdot [N]$ consists of all subsets of the cardinality $q(N + 1)$ set $\{ge_\alpha : g \in Q, 1 \leq \alpha \leq N + 1\}$, and $[N] * \dots * [N]$ of all faces of all N -simplices of the type $\{g_1e_1, \dots, g_{N+1}e_{N+1}\}$. So $[N] * \dots * [N]$ (q times) identifies with $E_N(Q) = Q \cdot \dots \cdot Q$ ($N + 1$ times).

Frequently we'll equip the set Q with a group structure G , and then let G act simplicially on $[N] \cdot \dots \cdot [N]$ by $h \bullet (ge_\alpha) = (hg)e_\alpha$. Note that this action preserves, and is free on, the subcomplex $[N] * \dots * [N]$. We recall that such free G -complexes $E_N(G) = G \cdot \dots \cdot G$ ($N + 1$ times), $EG = \cup_N E_N(G)$, go into Milnor's definition [14] of a *classifying space* BG of $G : BG = EG/G = \cup_N (B_N G)$, where $B_N G = E_N(G)/G$.

2.2. . We'll identify our affine d -space \mathbb{A}^d with the hyperplane $\sum_k x_k = 1$ of \mathbb{R}^{d+1} , and the q -fold product $\mathbb{R}^{d+1} \times \dots \times \mathbb{R}^{d+1}$ with the vector space \mathbb{L} of all real $(d + 1) \times q$ matrices, with \mathbb{L}^\perp denoting the $(q - 1)(d + 1)$ dimensional subspace consisting of all matrices having row sums zero. Note that \mathbb{L}^\perp is the orthogonal complement of the diagonal subspace Δ of matrices having all columns equal to each other.

We'll index the columns of our matrices by the cardinality q set Q . Frequently Q will be equipped with a group structure G , and then we'll permute the columns by left translations. The resulting representations of G will be denoted $\mathbb{L}(G)$ and $\mathbb{L}^\perp(G)$. Note that $\mathbb{L}(G) = \mathbb{R}^{d+1}[G]$, the $(d + 1)$ -fold direct sum of the *regular representation* $\mathbb{R}[G]$ provided by each row, and that $\mathbb{L}^\perp(G)$ contains no trivial representation. So the action of G on the unit sphere $S(\mathbb{L}^\perp)$ is always without fixed points. When $d + 1$ is even we'll identify $\mathbb{L}(G)$ with the representation $\mathbb{C}^{(d+1)/2}[G]$ provided by all $\frac{d+1}{2} \times q$ complex matrices by taking real and imaginary parts of each row, and we'll equip $\mathbb{L}(G)$ with the orientation prescribed by this complex structure.

For the case $G = \mathbb{Z}/q$ note that the action is free on $S(\mathbb{L}^\perp)$ iff q is prime, and that the action preserves the orientation of $\mathbb{L}^\perp(\mathbb{Z}/q)$ iff $(q - 1)(d + 1)$ is even.

2.3. Proof of theorems of Tverberg and Bárány-Shlosman-Szücs.

Let $s_\alpha, 1 \leq \alpha \leq N + 1, N = (q - 1)(d + 1)$, be the points of the given set $S \subset \mathbb{A}^d$ and consider the linear map $K = [N] \xrightarrow{f} \mathbb{A}^d$ such that $e_\alpha \mapsto s_\alpha \forall \alpha$. More generally consider any continuous map $[N] \xrightarrow{f} \mathbb{A}^d$. We want to show that there exist q disjoint faces $\sigma_1 \dots, \sigma_q$ of K such that $f(\sigma_1) \cap \dots \cap f(\sigma_q) \neq \emptyset$. Equivalently, if we compose the q -fold join $K * \dots * K \rightarrow \mathbb{A}^d \cdot \dots \cdot \mathbb{A}^d \subset \mathbb{R}^{d+1} \times \dots \times \mathbb{R}^{d+1} = \mathbb{L}$ of f with the orthogonal projection $\mathbb{L} \rightarrow \mathbb{L}^\perp$ to get a map

$$s : [N] * \dots * [N] \rightarrow \mathbb{L}^\perp,$$

then what we have to show is that $0 \in \text{Im}(s)$.

For this, note first that s commutes with the group actions, defined above. Now the linear case follows by applying the "linear Borsuk-Ulam" theorem 2.4. Likewise, for q prime, we see that the \mathbb{Z}/q -map s associated to a continuous f must have a zero, by using the generalization 2.5 of the usual continuous Borsuk-Ulam. □

2.4. “Linear Borsuk-Ulam”. *If \mathbb{E} does not contain the trivial representation, then any linear G -map $s : E_N(G) \rightarrow \mathbb{E}$ has a zero.*

Note that the condition on \mathbb{E} is obviously necessary.

Proof. This is a particular case of Bárány [3] the argument being as follows. If $\text{conv} \langle s(g_\alpha e_\alpha) : g_\alpha \in G, 1 \leq \alpha \leq N + 1 \rangle$ is at a distance $\delta > 0$ from $0 \in \mathbb{E}$, then its nearest point P is contained in the hyperplane H normal to $0P$ and out of the points $s(g_\alpha e_\alpha)$ we can choose $\leq N$ which all lie on H and are such that P is in their convex hull. The remaining points will be either on H or in the component of $\mathbb{E} \setminus H$ not containing $\{0\}$. Let $s(g_\beta e_\beta)$ be any of these points. Since s commutes with the G actions, and \mathbb{E} does not contain the trivial representation, we have $\sum_g s(g e_\beta) = \sum_g g(s e_\beta) = 0$. So some $s(g e_\beta)$ must be in the component of $\mathbb{E} \setminus H$ which contains $\{0\}$. Replacing g_β by such a g we can make δ still smaller. So the minimum δ must be zero. \square




Figure 1.

2.5. . Liulevicius [12], Dold [8]. *If $G \neq 1$ acts freely on $S(\mathbb{E})$ then \mathbb{E} has the Borsuk-Ulam property, i.e., every continuous G -map $s : E_N(G) \rightarrow \mathbb{E}$ has a zero.*

This generalizes Borsuk’s theorem [6] which says (because $E_N(\mathbb{Z}/2) =$ octahedral N -sphere equipped with the antipodal $\mathbb{Z}/2$ action) that the representation of $\mathbb{Z}/2$ in \mathbb{R}^N given by $x \mapsto -x$ has the B-U property.

Proof. It suffices to prove the result for complex representations, for if there were a G -map $E_N(G) \rightarrow S(\mathbb{E})$, then its 2-fold join would provide a G -map $E_{2N}(G) \subset E_{2N+1}(G) \rightarrow S(\mathbb{E}) \cdot S(\mathbb{E}) = S(\mathbb{E} \oplus \mathbb{E} \cong \mathbb{E} \otimes \mathbb{C})$ with G acting freely on $S(\mathbb{E} \otimes \mathbb{C})$.

Also it suffices to do just the prime cyclic case: for each G contains a subgroup $H \cong \mathbb{Z}/p$, and this case then gives us at least $|G| \div p$ zeros, one in each $E_N(H_g) = (H_g) \cdots (H_g)$. So the result follows from 2.6.3 which in fact gives for all q an explicit characterization of complex \mathbb{Z}/q representations having the Borsuk-Ulam property. \square

2.6. Characteristic classes of representations. Recall that the cohomology of $G \cong \pi_1(BG)$ is defined to be that of the classifying space BG . Likewise — see the appendix of Atiyah [1] — the characteristic classes of any representation \mathbb{E} of G are defined to be those of the corresponding vector bundle $\mathcal{E} = EG \times_G \mathbb{E} \rightarrow BG$.

2.6.1. . In dimensions $\leq N$ a characteristic class of \mathcal{E} vanishes iff its restriction to B_NG vanishes.

Proof. “Only if” is obvious. Using naturality of characteristic classes note that the restriction is the corresponding class of the bundle $E_NG \times_G \mathbb{E} \rightarrow B_NG$. Further the $(N+1)$ -fold join $E_NG = G \cdots G$ is $(N-1)$ -connected, so its identity map extends to a continuous G -map $(EG)_N \rightarrow E_NG$ from the N -skeleton $(EG)_N$ of EG to E_NG , thus giving us a bundle map $(EG)_N \times_G \mathbb{E} \rightarrow E_NG \times_G \mathbb{E}$. So, again by naturality, the corresponding class of $(EG)_N \times_G \mathbb{E} \rightarrow (BG)_N$ is also zero. This gives “if” because the inclusion induced map $H^i(BG) \rightarrow H^i((BG)_N)$ is injective for $i \leq N$. \square

We’ll equip \mathbb{E} with some orientation and let $\widehat{\mathbb{Z}}$ denote the integers equipped with the G -action $g \bullet n = \pm n$, the sign depending on whether $E \xrightarrow{g} \mathbb{E} \xrightarrow{\cong} \mathbb{E}$ preserves or reverses orientation. Now take any continuous G -map $s : EG \rightarrow \mathbb{E}$ with no zeros on the $(N-1)$ -skeleton and associate to any oriented N -simplex σ the *degree* of the map $s : \partial\sigma \rightarrow \mathbb{E} \setminus \{0\}$. This cochain $\sigma \mapsto \deg(s|\partial\sigma)$, which is equivariant with respect to the G -actions of EG and $\widehat{\mathbb{Z}}$, can be verified to be a cocycle, and its cohomology class $e(\mathbb{E}) \in H^N(G; \widehat{\mathbb{Z}})$ verified to be independent of the map s chosen. For these standard facts of *obstruction theory* see Steenrod [23, §35].

For example, we can choose s linear, when of course $\deg(s|\partial\sigma) \in \{-1, 0, +1\}$, and the “Linear Borsuk-Ulam” 2.4 tells us that this cocycle is nonzero for all \mathbb{E} not containing the trivial representation. The vanishing of its cohomology class interprets as follows.

2.6.2. .

Theorem 1. *The representation \mathbb{E} has the Borsuk-Ulam property iff the characteristic class $e(\mathbb{E}) \in H^N(G; \widehat{\mathbb{Z}})$ is nonzero.*

Proof. By 2.6.1 this class is zero iff the corresponding class of the bundle $E_N G \times_G \mathbb{E} \rightarrow B_N G$ is zero, but this happens (see [23, §35]) iff this vector bundle admits a continuous nonzero section, i.e., iff there is a continuous G -map $E_N G \rightarrow \mathbb{E}$ having no zeros. \square

It might be appropriate to call $e(\mathbb{E})$ the *van Kampen class* of \mathbb{E} because it can be traced back, for the case $G = \mathbb{Z}/2$ to [11]. In case the action of G on \mathbb{E} is orientation preserving, i.e., $\widehat{\mathbb{Z}} = \mathbb{Z}$, the integers equipped with the trivial action of G , then $e(\mathbb{E}) \in H^N(G; \mathbb{Z})$ identifies — see Milnor-Stasheff [15, p. 147] — with the *Euler class* of the oriented N -dimensional plane bundle $\mathcal{E} \rightarrow BG$. Thus, if N is even and \mathbb{E} is a complex $N/2$ -dimensional representation of G , then $e(\mathbb{E})$ coincides — see [15, p. 158] — with the $N/2$ -th *Chern class* $c_{N/2}(\mathbb{E})$ of this complex $N/2$ -dimensional bundle \mathcal{E} . Evens [9] has shown that $c_{N/2}(\mathbb{E})$ can always be computed purely algebraically, provided one knows the cohomology ring of G and the Brauer decomposition of \mathbb{E} . These computations can be quite hard, but the simple cases we need are easily dealt with directly.

We recall that \mathbb{Z}/q has q irreducible complex representations, all one-dimensional, being in fact the q homomorphisms $\mathbb{Z}/q \rightarrow \mathbb{C}^\times$, $\omega \mapsto \omega^\ell$, $1 \leq \ell \leq q$, where ω denotes the generator $\exp(2\pi i/q)$ of \mathbb{Z}/q .

2.6.3. .

Theorem 2. *Let m_ℓ denote the multiplicity of ω^ℓ in the irreducible decomposition of the complex $N/2$ -dimensional representation \mathbb{E} of \mathbb{Z}/q . Then \mathbb{E} has the Borsuk-Ulam property iff $q \nmid \prod_\ell (\ell)^{m_\ell}$.*

Proof. We'll use 2.6.2. The multiplicativity of Chern classes shows

$$e(\mathbb{E}) = c_{N/2}(\mathbb{E}) = \prod_\ell (c_1(\omega^\ell))^{m_\ell},$$

where $c_1(\omega^\ell) \in H^2(\mathbb{Z}/q; \mathbb{Z})$ denotes the first Chern class of the representation $\omega \mapsto \omega^\ell$ and multiplication is the cup product of $H^*(\mathbb{Z}/q; \mathbb{Z})$. Since $c_1 : \text{Hom}(G, \mathbb{C}^\times) \rightarrow H^2(G; \mathbb{Z})$ is always a group isomorphism — see Atiyah [1, (3), p. 62] — it follows that

$$e(\mathbb{E}) = \prod_\ell (\ell c_1(\omega))^{m_\ell} = \prod_\ell (\ell)^{m_\ell} (c_1(\omega))^{N/2} = u^{N/2} (\prod_\ell (\ell)^{m_\ell}),$$

where $u : H^i(\mathbb{Z}/q; \mathbb{Z}) \rightarrow H^{i+2}(\mathbb{Z}/q; \mathbb{Z})$ is the map given by taking cup product with the generator $c_1(\omega)$ of $H^2(\mathbb{Z}/q; \mathbb{Z})$ and $\prod_\ell (\ell)^{m_\ell} \in \mathbb{Z} = H^0(\mathbb{Z}/q; \mathbb{Z})$. This periodicity map u is an epimorphism for $i = 0$ and an isomorphism for $i \geq 1$ (and the remaining odd dimensional cohomology of \mathbb{Z}/q is zero): see Cartan-Eilenberg [7, p. 260]. So it follows that $e(\mathbb{E})$ vanishes iff $u(\prod_\ell (\ell)^{m_\ell}) = \prod_\ell (\ell)^{m_\ell} \cdot c_1(\omega)$ vanishes, i.e., iff q divides $\prod_\ell (\ell)^{m_\ell}$. \square

It seems one can give a similar explicit characterization of the complex Borsuk-Ulam representations of any finite Abelian group G .

2.7. . The last theorem gives rise to some remarks.

2.7.1. . Obviously, for a complex \mathbb{Z}/q representation \mathbb{E} , the *group action is free on $S(\mathbb{E})$ iff only those representations $\omega \mapsto \omega^\ell$ occur in it for which ℓ is relatively prime to q* . So 2.6.3 shows that the Borsuk-Ulam holds in many cases not covered by 2.5.

However 2.6.3 also shows that *if q is composite and $d + 1$ is an even number ≥ 4 , then there exist continuous \mathbb{Z}/q maps $E_N(\mathbb{Z}/q) \rightarrow \mathbb{L}^\perp$ having no zeros*. This follows because, for $\mathbb{L}^\perp = \mathbb{C}^{(d+1)/2}[\mathbb{Z}/q]$ the number $\Pi_\ell(\ell)^{m_\ell}$ equals $((q-1)!)^{(d+1)/2}$, and $q!(q-1)!$ unless q is prime or equal to 4. Thus to generalize the continuous version of the proof of 2.3 beyond the case q prime one needs non-cyclic groups G .

2.7.2. . Sierksma [22] has conjectured that a cardinality $(q-1)(d+1)+1$ subset of d -space has at least $((q-1)!)^d$ Tverberg partitions, i.e., that the linear map $s : E_N(Q) \rightarrow \mathbb{L}^\perp$ of 2.3 has at least $((q-1)!)^{d+1}$ zeros. It may in fact be possible to algebraically count these generic *Tverberg zeros* with appropriate *local degrees* ± 1 , so that one always get $((q-1)!)^{d+1}$. One cannot hope however for a similar index formula for *Tverberg partitions*, because this would imply, for $q = 3$ and $d = 2$, that the number of these partitions is always even, which is not so.

If one attempts such a signed counting by using finite group actions then one runs into problems. For example by taking $S \subset \mathbb{A}^d$ in a *general position* we can ensure that s has no zeros on the $(N-1)$ -skeleton of $E_N(\mathbb{Z}/q)$ — i.e., that no proper subset of S has a Tverberg partition into q parts — and then evaluate the cocycle $\sigma \mapsto \deg(s|\partial\sigma)$ of $e(\mathbb{L}^\perp)$ on some equivariant N -cycle of $E_N(\mathbb{Z}/q)$. However this algebraic counting does not give an *integer invariant* because $e(\mathbb{L}^\perp)$ lives in $H^N(\mathbb{Z}/q; \mathbb{Z}) \cong \mathbb{Z}/q$ and so is of finite order. Anyhow for the q prime case this method does suffice to give rough lower bounds for the number of Tverberg partitions: see Vučić-Zivaljević [26].

2.7.3. . Sierksma’s problem is *stable with respect to d* i.e., we can increase d by 1. To see this add, to a general position $S \subset \mathbb{A}^d \subset \mathbb{A}^{d+1}$, $q-1$ new points of $\mathbb{A}^{d+1} \setminus \mathbb{A}^d$, and at the same time perturb one of the old points v out of \mathbb{A}^d . In a Tverberg partition of this set $\hat{S} \subset \mathbb{A}^{d+1}$ the part containing v cannot contain any of the new $q-1$ points, for then some other part contains none and so is in \mathbb{A}^d , and thus restricting to \mathbb{A}^d we would have got a Tverberg partition of the proper subset $S \setminus \{v\}$ of the general position set $S \subset \mathbb{A}^d$. Thus \hat{S} has at most $(q-1)!$ times as many Tverberg partitions as S . A similar but simpler argument shows likewise that the “continuous” Tverberg problem is also stable with respect to d . So we can assume $d+1$ *even* (this we’ll do from here on), $d \gg q$, etc., with impunity in our proofs.

2.8. . The next result has also been proved independently by Ozaydin [16] and Volovikov [25].

2.8.1. .

Theorem 3. *The “continuous” Tverberg theorem is true for all prime powers $q = p^k$.*

Proof. We know from 2.7.1 that the argument of 2.3 can work for $k \geq 2$ only if we use the representation $\mathbb{L}^\perp(G)$ of some non-cyclic order q group G . By 2.8.2 below it does work for $G = (\mathbb{Z}/p)^k$. \square

2.8.2. .

Theorem 4. *The B-U property holds for any representation \mathbb{E} of $(\mathbb{Z}/p)^k$ not containing the trivial representation.*

Proof. Without loss of generality (cf. proof of 2.5) we can assume \mathbb{E} complex. So it is the direct sum of irreducible one-dimensional representations $(\mathbb{Z}/p)^k \rightarrow \mathbb{C}^\times$. These form a group, each member being of the type

$$(\omega_1, \dots, \omega_k) \mapsto \omega_1^{\ell_1} \cdots \omega_k^{\ell_k}$$

where ω_i 's denote copies of the generator $\omega = \exp(2\pi i/p)$, and $0 \leq \ell < p$ with not all ℓ_i 's zero. If, in the isomorphic group $H^2((\mathbb{Z}/p)^k; \mathbb{Z})$, x_i denotes the first Chern class of $(\omega_1, \dots, \omega_k) \mapsto \omega_i$, then $(\omega_1, \dots, \omega_k) \mapsto \omega_1^{\ell_1} \cdots \omega_k^{\ell_k}$ has first Chern class $\ell_1 x_1 + \cdots + \ell_k x_k$.

With mod p field coefficients the cohomology algebra of $(\mathbb{Z}/p)^k$ is isomorphic to the polynomial algebra $\mathbb{Z}/p[x_1, \dots, x_k]$ — this follows by using the case $k = 1$, $B(\mathbb{Z}/p) \times \cdots \times B(\mathbb{Z}/p) \simeq K((\mathbb{Z}/p)^k, 1)$, and the Kunneth formula for field coefficients — and so has no zero divisors. Therefore the cup product $e(\mathbb{E})$ of all these nonzero 2-dimensional classes $\ell_1 x_1 + \cdots + \ell_k x_k$ is nonzero, which by 2.6.2 is same as saying that \mathbb{E} has the B-U property. \square

2.8.3. . *Let \mathbb{E} be as above, and \mathbb{F} be any other representation of $(\mathbb{Z}/p)^k$ with $\dim(\mathbb{F}) > \dim(\mathbb{E})$. Then there does not exist a continuous $(\mathbb{Z}/p)^k$ -map from the sphere $S(\mathbb{F})$ to the sphere $S(\mathbb{E})$.*

This is another (known) generalization of Borsuk’s theorem [6] which is the case of $\mathbb{Z}/2$ acting on two Euclidean spaces via $x \mapsto -x$. See e.g., Atiyah-Tall [2] and Bartsch [5] for more on equivariant maps between representation spheres.

Proof. Since $\dim(\mathbb{F}) > \mathbb{N}$ the connectivity of the sphere $S(\mathbb{F})$ allows us to construct a continuous $(\mathbb{Z}/p)^k$ -map into it from the free N -dimensional $(\mathbb{Z}/p)^k$ -complex $E_N((\mathbb{Z}/p)^k)$. This and 2.8.2 rule out the possibility of any equivariant map $S(\mathbb{F}) \rightarrow S(\mathbb{E})$. \square

2.9. . Unfortunately one cannot extend the “continuous” Tverberg further by a similar use of other groups G of order $q \neq p^k$.

2.9.1. . For any finite group G whose order is not a prime power there exists a continuous G -map $E_N G \rightarrow \mathbb{L}^\perp(G)$ having no zeros.

One way of checking this is to note first that if $H < G$, and $\mathbb{L}^\perp(H)$ does not have the Borsuk-Ulam property, then $\mathbb{L}^\perp(G)$ also does not have the Borsuk-Ulam property. This follows because $\mathbb{L}(G)$ is induced by $\mathbb{L}(H)$, so allowing us to construct from a given H -map $E_{N'}(H) \rightarrow \mathbb{L}(H)$ whose image misses the diagonal, a G -map $E_N(G) \rightarrow \mathbb{L}(G)$ whose image also misses the diagonal. Hence by 2.6.3 we are only left to consider those G 's, of non-prime power order, which are such that all elements are of prime order. Some group theory shows that such a G must contain a subgroup H which is a non-Abelian extension of $(\mathbb{Z}/p)^k$ by a cyclic group of a different prime order. The proof can now be completed by checking that the Euler class of $\mathbb{L}^\perp(H)$ is zero.

We have omitted the details — cf. Bartsch [5] who proceeds as above (instead of Euler classes he uses a Burnside ring argument) to obtain a similar result about maps between representation spheres — because we'll see below that a simpler reasoning gives more.

2.9.2. . The point to note is that in 2.1 to 2.3 the natural group to use was the symmetric group Σ_q of all permutations of Q . It acts in the obvious way on $Q \cdot \dots \cdot Q$, and on \mathbb{L} , and the map $s : Q \cdot \dots \cdot Q \rightarrow \mathbb{L}^\perp$ of 2.3 commutes with these Σ_q actions. Further \mathbb{L}^\perp contains no trivial representation of Σ_q , or for that matter of any subgroup of Σ_q which acts transitively on Q . The only advantage in using the simply transitive subgroups G was that their action on $Q \cdot \dots \cdot Q$ is free.

When we consider \mathbb{L}^\perp as a Σ_q -representation its Euler class lives in $H^N(\Sigma_q; \mathbb{Z})$. We were previously looking at its restrictions to $H^N(G; \mathbb{Z})$ for some subgroups $G \subset \Sigma_q$, e.g., for $q = p^k$, $k \geq 2$, 2.6.3 and 2.8.2 show respectively that this restriction is zero for $G = \mathbb{Z}/p^k$ but nonzero for $G = (\mathbb{Z}/p^k)$. Could it not be that for a $q \neq p^k$ this class is nonzero despite the fact 2.9.1 that its restriction to all simply transitive subgroups G is zero? If so the “continuous” Tverberg would extend to such a q , because we obviously have a continuous Σ_q -map from the free and N -dimensional Σ_q -complex $E_N \Sigma_q$ to the $(N - 1)$ -connected Σ_q -complex $Q \cdot \dots \cdot Q$. Unfortunately the answer to this new question is also “no”.

2.9.3. .

Theorem 5. The Euler class of the Σ_q -representation \mathbb{L}^\perp is nonzero iff q is a prime power.

Proof. By 2.8.2 it only remains to look at the case $q \neq p^k$. One has $H^N(\Sigma_q; \mathbb{Z}) = \bigoplus_p H^N(\Sigma_q; \mathbb{Z}, p)$, where p runs over all primes, and $H^N(\Sigma_q; \mathbb{Z}, p)$ denotes the p -primary component of $H^N(\Sigma_q; \mathbb{Z})$. If $P \subset \Sigma_q$ is a p -Sylow subgroup then — see Cartan-Eilenberg [7, p. 259, Thm. 10.1] —

restriction gives us a monomorphism $H^N(\Sigma_q; \mathbb{Z}, p) \rightarrow H^N(P; \mathbb{Z})$. So it suffices to show that the restriction of our class to each $H^N(P; \mathbb{Z})$ is zero. To see this note that $|P|$ is not divisible by $q \neq p^k$, so P does not act transitively on Q , so there are trivial P -representations outside the diagonal of \mathbb{L} , i.e., in \mathbb{L}^\perp . \square

Note that, the Σ_q -action on $E_N(Q)$ being not free, this still leaves open the question whether, for $q \neq p^k$, one can have a continuous Σ_q -map $E_N(Q) \rightarrow \mathbb{L}^\perp$ having no zeros? It seems that $U(q)$ -actions are called for to settle this point, so we postpone it to a sequel which will deal with infinite group actions.

References

- [1] M.F. Atiyah, *Characters and cohomology of finite groups*, Publ. I.H.E.S., **9** (1961), 23-64.
- [2] M.F. Atiyah and D.O. Tall, *Group representations, λ -rings and the J-homomorphism*, Topology, **8** (1969), 253-297.
- [3] I. Bárány, *A generalization of Carathéodory's theorem*, Disc. Math., **40** (1982), 141-152.
- [4] I. Bárány, S.B. Shlosman and A. Szücs, *On a topological generalization of a theorem of Tverberg*, Jour. Lond. Math. Soc., **23** (1981), 158-164.
- [5] T. Bartsch, *On the existence of Borsuk-Ulam theorems*, Topology, **31** (1992), 533-543.
- [6] K. Borsuk, *Drei Sätze über die n-dimensionale Euklidische Sphäre*, Fund. Math., **20** (1933), 177-190.
- [7] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton, 1956.
- [8] A. Dold, *Simple proofs of some Borsuk-Ulam results*, Contemp. Math., **19** (1983), 65-69.
- [9] L. Evens, *On the Chern classes of representations of finite groups*, Trans. Amer. Math. Soc., **115** (1965), 180-193.
- [10] G. Kalai, *Combinatorics and Convexity*, Proc. I.C.M. Zürich, Birkhäuser, (1995), 1363-1374.
- [11] E.R. van Kampen, *Komplexe in euklidischen Räumen*, Abhand. Math. Sem. Hamburg, **9** (1932), 72-78, 152-153.
- [12] A. Liulevicius, *Borsuk-Ulam theorems for spherical space forms*, Contemp. Math., **19** (1983), 189-192.
- [13] M. de Longueville, *The topological Tverberg theorem for prime powers*, Seminar notes, Technisches Universität Berlin, 1998.
- [14] J.W. Milnor, *Construction of universal bundles*, II, Ann. of Math., **63** (1956), 430-436.
- [15] J.W. Milnor and J.D. Stasheff, *Characteristic Classes*, Princeton, 1974.
- [16] M. Ozaydin, *Equivariant maps for the symmetric group*, unpublished.
- [17] K.S. Sarkaria, *A generalized Kneser conjecture*, J. Combin. Th., **B 49** (1990), 236-340.
- [18] _____, *A generalized van Kampen-Flores Theorem*, Proc. Amer. Math. Soc., **111** (1991), 559-565.

- [19] ———, *Sierksma's Dutch Cheese Problem*, ms. circulated in 1988-89, unpublished.
- [20] ———, *Tverberg's theorem via number fields*, *Isr. Jour. of Math.*, **79** (1992), 317-320.
- [21] ———, *Sierksma's Dutch Cheese Problem*, revised version of M.S.R.I. Berkeley preprint, 1997-025.
- [22] G. Sierksma, *Convexity without linearity; the Dutch cheese problem*, Mimeographed Notes, University of Gröningen, 1979.
- [23] N.E. Steenrod, *The topology of fibre bundles*, Princeton, 1951.
- [24] H. Tverberg, *A generalization of Radon's theorem*, *Jour. Lond. Math. Soc.*, **41** (1966), 123-128.
- [25] A.Yu. Volovikov, *On a topological generalization of the Tverberg theorem*, *Math. Notes*, **59** (1996), 324-326; *Mat. Zametki*, **59** (1996), 454-456.
- [26] A. Vučić and R.T. Zivaljevic, *Note on a conjecture of Sierksma*, *Disc. Comput. Geom.*, **9** (1993), 339-349.

Received August 15, 1998.

PANJAB UNIVERSITY
CHANDIGARH 160014
INDIA