

*Pacific
Journal of
Mathematics*

AN ALGEBRAIC ANNULUS THEOREM

G.P. SCOTT AND G.A. SWARUP

Volume 196 No. 2

December 2000

AN ALGEBRAIC ANNULUS THEOREM

G.P. SCOTT AND G.A. SWARUP

We present an extension of Dunwoody's theory of tracks and use it to prove an analogue of the annulus theorem for hyperbolic groups.

1. Introduction.

We prove the following analogue for (word) hyperbolic groups of the Annulus Theorem for 3-manifolds:

Theorem 1.1. *Let G be a torsion free hyperbolic group with one end. Suppose that G has an infinite cyclic subgroup such that the number of ends of the pair (G, H) is greater than one. Then G splits over some infinite cyclic subgroup.*

Corollary 1.2. *Let G' be a subgroup of finite index in a torsion free hyperbolic group G . Then G splits over an infinite cyclic group if and only if G' splits over an infinite cyclic group.*

This result has also been proved by Bowditch [2], using very different methods. We discuss these differences at the end of this introduction.

The terminology used is standard (from [11] and [17]). The importance of splitting groups along infinite cyclic subgroups is well known from the work of Paulin [13], Rips and Sela [14] and Sela [18].

If G has an infinite cyclic subgroup H such that $e(G, H) \geq 3$, we will say that G is of *multi-band type*. Otherwise we will say that G is of *surface type*. The proofs of Theorem 1.1 are different in the two cases. We do not need the assumption that G is torsion free in the multi-band case. In this case we obtain the following result.

Theorem 1.3. *Let G be a one-ended hyperbolic group and let H be a two-ended subgroup with $e(G, H) \geq 3$. Then G splits over a subgroup commensurable with H .*

Both of the above results are closely related to the Annulus Theorem for 3-manifolds. There are several versions of this result and we state one of the most basic ones here. A map $f : S^1 \times I \rightarrow M$ of the annulus into a 3-manifold M is *essential* if it is π_1 -injective, proper in the sense that f maps

the boundary of the annulus into the boundary ∂M of M , and in addition f is not properly homotopic into ∂M .

Annulus Theorem: *Let M be a compact orientable irreducible 3-manifold with incompressible boundary. If M admits an essential map of the annulus $S^1 \times I$, then it admits an essential embedding of the annulus.*

The connections between the Annulus Theorem for 3-manifolds and our algebraic analogues are described briefly as follows. The assumptions that M be compact orientable and irreducible with non-empty boundary imply that $G = \pi_1(M)$ is torsion free. The additional assumption that M has incompressible boundary implies that G has one end. Also if $f : S^1 \times I \rightarrow M$ is a π_1 -injective proper map and H denotes the infinite cyclic subgroup of G carried by f , then f is essential if and only if $e(G, H) \geq 2$, and G splits over H if and only if f is properly homotopic to an embedding. Finally, if $e(G, H) \geq 3$, then f can be homotoped to cover an embedded essential annulus.

Our paper is organised as follows. In Section 2, we develop the general theory of patterns and tracks in 2-complexes and its connection with the number of ends of a 2-complex. This theory was introduced by Dunwoody in [6]. We extend Dunwoody's theory by adding the idea of a singular pattern and of the length of a pattern. Dunwoody introduced patterns and tracks in order to prove the accessibility of finitely presented groups. We show how our new ideas give an alternative version of his arguments. In Section 3, we discuss the use of tracks in a situation where one has a pair of groups such that $e(G, H) \geq 2$, and H is infinite cyclic. In Section 4, we discuss crossing of tracks and give examples of groups of multi-band type and of surface type. The proof of Theorem 1.3 on the multi-band case is completed in Section 5. The rest of the paper is devoted to the surface type case, and the main construction of the proof comes in Section 8. We use hyperbolicity of the group a bit more in the surface type case. On the whole our use of hyperbolicity seems weak and we expect that the 2-complex techniques set out in our proofs will be useful in the study of splittings of finitely presented groups. Another feature of the proof is the surface-like properties of a large class of one-ended torsion free hyperbolic groups, particularly those which we call surface type groups. In this case, we need to deal with orientation reversing elements in the group; we show that if a surface-like group has orientation reversing elements then it splits as an amalgamated free product over an infinite cyclic subgroup which has index two in one of the vertex groups which is also infinite cyclic. This is the analogue of the fact that a non-orientable surface must contain an embedded Mobius band. Our proof is suggested by the arguments of Dunwoody [6] and Tukia [23] and by the least area arguments of Freedman, Hass and Scott [8] and [9]. As in Tukia [23], we run into difficulties in the torsion case. With some

more work one can show without the torsion free assumption that either the conclusion of Theorem 1.1 holds or there is a subgroup of G which “looks like” a triangle group.

Brian Bowditch [2] recently developed a theory of JSJ-decompositions for one-ended hyperbolic groups with locally connected boundary, and Swarup then showed that any one-ended hyperbolic group has locally connected boundary [22]. Using an extension of the work of Rips, Bestvina and Feighn (see [3]), Bowditch showed that if one removes the torsion freeness hypothesis from Theorem 1.1, the conclusion remains true except for the case of triangle groups. His techniques are very different from ours, making heavy use of the boundary of a hyperbolic group, and as explained above we expect that our techniques will be useful in the study of splittings of finitely presented groups. It is for this reason that we have presented the theory of patterns in more generality and detail than we require in this paper. Perhaps Gabai’s techniques [10] may also be useful for extending Theorem 1.1. Another obvious question is whether there is an analogue of orientation covers for hyperbolic groups.

Acknowledgments. Much of this work was carried out while the first author was visiting the University of Melbourne in 1994 and the Mathematical Sciences Research Institute in Berkeley in 1994 and 1995. He is grateful for the partial support provided by the University of Melbourne, by NSF grant DMS-9306240, and by MSRI, which is partially supported by NSF grant DMS-9022140.

Note added after acceptance by the Pacific Journal in October 1998: Apart from corrections to two misprints and updating of references, this paper is unchanged from the version completed by the authors in July 1996. In the intervening two years, our result has been generalised to the case of arbitrary finitely presented groups and splittings over free abelian groups by Dunwoody and Swenson in a series of preprints each of which has been withdrawn and replaced by an updated version. We have further generalised their result to the case of splittings over virtually polycyclic groups, and we heard recently that Dunwoody and Swenson have done this also.

Note added in proof in September 2000: Dunwoody and Swenson’s paper appeared in *Inventiones* earlier this year.

2. Tracks and ends.

Throughout this section, Y will denote a connected locally finite 2-dimensional simplicial complex. A subset of Y will be called a *pattern* if it intersects each closed simplex of Y in a compact properly embedded submanifold of codimension one. This means that it avoids the 0-skeleton of Y ,

meets each 1-simplex in a finite set and meets each 2-simplex in a compact 1-dimensional submanifold whose boundary lies on the boundary of the simplex. See Figure 1a. A pattern t in Y locally has a collar neighbourhood, and globally there is a neighbourhood of t in Y which is an I -bundle over t . If this bundle is trivial, we say that t is *two-sided*. In this paper, we will mostly consider two-sided patterns but we will also need to consider one-sided patterns at times. A component of a pattern which is a circle in the interior of a 2-simplex will be called a *trivial circle*.




Figure 1.a.
(Pattern)




Figure 1.b.
(Normal Pattern)

Note that a pattern is compact if and only if it meets only a finite number of the simplices of Y . As we are mainly interested in the combinatorics described by a pattern, we will often say that a compact pattern is finite. Finite two-sided patterns typically arise from maps to the real line by taking inverse images of regular points. They can also be constructed using coboundaries of 0-cochains. A pattern will be called *normal* if it intersects each 2-simplex only in arcs, and the two ends of each arc lie on different edges of the simplex. See Figure 1b. A connected pattern is called a *track*. It should be noted that our definitions of the terms pattern and track are more general than Dunwoody's definitions in [6]. His patterns and tracks are all normal in our terminology. Note that if Y has a free edge e , which means that e is not a face of any 2-simplex, then a single point of the interior of e is a normal track in Y . We will not usually be interested in such tracks, but one does need to be aware of their existence.

As patterns have similar separation properties to those of a codimension-one submanifold of a manifold, we will say that a two-sided pattern t in a 2-complex Y *bounds* Z if t cuts Y into pieces such that the union of some of them has closure Z , and Z contains t but meets only one side of each component of t . A choice of normal direction for each component of a two-sided pattern t will be called a *transverse orientation* of t , and we will refer to t with a prescribed transverse orientation as an *oriented pattern*. Our interest in patterns and tracks in Y is because they give a good way of discussing the number of ends of Y . They are also a natural generalisation of the idea of a normal surface in a 3-manifold. For our purposes, we will need to consider patterns which are not normal, although these will play a small

role in our arguments. Consider a finite pattern t in Y . We will be interested in whether the closures of the components of $Y - t$ are compact or not. Again, because it is the combinatorics which are important, a component whose closure is compact will be called finite. This is equivalent to saying that the component contains only a finite number of vertices of Y . We will say that a finite pattern t in Y *splits* Y if it cuts Y into pieces at least two of which are infinite. Now consider an oriented finite pattern t . If f denotes a proper map of the line into Y or a map of the circle into Y such that f is transverse to t , we define the sign of an intersection point of f with t by comparing the orientation of the line or circle with the transverse orientation of t . (Here we use the definition that a map $f : X \rightarrow Y$ is *proper* if the pre-image of any compact set is compact.) By summing over all intersection points, we define the intersection number of f with t . As any proper map of the line into Y and any map of the circle can be properly homotoped to be transverse to t , and as the intersection number obtained is independent of the homotopy, this defines the intersection number of any such map with t . We say that an oriented finite pattern t is *essential* if any loop in Y has zero intersection number with t and there is a proper map of the line into Y which has non-zero intersection number with t . Note that if t is not connected, then the condition of essentiality may well depend on the choice of transverse orientation. We will say that a two-sided finite pattern t is *essential* if t admits an orientation which makes it essential. Clearly an essential pattern must split Y . The additional conditions which we impose for a pattern to be essential are needed because we want to consider patterns which are not connected. They are aimed at simplifying the discussion of essential patterns but they do not really restrict us in any way. To see this, consider a finite pattern t which splits Y . Next pick an infinite component of the complement of t and denote its closure by U . We define a sub-pattern t' of t to consist of those components of t which meet U but do not lie in the interior of U . Then t' bounds U and if we transversely orient t' to point into U , then t' is essential in Y . Thus a splitting pattern always contains an essential pattern. We will say that an essential pattern t is *elementary* if t bounds a subset U of Y such that all the orientations point into U or all point out of U . (Of course, such U must be infinite.) We have just seen that any splitting pattern contains an elementary essential pattern.

We will say that two oriented finite patterns t_1 and t_2 in Y are *equivalent* if, for any proper map h of the real line into Y and any map h of the circle into Y , the intersection number of h with t_1 equals the intersection number of h with t_2 . If t_1 and t_2 are disjoint, they are equivalent if and only if the union of t_1 and of t_2 with opposite orientation bounds a compact subset U of Y such that for each component V of U , either all the transverse orientations point into V or all point out of V . Thus the relation of equivalence could reasonably

be called oriented cobordism. We will say that an oriented pattern is *trivial* if it is equivalent to the empty set.

Before we discuss ends, we prove the following fundamental lemma about patterns. An invariant of a finite pattern t which will be used often is its *weight* $w(t)$, which we define to be the total number of points of $t \cap Y^{(1)}$, where $Y^{(1)}$ denotes the 1-skeleton of Y .

Lemma 2.1. *Any finite oriented pattern in Y is equivalent to a finite oriented normal pattern, possibly empty.*

Remark 2.2. If we apply the procedure below to a one-sided pattern, we will obtain a one-sided pattern, but equivalence is not defined for one-sided patterns.

Proof. (See [21].) Let t be a finite pattern in Y . If t is not normal, then there is a 2-simplex σ of Y and a component of $t \cap \sigma$ which is either a circle C or an arc λ with both endpoints on one edge e of σ . If there is such a circle C , we alter t by deleting C . If there is such an arc λ , let λ' denote the sub-arc of e with the same endpoints as λ . We alter t by removing a small neighbourhood of λ and, for each 2-simplex σ' , other than σ , which contains the edge e , we add to t an arc parallel to λ' . There is a natural choice of transverse orientation for the new arc which is compatible with the transverse orientation on the rest of t . See Figure 2.




Figure 2.a.




Figure 2.b.

Each of these moves replaces t by an equivalent pattern. Also each of these moves reduces the sum $w(t) + d(t)$, where $w(t)$ is the weight of t and $d(t)$ denotes the sum over all 2-simplices σ of Y of the number of components of $t \cap \sigma$. As this sum is a non-negative integer, this sequence of alterations to t must terminate, at which point we will have obtained an oriented normal pattern as required.

Note that even if t has no components which are trivial circles, such components can be introduced by the second type of alteration. Thus we may need to delete trivial circles at some stage. Now we consider the connection between ends and patterns.

Lemma 2.3. *The locally finite 2-dimensional simplicial complex Y has at least two ends if and only if it contains a finite essential pattern.*

Remark 2.4. If Y contains a finite essential pattern, the preceding lemma shows that it contains a finite normal pattern which is essential.

Proof. If Y contains a finite essential pattern, it is immediate that $Y - \{t\}$ has at least two infinite components, so that Y must have at least two ends. Now suppose that Y has at least two ends. This implies that we can find a proper map g of Y onto the real line \mathbb{R} . Without altering g on the vertices of Y , we can homotop it to be linear on the simplices of Y and this will be a proper homotopy. Now if z is a point of \mathbb{R} which is not the image of a vertex, then $g^{-1}(z)$ is a finite two-sided pattern t which we give the transverse orientation induced from the orientation of \mathbb{R} . Any loop in Y has zero intersection number with t , as any loop in \mathbb{R} has zero intersection number with z . Also if we choose a proper map λ of the line into Y such that $g \circ \lambda$ sends the ends of the line to distinct ends of \mathbb{R} , then λ will have non-zero intersection number with t as $g \circ \lambda$ has non-zero intersection number with z . This shows that t is essential, as required. Note that t will be normal automatically.

So far, we have not discussed anything really different from the ideas of Dunwoody. The first new concept we need to introduce is that of a singular pattern in Y . Note that a pattern t in Y has a natural induced structure as a 1-complex with vertex set consisting of $t \cap Y^{(1)}$. Any circle component of t which lies in the interior of a 2-simplex of Y will not have any vertices, but we will allow this in our definition of a 1-complex for the purposes of discussing singular patterns.

Definition 2.5. A singular pattern in Y is a proper map f of a 1-complex t into Y such that:

- 1) f maps vertices of t to interior points of edges of Y .
- 2) f maps non-vertex points of t to interior points of 2-simplices of Y .
- 3) If v is a vertex of t of valence d , it is mapped to an edge e of Y also of valence d , with distinct rays of the neighborhood of v being mapped to distinct 2-simplices of Y .

We define the *weight* $w(f)$ of a singular pattern f to equal the number of vertices of t . If f is an embedding this agrees with the original definition of weight. We also define a singular pattern f to be *normal* if t has no component which is a circle without vertices, and if, for each edge λ of t , the two endpoints of λ are mapped to distinct edges of Y . Again this agrees with the original definition of normality when f is an embedding.

The conditions in the above definition imply that a singular pattern has similar local separation properties to those of a pattern. In particular it has an induced normal bundle, which means that there is a unique bundle

over t with fibre the interval I such that f extends to the total space of this bundle and the extended map is injective on each fibre of the bundle. We will say that f is two-sided if its normal bundle is trivial. In this case, a choice of normal direction for each component of t will be called a transverse orientation of f and we will say that f is oriented. Our main interest in singular patterns is whether a finite singular pattern is essential in a sense analogous to that for an embedded pattern. We consider a finite oriented singular pattern $f : t \rightarrow Y$. We will say that f is *essential* if any loop in Y has zero intersection number with f and there is a proper map of the line into Y which has non-zero intersection number with f . This clearly agrees with the original definition of essentiality if f is an embedding. Finally we will say that two singular finite oriented patterns f and g are *equivalent* if, whenever h is a proper map of the real line into Y or a map of the circle into Y , the intersection number of h with f equals the intersection number of h with g . Again this agrees with the definition of equivalence of embedded oriented patterns, when f and g are embeddings.

Now we want to bring in ideas of Jaco and Rubinstein [12]. They defined a PL-analogue of area for a normal surface in a triangulated 3-manifold. We will follow their ideas to define an idea of length for a finite, possibly singular, pattern in Y . Jaco and Rubinstein [12] considered the case when the 2-complex is the 2-skeleton of a 3-manifold but their ideas extend to the general case. First one needs a “metric” on Y . To provide this, we simply choose an identification of each 2-simplex of Y with an ideal triangle in the hyperbolic plane. We then glue these triangles by isometries to obtain our “metric” on Y . We will call this a *hyperbolic structure* on Y . Really this structure is defined on $Y - Y^{(0)}$, where $Y^{(0)}$ denotes the set of vertices of Y . It does yield a metric on $Y - Y^{(0)}$, but we will not use this fact. Note that there is no requirement that this metric be complete, and completeness will be irrelevant for our arguments. For simplicity, we will give the definitions for embedded patterns only, with brief comments about how they extend to singular patterns. We define the complexity $c(t)$ of a finite pattern t in Y to be the ordered pair $(w(t), L(t))$, where $L(t)$ denotes the sum over all 2-simplices σ of Y of the length of the 1-manifold $t \cap \sigma$. (Note that $L(t)$ is non-zero unless t consists only of isolated points on free edges of Y .) For a singular pattern $f : t \rightarrow Y$, note that one does not want to measure the length of the image of f , but the length with appropriate multiplicity. This is done by pulling back the metric on Y to a metric on t . It is assumed that we are only considering finite patterns whenever we talk of complexity. These complexities are ordered lexicographically and they are added component-wise. Note that a hyperbolic structure on Y immediately induces a hyperbolic structure on every cover of Y . The idea now is that if we consider patterns whose complexity is as small as possible, then such patterns should have good properties. In particular they should intersect in

a reasonably simple way. This is by analogy with the very nice properties of surfaces of least area in 3-manifolds [8] and of shortest curves on surfaces [9]. It will be very convenient to refer to a pattern of least possible complexity as *shortest*. However, the reader should keep clear the fact that this term means least complexity $c(t)$ and not least length $L(t)$.

We say that two patterns t_1 and t_2 in Y intersect *transversely* if $t_1 \cap t_2$ does not meet the 1-skeleton of Y , and for each 2-simplex σ of Y the 1-manifolds $t_1 \cap \sigma$ and $t_2 \cap \sigma$ intersect transversely in the interior of σ . If t_1 and t_2 are finite patterns in Y which intersect transversely, we can define a new pattern t by performing cut and paste at each point of $t_1 \cap t_2$ and rounding corners. As rounding corners reduces length, we see that $c(t) \leq c(t_1) + c(t_2)$, with strict inequality so long as $t_1 \cap t_2$ is not empty. (For singular patterns which intersect transversely, the same construction can be made so long as they are in general position. This means that no point of $f_1(t_1) \cap f_2(t_2)$ is a multiple point of f_1 or f_2 .) This essentially trivial observation will have important consequences for us in the cases in which we are interested. Note that even if t_1 and t_2 are normal, the new pattern t will usually not be normal. In fact, for each point x of $t_1 \cap t_2$, exactly one of the two possible cuts and pastes at x will yield normal arcs, so that there is a unique cut and paste of t_1 and t_2 which yields a normal pattern. In Dunwoody's work [6], he used only this cut and paste which explains why he did not need to consider non-normal patterns. We will definitely need to use other cut and paste operations. We introduce one further piece of terminology. If t_1 and t_2 are finite patterns in Y which intersect transversely in a non-empty set, then $t_1 \cap t_2$ cuts t_1 and t_2 into pieces which are not patterns. We will call such pieces *partial patterns*. The complexity of a partial pattern is well defined in the same way as the complexity of a pattern, and if s_1, \dots, s_k denote the partial patterns which are contained in t_1 , we have $c(t_1) = \sum c(s_i)$. Note that this equality holds because $t_1 \cap t_2$ does not meet any edge of Y .

We want to use cut and paste constructions on shortest patterns to show that such patterns have nice intersections and we will also need to show that shortest patterns exist in suitable classes of patterns in order to apply these ideas. A good example to bear in mind is that of an essential simple closed curve C on an annulus A equipped with some metric. If this metric is reasonable, e.g., if A covers a closed surface F and the metric on A is lifted from that on F , we can find a shortest loop in the homotopy class of C . Of course, we do not want to think of the homotopy class of a pattern. For us a finite pattern is interesting if it is essential. Given a finite essential pattern t on a 2-complex Y , we will want to know that there is a shortest pattern equivalent to t and/or that there is a pattern which is shortest among all essential patterns in Y .

Before proving the existence results which we need, we will show how to use essential patterns of least complexity. The following result will be useful.

Lemma 2.6. *Suppose that Y has at least two ends, and let t be a finite essential pattern in Y of least possible weight and without trivial circles. Then t is normal and intersects each 2-simplex of Y in at most one arc.*

Proof. The fact that t is normal follows from the proof of Lemma 2.1 as any normalising moves will reduce the weight of t . If E, E^* are infinite disjoint sets of vertices of Y with $Y^{(0)} = E \cup E^*$, then we can construct from δE a pattern which has the required property. If we have an essential pattern t , we take the corresponding subdivision E, E^* of $Y^{(0)}$. Then δE gives an essential pattern s and any 2-simplex which s meets must also be met by t . It follows that $w(s) \leq w(t)$. The minimality of $w(t)$ implies that s and t have the same weight, and now it follows that t must meet each 2-simplex of Y in at most one arc.

Now we come to a key property of essential patterns of least complexity.

Lemma 2.7. *Suppose that Y has at least two ends and let s and t be essential patterns in Y which are shortest among all such patterns. Then either s and t coincide or they are disjoint.*

Proof. First note that s and t are two-sided and must have the same complexity. Next recall that each must contain an elementary essential pattern, so the fact that each is shortest implies that each is already elementary. Now suppose that s and t are not disjoint and intersect transversely. Denote the two pieces of Y bounded by s by S and S^* , and the two pieces of Y bounded by t by T and T^* . Thus $s \cap t$ cuts s into two partial patterns $s \cap T$ and $s \cap T^*$ which we denote s' and s'' , and it cuts t into $t \cap S$ and $t \cap S^*$ which we denote by t' and t'' . None of these four partial patterns need be connected even if s and t are connected. Note that $c(s') + c(s'') = c(s) = c(t) = c(t') + c(t'')$. Of these four partial patterns, we consider one of the shortest, i.e., of least complexity. Without loss of generality, s' is a shortest one. Then consider the patterns $s' \cup t'$ and $s' \cup t''$. After rounding corners, we obtain two patterns each of which has complexity strictly less than $c(t)$, which implies that both patterns are inessential in Y . The pattern $s' \cup t'$ bounds $S \cap T$ and its complement, and this complement must be infinite as it contains S^* . Hence $S \cap T$ must be finite or $s' \cup t'$ would be essential. Similarly $S^* \cap T$ must be finite. But this implies that T is finite so that t is inessential. This contradiction completes the proof of the lemma in the case when s and t intersect transversely.

If s and t intersect but not transversely, we use the trick of Meeks and Yau, see Lemma 1.3 of [8], as modified by Jaco and Rubinstein in Section 2.3 of [12]. Here is how the trick works in our situation. Note that the intersection of s and of t with each 2-simplex σ of Y must consist of geodesic arcs in the hyperbolic metric on σ . We will start by discussing the simplest case when no two geodesic arcs of s and t coincide, and $s \cap t$ meets the 1-skeleton of Y in exactly one point x which lies on an edge e of Y . The problem is

that cutting and pasting s and t at x does not yield a strict reduction in the length of $s \cup t$ as we cannot round corners. We resolve this as follows. First pick a direction along e and measure the angles at which the various geodesic arcs of s and of t meet e at x . The fact that s is a critical point for the length function implies that the sum of the cosines of the angles corresponding to the edges of s must be zero, and the analogous statement holds for the edges of t . In particular, it follows that s and t must locally cross at x . This means that any neighbourhood of x contains points of s on each side of t and vice versa. Now we perturb s slightly to a new track s_1 by moving the point $s \cap e$ a small distance ϵ along e . We will discuss how small to choose ϵ in a moment. The new track s_1 must cross t in at least one 2-simplex σ_1 which has e as an edge. Clearly $c(s_1) \geq c(s)$, and we can assume the inequality is strict as otherwise s_1 and t would be two shortest essential tracks on Y which intersect transversely, contradicting the result of the previous paragraph. Now $L(s_1) - L(s)$ is a function of ϵ which must be zero to the first order as s is a critical point of L . However, the reduction of length obtained by cutting and pasting s_1 and t in σ_1 is, to the first order, linear in ϵ with (non-zero) coefficient $|\cos \theta - \cos \phi|$ where θ and ϕ denote the original angles in σ_1 between e and the arcs of s and t which met e at x . It follows that if we choose ϵ suitably small, then the reduction will outweigh the increase so that the arguments of the preceding paragraph will yield a contradiction by applying them to s_1 and t . If no two geodesic arcs of s and t coincide, and $s \cap t$ meets the 1-skeleton of Y in more than one point, essentially the same argument will work by perturbing s at each of the points where $s \cap t$ meets the 1-skeleton of Y . If some geodesic arcs of s and t coincide, we need to perturb s along the entire 1-complex Γ of coincident arcs and make the same argument as before at each extreme point of Γ . Note that s and t must locally cross at each extreme point x of Γ , so that there is an intersection between s_1 and t in a 2-simplex σ_1 which contains x and contains non-coincident edges of s and t which end at x .

We will need one further related result for singular essential patterns.

Lemma 2.8. *Let $f : t \rightarrow Y$ be a finite two-sided essential singular pattern. Then either f factors through a covering of an essential embedded pattern in Y or there is an essential embedded pattern s in Y such that s and f are equivalent and $c(s) < c(f)$.*

Remark 2.9. This result means that if we can minimise complexity among all singular patterns in an equivalence class or collection of equivalence classes, then the least complexity pattern must be an embedding or a covering of an essential embedded pattern.

Proof. We will first consider the case where f is self-transverse. Then we can perform cut and paste at all the singular points of $f(t)$ to obtain a pattern s in Y such that $c(s) < c(f)$. In order to ensure that s is equivalent to f , we

choose all the cut and paste operations to be orientation preserving. This means that we choose the transverse orientation for $f(t)$ specified by the transverse orientation which makes f essential and then ensure that each cut and paste preserves this direction. The orientation preserving condition ensures that any line or loop which has algebraic intersection number d with f also has algebraic intersection number d with the new pattern, where we choose the obvious transverse orientation for the new pattern.

Now we need to consider the more general situation where f need not be self-transverse. Our aim is to apply the Meeks-Yau trick as in the proof of the preceding lemma. In other words, we will perturb f by a small amount to a self-transverse pattern f' , and then perform cut and paste on f' as in the preceding paragraph to obtain an embedded pattern s such that $c(s) < c(f)$. Note that if f is a covering map of an embedded pattern, there is no way to carry out this procedure. Suppose first that t is connected and that f is not a covering map of an embedded pattern. Then the methods apply to obtain an embedded pattern s equivalent to f such that $c(s) < c(f)$. If t is not connected, but f fails to be a covering map when restricted to some component t_1 of t , we can use the same methods to obtain an embedded pattern s_1 equivalent to f_1 , the restriction of f to t_1 , such that $c(s_1) < c(f_1)$. Let F denote the pattern obtained from f by replacing f_1 by the inclusion of s_1 and let ϵ denote $c(f) - c(F)$. We can perturb F on each of the other components of t , and perform more cut and paste to obtain an embedded pattern s which is equivalent to f . Further, this can be done while increasing the complexity of F by less than ϵ , so that $c(s) < c(f)$ as required.

In the first part of the proof of the preceding lemma, we showed that if f is a two-sided essential self-transverse pattern in Y , then oriented cut and paste yields an equivalent embedded two-sided pattern. A natural question is what can one say when one considers a one-sided pattern. The answer is the following little result which we will use later.

Lemma 2.10. *Let $f : t \rightarrow Y$ be a finite one-sided pattern in Y . Then either f is an embedding, or there is a one-sided embedded pattern in Y which has less complexity than f . In particular, Y cannot be simply connected.*

Proof. If f covers an embedded pattern s in Y , then s must also be one-sided. Thus either f is an embedding or s has less complexity than f as required. Without loss of generality we can assume that t is connected. Suppose also that f is in general position. Pick a maximal tree T in t , and let S denote a subset of t consisting of one point from the interior of each edge of $t - T$. Then there is a transverse orientation for $t - S$. By removing points from S , we can assume that this transverse orientation does not extend over any point of S . We can assume that none of the points of S are double points of f . This allows us to perform oriented cut and paste of f to obtain an embedded pattern s in Y which is transversely oriented except at the points

of S , and the orientation does not extend across the points of S . It follows that s is one-sided as required. If f is not in general position, we apply the Meeks-Yau trick as in the proof of the previous lemma.

Now we want to prove an existence result for patterns of least complexity. First we consider only patterns of the same combinatorial type as a given singular normal pattern f . This means that any such pattern is homotopic to f through singular patterns in Y , and so, in particular, has the same weight as t . We call such a homotopy a *normal* homotopy.

Lemma 2.11. *Let Y be a locally finite 2-dimensional simplicial complex equipped with some hyperbolic structure, and let $f : t \rightarrow Y$ be a finite singular normal pattern in Y such that the restriction of f to each component of t cannot be normally homotoped arbitrarily close to a vertex of Y . Then there is a singular pattern f' which is of the same combinatorial type as f and has least complexity among all such patterns.*

Remark 2.12. The hypothesis on the components of t implies that no component of t can be a single point.

Proof. This is proved exactly as on page 500 of Jaco and Rubinstein [12].

We will be interested in the existence of shortest patterns equivalent to a given essential singular pattern $f : t \rightarrow Y$ but not necessarily of the same combinatorial type. In order to apply the above result, we will need to know that the essentiality of f implies that no component of t can be homotoped arbitrarily close to a vertex of Y . But this need not be true unless we impose some condition on Y . For example, suppose that Y is obtained from two infinite 2-complexes Y_1 and Y_2 by glueing them at a vertex v and let t_i denote the pattern in Y_i which has one vertex on each edge which contains v and has one edge in each 2-simplex which contains v . Thus t_i is a copy of the link of v in Y_i . Then each t_i is essential in Y , as it separates Y into pieces which are essentially Y_1 and Y_2 , and each t_i can be isotoped arbitrarily close to v .

If Y is a locally finite 2-dimensional simplicial complex, we will say that a vertex v of Y is a *splitting vertex* if some component of the link of v in Y does not bound a compact subset of Y . The assumption which we will make for the rest of this paper is that the 2-complexes which we consider have no splitting vertices. This will not be a problem for our applications for the following reason. If Y does have a splitting vertex, so does its universal cover, and it follows that the universal cover has at least two ends. Thus we have the following result.

Lemma 2.13. *Let Y be a locally finite 2-dimensional simplicial complex which covers a finite simplicial complex Z with fundamental group G . If Y has a splitting vertex, then $e(G) > 1$.*

Now we can state and prove our existence result for shortest patterns.

Lemma 2.14. *Let Y be a locally finite 2-dimensional simplicial complex without splitting vertices, and suppose that Y has a hyperbolic structure lifted from a finite complex Z covered by Y . Let f denote an essential finite singular pattern in Y . Then:*

- 1) *There is a singular pattern f' which is equivalent to f and has least complexity among all such singular patterns. Further any such f' must be a covering map of an embedded normal pattern in Y .*
- 2) *There is a singular pattern f'' which has least complexity among all essential singular patterns in Y . Further, any such f'' must be an embedded normal pattern in Y .*

Remark 2.15. In the first part of this lemma, the situation where f' covers an embedded pattern but is not an embedding can certainly occur. Here is a simple example. Consider any essential track t embedded in Y and let f consist of two parallel copies of t . Then f' could map the two copies of t onto one copy of t .

Proof. Let (w_0, L_0) denote the infimum of the complexities of all singular patterns in Y which are equivalent to f , and let $f_i : t_i \rightarrow Y$ denote a sequence of such patterns such that $c(f_i) \rightarrow (w_0, L_0)$ as i tends to infinity. As remarked earlier, the proof of Lemma 2.1 shows that we can assume that each f_i is normal, as any normalising move reduces complexity. By passing to a subsequence, we can also assume that each f_i has weight w_0 . We consider the projections of the f_i 's into the finite complex Z . This gives us a sequence g_i of singular patterns in Z . Note that our hypothesis that the hyperbolic structure on Y is lifted from one on Z implies that the complexities of f_i and g_i are equal. As there are only finitely many combinatorial types of singular patterns of a fixed weight in a given finite 2-complex, we can arrange, by again passing to a subsequence, that all the g_i 's are of one combinatorial type. This means that each t_i can be identified with a fixed 1-complex t' and that for a given vertex v of t' each g_i maps v to the same edge of Z . Now the proof of Lemma 2.11 shows that there is a singular pattern $g : t' \rightarrow Z$ of the same combinatorial type as the g_i 's and with complexity (w_0, L_0) unless, for large values of i , there is a component u of t' whose image $g_i(u)$ lies arbitrarily close to a vertex of Z . This would imply that $f_i(u)$ also lies very close to a vertex of Y and so must cover some component of the link of this vertex. The fact that Y has no splitting vertices implies that this component bounds a compact subset of Y , and hence that the restriction of f_i to $t' - u$ is equivalent to f_i which contradicts our assumption that each f_i has the least possible weight w_0 . The fact that g has the same combinatorial type as each g_i , means that g can be lifted to a (possibly singular) pattern f'_i in Y , where f'_i is of the same combinatorial type as f_i . For each i , the singular pattern f'_i is equivalent to the original singular pattern f , and it has least possible complexity because its complexity is equal to the complexity

of g . Now Remark 2.9 shows that f'_i must be an embedding or a covering map of an embedded normal pattern.

For the second part of the lemma, the existence part of the proof is essentially the same as in the preceding argument, and is carried out by minimising over the class of all essential singular patterns in Y . Now consider a singular essential pattern $f'' : t'' \rightarrow Y$ of least possible complexity. We already know that f'' must be a covering map of an embedded normal pattern s in Y , so that $c(s) \leq c(f'')$ with equality if and only if f'' is an embedding. There must be a component of $Y - s$ with infinite closure U and with infinite complement as otherwise f'' could not be essential. Let s_1 denote the sub-pattern of s which is the union of those components of s which meet U but do not lie in the interior of U . Then s_1 is an essential pattern in Y . As $c(s_1) \leq c(s) \leq c(f'')$, the fact that f'' minimises complexity over all essential patterns in Y implies that these inequalities are equalities and so $c(s) = c(f'')$ which implies that f'' is an embedding as required.

Using the ideas of the preceding proof, one can give a somewhat different proof of Stallings' Theorem for finitely presented groups and of the accessibility of finitely presented groups by following Dunwoody's arguments in [6]. We sketch the proof of Stallings' Theorem. Consider a finitely presented group G which is the fundamental group of a finite 2-dimensional simplicial complex Y_G with universal cover Y . First we suppose that Y does not have any splitting vertices. We choose a hyperbolic structure on Y_G and give Y the induced hyperbolic structure. The hypothesis of Stallings' Theorem is that Y has at least two ends. Now we can apply Lemma 2.14 to obtain a shortest essential pattern t in Y . A crucial point which comes in here is that as Y is simply connected, each component of t must separate Y . It follows immediately, that some component of t must be essential, and the fact that t has least complexity shows that t must equal this component, so that t is connected. Now we consider the action of G on Y . Lemma 2.7 tells us that for each g in G the translate gt of t is disjoint from or coincides with t . It follows that t projects into Y_G as a covering of some embedded track s . This immediately gives a splitting of G over the finite group C which is the stabiliser of t . If Y does have a splitting vertex v , the existence result for shortest tracks fails but it is not needed. Let u denote the image of v in Y_G . It is immediate that u determines a splitting of G over the trivial group, which is the stabiliser of v . Thus in either case, Stallings' Theorem follows for finitely presented groups.

3. Ends of pairs of groups.

We will now fix some of the notation to be used in most of the rest of this paper. G is a torsion free hyperbolic group with one end and G has an infinite cyclic subgroup H such that $e(G, H) \geq 2$. Following Bowditch, we

sometimes call the number $e(G, H)$ the number of co-ends of H (in G). X denotes a simply-connected 2-dimensional simplicial complex on which G acts on the left freely and simplicially so that the quotient is compact and inherits a simplicial complex structure. If L is a subgroup of G then X_L denotes the quotient complex X/L . We will always use only the triangulation of X_L inherited from the given triangulation of X . The notation X_H usually means that H is infinite cyclic and if h is a generator of H we sometimes write X_h instead of X_H .

We will often be interested in the subgroup of $\pi_1(Y)$ represented by a track t in Y , where Y is any locally finite 2-dimensional simplicial complex. Of course, the fundamental group of any track is a free group, but the group in which we will be interested is the image of the natural map $\pi_1(t) \rightarrow \pi_1(Y)$. If K denotes this subgroup, we will say that t carries K . Note that K is only defined up to conjugacy in $\pi_1(Y)$, but this will not cause any problems as $\pi_1(Y)$ will usually be abelian in the cases in which we are interested.

Now we consider an infinite cyclic subgroup H of G and the corresponding 2-complex X_H . We will assume that H has infinite index in G , so that X_H is not compact. The reader should think of the special case where X_G is a closed orientable surface so that X_H is an annulus which contains an embedded essential simple closed curve. From our point of view, the fundamental property of this simple closed curve is that it separates the annulus into two infinite pieces. In addition, of course, it is connected and carries the entire group H . In the general situation, we want to find a track which will play the role of this simple closed curve on the annulus.

Lemma 3.1. *Let H be an infinite cyclic subgroup of G which has infinite index in G . If t is a finite two-sided track in X_H , then it must separate X_H . If, in addition, t carries the trivial group, then t must separate X_H into two pieces one of which is compact and carries the trivial group, so that, in particular, t is inessential.*

Proof. If t does not separate X_H , it follows that t must carry the trivial subgroup of H . Hence t must lift to X , and as X is simply connected this lift s must separate X into two pieces. Each of these pieces must be infinite, which contradicts our hypothesis that X has only one end. See [20] for a related argument. Now we consider the case when t is separating and carries the trivial group. Again t must lift to X , and this lift s must separate X into two pieces. If both the components of the complement of t in X_H are infinite, or if the finite component carries a non-trivial subgroup of H , then it follows that the complement of s in X consists of two infinite components contradicting our hypothesis that X has only one end. This completes the proof of the lemma.

Now we will concentrate on finite tracks t in X_H which carry a non-trivial subgroup of H .

Lemma 3.2. *Let H be an infinite cyclic subgroup of G which has infinite index in G . Then there is a finite two-sided track in X_H which carries a non-trivial subgroup of H .*

Proof. As X_H is not compact, we can take a proper map $\phi : X_H \rightarrow \mathbb{R}^+$, the non-negative real numbers, and consider the finite pattern $t = \phi^{-1}(c)$ where $\phi^{-1}([0, c])$ contains a loop representing h . At least one of the components of t must be a track of the required type, by the preceding lemma.

Remark 3.3. Another way to obtain tracks is to take suitable sets in X_H whose frontiers are disjoint from vertices.

Lemma 3.4. *If $e(X_H) \geq 2$, there is a finite essential track t in X_H . Any such track carries a non-trivial subgroup of H .*

Proof. As $e(X_H) \geq 2$, we can take a proper map φ of X_H onto the reals and consider the finite essential pattern $\phi^{-1}(c)$ for some c in \mathbb{R} . Now we apply Lemma 3.1 to show that some component of $\phi^{-1}(c)$ is essential. The same lemma shows that any essential track in X_H must carry a non-trivial subgroup of H .

Remark 3.5. If W is a subset of X_H whose projection to \mathbb{R} is not onto, we can choose t to be disjoint from W by choosing c in \mathbb{R} but not in the image of W .

The inverse image in X of such a track t in X has one component if t carries H . Otherwise t must carry a subgroup of H of finite index and we have a finite number of components in the inverse image of t . If we fix a generator h of H , then we can talk of the positive and negative end points of any one of these components in ∂X , the Gromov boundary of X or G .

We want to use tracks to subdivide ∂X . Note that any track in X is two-sided as X is simply connected. We will generally assume that any track A in X which we are considering has two ends; often it will have infinite cyclic stabilizer. The quotient by this stabiliser must be compact if A has two ends. We will use the term *axis* for any track in X with two ends and infinite cyclic stabilizer. We will use $\text{Stab}^0 A$ to denote the subgroup of $\text{Stab} A$ consisting of elements which do not interchange the sides of an axis A , and if h is a generator of $\text{Stab}^0 A$, then ∂A or $E(A)$ will denote the fixed points of h . Usually P_A denotes the positive or attracting fixed point of h , and N_A denotes the negative or repelling fixed point of h . These will also be called positive and negative end points of A . By Coornaert [4], $H = \text{Stab}^0 A$ acts properly discontinuously on $X - E(A)$ with compact quotient. If t is the image of A in X_H , then t carries H . We will say that the axis A is *essential* if t is an essential track in X_H . Note that if A is essential, then $e(X_H) \geq 2$, but the following example shows that the converse is false.

Example 3.6. Let X_H be obtained from an open annulus Z and a compact annulus Z' by identifying one component of $\partial Z'$ with an essential circle S of Z . Consider circles C and C' in Z and Z' respectively which are parallel to S and disjoint from S . By an isotopy, we can arrange that each is a track in X_H . Then C is an essential track in X_H , but C' is not essential. Thus the pre-images of C and C' in X are tracks A and A' with the same endpoints, with A essential and A' inessential.

If A is an essential axis in X , we denote the two parts of the image of $(\partial X - E(A))/H$ separated by t as $\partial_0(X_H)$ and $\partial_1(X_H)$ and think of these as the left and right parts of the boundary $\partial(X_H)$ of X_H . If p denotes the projection from $X_H \cup (\partial X - E(A))/H$ to $X_H \cup \partial(X_H)$, then the subdivision of $(\partial X - E(A))$ consists of $p^{-1}(\partial_0(X_H))$, $p^{-1}(\partial_1(X_H))$. If we take any finite cover X_K of X_H and the lift t' of t and repeat the above construction, then we get the same subdivision of $(\partial X - E(A))$. On the other hand, if we start with a track t whose fundamental group does not surject onto H , then the subdivision of $(\partial X - E(A))$ may depend on the lift of t that we take in X_K . Finally note that distinct essential axes A and A' with the same end points may yield different subdivisions of $(\partial X - E(A))$, as the following example shows.

Example 3.7. This example is the same as the preceding one except that we take Z' to be a half open annulus, so that X_H has three ends. Now C and C' are both essential tracks in X_H , but A and A' divide $(\partial X - E(A))$ in distinct ways.

The positive outcome of the above discussion is the following.

Lemma 3.8. *Given an essential axis A in X and an infinite cyclic $H \subseteq \text{Stab}^0 A$, then A/H defines a subdivision of ∂X_H and the corresponding subdivision of $(\partial X - E(A))$ is independent of the H chosen in $\text{Stab}^0 A$.*

If A, A' are disjoint and $H \subseteq \text{Stab}^0 A \cap \text{Stab}^0 A'$ and if A, A' bound a compact subset of X_H , we get the same subdivision of $(\partial X - E(A))$ whether we use A or A' (note that $E(A) = E(A')$). This generates an equivalence relation on essential axes with the same end points. We will denote these subsets of $(\partial X - E(A))$ by $\partial_0 A$ and $\partial_1 A$; thus implicitly we are using some transverse orientation for A . We will also often use the notation $\partial_L A$ and $\partial_R A$ for the above sets and L_A, R_A for the corresponding parts of X separated by A .

Remark 3.9. We will also use another description of the subdivision of ∂X . Taking a suitable base point x in R_A , we can describe $\partial_L A$ as the set of points of ∂X which can be joined by quasi-geodesic rays from x which cross A an odd number of times and $\partial_R A$ as the set of points of ∂X which can be joined by quasi-geodesic rays from x crossing A an even number of times.

4. Crossing of Tracks.

We next consider crossing of tracks in X . It is understood that all the tracks considered are two-ended and when we talk of subdivision of the boundary, the track considered has infinite cyclic stabilizer.

Definition 4.1. We say that B crosses A , if the end points of B are on different sides of A , that is, one end point of B is in $\partial_L A$ and the other in $\partial_R A$.

It is clear that if B crosses A and B' is a track with the same end points as B , then B' also crosses A . Also if A is equivalent to A' , meaning that A and A' induce the same splitting of ∂X , then B and B' cross A' . It is easy to see that crossing is not symmetric in general even if one restricts attention to axes (see the examples later in this section). This will cause us some difficulty, but it turns out that we will have symmetry in the cases in which we are interested. We note that if B crosses A , then any bi-infinite quasi-geodesic joining the end points of B crosses A an odd number of times. We can, if necessary, modify the quasi-geodesic to cross A exactly once.

We now examine the symmetry of crossing. The following proposition shows that we have symmetry in many cases.

Proposition 4.2. *Suppose that A, B are essential axes so that $e(X_H) \geq 2$ and $e(X_{H'}) \geq 2$, where $H = \text{Stab}^0 A$ and $H' = \text{Stab}^0 B$. Then A crosses B if and only if B crosses A .*

Proof. It will suffice to show that if B does not cross A , then A does not cross B . If B does not cross A , the picture in X_H is as shown in Figure 3. Remark 2.12 tells us that we can choose an essential track t' in X_H disjoint from $\pi(B)$ where π denotes the covering projection $X \rightarrow X_H$. This gives a track A' in X which is disjoint from B with the same end points as A . Thus A' cannot cross B . As A and A' have the same end points, it follows that A cannot cross B . This completes the proof of Proposition 4.2.




Figure 3.

The next proposition is crucial to our whole approach.

Proposition 4.3. *Let A be an essential axis, and suppose that for some $H \subseteq \text{Stab}^0 A$, $e(X_H) \geq 3$. Then the end points of A lie on one side of any axis B in X , i.e., A does not cross B .*

Proof. Consider $t = \pi(A)$ in X_H . There are three possibilities for the end points of $\pi(B)$: (1) Both end points are in $\partial_0 X_H$, (2) One end point is in $\partial_0 X_H$ and the other in $\partial_1 X_H$, (3) Both the end points are in $\partial_1 X_H$. (See Figure 4.)




Figure 4.

In each case, the fact that X_H has at least three ends while B has only two ends implies that there is a proper map $\phi : X_H \rightarrow \mathbb{R}$ such that the induced map from B to \mathbb{R} is not onto. Now Remark 3.5 implies that we can find a finite track s in X_H which carries a non-trivial subgroup of H and does not intersect $\pi(B)$ (see Figures 4a and 4b). This gives a track A' in X with the same end points as A which is disjoint from B .




Figure 4.a.




Figure 4.b.

It follows that A' , and hence A , does not cross B . This completes the proof.

Now we give the promised examples of asymmetry of crossing of axes.

Example 4.4. For any hyperbolic group G which is not elementary, if A is an essential axis with $H \subseteq \text{Stab}^0 A$, and if $e(X_H) \geq 3$, there is an axis B such that B crosses A but A does not cross B . This is immediate if we can find B as in case 2) in the proof of Proposition 4.3. To find such a B , we take $h \in G$ which has limit points as in case 2), (such h must exist by the density of endpoints of axes [24]), and take B to be the axis of h . By Proposition 4.2 we must have $e(X_{h^n}) = 1$ for any positive integer n . Similar examples can be constructed even when $e(G, H) \leq 2$ for all two-ended H in G . (See the definition and examples below.)

This example may give the impression that asymmetry in the crossing of axes is a surprising condition which can only occur for rather subtle reasons. The following simple example shows that asymmetry of crossing is not surprising and should have been expected.

Example 4.5. Let M be a compact orientable hyperbolic 3-manifold with an incompressible boundary component F which is not a sphere or torus. Identify the universal cover of F with the hyperbolic plane H^2 . Let C_1 and C_2 be essential simple closed curves on F such that C_1 and C_2 cannot be homotoped in F to be disjoint. Let G denote the fundamental group of M , let g_1 and g_2 denote generators of the cyclic subgroups of G carried by C_1 and C_2 and let l_1 and l_2 denote axes in H^2 for g_1 and g_2 which lie above C_1 and C_2 . The fact that C_1 and C_2 cannot be homotoped in F to be disjoint means that l_1 and l_2 must cross, in the sense of Definition 4.1.

Suppose that there is an essential annulus S_1 embedded in M with C_1 as a boundary component, and let S_2 denote an inessential annulus embedded in M with C_2 as a boundary component. This means that S_2 is parallel into F . Fix a triangulation of M such that S_1 and S_2 are normal surfaces in M . Let X_G denote the 2-skeleton of M , and let t_1 and t_2 denote the intersection of S_1 and S_2 with X_G . Thus t_1 and t_2 are tracks in X_G which contain C_1 and C_2 and hence carry the cyclic groups generated by g_1 and g_2 respectively. Let X denote the universal cover of X_G , and let A_1 and A_2 denote components of the pre-image in X of t_1 and t_2 , so that they contain l_1 and l_2 and are axes for g_1 and g_2 respectively. The fact that S_2 is inessential implies that $e(G, A_2) \leq 1$ so that A_1 cannot cross A_2 . The facts that S_1 is essential and that l_2 crosses l_1 implies that A_2 crosses A_1 .

Recall that a one-ended hyperbolic group G is said to have a multi-band or to be of multi-band type, if there is a two-ended subgroup H with $e(G, H) \geq 3$, and that G is of surface type if $e(G, H) \leq 2$ for every two-ended subgroup H . The proof of the main theorem is broken up into two cases using this division. We will prove the theorem in the multi-band case in the next section. The rest of the paper will be devoted to providing a proof in the surface type case. Here are some examples of both types. The first three are of multi-band type.

Example 4.6. Consider two orientable closed surfaces of genus ≥ 2 and identify them along a non-trivial simple loop on both. If G is the group so obtained and H the subgroup corresponding to the simple loop, then $e(G, H) = 4$ and G is hyperbolic by a theorem of Bestvina and Feighn [1].

Example 4.7. Let G_1, G_2 be hyperbolic groups, let g_1, g_2 be elements of G_1, G_2 respectively such that g_1 is indivisible in G_1 and g_2 has a root of order ≥ 3 in G_2 . Let G be the group obtained by amalgamating G_1 and G_2 along the cyclic subgroups generated by g_1 and g_2 , and let H denote this cyclic subgroup of G . Again, it is easy to verify that $e(G, H) \geq 3$ and that G is hyperbolic by [1].

Example 4.8. This is a special case of the previous example. We can take G_2 to be infinite cyclic and g_2 to be the n^{th} power of a generator of G_2 where $n \geq 3$. Of course, we take g_1 to be indivisible in G_1 and so that it does not generate the whole group.

We next construct surface type examples which are not surface groups.

Example 4.9. Let M_1, M_2 be two orientable hyperbolic 3-manifolds with incompressible boundary. We further assume that there are no essential annuli or tori in M_1, M_2 . Identify the manifolds along non-trivial embedded annuli $A_1 \subset \partial M_1, A_2 \subset \partial M_2$ to obtain M . Let G be the fundamental group of M and let H denote the subgroup corresponding to A_1, A_2 . We claim that G is of surface type. We have to show that $e(G) = 1$, and $e(G, K) \leq 2$ for any infinite cyclic K in G . It is easy to check that $e(G) = 1$, because M is irreducible with incompressible boundary. The claim about $e(G, K)$ is divided into a few cases. If X is the universal cover of M and X_K the cover corresponding to K , both X, X_K are built from various covers of M_1, M_2 . In both cases we need to consider only covers which are either simply connected or have infinite cyclic fundamental groups.

If N is any such cover we want to calculate $H_c^1(N; Z)$ and use the Mayer-Vietoris sequence to estimate $H_c^1(X_K; Z)$. We also have $H_c^1(N; Z) \cong H_2(N, \partial N; Z)$. The latter fits into the exact sequence

$$0 \rightarrow H_2(N, \partial N; Z) \rightarrow H_1(\partial N; Z) \rightarrow H_1(N; Z) \dots$$

Since each M_i is acylindrical, $H_1(\partial N; Z) \cong 0$ or Z and since M_i is boundary incompressible the map $H_1(\partial N; Z) \rightarrow H_1(N; Z)$ is injective. Thus $H_c^1(N; Z)$ is always zero in the cases we need. If N, N' are two such copies in X_K and if they intersect, they intersect along, say Y , which is an annulus or the product of an interval and a real line. Now the exact sequence

$$0 \rightarrow H_c^0(Y; Z) \rightarrow H_c^1(N \cup N'; Z) \rightarrow H_c^1(N; Z) \oplus H_c^1(N'; Z) \rightarrow H_c^1(Y; Z)$$

shows that $e(G, K) = 1$ when K is not conjugate to a subgroup of H , and that $e(G, K) = 2$ if K is conjugate to a subgroup of H .

Example 4.10. We can use a construction similar to the above to obtain a HNN-extension. This time, we start with M with similar properties to those of M_i above and identify two non-parallel annuli in the boundary of M . The proof that M is of surface type is similar.

5. Proof in the multi-band case.

We need the following variation of Theorem 2.2 of [15] stemming from the work of Stallings [19] and Dunwoody [5]. Let K be a finite simplicial 2-complex with universal cover X , and let A be a track in X with stabiliser H . Let π denote the projection $X \rightarrow X_H$ and let E, E^* be the two components of $X - A$.

Theorem 5.1. *Suppose that:*

- (a) A/H is finite.
- (b) Both the components of $X_H - A_H$ are infinite.
- (c) For every $g \in G = \pi_1(K)$, at least one of the four sets

$$gE \cap E, gE \cap E^*, gE^* \cap E, gE^* \cap E^*$$

has finite image in X_H , and:

- (d) *If two of the four sets in (c) have finite image in X_H , then at least one of them is empty.*
- (e) *The equation $gE = E^*$ does not hold for any element g of G .*

Then G splits over H .

Remark 5.2. Assumption (c) means that the axis A does not cross any of its translates. Assumption (d) means that if gA projects to a finite singular track in X_H then A and gA are disjoint or coincide. Assumption (e) means that no element of H interchanges the sides of A in X so that A/H is a two-sided track, and assumption (b) means that A/H is essential in X_H .

Ideally, we would like to prove this theorem by taking A/H to be shortest in some sense and showing that as A does not cross any of its translates, it must be disjoint from all its translates. This would immediately imply the result. However, there are some technical problems with this approach so we simply use the arguments in [15].

The theorem in [15] is stated in terms of 3-manifolds, but the proof of Theorem 2.2 in [15] carries over to a proof of Theorem 5.1 above. (The condition (d), is not explicitly stated in [15], though it is mentioned a few times that it is satisfied in the course of the proof.)

In this paragraph, we discuss the case of a general group G which can have a multi-band or be of surface type. We first examine the situation when one has two axes A, B which do not cross. We may assume by choosing appropriate transverse orientations that $N_A, P_A \in L_B$ and $N_B, P_B \in R_A$. Consider the infinite components of $X - (A \cup B)$. Let $H = \text{Stab}^0 A$ and

let $K = \text{Stab}^0 B$. Choosing a suitable base point near N_B or P_B in R_B (see Figure 9 in the proof of Lemma 8.1) we see that there is an infinite component X_1 of $X - (A \cup B)$ whose closure contains N_B, P_B but does not contain translates by K of N_A, P_A . Similarly there is a component X_2 whose closure contains N_A, P_A but does not contain translates by H of N_B, P_B . Finally, there are either one or two infinite components X_3, X_4 (one of them may be empty) such that their closures contain one of the end points of A and one of the end points of B (possibly all four). The closures of the X_i 's exhaust all of ∂X . Thus $X - \cup X_i$ is compact. With our notation, we have:

Lemma 5.3. *If A, B are axes which do not cross, then (with the above choices of transverse orientations) $L_A \cap L_B, R_A \cap R_B$ and $R_A \cap L_B$ are infinite, and $L_A \cap R_B$ is finite.*

We now assume that G has a multi-band and we want to prove Theorem 1.3 by using Theorem 5.1. We restate the result.

Theorem 1.3. *Let G be a one-ended hyperbolic group and let H_0 be a two-ended subgroup with $e(G, H_0) \geq 3$. Then G splits over a subgroup H commensurable with H_0 .*

Proof. The hypothesis that H_0 be two-ended implies that H_0 contains an infinite cyclic subgroup H_1 of finite index. Let g denote a generator of H_1 and let P and Q denote the fixed points of g in its action on ∂X . Let H_2 denote the stabiliser of $\{P, Q\}$. Then H_2 must contain H_1 as a subgroup of finite index so that H_2 also has two ends. The point of considering H_2 is that it is the unique maximal two-ended subgroup of G which contains H_0 . Let H denote the intersection of all the conjugates of H_1 in H_2 , so that H is normal in H_2 . As H_1 is of finite index in H_2 and is infinite cyclic, it follows that H is also of finite index in H_2 and infinite cyclic. Let X_i denote the quotient of X by H_i . We denote the quotient group H_2/H by K . Thus K is a finite group which acts on X_H by covering translations with quotient X_2 . As X_H is a finite cover of X_0 , and $e(X_0) \geq 3$ by assumption, it follows that $e(X_H) \geq 3$. We claim that there is a finite essential track t in X_H which is K -equivariant. This means that, for all k in K , the track kt is disjoint from or coincides with t . Let s denote a finite essential track in X_H , and let N denote a K -invariant neighbourhood of the union of all the translates of s by K . Then any component of the frontier of N in X_H is a K -equivariant finite track. If no component is essential in X_H , then all but one of the components of $X_H - N$ is finite, which would contradict the assumption that s is essential. Thus some component of the frontier of N is the required K -equivariant essential finite track t in X_H . The pre-image of t in X consists of a finite number of disjoint axes. Let A denote one of them. The K -equivariance of t implies that A is H_2 -equivariant.

Now we can verify the hypotheses of Theorem 5.1. Let h be an element of G . The fact that $e(X_H) \geq 3$ combined with Proposition 4.3 tells us that A , and hence also hA , cannot cross any axis. Hence if hA has different end points from A , then both the end points of hA are on one side of A , say in ∂R_A , and both the end points of A are on one side of hA , say in $h\partial L_A$. Lemma 5.3 tells us that exactly one of the four sets of 5.1.(c) is finite, namely $L_A \cap hR_A$. The other three sets have points near the boundary of X and thus project to unbounded sets under π . Hence exactly one of the four sets of 5.1.(c) has finite image in X_H . If hA has the same endpoints as A , then h lies in H_2 so that hA is equal to A or disjoint from A . This implies that the conditions (a-d) of 5.1 are satisfied. Condition (e) can fail only if t covers a one-sided track u in X_2 . If this occurs, we let v denote the boundary of a regular neighbourhood of u in X_2 and replace t by a parallel track t' which covers v . The new track t' satisfies condition (e) and automatically satisfies the other conditions also. This completes the proof of the theorem.

6. Orientation in surface type groups.

For the rest of this paper, we will concentrate on surface type groups. This time we cannot expect to split along a subgroup commensurable with H , as the example of the usual surface groups shows. Tukia [23] showed how to find better candidates if the original H does not work. We could combine this approach and Theorem 5.1 to complete the proof. Instead we follow the ideas of Freedman, Hass, Scott [8] and show that tracks of least complexity, like least area surfaces, result in the desired splittings.

We recall that we are considering a hyperbolic group G which is a one-ended torsion free group with $e(G, H) \leq 2$ for every infinite cyclic $H \subset G$. As mentioned above this case is very similar to the usual surface groups and we plan to adopt some of Tukia's ideas in [23]. We have the additional difficulty of orientation reversing elements which we discuss first. We start by observing that tracks have some nice properties in the case of groups of surface type.

Lemma 6.1. *If G is of surface type and $e(G, H) = 2$, and if t is a finite track in X_H , then t must be two-sided. If t is essential in X_H , then t must carry H .*

Proof. If t is one-sided, let W denote the space obtained from X_H by cutting along t . Then X_H has a double cover which is the union of two copies of W glued along a double cover of t . This double cover must have four ends which contradicts the hypothesis that G is of surface type. Hence t must be two-sided. If t is essential, then Lemma 3.1 shows that t must carry a subgroup H' of H of some finite index d . Thus t splits X_H into two infinite pieces C and D such that C carries H and D carries H' . It follows that the cover of X_H of degree d consists of a finite cover of C together with d copies

of D and so it must have at least $d + 1$ ends. The fact that G is of surface type implies that d equals 1, so that t must carry H as claimed.

Lemma 6.2. *Let G be of surface type and $e(G, \langle g \rangle) = 1$, but $e(G, \langle g^n \rangle) = 2$ for some n . Then $e(G, \langle g^{2m} \rangle) = 2$ and $e(G, \langle g^{2m+1} \rangle) = 1$.*

Proof. Let X_k denote the quotient of X by the cyclic group generated by g^k . Note that the finite cyclic group \mathbf{Z}_k acts on X_k with quotient X_1 . If X_k has two ends, the action of \mathbf{Z}_k on X_k induces an action on the two ends, yielding a homomorphism from \mathbf{Z}_k to \mathbf{Z}_2 . If this homomorphism were trivial, i.e., if the action preserved the two ends, it would follow that X_1 had two ends also. This contradiction shows that the homomorphism must be non-trivial, so that k must be even. If k is odd, we deduce that X_k must have one end. Hence if there is n such that X_n has two ends, then n is even and the analysis above shows that X_2 must also have two ends. Now it follows that X_{2m} has at least two ends, for all m , and hence that X_{2m} has exactly two ends, as G is of surface type. This proves the assertions of the lemma.

Definition 6.3. If G is of surface type and if $e(G, \langle g \rangle) = 1$ but $e(G, \langle g^n \rangle) = 2$ for some n , then we call g **orientation reversing**.

If g is not orientation reversing we call g **orientation preserving**.

In the Lemma above, we can map C onto a Mobius band and take the inverse image of the middle circle to obtain a finite one-sided pattern inside C . Any one-sided component t of this pattern will be a track in C whose pre-image in the double cover of C is essential. It follows that t carries $\langle g \rangle$. This gives an axis in X stabilized by $\langle g \rangle$ whose sides are interchanged by g . Conversely, if there is such an axis whose sides are interchanged by g , then g is clearly orientation reversing. Thus:

Lemma 6.4. *Let G be of surface type. An element $g \in G$ is orientation reversing if and only if there is an axis A in X stabilized by $\langle g \rangle$ such that the sides of A are interchanged by g .*

We will call A an axis for g if g stabilises A . We have:

Lemma 6.5. *If G is of surface type and $e(G, \langle g \rangle) = 2$, then all essential axes for g give the same separation of ∂X .*

7. Shortest tracks for groups of surface type.

It is simpler for many of the assertions that follow to assume that G is of surface type and even though some of the results of the following sections are valid more generally, **we will continue to confine ourselves to surface type groups**. Recall that Lemma 6.1 tells us that if $e(X_H) = 2$, then any finite track t in X_H is two-sided and that if t is essential in X_H then t must carry H . This is very similar to the situation of an annulus cover

of a surface as already discussed. We will use this analogy and use shortest tracks to obtain splittings in the surface type case. We will need the following existence result for shortest tracks.

Lemma 7.1. *Let G be a torsion free hyperbolic group of surface type and let H be an infinite cyclic subgroup. If $e(X_H) = 2$, there is an essential pattern t in X_H which is shortest among all such patterns, and any such pattern is normal, connected (hence a track), two-sided and carries H .*

Further, there is an infinite cyclic subgroup H of G and an essential track t in X_H such that t is shortest among all essential tracks in all quotients of X by an infinite cyclic subgroup of G .

Remark 7.2. The first part of this lemma is analogous to the second part of Lemma 2.14. There is no point discussing the analogue of the first part of Lemma 2.14, which is about the existence of a shortest pattern in a given equivalence class as all essential tracks in X_H are equivalent as X_H has two ends.

Proof. The second part of Lemma 2.14 shows that there is an essential singular pattern in X_H which is shortest among all such patterns and that any such shortest pattern is a normal embedding. Let t denote such a shortest pattern. As we remarked at the start of this section, Lemma 6.1 shows that each component of t must separate X_H . It follows that some component of t is essential. As t is shortest it follows that t equals this component so that t is connected. Of course, t must be two-sided as it is essential. Finally, we apply Lemma 6.1 again to show that t must carry H .

The last part of Lemma 7.1 is proved in much the same way as the existence result of Lemma 2.14 by projecting into X_G a sequence of tracks whose complexity approaches the infimum of all possible complexities. The fact that the tracks being considered are all essential in some quotient of X by an infinite cyclic subgroup of G ensures that some subsequence converges to a singular track in X_G , and this will yield a possibly singular track f in some X_H such that f has the required minimal complexity and is essential, and covers an embedded track t . Now the fact that f has minimal complexity implies that f must be an embedding and that t is the required track.

We will also need an analogous existence result for shortest one-sided tracks.

Lemma 7.3. *If X_H contains a one-sided track, then there is a shortest one-sided track in X_H . Further, there is an infinite cyclic subgroup H of G and a one-sided track t in X_H such that t is shortest among all one-sided tracks in all quotients of X by an infinite cyclic subgroup of G .*

Proof. This result has essentially the same proof as the existence results for shortest essential tracks in Lemma 2.14. One takes a minimising sequence of one-sided tracks in X_H . Remark 2.2 shows that we can assume that each

of these tracks is normal. The hypothesis of one-sidedness replaces the hypothesis of essentiality, and the key fact needed is that a one-sided track in a 2-complex cannot be very close to a vertex. For there is a canonical normal direction for a track close to a vertex v , in which the normal points towards v . One obtains a possibly singular one-sided track f in X_H of least possible complexity. Lemma 2.10 shows that f must be an embedding.

The second part of the lemma has essentially the same proof.

We next give analogues of results in Freedman, Hass and Scott [8] and [9]. The proofs are essentially identical. The first result is the analogue of the fact that two shortest simple closed curves on an annulus must coincide or be disjoint. It is a special case of Lemma 2.7, once one remembers that a shortest essential track in X_H is actually shortest among all essential patterns as proved in Lemma 7.1.

Lemma 7.4. *Suppose X_H has two ends and that s and t are shortest essential tracks in X_H . Then either s and t coincide or they are disjoint.*

Our next result is the analogue of the fact that a shortest loop on an annulus lifts to a shortest loop on any finite cover.

Lemma 7.5. *Suppose X_H has two ends and t is a shortest essential track in X_H . If X_K is any finite cover of X_H and t' denotes the lift of t to X_K , then t' is a shortest essential track in X_K .*

Proof. Let $p : X_K \rightarrow X_H$ be the covering projection, τ be a generator of the deck transformation group and let the order of the cover be d . Suppose there is an essential track s' in X_K with $c(s') < c(t')$. Lemma 6.1 shows that s' must carry K . Choose s' to be a shortest such track. Consider the translates of s' by the powers of τ . The preceding lemma implies that each $\tau^i s'$ must coincide with s' or be disjoint from s' . It follows that s' projects into X_H covering its image which we denote by s . Clearly s is an essential track in X_H , and so it must carry H by Lemma 6.1. It follows that each $\tau^i s'$ coincides with s' , so that s' covers s with degree d and hence $c(s') = d.c(s)$. As $c(t') = d.c(t)$, it follows that $c(s) < c(t)$. This contradiction completes the proof of the lemma.

Before continuing, we briefly consider the orientation reversing case. If g is orientation reversing, consider $p : X_{g^2} \rightarrow X_g$, let τ denote the non-identity covering translation and let t be a shortest essential track in X_{g^2} . Lemma 7.4 shows that t and $\tau(t)$ are either disjoint or coincide. In the first case $c(p(t)) = c(t)$, and in the second case $2c(p(t)) = c(t)$ and $p(t)$ is one-sided in X_g . Also recall from the discussion at the end of Section 5 that if one starts with a one-sided track t in X_g , then its pre-image in X_{g^2} must be an essential track. Thus:

Lemma 7.6. *If $c(g)$ is the minimum of $c(t)$ for one-sided tracks in X_g , and $c(g^2)$ is the minimum for essential tracks in X_{g^2} , then $2c(g) \geq c(g^2)$.*

Now we consider the analogue of the fact that if one has a shortest loop C on an annulus M then its pre-image in the universal cover \widetilde{M} of M is a length minimising line l . This means that any compact interval λ in l is the shortest path in \widetilde{M} connecting the two points which form $\partial\lambda$. The proof in [9] uses the facts that λ projects injectively into a finite cover M_1 of M and that the lift C_1 of C into M_1 is shortest and so any sub-arc is also shortest in its homotopy class. We need to replace the concept of homotopy class for our more general situation.

Let s and s' be oriented partial patterns in a 2-complex Y which have the same boundary. Thus $s \cup s'$ is naturally a singular pattern in Y . We will suppose that if we take the given transverse orientation on s and the opposite one on s' , this yields a transverse orientation on $s \cup s'$. We will say that s and s' are *equivalent* if every loop and proper map of the line into Y has zero intersection number with $s \cup s'$ equipped with this transverse orientation. Now the following result is clear.

Lemma 7.7. *Suppose that X_H has two ends and let t be a shortest essential track in X_H . If s is an oriented partial pattern contained in t which is equipped with a transverse orientation induced from a transverse orientation of t , then s is shortest among all oriented partial patterns in X_H which have the same boundary as s and are equivalent to s .*

It follows that if the pre-image of t in X is the axis A , then A is length minimising in the following sense.

Lemma 7.8. *Suppose that X_H has two ends and let t be a shortest essential track in X_H . Let the pre-image of t in X be the axis A . Then any finite pattern s contained in A is shortest among all partial patterns in X with the same boundary.*

Remark 7.9. Note that as X is simply connected, Lemma 2.10 shows that any singular pattern in X is two-sided. Thus the union of two oriented partial patterns in X with the same boundary is automatically orientable. As X has only one end, it follows that any two partial patterns in X with the same boundary are equivalent up to change of orientation so that this lemma does not need any reference to equivalence.

Proof. This follows from the preceding lemma because the finite pattern s in A projects injectively into X_{g^n} for some n .

Now we can prove the analogue of the fact that two length minimising lines in a plane must coincide, be disjoint or intersect transversely at a single point.

Lemma 7.10. *If A, B are minimal axes, then they must coincide or be disjoint, or cross. If they cross then they intersect in a finite graph such that both $A - (A \cap B)$ and $B - (A \cap B)$ consist of exactly two infinite components*

each. Further, given $\epsilon > 0$, there is an ϵ -isotopy of the track $A/\text{Stab}^0(A)$ which arranges that A and B intersect transversely in finitely many points and that both $A - (A \cap B)$ and $B - (A \cap B)$ consist of exactly two infinite components each.

Remark 7.11. This means that minimal axes in X behave very much like lines in the plane. Note that the lemma does not assert that A and B intersect transversely although we have no counterexample.

Proof. We start with the case when A and B have the same end points and hence must have a non-trivial common stabiliser H which preserves the two sides of each. The fact that A and B must coincide or be disjoint follows immediately by applying Lemma 7.4 to the quotient tracks A/H and B/H in X_H . For the rest of this proof we will assume that A and B do not have the same end points.

Suppose that A and B are distinct but intersect transversely. It is automatic that they intersect in a finite number of points. We will consider the possibility of non-transverse intersection later. If the lemma fails to hold, there must be a bounded component R of $X - (A \cup B)$. This is true whether A and B cross or not. Such a region will be bounded by compact pieces of A and B which will be partial patterns. In the case of lines intersecting in a plane, one can find R such that its boundary meets each of A and B in a connected set, but there seems no reason why this should be possible in general. However, we can still make what is in essence the usual cut and paste argument. We will perform cut and paste at each point of $A \cap B$ which lies in the boundary of R . At each such point one locally sees four regions exactly one of which is part of R . We choose the cut and paste which connects this region to the opposite one. We know that cut and paste operations reduce complexity but this does not mean much as both A and B have infinite complexity. We get round this problem as follows. Let A_+ and A_- denote the two infinite pieces of $A - (A \cap B)$, and similarly for B_+ and B_- . Let A_1 denote the partial pattern obtained by truncating A in A_+ and A_- and describe B_1 similarly. Lemma 7.8 tells us that A_1 and B_1 are each the shortest patterns in X with their boundary. The result of our cut and pastes is to replace A_1 and B_1 by A'_1 and B'_1 , where A'_1 is obtained from A_1 by removing $A \cap R$ and replacing it with $B \cap R$, and B'_1 is obtained similarly. Now we can say that $c(A'_1) + c(B'_1) < c(A_1) + c(B_1)$. This contradicts the fact that A_1 and B_1 are each the shortest partial patterns in X with their boundary, completing the proof of the lemma in the case when A and B intersect transversely.

If A and B have non-transverse intersection, we want to apply the Meeks-Yau trick as in the proof of Lemmas 2.7 and 2.8. If $X - (A \cup B)$ has a bounded component, we can use this trick and the arguments of the preceding paragraph to obtain a contradiction. Thus $X - (A \cup B)$ cannot have a bounded

component. Recall from the discussion in the proof of Lemma 2.7 that each extreme point of the finite graph $A \cap B$ is a local crossing point of A and B . This means that any neighbourhood of such a point contains points of A on each side of B and vice versa. If A and B do not cross, but do intersect, the local crossing property implies that A has points on both sides of B which in turn implies that $X - (A \cup B)$ has a finite component, which we know cannot occur. We conclude that if A and B do not cross, then they must be disjoint. If A and B do cross, the fact that there cannot be a finite component of $X - (A \cup B)$ implies that each of $A - (A \cap B)$ and $B - (A \cap B)$ cannot have a finite component, so that each consists of exactly two infinite components as required.

For the last part of the statement of the lemma, we consider the finite two-sided track $A/\text{Stab}^0(A)$ in $X/\text{Stab}^0(A)$. Denote the track by t_1 and the quotient of X by X_1 . As t_1 is two-sided in X_1 , it has a neighbourhood homeomorphic to $t_1 \times I$. This defines an isotopy of t_1 which moves it to a parallel copy, and all the nearby parallel copies will automatically be transverse to the image of B in X_1 , so that the new version of A will intersect B transversely in a finite set. Applying the Meeks-Yau trick again implies that $A \cap B$ must separate each of A and B into exactly two infinite pieces as required, so long as we move t_1 a small enough distance.

8. Subdivisions of the boundary of X .

We consider again subdivisions of the boundary, this time with respect to various crossing essential axes in the surface type case. We have already noted that if A is an essential axis in X , then $\partial L_A, \partial R_A$ can be described in terms of quasi-geodesic rays from some base point x . For example, in Figure 9 below,




Figure 9.




Figure 10.

∂L_A consists of points of the boundary for which the quasi-geodesic rays (hereafter called rays) joining them to x cross A an odd number of times and in fact we can even find rays which cross exactly once since A is connected. Similarly ∂R_A is described as the part of the boundary for which the rays cross an even number of times. We will use the convention that an arrow representing a transverse orientation of A always points towards to R_A . Since X has one end we can require the rays to avoid compact sets in X . Suppose C is a compact set in X such that $A - (A \cap C)$ has two infinite components, (there are automatically at most two). Let these be E_1, E_2 with $N_A \in E_1$ and $P_A \in E_2$. If R is a ray which avoids C and joins x to a point of ∂L_A , then the intersection of R with A is in $E_1 \cup E_2$. We can now further modify R to obtain a ray which intersects A exactly once. We next observe that L_A, R_A have exactly one end each. Thus:

Lemma 8.1. *We can join x to a point of ∂R_A by a ray which avoids any given compact set in X and does not intersect A . Similarly we can join x to a point of ∂L_A , by a ray which avoids a given compact set C and meets A exactly once either in E_1 or E_2 as required.*

Next suppose that two axes A and B cross. We can assume that P_B lies in ∂R_A by changing our choice of generator of $\text{Stab}^0 B$. Figure 10 shows the picture for one choice of transverse orientation of B . Given this choice, we subdivide ∂X into four sets $[P_A, P_B], [P_B, N_A], [N_A, N_B], [N_B, P_A]$ as follows: $[P_A, P_B] = \partial R_A \cap \partial R_B$, $(P_A, P_B) = (\partial R_A \cap \partial R_B) - \{P_A, P_B\}$ etc. Note that if we change the transverse orientation on B , then $[P_A, P_B] = \partial R_A \cap \partial L_B$. Choose a base point x in X which lies in R_A close to P_A and does not lie on B . Then (P_A, P_B) can be described as the set of points y in ∂X which can be joined to x by rays which cross both A and B an even number of times and $[P_A, P_B]$ is its closure. Now join x to y by a ray r which does not intersect A , and so lies in R_A . Such a ray, if it intersects B at all, will intersect B in the infinite component of $B - (A \cap B)$ in R_A an even number of times and thus can be replaced by a ray which does not intersect either A or B . If y_1, y_2 are two points of (P_A, P_B) , then the union of two such rays r_1, r_2 is a bi-infinite path joining y_1, y_2 which does not intersect $A \cup B$. Thus (P_A, P_B) is in the closure of one infinite component of $X - (A \cup B)$. Hence:

Lemma 8.2. *If A, B cross then $X - (A \cup B)$ has exactly four infinite components and their closures intersect ∂X in the subsets*

$$[P_A, P_B], [P_B, N_A], [N_A, N_B], [N_B, P_A].$$

Following Tukia, we sometimes denote these subsets of ∂X by I_1, I_2, I_3, I_4 in that order. See Figure 10. We next consider the situation when there is an axis C with $P_C \in (N_A, P_B)$, $N_C \in (N_B, P_A)$ and observe that we obtain subdivisions of ∂X as in the case of a circle. (See Figure 11 with transverse orientations as shown.)




Figure 11.

We have:

Lemma 8.3. *Suppose that C is an axis with $P_C \in (N_A, P_B)$, $N_C \in (N_B, P_A)$. Then we have with suitable choice of transverse orientation for C :*

- (1) $\partial R_C \supset [P_A, P_B], \partial L_C \supset [N_A, N_B]$,
- (2) $[N_B, P_A] = [N_C, P_A] \cup [N_B, N_C], [N_C, P_A] \cap [N_B, N_C] = \{N_C\}$,
- (3) $[N_A, P_B] = [N_A, P_C] \cup [P_C, P_B], [N_A, P_C] \cap [P_C, P_B] = \{P_C\}$.

Proof. We now take a base point x near P_A in L_A and a ray R joining x to a point of (P_A, P_B) . We assume that R avoids all the intersections of A, B, C and does not meet B and meets A exactly once near P_A . If R meets C an odd number of times, we modify R near the ends of C to R' so that R' meets C near P_C or N_C and still intersects A once and does not meet B . Since R' crosses A , R' must meet C near N_C . This means that N_C is in R_A and it is in R_B since R' does not meet B . Hence N_C should be in $R_A \cap R_B$ which contradicts the hypothesis that N_C is in (N_B, P_A) . This proves that $[P_A, P_B]$ is on one side of C . Similarly we conclude that $[N_A, N_B]$ is also on one side of C . We now choose transverse orientation on C so that $[N_A, N_B] \subset \partial L_C$ and $[P_A, P_B] \subset \partial R_C$. Since $\partial L_C \cap [P_A, P_B] = \emptyset$, we have

$$\begin{aligned} \partial L_C &= (\partial L_C \cap [N_A, N_B]) \cup (\partial L_C \cap (N_A, P_B)) \cup (\partial L_C \cap (N_B, P_A)) \\ &= [N_A, N_B] \cup (\partial L_C \cap (N_A, P_B)) \cup (\partial L_C \cap (N_B, P_A)). \end{aligned}$$

Since (N_A, P_B) is in ∂L_B , we have

$$\partial L_C \cap \partial R_B = (\partial R_B \cap [N_A, N_B]) \cup (\partial L_C \cap (N_B, P_A)) = (\partial L_C \cap [N_B, P_A]).$$

Hence $[N_B, N_C] \subset [N_B, P_A]$. Since $(P_A, P_B) \subset \partial R_C$, we have $[N_A, P_C] = \partial L_C \cap \partial R_A = \partial L_C \cap [N_A, P_B]$. Thus $[N_A, P_C] \subset [N_A, P_B]$. The other inclusions of Lemma 8.3 are proved similarly.

We note a consequence of the above lemma. Let $g \in \text{Stab}^0 A$ so that P_A is the attractive fixed point for g . We must have the end points of gB in $\partial(R_B)$. For otherwise, let $C = gB$. If C is as in Lemma 8.3, we see that $g[N_A, P_B] = [N_A, P_C]$ so that $g^i(P_B) \in [N_A, P_B]$ for all positive i contradicting the assumption that P_A is the positive fixed point of g . If $C = gB$ crosses B in the other direction, i.e., $P_C \in (P_A, P_B)$, $N_C \in (N_A, N_B)$, we can interchange the roles of B and C , and again arrive at a contradiction. Thus:

Corollary 8.4. *If A, B are essential axes which cross and if $g \in \text{Stab}^0 A$, then the end points of gB are in $\partial(R_B)$, the half containing P_A , the positive fixed point of g .*

Up to this point, we have discussed only one choice of configuration for crossing axes A and B as shown in Figures 10 and 11. If B equals hA , and we choose a transverse orientation for A and the induced one for B , then there are four possible configurations as shown in Figures 12a, 12b, 12c and 12d.




Figure 12.a.




Figure 12.b.




Figure 12.c.




Figure 12.d.

Note that configurations 12c and 12d can be obtained from 12a and 12b by interchanging A and B and replacing h by h^{-1} , so that we need only consider the configurations in 12a and 12b. We will say that hA crosses A in an *orientation preserving fashion* in case 12a and in an *orientation reversing fashion* in case 12b. More formally, in the first case $\{P_B, N_A\}$ (or $\{P_A, N_B\}$ in case 12c) is on the right or positive side of both A and B and in the second case $\{P_B, P_A\}$ (or $\{N_B, N_A\}$ in case 12d) is on the positive side of both A and B . We will use the abbreviations o.r. and o.p. We want to show that if the crossing is o.p. then h is o.p. Let A_h denote an axis for h . First observe that the fixed points of h must be in $I_2 \cup I_4$ as each must lie on the positive side of both A and hA or on the negative side of both. If A_h is not an essential axis, then h is o.p. by definition. So we may assume that A_h is essential. If both the fixed points of h lie in I_2 or if both lie in I_4 , then the endpoints of A and B lie on one side of A_h , so that h does not interchange the sides of A_h and h must be o.p. If one of the fixed points is in I_2 and the other in I_4 , then A_h crosses A . Since A_h is an essential axis, crossing is symmetric so that A crosses A_h . If h is o.p., the above corollary shows that $B = hA$ cannot cross A , a contradiction. If h is o.r., we also have a contradiction as P_A and $P_B = hP_A$ lie on the same side of A_h . Thus:

Lemma 8.5. *If hA crosses A in an o.p. fashion, then h is o.p. and the fixed points of h are in $[N_A, P_B] \cup [N_B, P_A]$. If h has an essential axis, then the fixed points of h are in either $[N_A, P_B]$ or in $[N_B, P_A]$ (that is, I_2 or I_4).*

We will prove later on (the main difficulty lies in showing that h has an essential axis) that if the crossing is o.r. then h is o.r. For the moment we will prove the following lemma.

Lemma 8.6. *If hA crosses A in an o.r. fashion, then the fixed points of h are in $I_1 \cup I_3$, one in I_1 and the other in I_3 .*

Proof. The fixed points of h must lie in $I_1 \cup I_3$, because if one of them is on the positive (negative) side of A it must lie on the negative (positive) side of hA . We consider the action of h on ∂X . The sets $[N_A, N_B]$ and $[N_B, N_C]$ must be one contained in the other. Thus h or h^{-1} sends one of these sets into itself and so this set must contain one of the fixed point of h . In particular h has a fixed point in I_3 . Similarly h must have a fixed point in I_1 .




Figure 13.

Now we consider another axis D which, like C in Lemma 8.3, has $N_D \in (N_B, P_A)$ and $P_D \in (N_A, P_B)$. See Figure 13. Then Lemma 8.3 tells us that

$$[N_B, P_A] = [N_B, N_C] \cup [N_C, P_A] = [N_B, N_D] \cup [N_D, P_A].$$

If $N_D \in (N_C, P_A)$, then N_D is above C as in Figure 13. If both endpoints of D lie above C then clearly $[N_B, N_C] \subset [N_B, N_D]$. In this case we use the notation $[N_C, N_D]$ for the closure of $[N_B, N_D] - [N_B, N_C]$. If D crosses C , then replacing A by D in Lemma 8.3, we see that again we have $[N_B, N_C] \subset [N_B, N_D]$. Similar statements hold if we reverse the roles of P_D and N_D . Thus:

Lemma 8.7. *If $D_i, 1 \leq i \leq n$ are a finite number of axes with one end point in $[N_B, P_A]$ and the other in $[N_A, P_B]$, then there is a D_i nearest to P_A in the sense that $[V_i, P_A]$ does not contain any of the end points of the other D_i 's, where V_i denotes the end point of D_i which lies in $[N_B, P_A]$.*

Remark 8.8. We apply the above lemma in the following situation. If $B = gA$ and A, B cross, we consider translates D of A with one end point

in (N_B, P_A) and the other in (N_A, P_B) . Freden [7] and Tukia [24] proved that G acts as a convergence group on ∂X . It follows as in Tukia [23] that there are only finitely many such D .

Following Tukia, we call any translate of A an A -axis. We also introduce the notion of canonical triple. We look at the situation when A and an A -axis B cross. Suppose that $B = gA$ and that the configuration is as in Figures 12a or 12b. Then $C = gB$ and B cross. We can vary g by an element in $\text{Stab}^0 B$ so that B is unchanged but $C = gB$ is moved up and down. As P_B lies in $\partial_R A$, it follows that P_C lies in $\partial_R B$. Whether C crosses A or not we can talk of the region $[N_A, P_C]$ in the first case and $[N_A, N_C]$ in the second case after moving the appropriate end point of C above A . We call the triple $(A, B = gA, C = gB)$ a **canonical triple** if $[N_A, P_C]$ (or $[N_A, N_C]$ in the o.r. case) does not contain the end point of any hC for $h \in \text{Stab}^0 B$. See Figures 14a and 14b.




Figure 14.a.




Figure 14.b.

We can always change g by an element of $\text{Stab}^0 B$ to achieve this, by using Lemma 8.7 above. We call a canonical triple a **good canonical triple** if in addition C is disjoint from A . (See Figures 15a and 15b.)




Figure 15.a.




Figure 15.b.

We will show that if an axis crosses its translates, then we can obtain good canonical triples starting with that axis. To handle the o.r. case, we need a lemma which will be proved in the next section.

Lemma 8.9. *If gA crosses A for some g , then there is a good canonical triple (A, D, E) .*

Proof. We proceed much as in Tukia [23]. Let $B = gA$ such that the triple (A, B, gB) is a canonical triple. If gA crosses A in an o.r. fashion, then Lemma 9.3 in the next section implies that g is o.r. and that an axis for g must cross A . Now Corollary 8.4 implies that g^2A and A must be disjoint, so that (A, B, gB) is good, and the lemma follows.

Otherwise gA crosses A in an o.p. fashion and we consider translates D of A such that N_D lies in $[N_B, P_A]$ and P_D lies in $[N_A, P_B]$. Note that gB is one such. Lemma 8.7 tells us that we can take such a D for which N_D is nearest to P_A . This means that there is h in G such that $D = hA$ and $(A, D = hA, E = h^2A)$ is a canonical triple. If h is o.p., then $P_E \in [N_A, P_D] \subset [N_A, P_B]$ and by the choice of D , N_E cannot be in $[N_D, P_A]$. Thus (A, D, E) is a good canonical triple. If h is o.r. then as in the preceding paragraph, E cannot cross A . Thus we again have a good canonical triple.

In the above situation if gA is o.r., then if necessary we can modify h by gA to obtain a new h which is o.r. Thus:

Corollary 8.10. *With the notation of the above lemma, if g_A is o.r. we can obtain a good canonical triple $(A, D = hA, hD)$ for which h is o.r.*

9. Construction of axes using good canonical triples.

We now describe a combinatorial version of a crucial construction of Tukia [23]. Tukia showed how given an element g of G such that $e(X, \langle g \rangle) = 2$ but G does not split over $\langle g \rangle$ one can find a “better” element. His improvement is in terms of intersection numbers. We could have adopted his procedure; instead we use shortest tracks and the least area ideas of Freedman, Hass, and Scott, [8] and [9], to obtain a splitting with the first choice of g . The proof that this works is based on Tukia’s idea of improving intersection numbers. The construction is also needed to show that if hA crosses A in an o.r. fashion, then h is o.r. To have clear pictures we want axes to be minimal (i.e., covering a shortest finite track) and by the results of §7, we can do this in the case when the axis stabiliser is o.p. Recall from Lemma 7.10, that minimal axes in X intersect like lines. This means that if A, B are minimal axes, then they are disjoint, or coincide, or cross. Further if they cross then they intersect in a finite graph and both $A - (A \cap B)$ and $B - (A \cap B)$ consist of exactly two infinite components each.

We now start with an essential axis A and let g_A be a generator of $\text{Stab}^0 A$. We assume that A is minimal. Consider the situation when $(A, gA, g^2A) = (A, B, C)$ is similar to a good canonical triple in the sense that A, gA cross but A, g^2A do not, but we do not assume that this triple is actually canonical. If g is o.p. with transverse orientations as shown in Fig. 12a, then $P_C \in [N_A, P_B] = I_2$ and $N_C \in [P_A, N_B] = I_1$. Recall from Lemma 8.5 that the fixed points of g lie in $I_2 \cup I_4$. However, no fixed point can lie in I_4 as all points of I_4 lie on the negative side of A and the positive side of C . Thus both the fixed points of g must lie in I_2 and in particular they lie on the positive side of A and all its translates. Denote $g^i A$ by A_i ; so that $A = A_0, B = A_1, C = A_2$. We will show now that A intersects A_i if and only if i equals 0, 1 or -1 . For otherwise, let k be the least integer greater than 1 such that A intersects A_k . Then the set of all points in X which lie on the positive side of each of A_0, \dots, A_k is compact, contradicting the fact that both the fixed points of g lie on the positive side of A and all its translates. This allows us to construct an axis for g as follows. By the lemmas above, one of the two infinite components of $A - A \cap B$ together with one of the infinite components of $B - A \cap B$ bounds the infinite component of $X - (A \cup B)$ which contains I_2 in its boundary. We denote this pattern by $A \# B = A_0 \# A_1$. Inductively we construct (see Figure 16) $A_{-n} \# \dots \# A_0 \# \dots \# A_n = L_n$, say. Since the collection $g^n A$ is locally finite, we obtain a limit pattern A_g of L_n such that $gA_g = A_g$ and the quotient of A_g by the action of g is a finite pattern t_g in X_g . We claim that some component of t_g is an essential track, so that, in particular, $\langle g \rangle$ has two co-ends.




Figure 16.

Recall that we are assuming that g is o.p. Thus if no component of t_g is essential in X_g , then every component of t_g bounds a compact subset of X_g . If we remove these compact sets from X_g and take the closure Y_g , then Y_g carries $\langle g \rangle$. This is because if a component of t_g carries the trivial group, then it bounds a compact subset of X_g which also carries the trivial group, by Lemma 3.1. Thus the pre-image of Y_g in X is its universal cover Y , and the number of ends of Y and of X are the same. Recall that the pattern A_g intersects A in the part of A between A_1 and A_{-1} . Now take a large enough N so that $g_A^N A_g$ is to the right of B . Then the portion of A between B and $g^N B$ is compact and divides Y into two infinite parts, which contradicts the assumption that X and hence Y has only one end. Thus we have:

Lemma 9.1. *If A is a minimal essential axis, if g is o.p. and if A, gA intersect but $A, g^2 A$ do not, then $\langle g \rangle$ has two co-ends.*

This is one place where the use of minimal axes seems convenient even though the lemma itself can be formulated purely algebraically. It should be possible to prove this lemma by more algebraic means and avoid minimal axes completely and proceed with intersection numbers as in Tukia [23]. The case of o.r. elements is somewhat easier.

Lemma 9.2. *If A is a minimal essential axis, if A and gA have o.r. crossing, and if $A, g^2 A$ do not cross, then g is o.r. and has fixed points in I_1 and I_3 , one in each.*

The construction is similar, but in this case the two sides of the boundary into which A_g separates ∂X are clearly non-empty, as they are $\bigcup g^{2i} [N_{-1}, P_A]$ and $\bigcup g^{2i+1} [N_{-1}, P_A]$ respectively, where P_{-1} and N_{-1} denote the positive and negative endpoints of A_{-1} . See Figure 17.




Figure 17.

We use this to prove:

Lemma 9.3. *If hA crosses A in an o.r. fashion, then h is o.r.*

Proof. We will assume that A is minimal. As the fixed points of h in ∂X must be distinct from the end points of A , there is a greatest power h^n of h such that $h^n A$ crosses A . If n is even, then this crossing must be o.p. As $h^{2n} A$ does not cross A , Lemma 9.1 shows that h^n has an essential axis. It follows that the same is true for h . Now it follows that h must be o.r. If n is odd, then the crossing of A and $h^n A$ must be o.r. and Lemma 9.2 shows that h^n is o.r. which also implies that h is o.r. This completes the proof of Lemma 9.3.

We will need the following consequence in the next section. This easily follows from the above since h and hence h^2 has an essential axis.

Corollary 9.4. *If A is a minimal essential axis and hA crosses A in an o.r. fashion, then $h^2 A$ is disjoint from A .*

Whether A is minimal or not, as long as it is essential, by replacing A by a minimal axis with the same end points we conclude:

Corollary 9.5. *If A is an essential axis and hA crosses A in an o.r. fashion, then the end points of $h^2 A$ are on the same side of A .*

10. Proof of the annulus theorem in the surface type case.

We start on the proof of the annulus theorem in the surface type case. We first consider the case when G has o.r. elements and prove the following special case of the annulus theorem.

Lemma 10.1. *Let G be a torsion free hyperbolic group with one end and suppose that G is of surface type. If G contains an o.r. element, then there is an o.r. element g of G such that G splits over $\langle g^2 \rangle$ as an amalgamated free product with $\langle g \rangle$ as one of the two factor groups.*

Proof. We choose g to be an o.r. element of G such that $c(g)$ is minimal, where $c(g)$ is the complexity of a shortest one-sided track s in X_g . Note that g will be indivisible. Let S denote the corresponding axis of g . Also let t be a shortest essential track in X_{g^2} , and recall from Lemma 7.6 that $2c(g) \geq c(g^2)$ where $c(g^2)$ is the complexity of t . Let A denote the axis of g^2 corresponding to t . Note that S and A are both stabilised by g^2 and so have the same end points in ∂X . Also g stabilises S but may not stabilise A , and A is minimal but S need not be minimal. If S and its translates never cross, then Theorem 5.1 shows that G must split over a subgroup commensurable with $\langle g \rangle$, and one can then show that G has the required splitting over $\langle g^2 \rangle$. However, as G is of surface type, we can use the properties of minimal axes to give a much simpler proof. As S and its translates never cross, the same holds for A and its translates. As A is minimal, Lemma 7.10 shows that A is disjoint from or coincides with each translate. It follows that A covers a finite track u in X_G which must carry $\langle g \rangle$ or $\langle g^2 \rangle$. If u carries $\langle g \rangle$ then g stabilises A . In this case, the projection of A into X_g must be a one-sided track so that u also must be one-sided. We replace u by the boundary of a regular neighbourhood in X_G . This will be a two-sided track v carrying $\langle g^2 \rangle$ and v separates X_G into two pieces one of which is a regular neighbourhood of u and so carries $\langle g \rangle$. It follows at once that G has the required splitting. Now suppose that u carries $\langle g^2 \rangle$. Thus u lifts to X_{g^2} and projects to an embedded two-sided track in X_g which we denote by u' . The region of X_{g^2} between u and its translate under the covering involution projects to a compact connected subset Z of X_g which carries $\langle g \rangle$ and is bounded by u' . We claim that Z projects into X_G by a homeomorphism so that u bounds a copy of Z in X_G . Again this implies that G has the required splitting. To prove the claim, let $p : X_g \rightarrow X_G$ denote the covering projection and consider $p^{-1}(u) \cap Z$. This consists of u' and possibly other finite covers of u , contained completely in Z . Let v denote a finite cover of u contained in Z . It must carry $\langle g^n \rangle$ for some positive n and its pre-image in X must be a translate kA of A which is stabilised by $\langle g^n \rangle$. It follows that kg^2k^{-1} equals g^n or g^{-n} . This is only possible if $n = 2$ and k is a power of g , as G is word hyperbolic. This implies that v equals u' , so that $p^{-1}(u) \cap Z = u'$. Let Σ denote the subset $\{z \in Z : \exists y \in Z, y \neq z, p(y) = p(z)\}$ of Z . Then Σ must be disjoint from u' . It follows that Σ is open and closed in Z , and hence equals Z or is empty. The fact that Σ is disjoint from u' implies that Σ must be empty, so that Z must project into X_G by a homeomorphism as claimed.

Now we consider the case where some translate of A crosses A . We will also assume that the intersection of any two translates of A is transverse. The general case can again be handled by using the Meeks-Yau trick. Some translate of S must also cross S , so that Corollary 8.10 tells us that there is a good canonical triple (S, hS, h^2S) with h o.r. It follows that there is a good canonical triple $(A, D = hA, hD)$ with h o.r. This means that there is no translate E of A having $P_E \in (N_D, P_A)$ and $N_E \in (N_A, P_D)$. In particular the axes $(g_D^{2m})hD$ do not cross A , for any value of m , where $g_D = hgh^{-1}$. Choose m to be the least value such that the end points of $(g_D^{2m})hD$ lie above A and replace h by $(g_D^{2m})h$. This does not change D , so we obtain a new triple $(A, D = hA, hD)$ and hD still does not cross A . We have the end points of hD above A and now the end points of $g_D^{-2}hD$ lie below A (see Figure 18).




Figure 18.

Let k denote $g_D^{-2}h$ and consider the triples (A, D, hD) and (kD, D, A) . The second triple is also like a good canonical triple since $k^{-1}D = A$, but it need not be canonical. Also kD and D cross o.r. so that k must be o.r. If D_1 denotes the portion of D between A and hD and D_2 the portion of D between $g_D^{-2}hD$ and A , we have $c(D_1) + c(D_2) = c(g_D^2) = c(g^2)$. Note that the proof of Lemma 9.2 implies that the image of D_1 in X_h contains a one-sided track carrying $\langle h \rangle$, and the image of D_2 in X_k contains a one-sided track carrying $\langle k \rangle$. Thus either $c(h)$ or $c(k)$ is, by rounding off corners, strictly less than half of $c(g^2)$. This contradicts our choice of $c(g)$. It follows that the translates of A cannot cross, so that G splits over $\langle g^2 \rangle$ as claimed. This completes the proof of the Annulus Theorem when G contains o.r. elements.

In order to complete our proof of the Annulus Theorem, we consider the case when G has no orientation reversing elements. In this case we choose an element g of G such that $c(g)$ is minimal among all elements of G with two co-ends, and let A denote the corresponding minimal essential axis for g . If hA intersects A for some h , then by the results of §8, we can choose h so that (A, hA, h^2A) is a good canonical triple. (See Figure 19. This is similar to Figure 5 in Tukia [23].)




Figure 19.

Note that h is automatically o.p. Now consider $g_B^{-1}C = D$. Since (A, B, C) is a good canonical triple $g_B^{-1}P_C = P_D$ is automatically below A . If N_D is also below A , we can argue as in the proof of Lemma 10.1, that either $c(h)$ or $c(g_B^{-1}h)$ is strictly less than $c(g)$, contradicting the choice of g . If N_D is above A we arrive at a contradiction as follows. Consider the fixed points of $g_B^{-1}h$. Since $B = hA = g_B^{-1}hA$, the fixed points of $g_B^{-1}h$ should be in $[N_A, P_B] \cup [N_B, P_A]$, by Lemma 8.5. Similarly, since $D = g_B^{-1}C = g_B^{-1}hB$, the fixed points of $g_B^{-1}h$ should be in $[P_D, N_B] \cup [N_D, P_B]$. But these sets are disjoint and this contradiction completes the proof of the annulus theorem in the surface type case.

References

- [1] M. Bestvina and M. Feighn, *A combination theorem for negatively curved groups*, Journal of Differential Geometry, **35** (1992), 85-101.
- [2] B. Bowditch, *Cut points and canonical splittings of hyperbolic groups*, Acta Math., **180** (1998), 145-186.

- [3] ———, *Group actions on trees and dendrons*, *Topology*, **37** (1998), 1275-1298.
- [4] M. Coornaert, *Sur les groupes proprement discontinus d'isometries des espaces hyperboliques au sens de Gromov*, Thesis, Universite Louis Pasteur, Strasbourg, 1990.
- [5] M.J. Dunwoody, *Accessibility and groups of cohomological dimension one*, *Proc. London Math. Soc.*, **38** (1979), 193-215.
- [6] ———, *The accessibility of finitely presented groups*, *Inv. Math.*, **81** (1985), 449-457.
- [7] E. Freden, *Negatively curved groups have convergence property*, *Ann. Acad. Sci. Fenn. Ser. A Math.*, **20** (1995), 333-348.
- [8] M.H. Freedman, J. Hass and G.P. Scott, *Least area incompressible surfaces in 3-manifolds*, *Inv. Math.*, **71** (1983), 609-642.
- [9] ———, *Closed Geodesics on Surfaces*, *Bull. London Math. Soc.*, **14** (1982), 385-391.
- [10] D. Gabai, *Convergence groups are Fuchsian groups*, *Annals of Maths.*, **136** (1992), 447-510.
- [11] E. Ghys and P. de la Harpe, *Sur les groupes hyperboliques d'apres Mikhail Gromov*, *Progress in Maths.*, No. 83, Birkhauser, (1990).
- [12] W. Jaco and J.H. Rubinstein, *PL minimal surfaces in 3-manifolds*, *J. Diff. Geom.*, **27** (1988), 493-524.
- [13] F. Paulin, *Outer automorphisms of hyperbolic groups and small actions on R-trees*, in 'Arboreal Group Theory' (edited by R. Alperin), *MSRI Publications*, **18** (1991), 331-343.
- [14] E. Rips and Z. Sela, *Structure and rigidity of hyperbolic groups*, *Geom. and Funct. Analysis*, **4** (1994), 337-371.
- [15] G.P. Scott, *A new proof of the annulus and torus theorems*, *American J. of Math.*, **102** (1980), 241-277.
- [16] ———, *Subgroups of surface groups are almost geometric*, *Journal of L.M.S.*, **17** (1978), 555-565.
- [17] ———, *Ends of pairs of groups*, *J. of Pure and Applied Algebra*, **11** (1977), 179-198.
- [18] Z. Sela, *Structure and rigidity in (Gromov) hyperbolic groups and discrete groups in rank 1 Lie groups II*, *Geom. Funct. Anal.*, **7** (1997), 561-594.
- [19] J.R. Stallings, *Group theory and 3-dimensional manifolds*, Yale University Press, New Haven, 1971.
- [20] G.A. Swarup, *On ends of pairs of groups*, *J. of Pure and Applied Algebra*, **11** (1993), 179-198.
- [21] ———, *A note on accessibility*, in 'Geometric Group Theory' (Edited by G. Niblo and M. Roller), Cambridge University Press, (1993), 204-207.
- [22] ———, *On the cut point conjecture*, *Electronic Res. Announc. Amer. Math. Soc.*, **2** (1996), 98-100.
- [23] P. Tukia, *Homeomorphic conjugates of Fuchsian groups*, *J. Reine Angew. Math.*, **391** (1988), 1-54.

- [24] ———, *Convergence groups and Gromov's metric hyperbolic spaces*, New Zealand J. Math., **23** (1994), 157-187.

Received July 25, 1998.

UNIVERSITY OF MICHIGAN
ANN ARBOR, MI 48109-1109
E-mail address: pscott@math.lsa.umich.edu

UNIVERSITY OF MELBOURNE
PARKVILLE VICTORIA 3052
AUSTRALIA
E-mail address: gadde@ms.unimelb.edu.au