

*Pacific
Journal of
Mathematics*

CAUSAL COMPACTIFICATION AND HARDY SPACES
FOR SPACES OF HERMITIAN TYPE

FRANK BETTEN AND GESTUR ÓLAFSSON

Volume 200 No. 2

October 2001

CAUSAL COMPACTIFICATION AND HARDY SPACES FOR SPACES OF HERMITIAN TYPE

FRANK BETTEN AND GESTUR ÓLAFSSON

Let G/H be a compactly causal symmetric space with causal compactification $\Phi : G/H \rightarrow \check{S}_1$, where \check{S}_1 is the Bergman-Šilov boundary of a tube type domain G_1/K_1 . The Hardy space $H_2(C)$ of G/H is the space of holomorphic functions on a domain $\Xi(C^\circ) \subset G_{\mathbb{C}}/H_{\mathbb{C}}$ with L^2 -boundary values on G/H . We extend Φ to imbed $\Xi(C^\circ)$ into G_1/K_1 , such that $\Xi(C^\circ) = \{z \in G_1/K_1 \mid \psi_m(z) \neq 0\}$, with ψ_m explicitly known. We use this to construct an isometry I of the classical Hardy space H_{cl} on G_1/K_1 into $H_2(C)$ or into a Hardy space $\widetilde{H}_2(C)$ defined on a covering $\widetilde{\Xi}(C^\circ)$ of $\Xi(C^\circ)$. We describe the image of I in terms of the highest weight modulus occurring in the decomposition of the Hardy space.

1. Introduction.

Hardy spaces on tube type domains $T_\Omega = \mathbb{R}^n + i\Omega$, associated to a homogeneous self dual cone $\Omega \subset \mathbb{R}^n$, are important objects in analysis. The Hardy space H_{cl} is by definition the space of holomorphic functions on T_Ω , such that the Hardy norm

$$\|f\|_2^2 := \sup_{y \in \Omega} \int_{\mathbb{R}^n} |f(x + iy)|^2 dx$$

is finite ([SW71, FK94]). The *boundary value map* $\beta : H_{cl} \rightarrow L^2(\mathbb{R}^n)$ is given by

$$\beta(f)(x) = \lim_{y \rightarrow 0} f(x + iy),$$

the limit taken in the $L^2(\mathbb{R}^n)$ -norm. The image of β is described by the *positivity condition*

$$\text{Im}(\beta) = \mathcal{F}(L^2(\Omega)),$$

where \mathcal{F} is the Fourier transform and $L^2(\Omega)$ the space of L^2 -functions supported on $\Omega \subset \mathbb{R}^n$. The evaluation map $H_{cl} \ni f \mapsto f(w) \in \mathbb{C}$, $w \in T_\Omega$, is continuous, and thus given by an element $K_w \in H_{cl}$. The function $K(z, w) := K_w(z)$ is the *Cauchy kernel* associated to the tube domain T_Ω . The Cauchy kernel is determined by a function of one variable $K(z, w) =$

$K(z - \bar{w})$, where $K(z)$ is the Laplace transform of the characteristic function of the cone. Finally the inverse of the boundary value map is

$$f(z) = \int_{\mathbb{R}^n} \beta(f)(y) K(z - y) dy.$$

For us the tube domain T_Ω is always the unbounded realization of a Hermitian symmetric space G_1/K_1 such that H_{cl} is also a G_1 -representation space ([FK94]).

The notation of Hardy spaces was generalized to compactly causal symmetric spaces G/H ([HÓØ91]). In this case a complex manifold $\Xi(C^o) \subset G_{\mathbb{C}}/H_{\mathbb{C}}$, depending on the causal structure $m \mapsto C(m)$ on G/H , was constructed. Three important properties of $\Xi(C^o)$ are:

- (1) The manifold $\Xi(C^o)$ is locally isomorphic to a tube domain $\mathfrak{q} + iC^o$, where \mathfrak{q} is the tangent space of G/H at $x_o = eH$ and C^o the interior of $C := C(x_o)$.
- (2) The homogeneous space G/H is a boundary component of $\Xi(C^o)$.
- (3) There is a semigroup Γ containing G and determined by C such that $\Xi(C^o) = (\Gamma^o)^{-1} \cdot x_o$.

The Hardy space $H_2(C)$ is defined as in the classical case to be the space of holomorphic functions on $\Xi(C^o)$ such that the Hardy norm

$$\|f\|_H^2 := \sup_{\gamma \in \Gamma^o} \int_{G/H} |f(\gamma^{-1} \cdot m)|^2 dm$$

is finite. $H_2(C)$ is a Hilbert space with norm $\|\cdot\|_H$, and — as in the classical case — there exists an isometry $\beta : H_2(C) \rightarrow L^2(G/H)$ given by

$$\beta(f) = \lim_{\Gamma^o \ni \gamma \rightarrow 1} \gamma \cdot f,$$

where the limit is in $L^2(G/H)$ and $\gamma \cdot f(\xi) := f(\gamma^{-1} \cdot \xi)$. The left action defines a *holomorphic* representation T of Γ on the Hardy space and a unitary representation λ of G on $L^2(G/H)$ such that β is an intertwining operator for the G -actions.

The parallel to the classical case goes further. In particular one can describe the image of β by a *positivity condition*: $\beta(H_2(C))$ is the direct sum of all the holomorphic discrete series from [ÓØ91] which are *C-admissible*. Point evaluation is also continuous and thus defines a kernel, the Cauchy-Szegö kernel $K(\cdot, \cdot)$. This kernel is determined by a holomorphic H -invariant function $\Theta_K : \Xi(C^o) \rightarrow \mathbb{C}$ such that

$$K(\gamma_1 \cdot x_o, \gamma_2 \cdot x_o) = \Theta_K(\gamma_2^* \gamma_1 \cdot x_o).$$

For the first variable fixed, the kernel extends in the second variable smoothly to the boundary of $\Xi(C^o)$. Then the inverse of the boundary

value map is given just as in the classical case

$$f(z) = \int_{G/H} \beta(f)(x) K(z, x) dx = \int_{G/H} f(\dot{g}) \Theta_K(g^{-1}\gamma \cdot x_o) d\dot{g},$$

where $z = \gamma \cdot x_o \in \Xi(C^o)$ and $\dot{g} = g \cdot x_o \in G/H$.

Let $H_2(C) = \oplus_\delta H_\delta$ be the decomposition of $H_2(C)$ into holomorphic discrete series. Then each of the representations ϵ_δ on H_δ give rise to a spherical distribution

$$\Theta_\delta(f) := \text{pr}_\delta(f)(x_o),$$

where pr_δ is the orthogonal projection onto H_δ . The distribution Θ_δ has an analytic continuation to Ξ^o , determined up to a constant by a spherical function ([**Ó97a**, **Ó97b**, **Ó00**]), with a well-known expansion formula in terms of elementary functions. On $\Xi(C^o)$ we have then the identity $\sum_\delta \Theta_\delta = \Theta_K$. It is still an open problem to evaluate this sum in general.

To calculate Θ_K independently, in a series of lectures at the University of Poitiers in 1990 by B. Ørsted and one of the authors the problem of relating the Hardy spaces H_{cl} and $H_2(C)$ via a *causal compactification* $G/H \rightarrow \check{S}_1$, with \check{S}_1 the Bergman-Šilov boundary of G_1/K_1 , was discussed. It was shown that for the *Cayley type* spaces those are actually isomorphic (modulo a double covering in some cases) ([**ÓØ99**]). In particular this result gives a formula for $K(\cdot, \cdot)$ in terms of the well-known classical Cauchy-Szegö kernel. The above summation formula can then be interpreted as a G -equivariant decomposition of the classical Cauchy-Szegö kernel or a generalized Heine formula ([**ÓØ99**]).

Further results in this direction for special cases were obtained by K. Koufany and B. Ørsted ([**KØ96**, **KØ97**]), G.I. Ol’shanskiĭ ([**O95**]) and V.F. Molchanov for $SO(2, n)/SO(1, n)$ ([**M97**]). The Cayley type spaces were also studied via a Jordan algebra approach by M. Chadli in his thesis ([**C96**]).

One of the obstacles for obtaining general results in this direction was the lack of general theory for the causal compactification of compactly causal symmetric spaces. This was finally obtained in [**B97**]. It was shown that there is a natural causal compactification of a “central extension” of most of those spaces in the Bergman-Šilov boundary of the bounded realization of a tube type domain T_Ω .

In this paper we extend the results form [**B97**] to the domain $\Xi(C^o)$ and show that there is a holomorphic function $\psi_m : G_1/K_1 \rightarrow \mathbb{C}$ such that

$$\Xi(C^o) = \{z \in G_1/K_1 \mid \psi_m(z) \neq 0\}.$$

The map I from the classical Hardy space into $H_2(C)$ is then given by $f \mapsto f \sqrt[m]{\psi_m}$. In general this has only meaning on a m -fold covering of $\Xi(C^o)$. In that case we get an isomorphism into a Hardy space $\tilde{H}_{2,\text{odd}}(C)$ on this covering of $\Xi(C^o)$. The image is a direct sum of highest weight moduls.

We describe the image in terms of the lowest K -type of the highest-weight modules.

We remark that similar results to ours can also be obtained with Jordan algebra methods ([Bal]).

The paper is organized as follows. In the first section we recall some structure theory for causal symmetric spaces and recollect the facts needed by us from [B97] on causal compactifications.

Section 3 is devoted to the description of the Hardy space $H_2(C)$ and here we introduce the semigroup Γ and the domain $\Xi(C^\circ)$.

We construct the function ψ_m in Section 4. In Section 5 we give an isometry from $L^2(\check{S}_1)$ onto $L^2(G/H)$, and construct the covering $\tilde{\Xi}(C^\circ) \rightarrow \Xi(C^\circ)$ of $\Xi(C^\circ)$ corresponding to the m -th root of ψ_m .

In Section 6 we show that that $\Xi(C^\circ) = \{z \in G_1/K_1 \mid \psi_m(z) \neq 0\}$. This gives us the necessary tool to analyze the covering $\tilde{\Xi}(C^\circ)$ in more detail in the next section. In particular we show that we need at most a double covering for describing the Hardy spaces. The final result of this paper is Theorem 7.8 where we characterize the image of I .

The authors would like to thank the Institute Mittag-Leffler for their hospitality during their stay there in the Spring of 1996. The first author would like to thank Louisiana State University and the second author for their hospitality during his stay in September, 1996.

2. Causal compactifications.

In this section we collect some standard facts on symmetric spaces. We also collect some newer results on causal compactifications. We use the monograph [HÓ96] and the original papers [ÓØ88, Ó91, B97] as references. We call (G, H, τ) a *symmetric space*, when G is a connected Lie group, $\tau : G \rightarrow G$ is an involutive automorphism, and $H \subset G$ a closed subgroup with

$$(G^\tau)_o \subset H \subset G^\tau,$$

where $G^\tau := \{a \in G \mid \tau(a) = a\}$ and the subscript $_o$ denotes the connected component containing the identity. By abuse of notation we then also call G/H a symmetric space. A *symmetric Lie algebra* is a triple $(\mathfrak{g}, \mathfrak{h}, \tau)$, where \mathfrak{g} is a Lie algebra, $\tau : \mathfrak{g} \rightarrow \mathfrak{g}$ is an involutive automorphism, and $\mathfrak{h} = \{X \in \mathfrak{g} \mid \tau(X) = X\}$ the subalgebra of τ -fixed elements. To every symmetric space (G, H, τ) there is associated the symmetric algebra $(\mathfrak{g}, \mathfrak{h}, \tau') := (\text{Lie}(G), \text{Lie}(H), d\tau)$. In the sequel we will denote the differential of τ (and similarly the differentials of all other group homomorphisms) always by the same letter. The symmetric algebra $(\mathfrak{g}, \mathfrak{h}, \tau)$ is *irreducible* if there is no nontrivial τ -stable ideal of \mathfrak{g} not contained in \mathfrak{h} . The symmetric algebra $(\mathfrak{g}, \mathfrak{h}, \tau)$ is called *reductive* respectively *semisimple* if \mathfrak{g} is reductive respectively semisimple.

Let $(\mathfrak{g}, \mathfrak{h}, \tau)$ be a reductive symmetric algebra. Choose a Cartan involution θ of \mathfrak{g} commuting with τ ([H78, p. 192] or [Ó84, Lemma 2.1]). Let $\mathfrak{q} := \{X \in \mathfrak{g} \mid \tau(X) = -X\}$, $\mathfrak{k} := \{X \in \mathfrak{g} \mid \theta(X) = X\}$, and $\mathfrak{p} := \{X \in \mathfrak{g} \mid \theta(X) = -X\}$, then

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q} = \mathfrak{k} \oplus \mathfrak{p} = \mathfrak{h}_k \oplus \mathfrak{h}_p \oplus \mathfrak{q}_k \oplus \mathfrak{q}_p,$$

where an index denotes the intersection with the corresponding subspace, i.e., $\mathfrak{h}_p := \mathfrak{h} \cap \mathfrak{p}$, etc.

Let $\mathfrak{g}_s := [\mathfrak{g}, \mathfrak{g}]$ denote the semisimple part of \mathfrak{g} and \mathfrak{z} the center of \mathfrak{g} . If \mathfrak{l} is a subspace of \mathfrak{g} such that $\mathfrak{l} = \mathfrak{l} \cap \mathfrak{z} \oplus \mathfrak{l} \cap \mathfrak{g}_s$, then we set $\mathfrak{l}_z := \mathfrak{l} \cap \mathfrak{z}$, and $\mathfrak{l}_s := \mathfrak{l} \cap \mathfrak{g}_s$. For (G, H, τ) associated to $(\mathfrak{g}, \mathfrak{h}, \tau)$, let G_s be the analytic subgroup corresponding to \mathfrak{g}_s , and $Z := \exp \mathfrak{z}$ those corresponding to \mathfrak{z} . Then $G = ZG_s$ and $D := Z \cap G_s$ is a discrete central subgroup in G_s . Moreover we have a (right) action

$$D \times (Z \times G_s) \rightarrow Z \times G_s, \quad (d, (z, g)) \mapsto (zd, d^{-1}g),$$

such that

$$G \simeq (Z \times G_s)/D =: Z \times_D G_s.$$

Let E be a finite dimensional vector space over the reals. A subset $C \subset E$ is called a *cone* if C is convex and closed under multiplication by \mathbb{R}^+ . The closed cone C is *pointed* if $C \cap -C = \{0\}$ and *generating* if $C - C = E$. A pointed generating cones is called *regular*. We remark that a cone C is generating if and only if its interior C° is nonempty.

Definition. Assume that (G, H, τ) is a symmetric space with associated reductive symmetric Lie algebra $(\mathfrak{g}, \mathfrak{h}, \tau)$.

- 1) The symmetric space is called *compactly causal* if there exists a H -invariant regular cone C in \mathfrak{q} such that $C^\circ \cap \mathfrak{k} \neq \emptyset$.
- 2) $(\mathfrak{g}, \mathfrak{h}, \tau)$ is called *compactly causal* if $(G, (G^\tau)_o, \tau)$ is compactly causal.
- 3) The symmetric space is called *noncompactly causal* if there exists a H -invariant regular cone $C \subset \mathfrak{q}$ such that $C^\circ \cap \mathfrak{p} \neq \emptyset$.
- 4) (G, H, τ) is called of *Cayley type* if it is semisimple and both compactly and noncompactly causal.
- 5) Let $\mathfrak{c}(\mathfrak{q}_k) := \{X \in \mathfrak{q}_k \mid [\mathfrak{q}_k, X] = \{0\}\}$. The symmetric space respectively the associated algebra are called of *weakly Hermitian type* if $\mathfrak{z}_{\mathfrak{q}}(\mathfrak{c}(\mathfrak{q}_k)) = \mathfrak{q}_k$.

Notice that in this definition we *do not* assume that G is noncompact. Thus every compact symmetric space is weakly Hermitian. On the other hand a compact symmetric space can only be compactly causal if it is reductive ([HÓ96]). The spaces we consider will by construction be both compactly causal and of weakly Hermitian type. We refer to [KN96] and [KNÓ97] for a general discussion of the connection between weakly Hermitian and compactly causal.

Let $(\mathfrak{g}_1, \mathfrak{h}_1, \tau_1)$ be a compactly causal irreducible symmetric algebra with \mathfrak{g}_1 simple and noncompact. Let $G_{1\mathbb{C}}$ the simply connected complex Lie group with Lie algebra $\mathfrak{g}_{1\mathbb{C}} := \mathfrak{g}_1 \otimes \mathbb{C}$. We extend τ_1 to a complex linear involution of $\mathfrak{g}_{1\mathbb{C}}$, and denote this involution and the corresponding involution on $G_{1\mathbb{C}}$ again by τ_1 . Then, as $G_{1\mathbb{C}}$ is assumed simply connected, it follows that $H_{1\mathbb{C}} := G_{1\mathbb{C}}^{\tau_1}$ is connected ([L69, p. 171]). Let G_1 be the analytic subgroup of $G_{1\mathbb{C}}$ with Lie algebra \mathfrak{g}_1 and

$$H_1 := G_1 \cap H_{1\mathbb{C}} = G_1^{\tau_1}.$$

Let θ_1 be a Cartan involution commuting with τ_1 , and let $\mathfrak{g}_1 = \mathfrak{k}_1 \oplus \mathfrak{p}_1$ be the corresponding Cartan decomposition. Recall that for $(\mathfrak{g}_1, \mathfrak{h}_1, \tau_1)$ irreducible compactly causal the Riemannian symmetric space G_1/K_1 is Hermitian symmetric ([HÓ96, Remark 3.19]) and that $(\mathfrak{g}_1, \mathfrak{h}_1, \tau_1)$ is of weakly Hermitian type ([HÓ96, Lemma 1.2.1, Lemma 1.3.5]). We will assume that G_1/K_1 is a tube type domain.

Let \mathfrak{t}_1 be a Cartan subalgebra of \mathfrak{k}_1 . Let $\Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$ be the set of roots of $\mathfrak{t}_{1\mathbb{C}}$ in $\mathfrak{g}_{1\mathbb{C}}$. We have the two subsets

$$\Delta(\mathfrak{k}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}}) = \{\alpha \in \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}}) \mid \mathfrak{g}_{1\mathbb{C},\alpha} \subset \mathfrak{k}_{1\mathbb{C}}\}$$

of *compact* and

$$\Delta(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}}) = \{\alpha \in \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}}) \mid \mathfrak{g}_{1\mathbb{C},\alpha} \subset \mathfrak{p}_{1\mathbb{C}}\}$$

of *noncompact* roots. We choose an ordering in $i\mathfrak{t}_1^*$ such that the positive noncompact dominate the positive compact roots and define

$$\mathfrak{p}_1^+ := \sum_{\alpha \in \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})} \mathfrak{g}_{1\mathbb{C},\alpha} \quad \text{respectively} \quad \mathfrak{p}_1^- := \sum_{\alpha \in \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})} \mathfrak{g}_{1\mathbb{C},-\alpha}.$$

Recall that two roots α and β are *strongly orthogonal* if $\alpha \neq \pm\beta$ and $\alpha \pm \beta \notin \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$. Let $\{\gamma_1, \dots, \gamma_r\} \subset \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$ be a maximal system of strongly orthogonal roots. Let $H_j \in [\mathfrak{g}_{1\mathbb{C},\gamma_j}, \mathfrak{g}_{1\mathbb{C},-\gamma_j}] \cap i\mathfrak{t}_1$ be such that $\gamma_i(H_j) = 2\delta_{i,j}$. Choose $E_{\pm j} \in \mathfrak{g}_{1\mathbb{C},\pm\gamma_j}$ such that

$$X_j := E_j + E_{-j}, \quad Y_j := iE_j - iE_{-j} \in \mathfrak{p}_1,$$

and

$$[E_j, E_{-j}] = H_j.$$

It is known that $\mathfrak{a}_p := \sum \mathbb{R}X_j$ is maximal abelian in \mathfrak{p}_1 ([H78, p. 387]). Let

$$X^0 := \sum X_j, \quad Y^0 := \sum Y_j, \quad \text{and} \quad Z^0 := -\frac{1}{2} \sum iH_j,$$

then $Z^0 \in \mathfrak{z}(\mathfrak{k}_1)$, with $\mathfrak{z}(\mathfrak{k}_1)$ the center of \mathfrak{k}_1 , and $\text{ad } X^0$ has eigenvalues 0, 2 and -2 . Let

$$\eta := \text{Ad} \left(\exp \frac{\pi}{2} iX^0 \right)$$

be the involution on $G_{1\mathbb{C}}$ respectively $\mathfrak{g}_{1\mathbb{C}}$ given by conjugation with $\exp \frac{\pi}{2} i X^0$. Then the symmetric space G_1/G_1^η is of Cayley type and

$$\mathfrak{q}_{1\eta} := \mathfrak{g}_1^{-\eta} = \{X \in \mathfrak{g}_1 \mid \eta(X) = -X\}$$

is the direct sum of the two eigenspaces $\mathfrak{q}_{1\eta}^\pm := \{X \in \mathfrak{q}_{1\eta} \mid [X^0, X] = \pm 2X\}$ of $\text{ad } X^0$, which are also G_1^η -invariant. As $\theta_1(X^0) = -X^0$ we get $\theta_1(\mathfrak{q}_{1\eta}^+) = \mathfrak{q}_{1\eta}^-$.

Define $Y_\pm \in \mathfrak{q}_{1\eta}^\pm$ by $Y^0 = Y_+ + Y_-$. The sets $C_\pm := \overline{\text{Ad}(G_1^\eta)Y_\pm}$ are regular cones in $\mathfrak{q}_{1\eta}^\pm$ and the cone $C_k := C_+ - C_- \subset \mathfrak{q}_{1\eta}$ defines the compactly causal structure on G_1/G_1^η . Let \mathfrak{p}' be the sum of the 0- and (-2)-eigenspaces of $\text{ad } X^0$. Then $P' := N_{G_1}(\mathfrak{p}')$ is a parabolic subgroup and

$$\check{S}_1 := G_1/P' = K_1/K_1^\eta$$

is the Bergman-Šilov boundary of G_1/K_1 . The compact symmetric space \check{S}_1 is causal with causal structure defined by $C_+ \subset \mathfrak{q}_{1\eta}^+ \simeq T_{eP'}\check{S}_1$. Moreover

$$G_1^\eta = Z_{G_1}(X^0) \subset P',$$

and the canonical projection $\Phi_1 : G_1/G_1^\eta \rightarrow \check{S}_1$ is causal.

Under conditions specified in [B97] there exists an involution σ , commuting with τ_1, θ_1 , and η , such that

$$(G, H, \tau) := ((G_1^\sigma)_o, H_1 \cap (G_1^\sigma)_o, \tau_1|(G_1^\sigma)_o)$$

is a compactly causal symmetric subspace of (G_1, G_1^η, η) and Φ_1 can be used to define a causal compactification of it. (For spaces of Cayley type $(\mathfrak{g}_1, \mathfrak{h}_1, \tau_1)$ in the preceding discussion has to be replaced by $(\mathfrak{g}_1 \times \mathfrak{g}_1, \mathfrak{h}_1 \times \mathfrak{h}_1, \tau_1 \times \tau_1)$ and $\sigma(g, h) = (h, g)$.) For the general theory we will not need the explicit form of σ but only the “axiomatic” properties of the construction that we collect in the remaining part of this section.

Define $\theta = \theta_1|_G$ then θ is a Cartan involution on G commuting with τ . We denote the corresponding Cartan decomposition by $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. Let $\mathfrak{g}_1 = \mathfrak{g} + \mathfrak{q}_{1\sigma}$ be the eigenspace decomposition of σ . We have $\eta|_G = \tau_1|_G$ — in fact, in many cases $\eta = \tau_1$ — such that $G \cap G_1^\eta = G \cap H_1$ and also $\mathfrak{q} := \mathfrak{q}_1 \cap \mathfrak{g} = \mathfrak{q}_{1\eta} \cap \mathfrak{g}$.

Lemma 2.1. *With notation as above the following holds:*

- (1) *The algebra $\mathfrak{b} := \mathfrak{a}_\mathfrak{p} \cap \mathfrak{q}_{1\sigma}$ is maximal abelian in $\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$, $X^0 \in \mathfrak{b}$, $Y^0 \in \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$, and $Z^0 \in \mathfrak{k} \cap \mathfrak{q}$.*
- (2) *The cone C_k is σ -invariant.*
- (3) *Let $\text{pr}_\mathfrak{q} : \mathfrak{q}_{1\eta} \rightarrow \mathfrak{q}$ be the projection with respect to the decomposition $\mathfrak{q}_{1\eta} = \mathfrak{q} \oplus \mathfrak{q}_{1\eta} \cap \mathfrak{q}_{1\sigma}$. Then $C_k \cap \mathfrak{q} = \text{pr}_\mathfrak{q}(C_k)$ and $(\text{pr}_\mathfrak{q}(C_k))^o \cap \mathfrak{k} \neq \emptyset$.*

Proof. (1) With $Z^0 = \frac{1}{4}[X^0, Y^0]$ the last claim follows from the preceding ones. These are prerequisites for the construction of σ in [B97, Theorem 5.1] respectively [B97, Theorem 5.9].

(2) From

$$Y_+ + Y_- = Y^0 = -\sigma(Y^0) = -\sigma(Y_+) - \sigma(Y_-)$$

and

$$\sigma(Y_{\pm}) = \sigma\left(\pm\frac{1}{2}[X^0, Y_{\pm}]\right) = \mp\frac{1}{2}[X^0, \sigma(Y_{\pm})]$$

we have $\sigma(Y_{\pm}) = -Y_{\mp}$. By the σ -invariance of G_1^η this implies the claim together with the definition of C_k .

(3) The σ -invariance of C_k implies that $C_k \cap \mathfrak{q} = \text{pr}_{\mathfrak{q}}(C_k)$. That $(\text{pr}_{\mathfrak{q}}(C_k))^o \cap \mathfrak{k}$ is nonempty follows since $Z^0 \in C_k^o$. \square

By the last part of the Lemma $C := C_k \cap \mathfrak{q}$ is a H -invariant regular cone in \mathfrak{q} which defines a compactly causal structure for (G, H, τ) . It is clear that this definition makes (G, H, τ) into a causal subspace of (G_1, G_1^η, η) .

Corollary 2.2. $C + \mathfrak{p}' = C_k + \mathfrak{p}' = C_+ \oplus \mathfrak{p}'$.

Proof. By definition we have

$$C + \mathfrak{p}' \subset C + \mathfrak{p}' = C_+ \oplus \mathfrak{p}'.$$

When we write $X \in C_+$ in the form $(X + \sigma(X)) - \sigma(X)$, where $X + \sigma(X) \in C_k \cap \mathfrak{g}$ and $\sigma(X) \in \sigma(\mathfrak{q}_{1\eta}^+) \subset \mathfrak{p}'$, the other inclusion is also obvious. \square

Definition ([$\acute{O}\acute{O}99$]). Let M and N be manifolds, with N compact, and with causal structure $M \ni m \mapsto C(m) \subset T_m M$ respectively $N \ni n \mapsto D(n) \subset T_n N$. Let $\Phi : M \rightarrow N$ be smooth. The pair (N, Φ) is called a causal compactification of M if Φ is a diffeomorphism onto an open dense subset of N , and $(d\Phi)_m(C(m)) = D(\Phi(m))$ for all $m \in M$. Let L be a Lie group acting on M and N such that the causal structures are L -invariant. Then the causal compactification is L -equivariant if Φ is L -equivariant.

Theorem 2.3. *The canonical inclusion $\iota : G/H \hookrightarrow G_1/G_1^\eta$ is causal and the map $\Phi := \Phi_1 \circ \iota : G/H \hookrightarrow \check{S}_1$ is a G -equivariant causal compactification of G/H .*

Proof. By [**B97**] we know that $\Phi : G/H \rightarrow \check{S}_1$ is an injective G -map with open and dense image. With the canonical identifications $T_{eH}G/H \simeq \mathfrak{q}$, $T_{eG_1^\eta}G/G_1^\eta \simeq \mathfrak{q}_{1\eta}$, and $T_{eP'}(\check{S}_1) \simeq \mathfrak{q}_{1\eta}^+$, the tangent map $d\Phi$ at the identity coset is given by first imbedding \mathfrak{q} into $\mathfrak{q}_{1\eta}$ and then projecting onto $\mathfrak{q}_{1\eta}^+$. Thus $d\Phi_{eH}(C) = C_+$ by Corollary 2.2. By the G -invariance of the causal structures on G/H respectively \check{S}_1 this is all to show. \square

As all groups are subgroups of $G_{1\mathbb{C}}$ it is immediate to see that Φ extends to a $G_{\mathbb{C}}$ -invariant map $G_{\mathbb{C}}/H_{\mathbb{C}} \hookrightarrow G_{1\mathbb{C}}/P'_{\mathbb{C}}$, which we denote again by Φ .

Example. Let $G_1 = SU(1, 1)$. Denote by $E_{i,j}$ the matrix with entry 1 in the i -th row and j -th column and otherwise zero. We choose $E_1 = E_{1,2}$, $E_{-1} = E_{2,1}$, and $H_1 = E_{1,1} - E_{2,2}$. Then

$$X^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \text{and} \quad Y^0 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}.$$

Thus η is given by conjugation by iX^0 . This is easily seen to be the same involution on $SU(1, 1)$ as $g \mapsto \bar{g}$, the complex conjugation. We will also need the *Cayley transform*, which is given by conjugation by

$$c := \exp \frac{\pi}{4} iY^0 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}.$$

As is well-known, the unit disk — identifying $\mathbb{C} \simeq \mathbb{C}E_1 = \mathfrak{p}_1^+$ — is the Harish-Chandra realization of G_1/K_1 , where G_1 operates by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot z = \frac{az + b}{cz + d}, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SU(1, 1).$$

The Bergman-Šilov boundary \check{S}_1 is now the unit circle and the Cayley transform $z \mapsto c \cdot z$ maps the origin to $-1 \in \check{S}_1$. We notice that $\text{Ad}(c)H_1 = X^0$. It follows in particular that $(\text{Ad}(c)(\mathfrak{k}_{1\mathbb{C}} + \mathfrak{p}_1^-)) \cap \mathfrak{g}_1 = \mathfrak{p}'$ and $\check{S}_1 = G_1/P' \simeq G_1 \cdot (-1)$. With \check{S}_1 in the Harish-Chandra realization the map Φ_1 is therefore given by $gG_1^\eta \mapsto g \cdot (-1)$.

We can choose the involution $\sigma = \theta_1$, i.e.,

$$(G, H) = \left(\left\{ \begin{pmatrix} a & 0 \\ 0 & \bar{a} \end{pmatrix} \mid a \in S^1 \right\}, \{\pm \text{id}\} \right),$$

and then

$$\Phi \left(\begin{pmatrix} a & 0 \\ 0 & \bar{a} \end{pmatrix} H \right) = -a^2.$$

We restrict us in the following to causal compactifications constructed as in the Theorem. (Indeed, as is shown in [B97], with one natural additional assumption all causal compactifications in the Bergman-Šilov boundary are of this form.) A list of the possible compactifications can be found in [B97], cf. also below.

For short we call $\langle A, B, C \rangle$ a $\mathfrak{sl}(2)$ -triple if

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \mapsto A, \quad \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \mapsto B, \quad \text{and} \quad \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \mapsto C$$

defines an isomorphism from $\mathfrak{sl}(2, \mathbb{C})$ onto the complex Lie algebra generated by $\{A, B, C\}$.

Proposition 2.4. *Let the notation be as above. Then the following holds:*

- (1) *There exist $\lambda_{j',j} \in \{0, 1\}$ such that with*

$$E'_{\pm j'} := \sum_j \lambda_{j',j} E_{\pm j},$$

$$H'_{j'} := [E'_{j'}, E'_{-j'}] = \sum_j \lambda_{j',j}^2 H_j$$

and

$$X'_{j'} := E'_{j'} + E'_{-j'} = \sum_j \lambda_{j',j} X_j$$

we have $\mathfrak{b} = \sum \mathbb{R}X'_{j'}$, and the triples $\langle H'_{j'}, E'_{j'}, E'_{-j'} \rangle$ are pairwise commuting $\mathfrak{sl}(2)$ -triples.

- (2) *With $c := \exp(\frac{\pi}{4}iY^0)$, the Cayley-transformed space $\mathfrak{a} := \text{Ad}(c)i\mathfrak{b}$ is maximal abelian in $\mathfrak{k} \cap \mathfrak{q}$.*
 (3) *Let $\mathfrak{t}_1^- := \sum \mathbb{R}iH_j$ then we have $\mathfrak{t}_1^- = \mathfrak{a} + \mathfrak{t}_1^- \cap \mathfrak{q}_{1\sigma}$.*
 (4) *We have $\mathfrak{z} \subset \mathfrak{k}$ and, if \mathfrak{g} is noncompact, then $(\mathfrak{g}, \mathfrak{h}, \tau)$, is weakly Hermitian. If $\mathfrak{l} \subset \mathfrak{h}$ is an ideal of \mathfrak{g} then $\mathfrak{l} \subset \mathfrak{k}$.*
 (5) *$C^\circ = \text{Ad}(H)(\mathfrak{a} \cap C^\circ)$ and $C = \text{Ad}(H)(\mathfrak{a} \cap C)$.*
 (6) *$\mathfrak{a} \cap C^\circ = -\sum \mathbb{R}^+ iH'_{j'}$, and $\mathfrak{a} \cap C = -\sum \mathbb{R}_0^+ iH'_{j'}$.*

Proof. (1) This is a direct consequence of [B97, Theorem 5.1] respectively [B97, Theorem 5.9].

- (2) The inner automorphism $\text{Ad}(c)$ maps the subalgebra

$$\mathfrak{g}_a := \mathfrak{k} + i(\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma})$$

onto itself, since $Y^0 \in \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma}$ by Lemma 2.1.

We consider first the causal compactifications described by [B97, Theorem 5.1]. Here we have $\tau_1 = \eta$ and $\mathfrak{b} \subset \mathfrak{h}_1$. Then $Y^0 = [X^0, Z^0] \in \mathfrak{p}_1 \cap \mathfrak{q}_1$ is $\tau_1\theta_1$ -fixed and consequently $\mathfrak{k} \cap \mathfrak{q} + i(\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} \cap \mathfrak{h}_1)$, the (-1) -eigenspace of the involution $\tau_1\theta_1$ in \mathfrak{g}_a , is $\text{Ad}(c)$ -invariant. Using $\theta_1 = \text{Ad}(\exp \pi Z^0)$, by a $\mathfrak{sl}(2)$ -calculation $\text{Ad}(c)\theta_1 = \tau_1\text{Ad}(c)$. Looking at the corresponding eigenspaces in $\mathfrak{k} \cap \mathfrak{q} + i(\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} \cap \mathfrak{h}_1)$ we get $\text{Ad}(c)i(\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} \cap \mathfrak{h}_1) = \mathfrak{k} \cap \mathfrak{q}$. As $\mathfrak{b} \subset \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} \cap \mathfrak{h}_1$, the claim for this case then follows.

For the compactifications given by [B97, Theorem 5.9] we have $\mathfrak{b} \subset \mathfrak{q}_1$. Similarly to the first case we get successively $Y^0 \in \mathfrak{h}_1$, the $\text{Ad}(c)$ -invariance of $\mathfrak{k} \cap \mathfrak{q} + i(\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} \cap \mathfrak{q}_1)$, and $\text{Ad}(c)\theta_1\tau_1 = \sigma\text{Ad}(c)$, where $\sigma := \eta \circ \tau_1$ as in [B97, Thm. 5.9] defined. Therefore $\text{Ad}(c)$ maps the $(+1)$ -eigenspace of $\theta_1\tau_1$ onto the $(+1)$ -eigenspace of σ , i.e., $\text{Ad}(c)i(\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} \cap \mathfrak{q}_1) = \mathfrak{k} \cap \mathfrak{q}$. Now the stated result for this case follows again from Lemma 2.1.

(3) For the orthogonal sum $\mathfrak{a}_p = \mathfrak{b} \oplus \mathfrak{b}^\perp$ (with respect to the inner product $(X, Y) = -B(X, \theta(Y))$) we have by Lemma 2.1 that $\mathfrak{b}^\perp \subset \mathfrak{p}_1 \cap \mathfrak{g}$. Now $\mathfrak{t}_1^- = \text{Ad}(c)i\mathfrak{a}_p = \mathfrak{a} \oplus \text{Ad}(c)\mathfrak{b}^\perp$. The second summand is contained in \mathfrak{k}_1 , since $Z^0 \in \mathfrak{a}$. But the Cayley transform maps with $\mathfrak{g}_a = \mathfrak{k} + i(\mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma})$ also

the orthocomplement \mathfrak{g}_a^\perp of \mathfrak{g}_a in $\mathfrak{k}_1 + i\mathfrak{p}_1$ onto itself. With $ib^\perp \subset \mathfrak{g}_a^\perp$ we have therefore $\text{Ad}(c)ib^\perp \subset \mathfrak{g}_a^\perp \cap \mathfrak{k}_1 = \mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma}$.

(4) As $Z^0 \in \mathfrak{g}$ by Lemma 2.1, and $\text{ad } Z^0|_{\mathfrak{p}_1}$ is regular, it follows that $\mathfrak{z}_{\mathfrak{g}}(Z^0) = \mathfrak{k}$. Hence

$$\mathfrak{z} \subset \mathfrak{z}_{\mathfrak{g}}(Z^0) = \mathfrak{k}.$$

Write $Z^0 = Z_1 + Z_2$, with $Z_1 \in \mathfrak{z}(\mathfrak{g})$, and $Z_2 \in [\mathfrak{g}, \mathfrak{g}]$. Assume that $Z_2 = 0$. Then the same argument shows that $\mathfrak{g} = \mathfrak{z}_{\mathfrak{g}}(Z^0) = \mathfrak{k}$. Hence \mathfrak{g} is compact. Thus, if \mathfrak{g} is not compact it follows that $Z_2 \neq 0$. Furthermore $\mathfrak{z}_{\mathfrak{g}}(Z_2) = \mathfrak{k}$, as Z_1 commutes with \mathfrak{g} . In particular it follows that $Z_2 \in \mathfrak{c}(\mathfrak{q}_k)$ and therefore

$$\mathfrak{z}_{\mathfrak{q}}(\mathfrak{c}(\mathfrak{q}_k)) \subset \mathfrak{z}_{\mathfrak{q}}(Z_2) = \mathfrak{q}_k.$$

Suppose that \mathfrak{l} is an ideal in \mathfrak{g} contained in \mathfrak{h} . As $Z_2 \in \mathfrak{q}$ it follows that $[Z_2, \mathfrak{l}] = \{0\}$. Hence

$$\mathfrak{l} \subset \mathfrak{z}_{\mathfrak{g}}(Z_2) = \mathfrak{k}$$

which shows that \mathfrak{l} is compact.

(5) Let $X \in C^o = (C_k^o \cap \mathfrak{g})$. Then $G^X = \{a \in G \mid \text{Ad}(a)X = X\}$ is compact. Therefore the elements in \mathfrak{g}^X are semisimple and with imaginary eigenvalues (see the proof of [HÓ96, Thm. 4.2.15]). Hence there is a $h \in H$ such that $\text{Ad}(h)X \in \mathfrak{q}_k$ ([M79, Thm. 1]). Let \mathfrak{a}_q be a maximal abelian algebra in \mathfrak{q}_k containing X , then (by [M79, Lemma 7]) we find $k \in H \cap K$ with $\text{Ad}(k)\mathfrak{a}_q = \mathfrak{a}$. In particular $\text{Ad}(kh)X \in \mathfrak{a}$. Therefore $C^o \subset \text{Ad}(H)(\mathfrak{a} \cap C^o)$. As the other inclusion is obvious we get $C^o = \text{Ad}(H)(\mathfrak{a} \cap C^o)$. Since the eigenvalues of $\text{ad } X$ are continuous in X , the same argument apply to X in the boundary of C . Thus the assertion for the whole cone follows.

(6) By [ÓØ99, Lemma 4.4] or [HÓ96, Chap. 4.2] we have $\mathfrak{t}_1^- \cap C_k^o = -\sum \mathbb{R}^+ iH_j$ respectively $\mathfrak{t}_1^- \cap C_k = -\sum \mathbb{R}_0^+ iH_j$. (Note that we have here a minus sign by our choice of Z^0 .) The claims then follow with $\mathfrak{t}_1^- \cap \mathfrak{a} = \sum \mathbb{R}iH'_j$. \square

Remarks. (1) The $\lambda_{j',j}$ are known by [B97, Thm. 5.1, Thm. 5.9] for every case. Either $\lambda_{j',j} = \delta_{j',j}$, for $j' = 1, \dots, r$, or $r = 2r'$ and $\lambda_{j',j} = \delta_{j',j} + \delta_{j'+r',j}$, for $j' = 1, \dots, r'$.
 (2) By the classification of causal compactifications the only $(\mathfrak{g}, \mathfrak{h}, \tau)$ with nontrivial center are those described by [B97, Corollary 5.3].

3. Hardy spaces.

We will from now on allways assume that \mathfrak{g} is noncompact. Proposition 2.4(4) asserts that $(\mathfrak{g}_s, \mathfrak{h}_s, \tau|_{\mathfrak{g}_s})$ is a symmetric Lie algebra of Hermitian type in the sense of [ÓØ88, Ó91, HÓ91] where we have a theory of Hardy spaces. In this section we generalize this to our reductive setting. Let \mathfrak{t} be a Cartan algebra of \mathfrak{g} , with $\mathfrak{c}(\mathfrak{q}_k) \subset \mathfrak{a} \subset \mathfrak{t} \subset \mathfrak{k}$, and $\Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$ respectively $\Delta := \Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{a}_{\mathbb{C}})$ the corresponding root systems. We may assume without loss of generality

$\mathfrak{t} \subset \mathfrak{t}_1$. Indeed, if $\mathfrak{t}_1^- \cap \mathfrak{q}_{1\sigma} = \{0\}$ by Proposition 2.4 then $[\mathfrak{t}, \mathfrak{t}_1^-] = \{0\}$ and (since the constructions in the last section needed only \mathfrak{t}_1^-) we can assume $\mathfrak{t} + \mathfrak{t}_1^- \subset \mathfrak{t}_1$. For the remaining (four) cases this is immediate from the realization of their causal compactifications ([B97]). We assume all introduced root systems to be lexicographically ordered with respect to the flag $\mathfrak{c}(\mathfrak{q}_k) \subset \mathfrak{a} \subset \mathfrak{t} \subset \mathfrak{t} + \mathfrak{t}_1^- \subset \mathfrak{t}_1$. Especially we have

$$\mathfrak{p}^\pm := \bigoplus_{\alpha \in \Delta^+(\mathfrak{p}_\mathbb{C}, \mathfrak{t}_\mathbb{C})} \mathfrak{g}_{\mathbb{C}, \pm\alpha} \subset \mathfrak{p}_1^\pm.$$

Let $\Delta_n := \Delta(\mathfrak{p}_\mathbb{C}, \mathfrak{a}_\mathbb{C})$, $\Delta_n^+ := \Delta(\mathfrak{p}^+, \mathfrak{a}_\mathbb{C})$, $\Delta_k := \Delta(\mathfrak{k}_\mathbb{C}, \mathfrak{a}_\mathbb{C})$, etc. Let $H_\alpha \in i\mathfrak{a}$ denote the co-root determined by the two conditions

$$H_\alpha \in [\mathfrak{g}_{\mathbb{C}, \alpha}, \mathfrak{g}_{\mathbb{C}, -\alpha}] \cap i\mathfrak{a} \quad \text{and} \quad \alpha(H_\alpha) = 2.$$

Assume for the moment that $(\mathfrak{g}_s, \mathfrak{h}_s, \tau|\mathfrak{g}_s)$ is irreducible. Recall that for this case in \mathfrak{q} we have the minimal closed cone C_{\min} and maximal closed cone C_{\max} given by

$$C_{\min} = \overline{\text{convAd}(H)Z_2} = C_{\min, s} \subset \mathfrak{g}_s$$

and

$$C_{\max} = C_{\min}^* = \mathfrak{z} + C_{\max, s}$$

where the subscript s indicates the corresponding cone in \mathfrak{g}_s . We have by [HÓ96, Chap. 4.2]

$$c_{\min} := C_{\min} \cap \mathfrak{a} = -i \overline{\sum_{\alpha \in \Delta_n^+} \mathbb{R}^+ H_\alpha}$$

and

$$c_{\max} := C_{\max} \cap \mathfrak{a} = \{H \in \mathfrak{a} \mid \forall \alpha \in \Delta_n^+ : -i\alpha(H) \geq 0\}.$$

Lemma 3.1. *Denote by pr_z respectively pr_s the projection onto \mathfrak{z} respectively \mathfrak{g}_s corresponding to the decomposition $\mathfrak{g} = \mathfrak{z} \oplus \mathfrak{g}_s$. Then the following holds:*

- (1) $\text{pr}_z(C) = \mathfrak{z}$.
- (2) $C_s := \text{pr}_s(C)$ is a regular H_s -invariant cone in \mathfrak{q}_s . In particular $C_{\min, s} \subset \text{pr}_s(C) \subset C_{\max, s}$.

Proof. (1) If $\mathfrak{z} \neq \{0\}$ the center is spanned by $Z'_1 = \sum \epsilon_j iH_j$, with $\epsilon_1 = \dots = \epsilon_p = 1 = -\epsilon_{p+1} = \dots = -\epsilon_r$ ([B97, Prop. 5.5]). In this case we also have $\lambda_{j',j} = \delta_{j',j}$. We get from Proposition 2.4(6) for elements in $\mathfrak{a} \cap C^o$

$$\sum a_j iH_j = aZ'_1 + X_s,$$

where $a_j < 0$ and $X_s \in \mathfrak{g}_s$. Since both sums are orthogonal it follows $a = \frac{1}{r} \sum \epsilon_j a_j$, which can take any value.

(2) Obviously C_s is a convex generating H_s -invariant cone in \mathfrak{q}_s because C is a convex generating H_s -invariant cone in \mathfrak{q} . Assume that C_s is not

pointed. Let $\mathfrak{q}_2 := C_s \cap -C_s$. Then $\mathfrak{g}_2 := [\mathfrak{q}_2, \mathfrak{q}_2] \oplus \mathfrak{q}_2$ is a subalgebra of \mathfrak{g}_s . As C is θ -stable it follows that \mathfrak{g}_2 is θ -stable and hence reductive. We have $[\mathfrak{h}_s, \mathfrak{g}_2] \subset \mathfrak{g}_2$ because C is H_s -stable and because of the Jacobi identity. Let \mathfrak{q}_2^\perp be the orthogonal complement to \mathfrak{q}_2 (with respect to the inner product $(X, Y) = -B(X, \theta(Y))$). Let $Z \in \mathfrak{q}_2$ and $Y \in \mathfrak{q}_2^\perp$. Then

$$-B([Z, Y], \theta([Z, Y])) = -B(Y, [Z, [\theta(Z), \theta(Y)]]) = 0$$

because $[\theta(Z), \theta(Y)] \in \mathfrak{h}_s$. It follows that \mathfrak{g}_2 is a τ -stable ideal in \mathfrak{g}_s . But $(\mathfrak{g}_s, \mathfrak{h}_s, \tau|_{\mathfrak{g}_s})$ is by [B97, Prop. 5.5] irreducible, hence $\mathfrak{q}_2 = \mathfrak{q}_s$. It follows in particular that $\pm H_\alpha \in C_s$ for all $\alpha \in \Delta_n$. Choose a Lie algebra homomorphism

$$\varphi_\alpha : \mathfrak{su}(1, 1) \rightarrow \mathfrak{g}_s$$

such that

$$\varphi_\alpha \left(\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \right) = iH_\alpha$$

and

$$T_\alpha := \varphi_\alpha \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \in \mathfrak{h}_s$$

([Ó088, p. 134]). Let $h_t := \exp tT_\alpha$ then a simple $SU(1, 1)$ -calculation shows that

$$\text{Ad}(h_t)H_\alpha + \text{Ad}(h_{-t})H_\alpha = 2 \cosh(2t)H_\alpha .$$

Choose $Z \in \mathfrak{z}$ such that $Z + H_\alpha \in C$. Then

$$\lim_{t \rightarrow \infty} \left(\frac{1}{\cosh(2t)} \left(\text{Ad}(h_t)(Z + H_\alpha) + \text{Ad}(h_{-t})(Z + H_\alpha) \right) \right) = 2H_\alpha \in C.$$

Similarly one shows that $-2H_\alpha \in C$. This contradicts the fact, that C is pointed. Hence C_s is also pointed. □

The cone C_k is minimal in $\mathfrak{q}_{1\eta}$ and is generated by $\text{Ad}(G_1^\eta)Z^0$. A minimal extension of C_k to a G_1 -invariant cone in \mathfrak{g}_1 is W_k , the minimal cone in \mathfrak{g}_1 generated by $\text{Ad}(G_1)Z^0$ ([HÓ96, Chap. 4.2]). As $Z^0 \in \mathfrak{q} \cap \mathfrak{k}$ it follows that W_k is σ -invariant. For a subset $D \subset W_k$ we define

$$\Gamma_1(D) := G_1 \exp iD \subset G_{1\mathbb{C}} .$$

It is known that the *Ol'shanskiiĭ semigroup* $\Gamma_1(W_k)$ is a closed semigroup in $G_{1\mathbb{C}}$ and that

$$\Gamma_1(W_k)^o = \Gamma_1(W_k^o) \simeq G_1 \times iW_k^o,$$

where the diffeomorphism is given by $(g, iX) \mapsto g \exp iX$.

Theorem 3.2. *For $W := W_k \cap \mathfrak{g}$ the following holds:*

- (1) $W = W_k^\sigma = \text{pr}_{\mathfrak{g}}(W_k)$, where $\text{pr}_{\mathfrak{g}} : \mathfrak{g}_1 \rightarrow \mathfrak{g}$ denotes the orthogonal projection.

- (2) W is a regular G -invariant cone in \mathfrak{g} such that $W \cap \mathfrak{q} = \text{pr}_{\mathfrak{q}}(W) = C$, where $\text{pr}_{\mathfrak{q}} : \mathfrak{g} \rightarrow \mathfrak{q}$ denotes the orthogonal projection, i.e., W is a G -invariant extension of C .
- (3) $W^\circ = \text{Ad}(G)(\mathfrak{t} \cap W^\circ)$.
- (4) Let $\Gamma(W) := G \exp iW$. Then

$$\Gamma(W) = \Gamma_1(W_k)^\sigma = \Gamma_1(W_k) \cap G_{1\mathbb{C}}^\sigma.$$

Thus $\Gamma(W)$ is a closed semigroup in $G_{\mathbb{C}} := G_{1\mathbb{C}}^\sigma$.

- (5) $\Gamma(W)^\circ = G \exp iW^\circ =: \Gamma(W^\circ)$, and

$$G \times iW^\circ \rightarrow \Gamma(W^\circ) \quad (g, iX) \mapsto g \exp iX$$

is a diffeomorphism.

Proof. (1), (2), and (3) follow in the same way as the corresponding claims for the cones C_k and C , using also the fact that W_k is an extension of C_k .

(4) Obviously $\Gamma(W) \subset \Gamma_1(W_k)^\sigma$, and $\Gamma(W) \subset \Gamma_1(W_k) \cap G_{\mathbb{C}}$. Let $\gamma = g \exp iX \in \Gamma_1(W_k)^\sigma$, $g \in G_1$ and $X \in W_k$. Then

$$\sigma(g) \exp i\sigma(X) = g \exp iX.$$

Because of the uniqueness of the decomposition in $\Gamma_1(W_k)$ we get $\sigma(g) = g$ and $\sigma(X) = X$. Hence $\Gamma(W) = \Gamma(W)^\sigma$. Assume now that $\gamma = g \exp iX \in G_{\mathbb{C}} \cap \Gamma_1(W_k)$, $g \in G_1$, $X \in W_k$. Then the same argument shows that $\sigma(g) = g$ and $\sigma(X) = X$. Hence $g \in G_1^\sigma$ and $X \in W_k^\sigma = W$. Hence $\gamma \in \Gamma(W)$. It follows that $\Gamma(W)$ is a closed semigroup.

(5) These claims follow by (4) and the corresponding facts for $\Gamma_1(W_k)$. \square

With these results it is straightforward to generalize the constructions of the holomorphic discrete series for G/H ([ÓØ88, ÓØ91]) and the Hardy spaces ([HÓØ91]) to our setting.

To be complete we collect here the results needed by us. A survey is also given in [HÓ96, Chap. 7]. Let

$$\rho := \frac{1}{2} \sum_{\alpha \in \Delta^+} \dim \mathfrak{g}_{\mathbb{C}, \alpha} \alpha.$$

We have that \mathfrak{p}^+ and \mathfrak{p}^- are abelian subalgebras, $\mathfrak{g}_{\mathbb{C}} = \mathfrak{p}^+ \oplus \mathfrak{k}_{\mathbb{C}} \oplus \mathfrak{p}^-$, and $[\mathfrak{k}_{\mathbb{C}}, \mathfrak{p}^\pm] \subset \mathfrak{p}^\pm$. With $P^\pm := \exp(\mathfrak{p}^\pm)$ we have $G \subset P^+ K_{\mathbb{C}} P^- \cap H_{\mathbb{C}} K_{\mathbb{C}} P^+$, and for $x \in P^+ K_{\mathbb{C}} P^-$ respectively $H_{\mathbb{C}} K_{\mathbb{C}} P^+$ we define $p^\pm(x) \in P^\pm$, $k_{\mathbb{C}}(x) \in K_{\mathbb{C}}$, and $k_H(x) \in K_{\mathbb{C}}$ (only the class in $K_{\mathbb{C}} \cap H_{\mathbb{C}} \backslash K_{\mathbb{C}}$ is well-defined) by

$$x = p^+(x) k_{\mathbb{C}}(x) p^-(x)$$

respectively

$$x \in H_{\mathbb{C}} k_H(x) P^+.$$

Let (δ, V_δ) be a holomorphic irreducible representation of $K_{\mathbb{C}}$ with nonzero $K_{\mathbb{C}}^\tau$ -fixed vector ν and $\delta|K$ unitary. For $\check{\delta}$ the contragredient representation, $\nu^0 \in V_{\check{\delta}}^*$ a $K_{\mathbb{C}}^\tau$ -invariant vector with $\langle \nu, \nu^0 \rangle = 1$, we define

$$\Phi_\delta : P^+ K_{\mathbb{C}} H_{\mathbb{C}} / H_{\mathbb{C}} \rightarrow \mathbb{C}, \quad gH \mapsto \langle \nu, \check{\delta}(k_H(g^{-1})^{-1})\nu^0 \rangle.$$

This is a well-defined holomorphic function because ν^0 is $K_{\mathbb{C}} \cap H_{\mathbb{C}}$ -invariant. We denote the restriction of Φ_δ to G/H simply by Φ_δ . With this convention notice that $\Phi_\delta|(G_s/H_s) = \Phi_{\delta_s}$ with $\delta_s = \delta|(K_{s,\mathbb{C}})$. Let $Z_{\mathbb{C}}$ be the complex torus given by $\exp \mathfrak{z}_{\mathbb{C}}$. Define $\chi_\delta : Z_{\mathbb{C}} \rightarrow \mathbb{C}$ by

$$\delta(z) = \chi_\delta(z) \text{id}.$$

Theorem 3.3 ([**ÓØ91**]). *Let μ be the highest weight of δ . Then the following holds:*

- (1) *The function Φ_δ is in $L^2(G/H)$ if and only if $\langle \mu + \rho, \alpha \rangle < 0$ for all $\alpha \in \Delta_n^+$.*
- (2) *Suppose that $\langle \mu + \rho, \alpha \rangle < 0$ for all $\alpha \in \Delta_n^+$. Let $H_\delta \subset L^2(G/H)$ be the G -module generated by Φ_δ . Then H_δ is an irreducible highest weight module.*
- (3) *Let \mathbb{C}_δ be the $Z_{\mathbb{C}}$ -module corresponding to χ_δ and let H_{δ_s} be the G_s -module generated by Φ_{δ_s} , then $H_\delta \simeq \mathbb{C}_\delta \otimes H_{\delta_s}$.*
- (4) *Let $E \subset L^2(G/H)$ be an irreducible highest weight module. Then there exists a representation δ such that $E \simeq H_\delta$ and the multiplicity of E in $L^2(G/H)$ is one.*

Proof. (1) For $g \in G$ choose $z \in Z$ and $g_s \in G_s$ such that $g = zg_s$. Then $\Phi_\delta(gH) = \chi_\delta(z)\Phi_{\delta_s}(g_sH_s)$, in particular

$$|\Phi_\delta(gH)| = |\Phi_{\delta_s}(g_sH)|.$$

Hence $\Phi_\delta \in L^2(G/H)$ if $\langle \mu + \rho, \alpha \rangle < 0$ for all $\alpha \in \Delta_n^+$ by [**ÓØ91**, Theorem 5.2] and [**HÓØ91**, Theorem 3.3].

(2), (3) Denote the G_s -module generated by Φ_{δ_s} in $L^2(G_s/H_s)$ by H_{δ_s} . Then

$$H_\delta \simeq \mathbb{C}_{\chi_\delta} \otimes H_{\delta_s}$$

as a $Z \times G_s$ module. The module on the right hand side is an irreducible highest weight module by [**ÓØ91**, Theorem 5.2]. Hence H_δ must be irreducible.

(4) Assume that $E \subset L^2(G/H)$ is an irreducible highest weight module. Then $E \simeq \mathbb{C}_\chi \otimes E_s$, where E_s is the restriction of E to G_s and \mathbb{C}_χ is the central character of E . The module E_s is an irreducible highest weight module for G_s , and $E_s \subset L^2(G_s/H_s)$. By [**HÓØ91**, Theorem 3.3] it follows that $E_s \simeq H_{\delta_s}$ for some irreducible $H_{s\mathbb{C}} \cap K_{s\mathbb{C}}$ -spherical representation of $K_{s\mathbb{C}}$. Let $\delta = \chi \otimes \delta_s$. Then δ is an irreducible $K_{\mathbb{C}} \cap H_{\mathbb{C}}$ -spherical representation of $K_{\mathbb{C}}$ and $E \simeq H_\delta$. The statement about the multiplicity follows now from [**ÓØ91**, Theorem 7.2]. □

These representations ϵ_δ on H_δ form the holomorphic discrete series of G/H . Let π be a unitary representation of G on the Hilbert space V . Denote the space of smooth vectors by V^∞ . Denote by π^∞ the derived representation of \mathfrak{g} on V^∞ . Define the *cone of negative elements* by

$$C(\pi) := \{X \in \mathfrak{g} \mid \forall u \in V^\infty : (i\pi^\infty(X)u \mid u) \leq 0\}.$$

Then $C(\pi)$ is a G -invariant cone in \mathfrak{g} . The representation π is called *W-admissible* if $C(\pi) \supset W$. If π is W -admissible then π extends to a holomorphic representation of $\Gamma(W)$. (In fact, this is true for W replaced by an arbitrary invariant cone.)

Proposition 3.4. *For $W = W_k \cap \mathfrak{g}$ the representation $(\epsilon_\delta, H_\delta)$ is W -admissible if and only if*

$$i\mu(\mathfrak{a} \cap C^o) = (i d\chi_\delta + i\mu_s)(\mathfrak{a} \cap C^o) \leq 0.$$

Proof. Since $W^o = \text{Ad}(G)(\mathfrak{t} \cap W^o)$ and ϵ_δ is unitary, W -admissibility is equivalent to

$$(*) \quad (i\epsilon_\delta^\infty(\mathfrak{t} \cap W^o)u \mid u) \leq 0$$

for all $u \in H_\delta^\infty$. Let $W(\mathfrak{k}, \mathfrak{t})$ be the Weyl group of $\Delta(\mathfrak{k}, \mathfrak{t})$. As H_δ is an irreducible highest weight module we have by Poincaré-Birkhoff-Witt $\mu_\lambda = \mu_+ - \sum_{\alpha \in \Delta^+(\mathfrak{p}_\mathbb{C}, \mathfrak{t}_\mathbb{C})} n_\alpha \alpha$, with $n_\alpha \geq 0$ and $\mu_+ \in \text{conv}(W(\mathfrak{k}, \mathfrak{t})\mu)$, for all occurring weights μ_λ . Since $\mathfrak{t} \cap W \subset \mathfrak{t}_1 \cap W_k = -\sum_{\alpha \in \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})} \mathbb{R}_0^+ iH_\alpha$ (the $H_\alpha \in \mathfrak{t}_{1\mathbb{C}}$ to $\alpha \in \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$ defined as the H_α to $\alpha \in \Delta$) ([HÓ96, p. 102]) such that $i\alpha(\mathfrak{t} \cap W) \geq 0$ for all noncompact roots $\alpha \in \Delta^+(\mathfrak{p}_\mathbb{C}, \mathfrak{t}_\mathbb{C})$, the condition $(*)$ remains to be checked for the vectors in $\text{conv}(W(\mathfrak{k}, \mathfrak{t})\mu)$. Now, since W is G -invariant, $\mathfrak{t} \cap W^o$ is $W(\mathfrak{k}, \mathfrak{t})$ -invariant, and we have to check $(*)$ only for μ . Finally, since δ is $K_\mathbb{C} \cap H_\mathbb{C}$ -spherical, we have $\mu|_{\mathfrak{h}} = 0$ and $(*)$ reduces to the claimed condition. \square

Remark. For \mathfrak{g} semisimple and $\mathfrak{a} \cap C = c_{\min}$ it is known by [HÓ091] that all $(\epsilon_\delta, V_\delta)$ are admissible.

Let $\Xi(C) := \Gamma(-W)x_o \subset G_\mathbb{C}/H_\mathbb{C}$, where $x_o = eH_\mathbb{C}$. As in [HÓ091, Lemma 1.3] we have $\Xi(C) \simeq G \times_H (-iC)$ and for $\Xi(C^o) := \Gamma(-W^o)x_o$ we have $\Xi(C)^o = \Xi(C^o)$. Thus $\Xi(C^o)$ is an open complex submanifold of $G_\mathbb{C}/H_\mathbb{C}$. Notice that $\Gamma(-W) = \Gamma(W)^{-1}$. As $\Gamma(W)\Gamma(W^o) \subset \Gamma(W^o)$ it follows that $\Gamma(-W^o)\Xi(C) \subset \Xi(C^o)$. Thus $\Gamma(W)$ acts on functions defined on $\Xi(C)$ respectively $\Xi(C^o)$ by

$$(\gamma \cdot f)(x) = f(\gamma^{-1}x)$$

and $(\gamma \cdot f)|_{G/H}$ is well-defined for $\gamma \in \Gamma(W^o)$. Define the *Hardy norm* on holomorphic functions on $\Xi(C^o)$ by

$$\|f\|_H := \sup_{\gamma \in \Gamma^o} \|\gamma \cdot f\|_{L^2(G/H)}.$$

Definition. The Hardy space $H_2(C)$ on G/H is

$$H_2(C) := \{f : \Xi(C^o) \rightarrow \mathbb{C} \mid f \text{ is holomorphic and } \|f\|_H < \infty\}.$$

Denote by T the representation of $\Gamma(W)$ in $H_2(C)$ induced by the left-regular action and by λ the left-regular representation of G in $L^2(G/H)$. The proof of the following theorem then generalizes without any change (or simply using again that $G = ZG_s$).

Theorem 3.5 ([HÓØ91]). (1) *The Hardy space $H_2(C)$ is a Hilbert space with norm $\|\cdot\|_H$.*

(2) *There is an isometry $\beta : H_2(C) \rightarrow L^2(G/H)$ given by*

$$\beta(f) = \lim_{\Gamma(W^o) \ni \gamma \rightarrow 1} \gamma \cdot f,$$

where the limit is in $L^2(G/H)$.

(3) *The boundary value map β is an intertwining operator for the G -actions, i.e., $\beta T(g) = \lambda(g) \beta$ for $g \in G$.*

(4) *The representation T is a holomorphic representation of $\Gamma(W)$ in $H_2(C)$.*

(5) *The image of β in $L^2(G/H)$ is the direct sum of all the holomorphic discrete series that are W -admissible and each occurs with multiplicity one.*

(6) *Assume that $w \in \Xi(C^o)$. Then the evaluation map $H_2(C) \ni f \mapsto f(w) \in \mathbb{C}$ is continuous. Let $K_w \in H_2(C)$ be such that $f(w) = (f \mid K_w)$, where $(\cdot \mid \cdot)$ denotes the scalar product on $H_2(C)$. Then the map $(z, w) \mapsto K(z, w) := K_w(z)$ is holomorphic in the first variable and antiholomorphic in the second variable. We have:*

a) $K(w, z) = K(z, w)$.

b) Let $\gamma \in \Gamma(-W)$, and let $z, w \in \Xi(C^o)$. Then

$$K(\gamma \cdot z, w) = K(z, \gamma^* \cdot w),$$

with $(g \exp iX)^ := \exp(iX)g^{-1}$.*

c) *There exists a holomorphic H -invariant function $\Theta_K : \Xi(C^o) \rightarrow \mathbb{C}$ such that*

$$K(\gamma_1 \cdot x_o, \gamma_2 \cdot x_o) = \Theta_K(\gamma_2^* \gamma_1 \cdot x_o).$$

(7) *Suppose that $z = \gamma \cdot x_o \in \Xi(C^o)$. Then $w \mapsto K(z, w)$ extends to a smooth map on $\Xi(C)$, and the inverse of β is given by*

$$F(z) = \int_{G/H} f(x) K(z, x) dx = \int_{G/H} f(\dot{g}) \Theta_K(g^{-1} \gamma \cdot x_o) d\dot{g},$$

where $\dot{g} = g \cdot x_o$, and $d\dot{g}$ is the G -invariant measure on G/H normalized by $d\dot{g} = dh d\dot{g}$.

We note for later use that the extension of Φ_δ from G/H to $\Xi(C^o)$ is given by the defining formula, as $\Gamma(-W) \subset H_{\mathbb{C}} K_{\mathbb{C}} P^+$ ([HÓØ91, Lemma 3.6]).

Example (continued). Since $\mathfrak{g} = \mathfrak{k} \cap \mathfrak{q} = \mathfrak{a}$ we have by Proposition 2.4

$$C^o = \mathbb{R}^+ Z^0 = \mathbb{R}^+ \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} = W^o$$

and further

$$\Gamma(-W) = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \mid |a| \leq 1 \right\}.$$

The extension of $\Phi : G/H \rightarrow \check{S}_1$ to a $G_{\mathbb{C}}$ -invariant holomorphic map is

$$\Phi : G_{\mathbb{C}}/H_{\mathbb{C}} \rightarrow \mathbb{C} \quad \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} H_{\mathbb{C}} \mapsto -a^2,$$

such that

$$\Phi(\Xi(C)^o) = \Phi(\Gamma(-W^o)x_o) = \{z \in \mathbb{C} \mid 0 < |z| < 1\}.$$

We describe the Hardy space using this explicit realization of $\Xi(C^o)$. Since G operates by rotations, the Hardy norm is given by

$$\|f\|_H = \sup_{0 < |z| < 1} \frac{1}{\pi} \int_0^\pi |f(e^{2i\varphi}z)|^2 d\varphi$$

and, using the Laurent series representation of f , we get

$$H_2(C) = \left\{ f : \{z \in \mathbb{C} \mid 0 < |z| < 1\} \rightarrow \mathbb{C} \mid f(z) = \sum_{n=0}^\infty a_n z^n \right. \\ \left. \text{with } \sum_{n=0}^\infty |a_n|^2 < \infty \right\}.$$

We can describe this space also as sum of W -admissible representations. Identifying G/H via Φ with S^1 , the holomorphic discrete series representations are the characters

$$\Phi_n : S^1 \rightarrow \mathbb{C}, \quad z \mapsto z^n, \quad n \in \mathbb{Z}.$$

All these functions have obviously a holomorphic continuation to $\Xi(C^o)$. Checking which of these representations are W -admissible, we get the condition $n \geq 0$, proving Theorem 3.5(5) for this case.

4. Characterization of $\Phi(G/H)$.

The Weyl group for the root system $\Delta(\mathfrak{k}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$ of compact roots acts by permutations on the positive noncompact roots $\Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$. Therefore

$$\rho_{1,n} := \frac{1}{2} \sum_{\alpha \in \Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})} \dim_{\mathbb{C}} \mathfrak{g}_{1\mathbb{C},\alpha} \alpha \in i\mathfrak{t}_1^*$$

is zero on the orthogonal complement of $\mathfrak{z}(\mathfrak{k}_1)$ in \mathfrak{t}_1 . We define $\Delta_1 := \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}}^-)$ and the positive roots Δ_1^+ by restricting the roots $\Delta^+(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$

to $\mathfrak{t}_{1\mathbb{C}}^-$. Since $(\mathfrak{g}_1, \mathfrak{k}_1, \theta)$ is assumed to be Hermitian symmetric of tube type, and, by abuse of notation, identifying the strongly orthogonal roots γ_i with elements of $(\mathfrak{t}_{1\mathbb{C}}^-)^*$, by the Theorem of Moore ([H78, p. 528] or [H94, p. 460])

$$\Delta_1 = \left\{ \pm \frac{1}{2}(\gamma_i + \gamma_j) \mid 1 \leq i \leq j \leq r \right\} \cup \left\{ \pm \frac{1}{2}(\gamma_i - \gamma_j) \mid 1 \leq i < j \leq r \right\}.$$

Thereby the positive noncompact roots are

$$\Delta_{1,n}^+ = \left\{ \frac{1}{2}(\gamma_i + \gamma_j) \mid 1 \leq i \leq j \leq r \right\},$$

the root spaces $\mathfrak{g}_{1\mathbb{C}, \gamma_i}$ are one dimensional, and root spaces corresponding to roots $\frac{1}{2}(\gamma_i \pm \gamma_j)$, $i \neq j$, have all the same dimension d . As the restriction of a noncompact root to $\mathfrak{t}_{1\mathbb{C}}^-$ is always nonzero, we get immediately

$$\rho_{1,n} = \frac{1}{2} \left(1 + \frac{d(r-1)}{2} \right) (\gamma_1 + \dots + \gamma_r).$$

For the moment we prefer to work with another Cartan algebra than \mathfrak{t}_1 . Therefore, let \mathfrak{t}_1^+ be the orthogonal complement of \mathfrak{t}_1^- in \mathfrak{t}_1 , then the Cayley transform is the identity on $\mathfrak{t}_{1\mathbb{C}}^+$ and maps $\mathfrak{t}_{1\mathbb{C}}^-$ onto $\mathfrak{a}_{p\mathbb{C}}$. For $\alpha \in \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$ we define the Cayley transformed root α^c by

$$\alpha^c := \alpha \circ \text{Ad}(c^{-1}) \in \Delta(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}}^+ \oplus \mathfrak{a}_{p\mathbb{C}}) =: \Delta_1^c,$$

and the positive system $\Delta_1^{c,+}$ by the transformed roots of $\Delta^+(\mathfrak{g}_{1\mathbb{C}}, \mathfrak{t}_{1\mathbb{C}})$.

We define

$$\rho_+ := \text{Ad}(c^{-1})^* \rho_{1,n} = \frac{1}{2} \left(1 + \frac{d(r-1)}{2} \right) (\gamma_1^c + \dots + \gamma_r^c) \in (\mathfrak{t}_{1\mathbb{C}}^+ + \mathfrak{a}_{p\mathbb{C}})^*,$$

which is zero on the orthogonal complement of $\mathbb{C}X^0$. We denote by (π_m, V_m) the irreducible finite-dimensional representation of $G_{1\mathbb{C}}$ with lowest weight $-m\rho_+$, if it exists.

We will need the *Riemannian dual* algebra \mathfrak{g}_1^d associated to $(\mathfrak{g}_1, \mathfrak{g}, \sigma)$ for the proof of the next proposition

$$\mathfrak{g}_1^d := \mathfrak{k}_1 \cap \mathfrak{g} + i(\mathfrak{p}_1 \cap \mathfrak{g}) + \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} + i(\mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma}).$$

The involution σ restricted to \mathfrak{g}_1^d is a Cartan involution. Hence $\mathfrak{k}_1^d := \mathfrak{k}_1 \cap \mathfrak{g} + i(\mathfrak{p}_1 \cap \mathfrak{g})$ is a maximal compactly embedded subalgebra of \mathfrak{g}_1^d and the corresponding orthogonal complement is

$$\mathfrak{p}_1^d := \mathfrak{p}_1 \cap \mathfrak{q}_{1\sigma} + i(\mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma}).$$

Let $\hat{\mathfrak{a}} \subset \mathfrak{k}_1 \cap \mathfrak{q}_{1\sigma} \cap \mathfrak{z}_{\mathfrak{g}_1}(\mathfrak{b})$ be maximal abelian, then $\mathfrak{a}^d := i\hat{\mathfrak{a}} + \mathfrak{b}$ is maximal abelian in \mathfrak{p}_1^d . For the root system $\Delta_1^d := \Delta(\mathfrak{g}_1^d, \mathfrak{a}^d)$ we choose the positive roots $\Delta_1^{d,+} := \Delta^+(\mathfrak{g}_1^d, \mathfrak{a}^d)$ compatible with $\Delta^+(\mathfrak{g}_1^d, \mathfrak{b}) = \Delta^+(\mathfrak{g}_1, \mathfrak{b}) = \{\alpha|_{\mathfrak{b}} \mid \alpha \in \Delta_1^{c,+}, \alpha|_{\mathfrak{b}} \neq 0\}$.

Proposition 4.1. *Let the notation be as above, then the following holds:*

- (1) *There exists an irreducible finite-dimensional representation (π_2, V_2) of $G_{1\mathbb{C}}$ with lowest weight $-2\rho_+$.*
- (2) *Assume that $\mathfrak{g}_1 \neq \mathfrak{sp}(2n, \mathbb{R}), \mathfrak{so}(2, 2k + 1)$, $n, k \geq 1$. Then there exists an irreducible finite-dimensional representation (π_1, V_1) of $G_{1\mathbb{C}}$ with lowest weight $-\rho_+$.*
- (3) *The weight space $V_{m, -m\rho_+}$ is left pointwise fixed by the identity component of $M'_\mathbb{C}$, where $M'_\mathbb{C}$ is defined by the Langlands decomposition $P' \simeq M' \times \exp(\mathbb{R}X^0) \times Q_{1\eta}^-$. For $m = 4$ it is fixed pointwise by $M'_\mathbb{C}$ and, if $\mathfrak{g}_1 \neq \mathfrak{sp}(2n, \mathbb{R}), \mathfrak{so}(2, 2k + 1)$, $n, k \geq 1$, the same is true for $m = 2$.*
- (4) *There is a $m \in \{1, 2, 4, 8\}$ such that π_m is $G_\mathbb{C}$ -spherical. In this case, let $(\cdot | \cdot)$ be a scalar product, invariant under the analytic subgroup to $\mathfrak{k}_1 + \mathfrak{ip}_1$, then we can choose $v_m \in V_{m, -m\rho_+}$ and $\xi_m \in V_m^{G_\mathbb{C}}$ such that $(v_m | \xi_m) = 1$.*

Proof. (1), (2) The first two assertions are [ÓØ99, Theorem 2.6].

(3) Let $\tilde{\mathfrak{a}}$ be the orthogonal complement of $\mathbb{R}X^0$ in \mathfrak{a}_p and $\Delta_0 := \{\alpha \in \Delta_1^c \mid \alpha(X^0) = 0\}$ then

$$\mathfrak{m}_\mathbb{C} = \mathfrak{t}_{1\mathbb{C}}^+ + \tilde{\mathfrak{a}}_\mathbb{C} + \sum_{\alpha \in \Delta_0} \mathfrak{g}_{1\mathbb{C}, \alpha}.$$

Every lowest weight vector is invariant under $\mathfrak{t}_{1\mathbb{C}}^+ + \tilde{\mathfrak{a}}$ and elements of root spaces $\mathfrak{g}_{1\mathbb{C}, \alpha}$ with $-\alpha \in \Delta_1^{c,+}$. By the invariance of the weights under the Weyl group, for $\alpha \in \Delta_1^{c,+}$ with $\langle \alpha, \rho_+ \rangle = 0$ the weight space to $-m\rho_+ + \alpha = s_\alpha(-m\rho_+ - \alpha)$ is trivial, hence $\mathfrak{m}_\mathbb{C}$ operates trivial on $V_{m, -m\rho_+}$. By the Theorem of Helgason $\pi_m(M_{\min})$, with $M_{\min} := Z_{K_1}(\mathfrak{a}_p)$, leaves the weight space pointwise fixed if and only if $\frac{\langle m\rho_+, \alpha \rangle}{\langle \alpha, \alpha \rangle} \in \mathbb{Z}^+$ for all $\alpha \in \Delta_1^{c,+}$. For $\alpha = \gamma_j^c$ this quotient is $\frac{m}{2} \left(1 + \frac{d(r-1)}{2}\right)$, for $\alpha = \frac{1}{2}(\gamma_i^c + \gamma_j^c)$ it is $m \left(1 + \frac{d(r-1)}{2}\right)$, and for $\alpha = \frac{1}{2}(\gamma_i^c - \gamma_j^c)$ it vanishes. This proves the invariance under M_{\min} , for m as in the assertion, by looking at the possible combinations for d and r ([H78, p. 530ff] or the table in [ÓØ99]). By $M_\mathbb{C} = M_{\min}\langle \exp(\mathfrak{m}_\mathbb{C}) \rangle$ the result now follows.

(4) By Helgason’s Theorem π_m is $G_\mathbb{C}$ -spherical if and only if

$$\frac{\langle m\rho_+, \alpha \rangle}{\langle \alpha, \alpha \rangle} \in \mathbb{Z}^+, \quad \forall \alpha \in \Delta_1^{d,+}.$$

By definition $2\rho_{1,n} | \mathfrak{b}$ is a linear combination of elements of $\Delta^+(\mathfrak{p}_{1\mathbb{C}}, \mathfrak{b})$ with positive integer coefficients. Therefore $2\rho_+$ is a linear combination with positive integer coefficients of elements of $\Delta^+(\mathfrak{g}_1, \mathfrak{b})$. Now by [B88, p. 362]

the system $\Delta(\mathfrak{g}_1, \mathfrak{b})$ is a root system in the axiomatic sense such that

$$2 \frac{\langle 2\rho_+, \tilde{\alpha} \rangle}{\langle \tilde{\alpha}, \tilde{\alpha} \rangle} \in \mathbb{Z} \quad \text{for } \tilde{\alpha} \in \Delta(\mathfrak{g}_1, \mathfrak{b}).$$

For $\tilde{\alpha} := \alpha|_{\mathfrak{b}}$, with $\alpha \in \Delta_1^d$, and $\tilde{\alpha} \neq 0$ by [Ó87, Lemma 2.3] we have

$$\langle \alpha, \alpha \rangle = n \langle \tilde{\alpha}, \tilde{\alpha} \rangle, \quad n \in \{1, 2, 4\},$$

and $2\alpha \in \Delta_1^d$ if $n = 4$. Since a root 2α gives a more severe restriction than α we can assume $n \in \{1, 2\}$, which implies the first assertion. The second assertion then is a consequence of the proof of Helgason’s Theorem [H84, p. 534ff]. \square

Example. We consider the causal compactification of $SO(2, n)/SO(1, n)$, as treated in [B97, Example 3.3]. Here $G_1 = SO(2, n + 1)$, i.e.,

$$G_1 = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in SL(n + 3, \mathbb{R}) \left| \begin{array}{l} {}^tAA - {}^tCC = I, \quad {}^tAB = {}^tCD \\ {}^tDD - {}^tBB = I \end{array} \right. \right\},$$

with Cartan involution $\theta(g) = {}^t g^{-1}$, $X^0 = 2(E_{1, n+3} + E_{n+3, 1})$, and

$$\sigma = \text{Ad} \begin{pmatrix} \text{id}_{n+2} & 0 \\ 0 & -1 \end{pmatrix}.$$

Then $\mathfrak{q}_{1\sigma}$ consists of matrices with nonzero entries only in the last row respectively column such that $\mathfrak{b} = \mathbb{R}X^0$ is maximal abelian in \mathfrak{p}_1^d , i.e., $\mathfrak{a}^d = \mathfrak{b}$. For the Riemannian dual algebra \mathfrak{g}_1^d we get therefore, restricting the roots of $\Delta^+(\mathfrak{g}_1, \mathfrak{a}_p) = \{\gamma_1^c, \gamma_2^c, \frac{1}{2}(\gamma_1^c \pm \gamma_2^c)\}$ to \mathfrak{b} , the positive system $\Delta^+(\mathfrak{g}_1^d, \mathfrak{b}) = \Delta^+(\mathfrak{g}_1, \mathfrak{b}) = \{\frac{1}{2}(\gamma_1^c + \gamma_2^c)\}$. By Helgason’s Theorem, applied to $(\mathfrak{g}_1^d, \mathfrak{b})$, the condition for π_m to be G -spherical is now

$$m \left(1 + \frac{d(r - 1)}{2} \right) \in \mathbb{Z}^+.$$

For our \mathfrak{g}_1 here $r = 2$ and $d = n - 1$ ([H78, p. 530ff] or the table in [Ó099]) such that π_2 is always G -spherical, whereas π_1 only for n odd.

In a similar fashion one calculates the minimal m ’s for the other causal compactifications as given in the following table. Here p and q are nonnegative integers with $p + q = n$. In the case $(\mathfrak{so}(2, q) \oplus \mathfrak{so}(p + 1), \mathfrak{so}(1, q) \oplus \mathfrak{o}(p), \mathfrak{so}(2, n + 1))$ we also assume that $p \geq 2$:

$(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1)$	minimal m
$(\mathfrak{su}(p, q) \oplus \mathfrak{u}(p, q), \mathfrak{su}(p, q), \mathfrak{su}(n, n))$	1 for n even 2 for n odd
$(\mathfrak{u}(2p, 2q), \mathfrak{sp}(p, q), \mathfrak{so}^*(4n))$	2
$(\mathfrak{u}(p, q), \mathfrak{so}(p, q), \mathfrak{sp}(n, \mathbb{R}))$	1 for $n \equiv 3 \pmod{4}$ 2 for $n \equiv 1 \pmod{4}$ 4 for n even
$(\mathfrak{so}(2, n-1) \oplus \mathfrak{so}(2), \mathfrak{so}(1, n-1), \mathfrak{so}(2, n+1))$	1 for $n \equiv 3 \pmod{4}$ 2 for $n \equiv 1 \pmod{4}$ 4 for n even
$(\mathfrak{e}_{6(-14)} \oplus \mathfrak{so}(2), \mathfrak{f}_{4(-20)}, \mathfrak{e}_{7(-25)})$	2
$(\mathfrak{so}^*(2n) \oplus \mathfrak{so}^*(2n), \mathfrak{so}^*(2n), \mathfrak{so}^*(4n))$	2
$(\mathfrak{sp}(n, \mathbb{R}) \oplus \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(2n, \mathbb{R}))$	4
$(\mathfrak{so}^*(2n), \mathfrak{so}(n, \mathbb{C}), \mathfrak{su}(n, n))$	1 for n even 2 for n odd
$(\mathfrak{sp}(2n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{C}), \mathfrak{su}(2n, 2n))$	1
$(\mathfrak{so}(2, q) \oplus \mathfrak{so}(p+1), \mathfrak{so}(1, q) \oplus \mathfrak{o}(p), \mathfrak{so}(2, n+1))$	1 for $n \equiv 3 \pmod{4}$ 2 for $n \equiv 1 \pmod{4}$ 4 for n even
$(\mathfrak{g}, \mathfrak{h}, \mathfrak{g} \times \mathfrak{g})$, for $(\mathfrak{g}, \mathfrak{h})$ of Cayley type	1 for $\mathfrak{g} \neq \mathfrak{so}(2, 2k+1), \mathfrak{sp}(2n, \mathbb{R})$ 2 for $\mathfrak{g} = \mathfrak{so}(2, 2k+1), \mathfrak{sp}(2n, \mathbb{R})$
$(\mathfrak{so}(2, n), \mathfrak{so}(1, n), \mathfrak{so}(2, n+1))$	1 for n odd 2 for n even

For G_1 we define $P_1^\pm, K_{1\mathbb{C}}$, and the functions p_1^\pm and $k_{1\mathbb{C}}$ in the same manner as for G . Since the map $P_1^+ \times K_{1\mathbb{C}} \times P_1^- \rightarrow G_{1\mathbb{C}}, (p^+, k, p^-) \mapsto p^+kp^-$, is a diffeomorphism onto a dense subset we can define

$$\zeta_1 : P_1^+ K_{1\mathbb{C}} P_1^- \rightarrow \mathfrak{p}_1^+ \quad g \mapsto \exp^{-1} p_1^+(g).$$

We will use the Harish-Chandra realization $D_1 := \zeta_1(GK_{1\mathbb{C}}P_1^-)$ of G_1/K_1 in \mathfrak{p}_1^+ and ζ_1 to state our results in a particularly nice form and note therefore:

Proposition 4.2. *In the Harish-Chandra realization $\check{S}_1 = \zeta_1(G_1c)$ and the image $\Phi(G/H)$ of the causal compactification is given by $\zeta_1(Gc)$.*

Proof. The first assertion is [KW65, Theorem 3.6], the second then follows from Theorem 2.3. □

To avoid a clumsy notation we will assume from now on, using the Proposition, that $\Phi(G/H) \subset \mathfrak{p}_1^+$. Define the *canonical cocycle*

$$J_1(g, z) = J_1^1(g, z) := k_{1\mathbb{C}}(g \exp z)^{2\rho_{1,n}},$$

where $(\cdot)^{2\rho_{1,n}}$ is the central character of $K_{1\mathbb{C}}$ defined by $2\rho_{1,n}$. More general, assume $m \geq 1$ such that (π_m, V_m) exists then

$$J_1^{m/2}(g, z) := k_{1\mathbb{C}}(g \exp z)^{m\rho_{1,n}}$$

is defined. If (π_m, V_m) is G -spherical define

$$\psi_m : \mathfrak{p}_1^+ \rightarrow \mathbb{C} \quad z \mapsto (\pi_m(\exp(z)c^{-1})v_m|\xi_m).$$

Theorem 4.3. *Assume $m \geq 1$ such that (π_m, V_m) exists and is G -spherical. Then the function $\psi_m(z)$ is holomorphic on \mathfrak{p}_1^+ and has the following properties:*

- (1) $\Phi(G/H) = \{z \in \check{S}_1 \mid \psi_m(z) \neq 0\}$.
- (2) For $g \in G_{\mathbb{C}}, z \in \mathfrak{p}_1^+$, such that $g \cdot z$ is defined, we have

$$\psi_m(g \cdot z) = J_1^{m/2}(g, z) \psi_m(z).$$

Proof. (1) For $z = \zeta_1(gc) \in \check{S}_1$, with $g \in G_1$, we have

$$\begin{aligned} \psi_m(z) &= (\pi_m(p_1^+(gc)c^{-1})v_m|\xi_m) \\ &= (\pi_m(gc p_1^-(gc)^{-1}k_{1\mathbb{C}}(gc)^{-1}c^{-1})v_m|\xi_m). \end{aligned}$$

Since $\text{Ad}(c)iZ^0 = \frac{1}{2}X^0$, the Cayley transformed group $\text{Ad}(c)(K_{1\mathbb{C}}P_1^-)$ is $P'_{\mathbb{C}}$, where we denote by $\text{Ad}(c)$ the inner automorphism $x \mapsto cxc^{-1}$ of $G_{1\mathbb{C}}$. In [B97] it is shown that GP' is the only open (G, P') -double coset in G_1 . Then by [BD92, Lemme 4] we have $(\pi_m(gp)v_m|\xi_m) \neq 0$, for $p \in P'_{\mathbb{C}}$, if and only if $g \in G$. (As M'/M'_0 is finite, and using that every component contains an element of the form $\exp i\lambda X^0$, the cited result is still valid with obvious modifications in its proof, even if v_m is not $M'_{\mathbb{C}}$ -fixed. Note also, that we use a lowest weight vector v_m , since the nilradical $\mathfrak{q}_{1\eta}^-$ of our parabolic algebra \mathfrak{p}' is the sum of root spaces for negative roots.)

(2) Since \mathfrak{p}_1^- is $K_{1\mathbb{C}}$ -invariant and mapped by $\text{Ad}(c)$ onto $\mathfrak{q}_{1\eta\mathbb{C}}^-$, we have

$$\begin{aligned} \psi_m(g \cdot z) &= (\pi_m(g \exp z p_1^-(g \exp z)^{-1}k_{1\mathbb{C}}(g \exp z)^{-1}c^{-1})v_m|\xi_m) \\ &= (\pi_m(\exp z p_1^-(g \exp z)^{-1}k_{1\mathbb{C}}(g \exp z)^{-1}c^{-1})v_m|\pi_m(\theta(g)^{-1})\xi_m) \\ &= (\pi_m(\exp z k_{1\mathbb{C}}(g \exp z)^{-1}c^{-1}\text{Ad}(ck_{1\mathbb{C}}(g \exp z)))(p_1^-(g \exp z)^{-1})v_m|\xi_m) \\ &= (\pi_m(\exp z c^{-1}\text{Ad}(c)(k_{1\mathbb{C}}(g \exp z)^{-1}))v_m|\xi_m) \\ &= J_1^{m/2}(g, z)(\pi_m(\exp zc^{-1})v_m|\xi_m). \end{aligned}$$

□

Example. We want to describe the causal compactification for G/H of Cayley type. Let $G_1 = \tilde{G} \times \tilde{G}$, with $\tilde{G} \subset \tilde{G}_{\mathbb{C}}$ simply connected and $(\tilde{\mathfrak{g}}, \tilde{\mathfrak{k}}, \tilde{\theta})$ Hermitian symmetric of tube type. With $\sigma(g, h) := (h, g)$ the fixpoint group G of this involution is isomorphic to \tilde{G} . For $\tilde{\mathfrak{a}}_p \subset \tilde{\mathfrak{p}}$ maximal abelian and

$\theta = \tilde{\theta} \times \tilde{\theta}$, we have $\mathfrak{a}_p = \tilde{\mathfrak{a}}_p \times \tilde{\mathfrak{a}}_p$ and $\mathfrak{b} = \{(X, -X) \mid X \in \tilde{\mathfrak{a}}_p\}$. With $\tilde{\Delta} := \Delta(\tilde{\mathfrak{g}}, \tilde{\mathfrak{a}}_p)$ we can define $\tilde{\rho}_+$ and get then $\rho_+ = (\tilde{\rho}_+, -\tilde{\rho}_+)$. Given $(\tilde{\pi}_m, \tilde{V}_m)$, a representation of $\tilde{G}_{\mathbb{C}}$ with lowest weight $-m\tilde{\rho}_+$, the representation $\pi_m := \tilde{\pi}_m \otimes \tilde{\pi}_m^*$, where $\tilde{\pi}_m^*$ is the contragredient representation to $\tilde{\pi}_m$, has lowest weight $-m\rho_+$. Moreover every such representation has the G -fixed vector $\sum w_i \otimes w_i^*$, where $\{w_i \mid i = 1, \dots, \dim \tilde{V}_m\}$ is a base of \tilde{V}_m and $\{w_i^*\}$ its dual. We have now $c = (\tilde{c}, \tilde{c}^{-1})$ and for $(z, w) \in \mathfrak{p}_1^+$

$$\begin{aligned} \psi_m((z, w)) &= \left(\tilde{\pi}_m(\exp z \tilde{c}^{-1}) \otimes \tilde{\pi}_m^*(\exp w \tilde{c}) \cdot \tilde{v}_m \otimes \tilde{v}_m^* \mid \sum w_i \otimes w_i^* \right) \\ &= \sum (\tilde{\pi}_m((\exp w \tilde{c})^{-1} \exp z \tilde{c}^{-1}) \otimes 1 \cdot \tilde{v}_m \otimes \tilde{v}_m^* \mid w_i \otimes w_i^*) \\ &= (\pi_m(\tilde{c}^{-1} \exp(z - w) \tilde{c}^{-1}) v_m \mid v_m) \\ &= (\pi_m(\tilde{c}^{-2} \exp(z - w)) \tilde{\pi}_m(\tilde{c}^{-1}) v_m \mid \tilde{\pi}_m(\tilde{c}^{-1}) v_m). \end{aligned}$$

Here we used $\tilde{\mathfrak{t}}^+ + \tilde{\mathfrak{a}}_p$ as a Cartan algebra contrary to $\tilde{\mathfrak{t}}^+ + \tilde{\mathfrak{t}}^-$ used in [ÓØ99]. Therefore the lowest weight vectors are related by the Cayley transform and our ψ_m is the same function as those constructed in [ÓØ99].

5. Some L^2 -isometries.

We know that $\Phi(G/H) \subset \check{S}_1$ is open and dense. In this section we relate the L^2 -spaces with respect to invariant measures on these manifolds. The following results can be found in [ÓØ99]:

For \check{S}_1 in the Harish-Chandra realization the quasi-invariant measure is given by

$$\int_{\check{S}_1} f(z) d\mu(z) = \int_{K_1/K_1 \cap Z_{G_1}(X^0)} f(k \cdot \zeta_1(c)) dk.$$

With $P' = Z_{G_1}(X^0)Q_{1\eta}^-$ we can decompose every $g \in G_1$ in the form

$$g = k_1(g)h(g)q^-(g),$$

where $k_1(g) \in K_1$, $h(g) \in Z_{G_1}(X^0)$, and $q^-(g) \in Q_{1\eta}^-$. To $z \in \check{S}_1$ choose $k \in K_1$ such that $z = k \cdot \zeta_1(c)$ and define

$$J_R(g, z) := h(gk)^{2\rho_+} = \det \left(\text{Ad}(h(gk))|_{\mathfrak{q}_{1\eta}^-} \right)^{-1},$$

then

$$\int_{\check{S}_1} f(g \cdot z) J_R(g, z) d\mu(z) = \int_{\check{S}_1} f(z) d\mu(z)$$

and the relation

$$k_{1\mathbb{C}}(g \exp z)^{2\rho_{1,n}} = k_1(gk)^{2\rho_{1,n}} k^{-2\rho_{1,n}} h(gk)^{2\rho_+}$$

holds. Note that the two first factors on the right hand side have modulus one by the compactness of K_1 .

For $k_{1\mathbb{C}}$ we also have the cocycle property

$$k_{1\mathbb{C}}(gh \exp z) = k_{1\mathbb{C}}(g \exp(h \cdot z)) k_{1\mathbb{C}}(h \exp z),$$

which gives

$$J_1^{m/2}(gh, z) = J_1^{m/2}(g, h \cdot z) J_1^{m/2}(h, z)$$

when $J_1^{m/2}$ is defined. The following theorem now follows from the cocycle property for $J_1(\cdot, \cdot)$.

Theorem 5.1. (1) *The G -invariant measure on $\Phi(G/H) \subset \check{S}_1$ is up to normalization given by*

$$\int_{\check{S}_1} f(z) |\psi_m(z)|^{-2/m} d\mu(z).$$

(2) *A unitary representation of G_1 in $L^2(\check{S}_1)$ is given by*

$$(\lambda_0(g)f)(z) := \sqrt{|J_1(g^{-1}, z)|} f(g^{-1} \cdot z).$$

(3) *Identify G/H with $\zeta_1(\Phi(G/H))$ and let λ denote the left regular representation on G/H , then $f \mapsto f|\psi_m|^{1/m}$ is a G -equivariant isometry of $(L^2(\check{S}_1), \lambda_0|G)$ onto $(L^2(G/H), \lambda)$.*

If $\mathfrak{g}_1 \neq \mathfrak{sp}(2n, \mathbb{R}), \mathfrak{so}(2, 2k + 1)$, the holomorphic square root $\sqrt{J_1} = J_1^{1/2}$ is well-defined. We get then in the same way as the preceding theorem.

Theorem 5.2. (1) *If $\mathfrak{g}_1 \neq \mathfrak{sp}(2n, \mathbb{R}), \mathfrak{so}(2, 2k + 1)$, then*

$$(\lambda_1(g)f)(z) := \sqrt{J_1(g^{-1}, z)} f(g^{-1} \cdot z)$$

is a unitary representation of G_1 in $L^2(\check{S}_1)$.

(2) *If further π_1 is G -spherical then $f \mapsto f\psi_1$ is a G -equivariant isometry of $(L^2(\check{S}_1), \lambda_1|G)$ onto $(L^2(G/H), \lambda)$.*

In the general case, to define the roots $\sqrt{J_1(\cdot, \cdot)}$ and $\sqrt[m]{\psi_m}$, we have to replace G_1 and $\Phi(G/H)$ by appropriate coverings. For

$$\mathcal{D} := \{z \in \bar{D}_1 \mid \psi_m(z) \neq 0\},$$

which is independent of m by the last section, define a m -fold covering by

$$\tilde{\mathcal{D}} := \{(z, x) \in \mathcal{D} \times \mathbb{C} \mid \psi_m(z) = x^m\}.$$

We define then a holomorphic m -th root $\sqrt[m]{\psi_m}$ on $\tilde{\mathcal{D}}$ by

$$\sqrt[m]{\psi_m} : \tilde{\mathcal{D}} \rightarrow \mathbb{C} \quad (z, x) \mapsto x.$$

In the same manner we define a double covering \tilde{G}_1 of G_1 such that $\sqrt{J_1(\cdot, \cdot)}$ is defined on $\tilde{G}_1 \times \bar{D}_1$ and equal to 1 on $[\tilde{K}_1, \tilde{K}_1] \times \{0\}$. More general, as the Ol'shanskii semigroup $\Gamma_1(W_k)$ is homeomorphic to $G_1 \times iW_k$ and $\Gamma_1(W_k) \subset \{\gamma \in G_{1\mathbb{C}} \mid \gamma^{-1}\bar{D}_1 \subset \bar{D}_1\}$ ([HÓØ91]), there is a double

covering $\tilde{\Gamma}_1(W_k)$ with $\sqrt{J_1(\cdot, \cdot)}$ defined on $\tilde{\Gamma}_1(-W_k) \times \overline{D}_1$ and holomorphic on $\tilde{\Gamma}_1(-W_k)^o \times D_1$.

Let $\kappa : \tilde{\Gamma}_1(-W_k) \rightarrow \Gamma_1(-W_k)$ be the canonical projection, then the covering semigroup operates on \overline{D}_1 by projecting first with κ . Moreover the cocycle relation

$$\sqrt{J_1(\tilde{\gamma}\tilde{\delta}, z)} = \sqrt{J_1(\tilde{\gamma}, \tilde{\delta} \cdot z)}\sqrt{J_1(\tilde{\delta}, z)},$$

with $\tilde{\gamma}, \tilde{\delta} \in \tilde{\Gamma}_1(-W_k)$ and $z \in \overline{D}_1$ is still valid. Let $\tilde{\Gamma}(\pm W) := \kappa^{-1}(\Gamma(\pm W)) \subset \tilde{\Gamma}_1(\pm W_k)$ be the set lying above $\Gamma(\pm W)$ then by Theorem 4.3 we get the $\Gamma(-W)$ - respectively $\tilde{\Gamma}(-W)$ -invariance of $\mathcal{D} \subset \overline{D}_1$. We have an operation of $\tilde{\Gamma}(-W)$ respectively \tilde{G} on $\tilde{\mathcal{D}}$ by setting

$$\tilde{\gamma} \cdot (z, x) := (\tilde{\gamma} \cdot z, \sqrt{J_1(\tilde{\gamma}, z)}x).$$

Therefore we can define as usual the left action $\tilde{\lambda}$ by

$$\tilde{\lambda}(\tilde{\gamma})f(z, x) := f(\tilde{\gamma}^{-1} \cdot (z, x))$$

for functions defined on $\tilde{\mathcal{D}}$.

For $X \subset \mathcal{D}$ let $\tilde{X} \subset \tilde{\mathcal{D}}$ be the subset lying above X .

Definition. Let $X \subset \mathcal{D}$ and $f : \tilde{X} \rightarrow \mathbb{C}$ be a function.

- (1) The function f is odd if $f(z, \xi x) = \xi f(z, x)$ for all m -th roots of unity ξ and $(z, x) \in \tilde{X}$.
- (2) The function f is even if $f(z, \xi x) = f(z, x)$ for all m -th roots of unity ξ and $(z, x) \in \tilde{X}$.

Every even function corresponds to a unique function on X and, by abuse of notation, we will denote both by the the same letter. We note also that we have a \tilde{G} -invariant measure on $\widetilde{G/H}$, defined as the pullback of the G -invariant measure on G/H (and normalized by m^{-1}). Therefore the L^2 -space of odd functions $L^2_{\text{odd}}(\widetilde{G/H})$ is a well-defined object.

With these definitions the proof of the last theorem can be generalized now and we get:

Theorem 5.3. (1) *We have a unitary representation of \tilde{G}_1 in $L^2(\check{S}_1)$ given by*

$$(\tilde{\lambda}_1(\tilde{g})f)(z) := \sqrt{J_1(\tilde{g}^{-1}, z)}f(\tilde{g}^{-1} \cdot z).$$

- (2) *If π_m is G -spherical then $L^2(\check{S}_1, \tilde{\lambda}_1|_{\tilde{\mathcal{G}}}) \rightarrow L^2_{\text{odd}}(\widetilde{G/H}, \tilde{\lambda})$, $f \mapsto f \sqrt[m]{\psi_m}$, is a surjective \tilde{G} -equivariant isometry.*

6. Characterization of $\Phi(\Xi(C^0))$.

Remember that $D_1 = \zeta_1(G_1 K_{1\mathbb{C}} P_1^-)$ is the Hermitian symmetric space in the Harish-Chandra realization. Since $\mathfrak{b} \subset \mathfrak{q}_{1\sigma} \cap \mathfrak{p}_1$ is maximal abelian, we have $G_1 = G \exp(\mathfrak{b}) K_1$. This, together with Proposition 2.4(1), gives now

$$D_1 = G \cdot \left\{ \sum x_{j'} E'_{j'} \mid |x_{j'}| < 1 \right\}$$

by a $\mathfrak{sl}(2)$ -calculation, cf. [H78, p. 387].

Proposition 6.1. (1) $\Phi(\Xi(C^o)) = \{z \in D_1 \mid \psi_m(z) \neq 0\}$.

(2) *There is a constant $c \neq 0$ such that*

$$\psi_m \left(\sum x_{j'} E'_{j'} \right) = c \prod_{j'} x_{j'}^{m\rho_{1,n}(H'_{j'}/2)} = c \prod_{j'} x_{j'}^{\frac{m}{2} \left(1 + \frac{d(r-1)}{2}\right) \sum_j \lambda_{j',j}}.$$

Proof. First we note that $\Gamma(-W^o) Z_{G_{1\mathbb{C}}}(X^0) = G \exp(-iC^o) Z_{G_{1\mathbb{C}}}(X^0)$, which can be proven as in [HÓØ91, Prop. 1.4]. Using this, we get

$$\Phi(\Xi(C^o)) = \zeta_1(\Gamma(-W)^o \cdot c) = \zeta_1(G \exp(-i(\mathfrak{a} \cap C^o)) \cdot c),$$

since $H \subset Z_{G_{1\mathbb{C}}}(X^0) \subset \text{Stab}_{G_{1\mathbb{C}}}(c) = P'_\mathbb{C}$ and using Proposition 2.4(5). By a $\mathfrak{sl}(2)$ -calculation

$$(**) \quad \exp(\lambda H_j) \cdot E_k = \begin{cases} E_k, & \text{if } j \neq k \\ e^{2\lambda} E_k & \text{if } j = k \end{cases},$$

and, as $\zeta_1(c) = -\sum E_j = -\sum E'_{j'}$, we finally get from Proposition 2.4(1) and (6) that

$$\Phi(\Xi(C^o)) = G \cdot \left\{ \sum x_{j'} E'_{j'} \mid 0 < |x_{j'}| < 1 \right\} \subset D_1.$$

Conversely, to determine the set $\{z \in D_1 \mid \psi_m(z) \neq 0\}$ by Theorem 4.3 it is enough to consider $\{\psi_m(\sum x_{j'} E'_{j'}) \mid |x_{j'}| < 1\}$. Using Proposition 2.4(1) we can calculate $\psi_m(\sum x_{j'} E'_{j'})$ via $\mathfrak{sl}(2)$ -reductions. Thus we assume for the moment $G_{1\mathbb{C}} = SL(2, \mathbb{C})$,

$$E'_{j'} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad E'_{-j'} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \text{and } H'_{j'} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

for j' fixed. Then $G_{1\mathbb{C}}^\sigma \exp(\mathbb{C} X'_{j'}) Q_{1\eta\mathbb{C}}^- \subset G_{1\mathbb{C}}$ is open and dense, defining a $r_{j'} \in \mathbb{C}$ by $\exp(x_{j'} E'_{j'}) c^{-1} \in G_{1\mathbb{C}}^\sigma \exp(r_{j'} X'_{j'}) Q_{1\eta\mathbb{C}}^-$ for almost all $x_{j'}$. To determine $r_{j'}$, choose $(X|Y) := \text{tr}(XY^*)$ as an invariant scalar product on $\mathfrak{g}_{1\mathbb{C}}$ and $v \in \mathfrak{q}_{1\eta\mathbb{C}}^+$ nonzero. Let $g \in G_{1\mathbb{C}}^\sigma$ and $q \in Q_{1\eta\mathbb{C}}^-$ be chosen such that $\exp(x_{j'} E'_{j'}) c^{-1} = g \exp(r_{j'} X'_{j'}) q$. Then we have

$$(\exp(x_{j'} E'_{j'}) c^{-1})^{-1} \sigma(\exp(x_{j'} E'_{j'}) c^{-1}) = q^{-1} \exp(-2x_{j'} X'_{j'}) \sigma(q)$$

and consequently, using $\sigma(Q_{1\eta\mathbb{C}}^-) = \theta(Q_{1\eta\mathbb{C}}^-) = Q_{1\eta\mathbb{C}}^+$, we get

$$(\text{Ad}((\exp(x_{j'}E'_{j'})c^{-1})^{-1}\sigma(\exp(x_{j'}E'_{j'})c^{-1}))v|v) = e^{-4r_{j'}}(v|v).$$

Inserting the appropriate matrices it follows easily $r_{j'} = -\frac{1}{2} \log 2x_{j'}$ and further

$$\psi_m \left(\sum x_{j'} E'_{j'} \right) = \left(\prod_{j'} e^{-m\rho_+(r_{j'} X'_{j'})} \right) (v_m | \xi_m) = c \prod_{j'} x_{j'}^{\frac{m}{2} (1 + \frac{d(r-1)}{2}) \sum_j \lambda_{j',j}}.$$

□

Remark. Using $C = \text{Ad}(H)(\mathfrak{a} \cap C)$ together with $\mathfrak{a} \cap C = -\sum \mathbb{R}_0^+ iH'_{j'}$, the arguments of the proof show that $\Xi(C) \subset \mathcal{D}$.

7. Some Hardy space isometries.

The classical Hardy space for G_1 respectively the covering group \tilde{G}_1 in case $\mathfrak{g}_1 = \mathfrak{sp}(2n, \mathbb{R})$, $\mathfrak{so}(2, 2k + 1)$, is defined as the Hilbert space

$$H_{cl} := \left\{ f : D_1 \rightarrow \mathbb{C} \mid f \text{ is holomorphic and } \sup_{\{0 < r < 1\}} \int_{K_1} |f(rk \cdot c)|^2 dk < \infty \right\}$$

with G_1 -action

$$(\lambda(g)f)(z) := \sqrt{J_1(g^{-1}, z)} f(g^{-1} \cdot z)$$

respectively \tilde{G}_1 -action

$$(\tilde{\lambda}(\tilde{g})f)(z) := \sqrt{J_1(\tilde{g}^{-1}, z)} f(\tilde{g}^{-1} \cdot z)$$

([FK94, XIII,3]). By Theorem 5.2(1) respectively Theorem 5.3(1), the operation of taking boundary values is a G_1 - respectively \tilde{G}_1 -equivariant isometry into $L^2(\check{S}_1)$.

Theorem 7.1. *We identify $\Xi(C^o)$ with $\Phi(\Xi(C^o))$. If $\mathfrak{g}_1 \neq \mathfrak{sp}(2n, \mathbb{R})$, $\mathfrak{so}(2, 2k + 1)$ and π_1 is G -spherical then $I : H_{cl} \rightarrow H_2(C)$ $f \mapsto f\psi_1|_{\Xi(C^o)}$ is a G -equivariant isometry.*

Proof. In $H_2(C)$ the action is given by left translation, i.e., for $f \in H_{cl}$ we get

$$\begin{aligned} (\lambda(g)I(f))(z) &= f(g^{-1} \cdot z) \psi_1(g^{-1} \cdot z) \\ &= \sqrt{J_1(g^{-1}, z)} f(g^{-1} \cdot z) \psi_1(z) = I(\lambda(g)f)(z), \end{aligned}$$

the second equality by Theorem 4.3. The map is also an isometry by Theorem 5.2(2), as taking boundary values is an isometry for the Hardy spaces. □

For the general case we define first $\tilde{H}_2(C)$ as covering space of $H_2(C)$ with a $\tilde{\Gamma}(W)$ -action by

$$\tilde{H}_2(C) := \{f : \tilde{\Xi}(C^o) \rightarrow \mathbb{C} \mid f \text{ is holomorphic and } \|f\|_{\tilde{H}} < \infty\},$$

where $\|f\|_{\tilde{H}} := \sup_{\tilde{\gamma} \in \tilde{\Gamma}^o} \|\tilde{\gamma} \cdot f\|_{L^2(\widetilde{G/H})}$. For this space the analogous results as stated in Section 3 hold. One proves now in the same manner as the theorem above:

Theorem 7.2. *If π_m is G -spherical then*

$$\tilde{I} : H_{cl} \rightarrow \tilde{H}_{2,\text{odd}}(C) \quad f \mapsto f \sqrt[m]{\psi_m} |_{\tilde{\Xi}(C^o)}$$

is a \tilde{G} -equivariant isometry.

Let us look closer at the different cases $m = 1, 2, 4$.

m=1: In this case both ψ_1 and $\sqrt{J_1}$ exists, so all coverings split in a trivial way. In particular $L^2_{\text{odd}}(\widetilde{G/H}) \simeq L^2_{\text{even}}(\widetilde{G/H}) \simeq L^2(G/H)$ and similarly also $\tilde{H}_{2,\text{odd}}(C) \simeq \tilde{H}_{2,\text{even}}(C) \simeq H_2(C)$. By the table before Proposition 4.2 the following $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1)$ -triples correspond to this case:

$(\mathfrak{su}(p, q) \oplus \mathfrak{u}(p, q)), \mathfrak{su}(p, q), \mathfrak{su}(n, n))$	n even
$(\mathfrak{u}(p, q), \mathfrak{so}(p, q), \mathfrak{sp}(n, \mathbb{R}))$	$n \equiv 3 \pmod{4}$
$(\mathfrak{so}(2, n-1) \oplus \mathfrak{so}(2), \mathfrak{so}(1, n-1), \mathfrak{so}(2, n+1))$	$n \equiv 3 \pmod{4}$
$(\mathfrak{so}^*(2n), \mathfrak{so}(n, \mathbb{C}), \mathfrak{su}(n, n))$	n even
$(\mathfrak{sp}(2n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{C}), \mathfrak{su}(2n, 2n))$	
$(\mathfrak{so}(2, q) \oplus \mathfrak{so}(p+1), \mathfrak{so}(1, q) \oplus \mathfrak{o}(p), \mathfrak{so}(2, n+1))$	$n \equiv 3 \pmod{4}$
$(\mathfrak{g}, \mathfrak{h}, \mathfrak{g} \times \mathfrak{g})$ for $(\mathfrak{g}, \mathfrak{h})$ of Cayley type	$\mathfrak{g} \neq \mathfrak{so}(2, 2k+1),$ $\mathfrak{sp}(2n, \mathbb{R})$
$(\mathfrak{so}(2, n), \mathfrak{so}(1, n), \mathfrak{so}(2, n+1))$	n odd

m=2: Let us first assume $\mathfrak{g}_1 \neq \mathfrak{sp}(2n, \mathbb{R}), \mathfrak{so}(2, 2k+1)$ such that $\sqrt{J_1}$ exists as a holomorphic function on $\Gamma_1(-W_k^o) \times D_1$. Those are the $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1)$ -triples:

$(\mathfrak{su}(p, q) \oplus \mathfrak{u}(p, q)), \mathfrak{su}(p, q), \mathfrak{su}(n, n))$	n odd
$(\mathfrak{u}(2p, 2q), \mathfrak{sp}(p, q), \mathfrak{so}^*(4n))$	
$(\mathfrak{u}(p, q), \mathfrak{so}(p, q), \mathfrak{sp}(n, \mathbb{R}))$	$n \equiv 1 \pmod{4}$
$(\mathfrak{so}(2, n-1) \oplus \mathfrak{so}(2), \mathfrak{so}(1, n-1), \mathfrak{so}(2, n+1))$	$n \equiv 1 \pmod{4}$
$(\mathfrak{e}_{6(-14)} \oplus \mathfrak{so}(2), \mathfrak{f}_{4(-20)}, \mathfrak{e}_{7(-25)})$	
$(\mathfrak{so}^*(2n) \oplus \mathfrak{so}^*(2n), \mathfrak{so}^*(2n), \mathfrak{so}^*(4n))$	
$(\mathfrak{so}^*(2n), \mathfrak{so}(n, \mathbb{C}), \mathfrak{su}(n, n))$	n odd
$(\mathfrak{so}(2, q) \oplus \mathfrak{so}(p+1), \mathfrak{so}(1, q) \oplus \mathfrak{o}(p), \mathfrak{so}(2, n+1))$	$n \equiv 1 \pmod{4}$

Let $E := \sum E_j = -\zeta_1(c)$. By the equation

$$\psi_2(g \cdot z) = J_1(g, z) \psi_2(z)$$

and the fact that $\Gamma(-W)$ acts transitively on $\Xi(C)$ we can define a global square root of ψ_2 on $\Xi(C)$ by

$$\sqrt{\psi_2}(z) := \sqrt{J_1(\gamma^{-1}, -E)},$$

with γ chosen such that $z = \gamma^{-1} \cdot (-E)$. Clearly this square root is holomorphic on $\Xi(C^o)$.

Lemma 7.3. *For*

$$\Xi(C)_+ := \left\{ \left(\gamma \cdot (-E), \sqrt{J_1(\gamma, -E)} \right) \mid \gamma \in \Gamma(-W) \right\}$$

and

$$\Xi(C^o)_- := \left\{ \left(\gamma \cdot (-E), -\sqrt{J_1(\gamma, -E)} \right) \mid \gamma \in \Gamma(-W) \right\}$$

we have:

- (1) $\Xi(C)_+ = \Gamma(-W) \cdot (-E, 1)$ and $\Xi(C)_- = \Gamma(-W) \cdot (-E, -1)$.
- (2) $\tilde{\Xi}(C) = \Xi(C)_+ \dot{\cup} \Xi(C)_-$ and the canonical projection $\kappa_2 : \Xi(C)_+ \rightarrow \Xi(C)$ is a diffeomorphism, biholomorphic on $\Xi(C)_+^o$.

Proof. It is clear that $\Xi(C)_+, \Xi(C)_- \subset \tilde{\Xi}(C)$ and that $\Xi(C)_+ \cap \Xi(C)_- = \emptyset$. Let $\xi = (z, x) \in \tilde{\Xi}(C)$ and $\gamma \in \Gamma(-W)$ such that $z = \gamma \cdot (-E)$ then

$$x^2 = \psi_2(z) = \psi_2(\gamma \cdot (-E)) = J_1(\gamma, -E) \psi_2(-E) = J_1(\gamma, -E).$$

Thus $\xi = (z, \sqrt{J_1(\gamma, -E)}) \in \Xi(C)_+$ or $\xi = (z, -\sqrt{J_1(\gamma, -E)}) \in \Xi(C)_-$ and in particular $\Xi(C) = \Xi(C)_+ \cup \Xi(C)_-$.

To define the inverse map to κ_2 , choose for $z \in \Xi(C)$ a $\gamma \in \Gamma(-W)$ such that $z = \gamma \cdot (-E)$. Then κ_2^{-1} is given by

$$z \mapsto (z, \sqrt{J_1(\gamma, -E)}),$$

which is a holomorphic map on $\Xi(C^o) = \Gamma(-W^o) \cdot (-E)$, since locally we may choose γ using a holomorphic section. □

The following is now clear:

Proposition 7.4. (1) *The restriction from $\tilde{\Xi}(C^o)$ to $\Xi(C)_+^o$ together with the canonical projection $\kappa_2|_{\Xi(C)_+^o} : \Xi(C)_+^o \rightarrow \Xi(C^o)$ induces an isomorphism of $\Gamma(W)$ -modules*

$$\tilde{H}_{2,\text{odd}}(C) \simeq \tilde{H}_{2,\text{even}}(C) \simeq H_2(C).$$

- (2) *The restriction from $\widetilde{G/H}$ to $G/H_+ := \widetilde{G/H} \cap \Xi(C)_+$ together with the canonical projection $\kappa_2|_{G/H_+} : G/H_+ \rightarrow G/H$ induces an isomorphism of G -modules*

$$L_{\text{odd}}^2(\widetilde{G/H}) \simeq L_{\text{even}}^2(\widetilde{G/H}) \simeq L^2(G/H).$$

- (3) *Taking boundary values in the L^2 -norm intertwines the G -actions on the respective modules.*

If $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1) = (\mathfrak{so}(2, 2k), \mathfrak{so}(1, 2k), \mathfrak{so}(2, 2k + 1))$ or $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g} \times \mathfrak{g})$ is one of the remaining Cayley type cases the preceding construction does not work. However, using the fact that $\mathfrak{g}_1 = \mathfrak{g} \times \mathfrak{g}$, for the Cayley type cases it is possible to define a global square root $\sqrt{J_1(\cdot, \cdot)}$ on $\Gamma(-W)$. Therefore analogous results to Lemma 7.3 and Proposition 7.4 hold, cf. [ÓØ99, Lemma 5.5, Lemma 6.5].

m=4: For these cases $\sqrt{J_1(\gamma, z)}$ does not exist as a globally defined map in the variable $\gamma \in \Gamma_1(-W)$, as can be seen from the occurring \mathfrak{g}_1 for the corresponding $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1)$ -triples:

$(\mathfrak{u}(p, q), \mathfrak{so}(p, q), \mathfrak{sp}(n, \mathbb{R}))$	n even
$(\mathfrak{so}(2, n - 1) \oplus \mathfrak{so}(2), \mathfrak{so}(1, n - 1), \mathfrak{so}(2, n + 1))$	n even
$(\mathfrak{sp}(n, \mathbb{R}) \oplus \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(2n, \mathbb{R}))$	
$(\mathfrak{so}(2, q) \oplus \mathfrak{so}(p + 1), \mathfrak{so}(1, q) \oplus \mathfrak{o}(p), \mathfrak{so}(2, n + 1))$	n even

From the equation

$$\psi_4(\gamma \cdot (-E)) = J_1^2(\gamma, -E) \psi_4(-E) = (J_1(\gamma, -E))^2,$$

valid for $\gamma \in \Gamma(-W)$, it follows that we can define a holomorphic square root of ψ_4 on $\Xi(C)$. But to define the needed function $\sqrt[4]{\psi_4}$ we have to go to a double covering. By the above, for $(\tilde{\gamma} \cdot (-E), x) \in \tilde{\Xi}(C)$, i.e., $x^4 = \psi_4(\tilde{\gamma} \cdot (-E))$, we have $x^2 = \pm J_1(\tilde{\gamma}, -E)$. Therefore we define in this case

$$\tilde{\Xi}(C)_+ := \{(\tilde{\gamma} \cdot (-E), x) \mid \tilde{\gamma} \in \Gamma(-W), x^2 = J_1(\tilde{\gamma}, -E)\} \subset \tilde{\Xi}(C)$$

and

$$\tilde{\Xi}(C)_- := \{(\tilde{\gamma} \cdot (-E), x) \mid \tilde{\gamma} \in \Gamma(-W), x^2 = -J_1(\tilde{\gamma}, -E)\} \subset \tilde{\Xi}(C).$$

Then $\tilde{\Gamma}(-W)$ acts on $\tilde{\Xi}(C)_\pm$ as subsets of $\tilde{\Xi}(C) \subset \tilde{\mathcal{D}}$ by

$$\tilde{\gamma} \cdot (z, x) = (\tilde{\gamma} \cdot z, \sqrt{J_1(\tilde{\gamma}, z)} x).$$

Note that for $z \in \Xi(C)$ there are two elements $\tilde{\gamma}, \tilde{\delta} \in \tilde{\Gamma}(-W)$ with $z = \tilde{\gamma} \cdot (-E) = \tilde{\delta} \cdot (-E)$ and $\sqrt{J_1(\tilde{\gamma}, -E)} = -\sqrt{J_1(\tilde{\delta}, -E)}$. Using this, the proof of Lemma 7.3 now generalizes word-by-word to give:

Lemma 7.5.

- (1) $\tilde{\Xi}(C)_+ = \tilde{\Gamma}(-W) \cdot (-E, 1)$ and $\tilde{\Xi}(C)_- = \tilde{\Gamma}(-W) \cdot (-E, -1)$.
- (2) We have

$$\tilde{\Xi}(C) = \tilde{\Xi}(C)_+ \dot{\cup} \tilde{\Xi}(C)_-$$

and the canonical projection $\kappa_4 : \tilde{\Xi}(C)_+ \rightarrow \Xi(C)$ is a double covering, holomorphic on $\tilde{\Xi}(C)_+$.

We have on $\widetilde{G/H}_+ := \kappa_4^{-1}(G/H)$ the space of L^2 -functions $L^2(\widetilde{G/H}_+)$ and on $\widetilde{\Xi}(C)_+^o = \kappa_4^{-1}(\Xi(C^o))$ the Hardy space $\widetilde{H}_2(C)_+$, as a space of holomorphic functions defined in the same way as $\widetilde{H}_2(C)$. Using roots of unity we can define the subspaces $L^2_{\text{odd}}(\widetilde{G/H}_+)$ respectively $\widetilde{H}_{2,\text{odd}}(C)_+$ of odd functions.

Proposition 7.6. (1) *The restriction from $\widetilde{\Xi}(C^o)$ to $\widetilde{\Xi}(C)_+^o$ induces an isomorphism of $\widetilde{\Gamma}(W)$ -modules*

$$\widetilde{H}_{2,\text{odd}}(C) \simeq \widetilde{H}_{2,\text{odd}}(C)_+.$$

(2) *The restriction from $\widetilde{G/H}$ to $\widetilde{G/H}_+$ induces an isomorphism of G -modules*

$$L^2_{\text{odd}}(\widetilde{G/H}) \simeq L^2_{\text{odd}}(\widetilde{G/H}_+).$$

The proof is obvious. We can in particular draw the conclusion from those considerations that we will need maximally a double covering of $\Xi(C)$ for identifying the Hardy space.

For the characterization of the image we give first the extension of Φ_δ to a function on $\Phi(\Xi(C^o)) \subset D_1 = G \cdot \{\sum x_{j'} E'_{j'}\}$.

Lemma 7.7. *For $g \in G$, define $g = k(g) \exp p(g)$ by the Cartan decomposition. Let $\{\nu_{\mu',r}\}$ be a base of weight vectors of V_δ , with $\nu_{\mu',r}$ of weight μ' , and define $a_{\mu',r}$ by $\delta(k)\nu = \sum a_{\mu',r}(k^{-1})\nu_{\mu',r}$. Identifying $\Xi(C^o)$ with $\Phi(\Xi(C^o))$, there is a constant $c \neq 0$ such that*

$$\begin{aligned} &\Phi_\delta \left(g \cdot \sum x_{j'} E'_{j'} \right) \\ &= c \sum_{\mu',r} a_{\mu',r}(k(g)) \prod_{j'} x_{j'}^{-\mu'(H'_{j'}/2)} \left\langle \delta \left(f \left(g, \sum x_{j'} E'_{j'} \right) \nu_{\mu',r}, \nu^0 \right) \right\rangle, \end{aligned}$$

with

$$f \left(g, \sum x_{j'} E'_{j'} \right) = k_H \left(\text{Ad} \left(\exp \left(-\frac{1}{2} \sum (\log x_{j'} + \pi i) H'_{j'} \right) p(g)^{-1} \right) \right)$$

and \log the principal branch continued continuously to $-\mathbb{R}^+$ from the upper half plane.

Proof. By Equation (**) and Proposition 2.4(1) we have

$$g \cdot \sum x_{j'} E'_{j'} = \zeta_1 \left(g \exp \left(\frac{1}{2} \sum (\log x_{j'} + \pi i) H'_{j'} \right) c \right).$$

Therefore

$$\begin{aligned} & \Phi_\delta \left(g \cdot \sum x_{j'} E'_{j'} \right) \\ &= \left\langle \nu, \check{\delta} \left(k_H \left(\exp \left(-\frac{1}{2} \sum (\log x_{j'} + \pi i) H'_{j'} \right) g^{-1} \right)^{-1} \right) \nu^0 \right\rangle \\ &= \left\langle \delta(k(g)^{-1}) \nu, \check{\delta} \left(\exp \left(\frac{1}{2} \sum (\log x_{j'} + \pi i) H'_{j'} \right) f \left(g, \sum x_{j'} E'_{j'} \right)^{-1} \right) \nu^0 \right\rangle \\ &= c \sum_{\mu', r} a_{\mu', r}(k(g)) \prod_{j'} x_{j'}^{-\mu'(H'_{j'}/2)} \left\langle \delta \left(f \left(g, \sum x_{j'} E'_{j'} \right) \right) \nu_{\mu', r}, \nu^0 \right\rangle. \end{aligned}$$

□

For $f \in H_2(C)$, the inverse map to I is given by extending the holomorphic function f/ψ_1 to D_1 . This extension is possible if and only if f/ψ_1 is locally bounded near $D_1 \setminus \Xi(C^o)$. By Theorem 3.5(5) it is enough to check this for the generating functions $f = \Phi_\delta$. Now by Theorem 4.3, Proposition 6.1, and the last lemma

$$(\Phi_\delta/\Psi_1) \left(g \cdot \sum x_{j'} E'_{j'} \right) = \sum_{\mu'} b_{\mu'} \left(g, \sum x_{j'} E'_{j'} \right) \prod_{j'} x_{j'}^{-(\mu' + \rho_{1,n})(H'_{j'}/2)},$$

where $b_{\mu'}$ is locally bounded on $G \times D_1$.

Theorem 7.8. $H_\delta \subset I(H_{cl}) \iff (\mu + \rho_{1,n})(H'_{j'}) \leq 0$ for all j' .

Proof. Equating $\Phi_\delta(\sum x_{j'} E'_{j'})$ by Lemma 7.7, we get the necessary condition $(\mu' + \rho_{1,n})(H'_{j'}) \leq 0$ for all μ' with $a_{\mu', r}(1) \neq 0$ and $\langle \nu_{\mu', r}, \nu^0 \rangle \neq 0$. By [H84, p. 535ff] for the highest weight μ of δ both $\nu_{\mu, 1} \neq 0$ and $\langle \nu_{\mu, 1}, \nu^0 \rangle \neq 0$, showing that the given condition is necessary.

For sufficiency, let μ' be a weight of (δ, V_δ) then $\mu' \in \text{conv}(W(\mathfrak{k}, \mathfrak{t}) \cdot \mu) \subset i\mathfrak{t}^*$. Applying [B88, Lemma 1.1] to the Lie algebra $\mathfrak{k}_{1\mathbb{C}}$ with involution $-\sigma$, we can assume $W(\mathfrak{k}, \mathfrak{t}) \subset W(\mathfrak{k}_1, \mathfrak{t}_1)$, where $W(\mathfrak{k}_1, \mathfrak{t}_1)$ is the Weyl group of the root system $\Delta(\mathfrak{k}_1, \mathfrak{t}_1)$. As we already have seen $\rho_{1,n} \in (\mathfrak{z}(\mathfrak{k}_1)_{\mathbb{C}})^* \subset (\mathfrak{t}_{1\mathbb{C}})^*$ such that now $W(\mathfrak{t}, \mathfrak{k})\rho_{1,n} = \rho_{1,n}$ and further

$$(w \cdot \mu + \rho_{1,n})(H'_{i'}) = w \cdot (\mu + \rho_{1,n})(H'_{i'}) = (\mu + \rho_{1,n})(w^{-1}H'_{i'}).$$

We note that $i(\mathfrak{a} \cap C^o) = \sum \mathbb{R}^+ H'_{j'} \subset i\mathfrak{a}$ is the Weyl chamber for the roots $\Delta_{1,n}^+ | \mathfrak{a}$. Indeed, since $\gamma_i(H_j) = 2\delta_{i,j}$, we deduce from the Remark following Proposition 2.4 that the co-roots for $\Delta_{1,n}^+ | \mathfrak{a}$ are the vectors $\{ \frac{1}{4}(H'_{i'} + H'_{j'}) \mid 1 \leq i' \leq j' \leq r^{(l)} \}$. This means that $i(\mathfrak{a} \cap C^o)$ is the set of all linear combinations of co-roots with positive coefficients, which — together with $\langle \alpha, \beta \rangle \geq 0$ for all $\alpha, \beta \in \Delta_{1,n}^+ | \mathfrak{a}$ — implies the assertion.

Now, as $W(\mathfrak{k}_1, \mathfrak{t}_1)$ permutes the noncompact roots, we get

$$\left(W(\mathfrak{k}, \mathfrak{t})(\Delta_{1,n}^+ | \mathfrak{a}) \right) | \mathfrak{a} \subset \Delta_{1,n}^+ | \mathfrak{a}.$$

Expressing this with co-roots we get $w^{-1}H'_{i'} \in \sum \mathbb{R}_0^+ H'_{j'}$, such that by our equation above the given condition is sufficient for the boundedness of Φ_δ/Ψ_1 . \square

Of course, the same condition determines the image of \tilde{I} in $\tilde{H}_{2,\text{odd}}(C)$.

This theorem allows us now to determine the image of the classical Hardy space in $H_2(C)$ respectively $\tilde{H}_{2,\text{odd}}(C)$ as described by the root inequalities of Theorem 3.3 and Proposition 3.4.

Example. We consider $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1) = (\mathfrak{sp}(n, \mathbb{R}) \oplus \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(2n, \mathbb{R}))$ as treated in [B97, Section 7.4]. Here

$$\mathfrak{g}_1 = \left\{ \left(\begin{array}{cc} A & B \\ \overline{B} & \overline{A} \end{array} \middle| \begin{array}{l} A, B \in M(2n \times 2n, \mathbb{C}) \\ -{}^t\overline{A} = A, \quad {}^tB = B \end{array} \right) \right\}$$

and

$$\mathfrak{g} = \left\{ \left(\begin{array}{cccc} A_1 & 0 & B_1 & 0 \\ 0 & A_4 & 0 & B_4 \\ \overline{B}_1 & 0 & \overline{A}_1 & 0 \\ 0 & \overline{B}_4 & 0 & \overline{A}_4 \end{array} \right) \in \mathfrak{g}_1 \mid A_i, B_i \in M(n \times n, \mathbb{C}) \right\}.$$

Further, $\mathfrak{t}_{1\mathbb{C}}^-$ is spanned by

$$H_j = \frac{1}{2} \begin{pmatrix} E_{jj} & E_{jj} & 0 & 0 \\ E_{jj} & E_{jj} & 0 & 0 \\ 0 & 0 & -E_{jj} & -E_{jj} \\ 0 & 0 & -E_{jj} & -E_{jj} \end{pmatrix},$$

$$H_{j+n} = \frac{1}{2} \begin{pmatrix} E_{jj} & -E_{jj} & 0 & 0 \\ -E_{jj} & E_{jj} & 0 & 0 \\ 0 & 0 & -E_{jj} & E_{jj} \\ 0 & 0 & E_{jj} & -E_{jj} \end{pmatrix}$$

for $j = 1, \dots, n$, whereas a base for $\mathfrak{a}_{\mathbb{C}}$ is given by the vectors $H'_j = H_j + H_{j+n}$. Since \mathfrak{g}_1 is a normal Lie algebra the multiplicity of all roots in Δ_1 is one. Defining $\gamma'_j := \frac{1}{2}(\gamma_j + \gamma_{j+n})$ we get by restricting the roots of Δ_1^+ to $\mathfrak{a}_{\mathbb{C}}$

$$\Delta^+ = \left\{ \frac{1}{2}(\gamma'_i + \gamma'_j) \mid 1 \leq i \leq j \leq n \right\} \cup \left\{ \frac{1}{2}(\gamma'_i - \gamma'_j) \mid 1 \leq i < j \leq n \right\},$$

and all these roots have multiplicity 2. With $\Delta_n^+ = \{ \frac{1}{2}(\gamma'_i + \gamma'_j) \mid 1 \leq i \leq j \leq n \}$ by Proposition 2.4(6) and the definition of c_{\min} we get $\mathfrak{a} \cap C = c_{\min}$. Therefore, by the Remark following Proposition 3.4 the Hardy space $\tilde{H}_2(C)$ comprises the whole holomorphic discrete series. To describe these in more

detail, with μ the highest weight of δ with respect to $\Delta_k^+ = \{\frac{1}{2}(\gamma'_i - \gamma'_j) \mid 1 \leq i < j \leq n\}$ we have for $H_\delta \subset \tilde{H}_{2,\text{odd}}(C)$

$$\mu = \sum m_j \gamma'_j, \quad m_1 \geq m_2 \geq \dots \geq m_n, \quad m_j \in \mathbb{Z} + \frac{1}{2}.$$

To apply Theorem 3.3(1) we calculate

$$\begin{aligned} \rho &= \frac{1}{2} \sum_{\alpha \in \Delta^+} \dim \mathfrak{g}_{\mathbb{C},\alpha} \alpha = \sum_{1 \leq i \leq j \leq n} \frac{1}{2}(\gamma'_i + \gamma'_j) + \sum_{1 \leq i < j \leq n} \frac{1}{2}(\gamma'_i - \gamma'_j) \\ &= (n+1) \sum_{j=1}^n \gamma'_j - \sum_{j=1}^n j \gamma'_j. \end{aligned}$$

Inserting into $\langle \mu + \rho, \alpha \rangle < 0$, for $\alpha \in \Delta_n^+$, we derive the condition

$$H_\delta \subset \tilde{H}_{2,\text{odd}}(C) \iff -n > m_1 \geq m_2 \geq \dots \geq m_n.$$

We show next that \tilde{I} is actually surjective. In fact, with

$$\rho_{1,n} = \frac{1}{2} \left(\frac{2n+1}{2} \right) (\gamma_1 + \dots + \gamma_{2n})$$

the condition of Theorem 7.8 reduces to

$$\langle \mu, \gamma'_j \rangle \leq -\frac{2n+1}{4} \quad \text{or} \quad m_j \leq -\frac{2n+1}{2}$$

such that, $m_j + \frac{1}{2}$ being an integer, the embedding \tilde{I} is onto. Another discussion of this example can be found in [KØ97].

Similar to the given example this “best possible” case, with $H_2(C)$ respectively $\tilde{H}_{2,\text{odd}}(C)$ being the direct sum of all holomorphic discrete series representations and I respectively \tilde{I} onto, occurs for the $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1)$ -triples

$$\begin{aligned} &(\mathfrak{sp}(n, \mathbb{R}) \oplus \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{R}), \mathfrak{sp}(2n, \mathbb{R})) \\ &(\mathfrak{sp}(2n, \mathbb{R}), \mathfrak{sp}(n, \mathbb{C}), \mathfrak{su}(2n, 2n)) \\ &(\mathfrak{g}, \mathfrak{h}, \mathfrak{g} \times \mathfrak{g}) \text{ for } (\mathfrak{g}, \mathfrak{h}) \text{ of Cayley type} \\ &(\mathfrak{so}(2, n), \mathfrak{so}(1, n), \mathfrak{so}(2, n+1)). \end{aligned}$$

Note that these are exactly the cases with $\mathfrak{a} \neq \mathfrak{t}_1^-$.

Example. We discuss next the compactifications described by [B97, Cor. 5.3]. These are the $(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1)$ -triples

$$\begin{aligned} &(\mathfrak{su}(p, q) \oplus \mathfrak{u}(p, q), \mathfrak{su}(p, q), \mathfrak{su}(n, n)), \quad (\mathfrak{u}(2p, 2q), \mathfrak{sp}(p, q), \mathfrak{so}^*(4n)), \\ &(\mathfrak{u}(p, q), \mathfrak{so}(p, q), \mathfrak{sp}(n, \mathbb{R})), \quad (\mathfrak{so}(2, n-1) \oplus \mathfrak{so}(2), \mathfrak{so}(1, n-1), \mathfrak{so}(2, n+1)), \\ &(\mathfrak{e}_{6(-14)} \oplus \mathfrak{so}(2), \mathfrak{f}_{4(-20)}, \mathfrak{e}_{7(-25)}). \end{aligned}$$

Here $\gamma'_{j'} = \gamma_{j'}$, i.e., $H'_{j'} = H_{j'}$, and $\mathfrak{z}(\mathfrak{g}) = \sum \epsilon_j i H_j$, where without loss of generality $\epsilon_j = 1$ for $j \leq p$ and $\epsilon_j = -1$ for $j > p$. By the explicit form of

$$\sigma = \text{Ad} \left(\exp \frac{\pi}{2} \sum \epsilon_j i H_j \right)$$

the action of this involution on the root spaces $\mathfrak{g}_{1\mathbb{C},\alpha}$ can be determined easily and gives us

$$\Delta_k = \left\{ \pm \frac{1}{2}(\gamma_i - \gamma_j) \mid i < j \leq p \text{ or } p < i < j \right\} \text{ and}$$

$$\Delta_n = \left\{ \pm \frac{1}{2}(\gamma_i + \gamma_j) \mid i \leq p < j \right\}.$$

The weight spaces for $\gamma \in \Delta$ are the same as for $\gamma \in \Delta_1$, especially all have the same dimension d , such that

$$\rho = \frac{d}{4} \left[\sum_{i < j \leq p} (\gamma_i - \gamma_j) + \sum_{p < i < j} (\gamma_i - \gamma_j) + \sum_{i \leq p < j} (\gamma_i + \gamma_j) \right] = \frac{d}{4}(r+1) \sum_{j=1}^r \gamma_j.$$

With $\Delta_k^+ = \{\frac{1}{2}(\gamma_i - \gamma_j)\}$ the highest weight has the form $\mu = \sum n_j \gamma_j$ with $n_1 \geq n_2 \geq \dots \geq n_p$ and $n_{p+1} \geq \dots \geq n_r$. Alternatively, with $\mu = d\chi_\delta + \mu_s$ we can describe the highest weight with $d\chi_\delta = \lambda(\frac{1}{2r} \sum \epsilon_j \gamma_j)$ and $\mu_s = \sum s_j \gamma_j$, where $\lambda \in \mathbb{R}$ and the s_j are integers with $s_1 \geq s_2 \geq \dots \geq s_p$ and $s_{p+1} \geq \dots \geq s_r$. (The s_j are also restricted by $\sum \epsilon_j s_j = 0$, since μ_s is zero on $\mathfrak{z}(\mathfrak{g})$.) As $d\chi_\delta$ integrates to a representation $\chi_\delta \in \hat{Z}$ respectively $\tilde{\chi}_\delta \in \tilde{Z}$ we have $\lambda \in \mathbb{Z}$ respectively $\lambda \in \frac{1}{2}\mathbb{Z}$. Finally, to get a representation $\delta = \chi_\delta \otimes \delta_s$ the representations χ_δ and δ_s must be equal on the central subgroup $D = Z \cap G_s$ respectively \tilde{D} for the covering group. We call this the (D) -condition.

With ρ as above we get from Theorem 3.3(1)

$$n_i + n_j < -\frac{d}{2}(r+1), \quad \text{for } i \leq p < j,$$

which is by the form of μ equivalent to $n_1 + n_{p+1} < -\frac{1}{2}(r+1)$, and from Proposition 3.4 the condition of W -admissibility

$$n_j \leq 0, \quad \text{for } j = 1, \dots, r.$$

To summarize, the Hardy space $H_2(C)$ ($\tilde{H}_2(C)$) is determined by $(n_1, \dots, n_r) \in \mathbb{Z}^r + (\frac{1}{2}\mathbb{Z}(1, \dots, 1))$ with

- (1) the (D) -condition,
- (2) $0 \geq n_1 \geq \dots \geq n_p$ and $0 \geq n_{p+1} \geq \dots \geq n_r$,
- (3) $n_1 + n_{p+1} < -\frac{d}{2}(r+1)$.

For the covering case $\tilde{H}_{2,\text{odd}}(C)$ is the subspace spanned by those holomorphic discrete series representations which are not representations for G .

We remark that this Hardy space is strictly smaller than the Hardy space $H_2(C_{\min})$ respectively $\tilde{H}_{2,\text{odd}}(C_{\min})$. Indeed, by Theorem 3.3, Proposition 3.4, and the following Remark, it is clear that this “maximal” Hardy space is the direct sum $\bigoplus_{\delta_s, \chi \in \hat{Z}(\delta_s)} \mathbb{C}_\chi \otimes H_{\delta_s}$, where the δ_s exhaust the holomorphic discrete series and $\hat{Z}(\delta_s)$ is the set of unitary characters which fulfill with δ_s the (D) -condition, i.e., $\chi \otimes \delta_s$ is a representation of G respectively \tilde{G} . To see that this is a bigger space than $H_2(C)$ we calculate, with Δ_n^+ as above,

$$\mathfrak{a} \cap C_{\min}^o = -i \sum_{a \leq p < b} \mathbb{R}^+(H_a + H_b).$$

Therefore the condition of admissibility for the minimal cone is $n_a + n_b \leq 0$, for $a \leq p < b$, or equivalently $n_1 + n_{p+1} \leq 0$ compared with $n_1 \leq 0$ and $n_{p+1} \leq 0$ for the cone C .

The image of the classical Hardy space is described by

$$n_j \leq -\frac{1}{2} \left(1 + \frac{d(r-1)}{2} \right), \quad \text{for } j = 1, \dots, r.$$

To give a more explicit example, we specialize further to the case $G/H = S(U(1, 1) \times U(1, 1))/SU(1, 1)$, cf. [B97, Section 6.5] for the realization. Here $H_j = (E_{j,j} + E_{j+2,j+2}) \in \mathfrak{su}(2, 2)_{\mathbb{C}} = \mathfrak{sl}(4, \mathbb{C})$ and

$$\mathfrak{a} = \mathfrak{t} \cap \mathfrak{q} = \mathbb{R} \begin{pmatrix} i & & & \\ & -i & & \\ & & -i & \\ & & & i \end{pmatrix} + \mathbb{R} \begin{pmatrix} i & & & \\ & i & & \\ & & -i & \\ & & & -i \end{pmatrix},$$

where the first summand is equal to $\mathfrak{z}(\mathfrak{g})$. Let $\mu \in i(\mathfrak{t} \cap \mathfrak{q})^*$ be the highest weight for a holomorphic discrete series representation of G/H . Identifying $\mathfrak{t} \cap \mathfrak{q}$ with its dual, we write

$$\mu = a \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix} + b \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix},$$

such that the first summand gives the central character $d\chi_\delta$ and the second μ_s . We have the integrability conditions $a, b \in \mathbb{Z}$, and the (D) -condition gives $a + b \in 2\mathbb{Z}$. Further

$$\Delta^+ = \Delta_n^+ = \left\{ \frac{1}{2}(\gamma_1 + \gamma_2) \right\}$$

and $\rho = \frac{1}{2}(\gamma_1 + \gamma_2)$. Thus Theorem 3.3(1) gives us $b < -1$ and of course no restriction on the “central parameter” a .

However we have also the condition of W -admissibility. Rewriting this in terms of the parameters a and b shows that W -admissibility is equivalent to $|a| \leq -b$.

For the determination of the image of the classical Hardy space in $H_2(C)$ it is straightforward from the definition that $\rho_{1,n} = H_1 + H_2$ (with our identification of \mathfrak{t} and \mathfrak{t}^*). Then the determining equations $(\mu + \rho_{1,n})(H_i) \leq 0$ are equivalent to $|a| \leq -b - 1$. The image of the classical Hardy space therefore misses the two half lines $a + b = 0$ and $a - b = 0$. We remark that this differs from [KØ96, Figure 3] where the missing half lines lie on one side of the cone.

One can deal with the remaining cases as above. With notation as in the preceding examples, we summarize the results as follows:

$(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1) = (\mathfrak{so}^*(2n) \oplus \mathfrak{so}^*(2n), \mathfrak{so}^*(2n), \mathfrak{so}^*(4n))$: The Hardy space $H_2(C) = H_2(C_{\min})$ is parametrized by $0 \geq n_1 \geq \dots \geq n_n$ with $n_i + n_j < -2n + i + j$, for $i \neq j$, and $I(H_{cl})$ is given by $n_1 \leq -n$.

$(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1) = (\mathfrak{so}^*(2n), \mathfrak{so}(n, \mathbb{C}), \mathfrak{su}(n, n))$: The Hardy space $H_2(C) = H_2(C_{\min})$ is parametrized by $0 \geq n_1 \geq \dots \geq n_n$ with $n_i + n_j < -n + \frac{i+j}{2}$, for $i \neq j$, and $I(H_{cl})$ is given by $n_1 \leq -\frac{n}{2}$.

$(\mathfrak{g}, \mathfrak{h}, \mathfrak{g}_1) = (\mathfrak{so}(2, q) \oplus \mathfrak{so}(p + 1), \mathfrak{so}(1, q) \oplus \mathfrak{o}(p), \mathfrak{so}(2, n + 1))$: $H_2(C)$ respectively $\tilde{H}_{2,\text{odd}}(C)$ is parametrized by $0 \geq n_1 \geq n_2$ with $n_1 + n_2 \leq -\frac{1}{4}(n + q - p)$ and $I(H_{cl})$ or $\tilde{I}(H_{cl})$ is given by $n_1 \leq -\frac{1}{4}(n + 1)$. Note that for this case the decomposition of $(\mathfrak{g}, \mathfrak{h}, \tau)$ into irreducible symmetric algebras gives not a decomposition of the causal structure.

We remark at the end that our results give now the Cauchy-Szegö kernel $K_I(\cdot, \cdot)$ for the subspace $I(H_{cl}) \subset H_2(C)$ respectively $\tilde{I}(H_{cl}) \subset \tilde{H}_{2,\text{odd}}(C)$. Indeed, with $\{\varphi_n\}$ an orthonormal base of $H_2(C)$ the kernel is generally given by

$$K(z, w) = \sum \varphi_n(z) \overline{\varphi_n(w)},$$

such that

$$K_I(z, w) = \sqrt[m]{\psi_m(z)} K_{cl}(z, w) \overline{\sqrt[m]{\psi_m(w)}},$$

where the kernel K_{cl} for the classical Hardy space is known ([FK94, Prop. X.1.3]). Note also that this kernel is invariant by deck transformations of the covering $\tilde{\Xi}(C) \rightarrow \Xi(C)$. Therefore, when we have an isomorphism of Hardy spaces as in Proposition 7.4, we get the corresponding image of the kernel by simply projecting down.

References

[B88] E.P. van den Ban, *The principal series for a reductive symmetric space I. H-fixed distribution vectors*, Ann. Scient. Éc. Norm. Sup., **21** (1988), 359-412, MR 90a:22016, Zbl 714.22009.

- [Bal] W. Bertram and J. Hilgert, *Hardy spaces and analytic continuation of Bergman spaces*, Bull. Soc. Math. France, **126** (1998), 435-482, MR 2000a:32012, Zbl 920.22006.
- [B97] F. Betten, *Causal compactification of compactly causal spaces*, Trans. A.M.S., to appear.
- [BD92] J.-L. Brylinski and P. Delorme, *Vecteurs distributions H -invariants pour les séries principales généralisées d'espaces symétriques réductifs et prolongement méromorphe d'intégrales d'Eisenstein*, Invent. Math., **109** (1992), 619-664, MR 93m:22016, Zbl 785.22014.
- [C96] M. Chadli, *Espace de Hardy d'un Espace Symétrique de Type Cayley*, Thesis, Paris, 1996, MR 97f:43013, Zbl 860.46036.
- [FK94] J. Faraut and A. Korányi, *Analysis on Symmetric Cones*, Clarendon Press, Oxford, 1994, MR 98g:17031, Zbl 841.43002.
- [H78] S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Academic Press, New York, 1978, MR 80k:53081, Zbl 451.53038.
- [H84] ———, *Groups and Geometric Analysis. Integral Geometry, Invariant Differential Operators and Spherical Functions*, Academic Press, New York, 1984, MR 86c:22017, Zbl 543.58001.
- [H94] ———, *Geometric Analysis on Symmetric Spaces*, A.M.S., Providence, Rhode Island, 1994, MR 96h:43009, Zbl 809.53057.
- [HÓ92] J. Hilgert and G. Ólafsson, *Analytic extension of representations, the solvable case*, Japan J. Math., **18** (1992), 213-290, MR 94f:22011.
- [HÓ96] ———, *Causal Symmetric Spaces, The Geometry and Harmonic Analysis*, Perspectives in Math., **18**, Academic Press, New York, 1996, MR 97m:43006.
- [HÓØ91] J. Hilgert, G. Ólafsson and B. Ørsted, *Hardy spaces on affine symmetric spaces*, J. Reine Angew. Math., **415** (1991), 189-218, MR 92h:22030, Zbl 716.43006.
- [KW65] A. Korányi and J.A. Wolf, *Realization of Hermitian symmetric spaces as generalized half planes*, Ann. of Math., **81** (1965), 265-288, MR 30 #4980, Zbl 137.27402.
- [KØ96] K. Koufany and B. Ørsted, *Function spaces on the Ol'shanskiĭ semigroup and the Gelf'and-Gindikin program*, Anal. Inst. Fourier, **46** (1996), 689-722, MR 97k:22021, Zbl 853.22010.
- [KØ97] ———, *Hardy spaces on two sheeted covering semigroups*, J. Lie Theory, **7** (1997), 245-267, MR 98k:22060, Zbl 884.22006.
- [KN96] B. Krötz and K.-H. Neeb, *On hyperbolic cones in mixed symmetric spaces*, J. of Lie Theory, **6** (1996), 69-146, MR 97k:17007, Zbl 860.22004.
- [KNÓ97] B. Krötz, K.-H. Neeb and G. Ólafsson, *Spherical representations and mixed symmetric spaces*, Representation Theory, **1** (1997), 424-461, MR 99a:22031, Zbl 887.22022.
- [L69] O. Loos, *Symmetric Spaces, I: General Theory*, W.A. Benjamin Inc., New York, 1969, MR 39 #365b, Zbl 175.48601.
- [M79] T. Matsuki, *The orbits of affine symmetric spaces under the action of minimal parabolic subgroups*, J. Math. Soc. Japan, **31** (1979), 331-357, MR 81a:53049, Zbl 396.53025.

- [M97] V.F. Molchanov, *Holomorphic discrete series for hyperboloids of Hermitian type*, J. Funct. Anal., **147** (1997), 26-50, MR 99b:22026, Zbl 881.43005.
- [Ó84] G. Ólafsson, *Die Langlands-Parameter für die Flensted-Jensensche fundamentale Reihe*, Math. Scand., **55** (1984), 229-244, MR 86k:22034, Zbl 561.22008.
- [Ó87] ———, *Fourier and Poisson transformation associated to a semisimple symmetric space*, Invent. Math., **90** (1987), 605-629, MR 89d:43011, Zbl 665.43004.
- [Ó91] ———, *Symmetric spaces of Hermitian type*, Diff. Geom. and Appl., **1** (1991), 195-233, MR 94g:22034, Zbl 785.22021.
- [Ó97a] ———, *Spherical functions and spherical Laplace transform on ordered symmetric spaces*, submitted, preprint: <http://www.math.lsu.edu/~olafsson>.
- [Ó97b] ———, *Open problems in harmonic analysis on causal symmetric spaces*, in 'Positivity in Lie Theory: Open Problems', eds. J. Hilgert, J.D. Lawson, K.-H. Neeb and E.B. Vinberg, Expositions in Mathematics, De Gruyter, Berlin, 1998, 249-270, MR 99h:43021, Zbl 919.43006.
- [Ó00] ———, *Analytic continuation in representation theory and harmonic analysis*, in 'Seminaires et Congres', **4**, Global Analysis and Harmonic Analysis, eds. J.P. Bourguignon, T. Branson and O. Hijazi, French Mathematical Society, 2000, 201-233.
- [ÓØ88] G. Ólafsson and B. Ørsted, *The holomorphic discrete series for affine symmetric spaces*, I, J. Funct. Anal., **81** (1988), 126-159, MR 89m:22021, Zbl 678.22008.
- [ÓØ91] ———, *The holomorphic discrete series of an affine symmetric space and representations with reproducing kernels*, Trans. A.M.S., **326** (1991), 385-405, MR 91j:22014, Zbl 760.22020.
- [ÓØ99] ———, *Causal compactification and Hardy spaces*, Trans. A.M.S., **351** (1999), 3771-3792, MR 99m:22005, Zbl 928.43007.
- [O95] G.I. Ol'shanskiĭ, *Cauchy-Szegő kernels for Hardy spaces on simple Lie groups*, J. Lie Theory, **5** (1995), 241-273, Zbl 860.43008.
- [SW71] E.M. Stein and G. Weiss, *Fourier Analysis on Euclidean Spaces*, Princeton University Press, Princeton, New Jersey, 1971, MR 46 #4102, Zbl 232.42007.

Received December 29, 1998. The second author was supported by NSF grant DMS-9626541, DMS-0070607, and LEQSF (1996-99)-RD-A-12.

UNIVERSITÄT GÖTTINGEN
 MATHEMATISCHES INSTITUT
 BUNSENSTRASSE 3-5, D-37073 GÖTTINGEN
 GERMANY
E-mail address: betten@cfgauss.uni-math.gwdg.de

DEPARTMENT OF MATHEMATICS
 LOUISIANA STATE UNIVERSITY
 BATON ROUGE, LA 70803
E-mail address: olafsson@math.lsu.edu