

*Pacific
Journal of
Mathematics*

ENTROPY IN TYPE I ALGEBRAS

SERGEY NESHVEYEV AND ERLING STØRMER

Volume 201 No. 2

December 2001

ENTROPY IN TYPE I ALGEBRAS

SERGEY NESHVEYEV AND ERLING STØRMER

It is shown that if (M, ϕ, α) is a W^* -dynamical system with M a type I von Neumann algebra then the entropy of α w.r.t. ϕ equals the entropy of the restriction of α to the center of M . If furthermore (N, ψ, β) is a W^* -dynamical system with N injective then $h_{\phi \otimes \psi}(\alpha \otimes \beta) = h_{\phi}(\alpha) + h_{\psi}(\beta)$.

1. Introduction.

In the theory of non-commutative entropy the attention has almost exclusively been concentrated on non-type I algebras. We shall in the present paper remedy this situation by proving the basic facts on entropy of automorphisms of type I C^* - and von Neumann-algebras. The results are as nice as one can hope. The CNT-entropy of an automorphism of a von Neumann algebra of type I with respect to an invariant normal state is the classical entropy of the restriction of the automorphism to the center of the algebra. If one factor of a tensor product of two von Neumann algebras is of type I and the other injective, then the entropy of a tensor product automorphism with respect to an invariant product state is the sum of the entropies. The results have obvious corollaries to type I C^* -algebras. The main idea behind our proofs is the use of conditional expectations of finite index, as employed in [GN].

We shall use the notation $h_{\phi}(\alpha)$ for the CNT-entropy of a C^* -dynamical system as defined by Connes, Narnhofer and Thirring in [CNT], and $h'_{\phi}(\alpha)$ for the ST-entropy defined by Sauvageot and Thouvenot in [ST].

2. Main results.

We first prove a general result for the Sauvageot-Thouvenot entropy for the restriction of an automorphism to a globally invariant C^* -subalgebra of finite index.

Proposition 1. *Let (A, ϕ, α) be a unital C^* -dynamical system. Let $B \subset A$ be an α -invariant C^* -subalgebra (with $1 \in B$). Suppose there exists a conditional expectation $E: A \rightarrow B$ such that $E \circ \alpha = \alpha \circ E$, $\phi \circ E = \phi$ and $E(x) \geq cx$ for all $x \in A^+$ for some $c > 0$. Then $h'_{\phi}(\alpha) = h'_{\phi}(\alpha|_B)$.*

Proof. Let (C, μ, β) be a C^* -dynamical system with C abelian. Using E we can lift any stationary coupling on $B \otimes C$ to a stationary coupling on $A \otimes C$. This, together with the property of monotonicity of relative entropy, shows that $h'_\phi(\alpha) \geq h'_\phi(\alpha|_B)$.

Conversely, suppose λ is a stationary coupling of (A, ϕ, α) with (C, μ, β) , and P is a finite-dimensional subalgebra of C with atoms p_1, \dots, p_n . Let

$$\phi_i(a) = \frac{\lambda(a \otimes p_i)}{\mu(p_i)} \text{ for } a \in A.$$

Then in the notations of [ST]

$$h'_\phi(\alpha) = \sup \left\{ H_\mu(P|P^-) - H_\mu(P) + \sum_{i=1}^n \mu(p_i) S(\phi, \phi_i) \mid (C, \mu, \beta, \lambda, P) \right\}.$$

Since $\phi_i \leq \frac{1}{\mu(p_i)}\phi$, ϕ_i is normal in the GNS-representation of ϕ . Since E is ϕ -invariant, it extends to a normal conditional expectation of the closure of A in the GNS-representation onto the closure of B . Thus we can apply [OP, Theorem 5.15] to ϕ and ϕ_i , and (as in the proof of Lemma 1.5 in [GN]) get

$$\begin{aligned} \sum_{i=1}^n \mu(p_i) S(\phi, \phi_i) &= \sum_{i=1}^n \mu(p_i) (S(\phi|_B, \phi_i|_B) + S(\phi_i \circ E, \phi_i)) \\ &\leq \sum_{i=1}^n \mu(p_i) S(\phi|_B, \phi_i|_B) - \log c. \end{aligned}$$

It follows that $h'_\phi(\alpha) \leq h'_\phi(\alpha|_B) - \log c$. Then for each $m \in \mathbb{N}$

$$h'_\phi(\alpha) = \frac{1}{m} h'_\phi(\alpha^m) \leq \frac{1}{m} h'_\phi(\alpha^m|_B) - \frac{1}{m} \log c = h'_\phi(\alpha|_B) - \frac{1}{m} \log c.$$

Thus $h'_\phi(\alpha) \leq h'_\phi(\alpha|_B)$. □

By [ST, Proposition 4.1] the Sauvageot-Thouvenot entropy coincides with the CNT-entropy for nuclear C^* -algebras. In fact, what is really necessary for the coincidence of the entropies, is the existence of a net of unital completely positive mappings γ_i of finite-dimensional C^* -algebras into A such that $S(\phi, \psi) = \lim_i S(\phi \circ \gamma_i, \psi \circ \gamma_i)$ for any positive linear functional ψ on A , $\psi \leq \phi$. We therefore have:

Corollary 2. *If in the above proposition A and B are injective von Neumann algebras and ϕ is normal then $h_\phi(\alpha) = h_\phi(\alpha|_B)$.*

To prove our main result we need also two simple lemmas. The first lemma is more or less well-known.

Lemma 3. *Let (M, ϕ, α) be a W^* -dynamical system. Then*

- (i) *if p is an α -invariant projection in M such that $\text{supp } \phi \leq p$, then $h_\phi(\alpha) = h_\phi(\alpha|_{M_p})$;*

- (ii) if $\{p_i\}_{i \in I}$ is a set of mutually orthogonal α -invariant central projections in M , $\sum_i p_i = 1$, then

$$h_\phi(\alpha) = \sum_i \phi(p_i)h_{\phi_i}(\alpha_i),$$

where $\phi_i = \frac{1}{\phi(p_i)}\phi$ is the normalized restriction of ϕ to Mp_i , and $\alpha_i = \alpha|_{Mp_i}$.

Proof.

(i) easily follows from the definitions; (ii) follows from [CNT, VII.5(iii)], (i) and [SV, Lemma 3.3] applied to the subalgebras $M(p_{i_1} + \dots + p_{i_n}) + \mathbb{C}(1 - p_{i_1} - \dots - p_{i_n})$. □

The proof of the following lemma is left to the reader.

Lemma 4. *Let T be an automorphism of a probability space (X, μ) , $f \in L^\infty(X, \mu)$ a T -invariant function such that $f \geq 0$ and $\int_X f d\mu = 1$. Let μ_f be the measure on X such that $d\mu_f/d\mu = f$. Then $h_{\mu_f}(T) \leq \|f\|_\infty h_\mu(T)$.*

Theorem 5. *Let (M, ϕ, α) be a W^* -dynamical system with M a von Neumann algebra of type I. Let Z denote the center of M . Then $h_\phi(\alpha) = h_\phi(\alpha|_Z)$.*

Proof. By Lemma 3(i) we may suppose that ϕ is faithful. Then M is a direct sum of homogeneous algebras of type I_n , $n \in \mathbb{N} \cup \{\infty\}$. By Lemma 3(ii) we may assume that M is homogeneous of type I_n . We first assume that $n \in \mathbb{N}$. Then $Z = L^\infty(X, \mu)$, where (X, μ) is a probability space and $\phi|_Z = \mu$. Thus

$$M \cong Z \otimes \text{Mat}_n(\mathbb{C}) = L^\infty(X, \text{Mat}_n(\mathbb{C})), \quad \phi = \int_X^\oplus \phi_x d\mu(x),$$

where $\phi_x = \text{Tr}(\cdot Q_x)$ is a state on $\text{Mat}_n(\mathbb{C})$, Tr the canonical trace on $\text{Mat}_n(\mathbb{C})$. We first assume $Q_x \geq c > 0$ for all x .

If $s \in M^+$, s is a function in $L^\infty(X, \text{Mat}_n(\mathbb{C}))$. Define the ϕ -preserving conditional expectation $E: M \rightarrow Z$ by $E(s)(x) = \phi_x(s(x))$. Then

$$E(s)(x) = \text{Tr}(s(x)Q_x) \geq c\text{Tr}(s(x)) \geq cs(x),$$

so $E(s) \geq cs$, and it follows from Corollary 2 that $h_\phi(\alpha) = h_\phi(\alpha|_Z)$.

If there is no $c > 0$ such that $Q_x \geq c$ for all x , let $X_c = \{x \in X \mid Q_x \geq c\}$, ($c > 0$),

$$N_c = L^\infty(X_c, \text{Mat}_n(\mathbb{C})) \quad \text{and} \quad M_c = N_c + \mathbb{C}\chi_{X \setminus X_c},$$

where $\chi_{X \setminus X_c}$ is the characteristic function of $X \setminus X_c$. Since ϕ is α -invariant so is M_c , so by the above argument and Lemma 3, letting $\phi_c = \frac{1}{\mu(X_c)}\phi|_{N_c}$ and $\mu_c = \frac{1}{\mu(X_c)}\mu|_{X_c}$, we obtain

$$h_\phi(\alpha|_{M_c}) = \mu(X_c)h_{\phi_c}(\alpha|_{N_c}) = \mu(X_c)h_{\mu_c}(T|_{X_c}) \leq h_\mu(T),$$

where T is the automorphism of (X, μ) induced by α . Letting $c \rightarrow 0$ and using [SV, Lemma 3.3] we obtain the Theorem when M is finite.

If M is homogeneous of type I $_{\infty}$, we have $M \cong L^{\infty}(X, \mu) \otimes B(H)$, where H is a separable Hilbert space. Let Tr denote the canonical trace on $B(H)$. Write again

$$\phi = \int_X^{\oplus} \phi_x d\mu(x), \quad \phi_x = \text{Tr}(\cdot Q_x),$$

and let $E_x(U)$ denote the spectral projection of Q_x corresponding to a Borel set U . Let $P_c \in M = L^{\infty}(X, B(H))$ be the projection defined by $P_c(x) = E_x([c, +\infty))$, where $c > 0$. Then P_c is an α -invariant finite projection. Let

$$M_c = P_c M P_c + \mathbb{C}(1 - P_c).$$

Then M_c is a finite type I von Neumann algebra. Its center is isomorphic to $L^{\infty}(X_c, \mu_c) \oplus \mathbb{C}$, and the restriction of ϕ to it is $\phi(P_c)\mu_c \oplus \phi(1 - P_c)$, where $X_c = \{x \in X \mid P_c(x) \neq 0\}$ and

$$\int_{X_c} f(x) d\mu_c(x) = \frac{1}{\phi(P_c)} \int_{X_c} f(x) \phi_x(P_c(x)) d\mu(x).$$

So we can apply the first part of the proof to M_c . Since $d\mu_c/d\mu \leq \frac{1}{\phi(P_c)}$, applying Lemma 4 we get

$$h_{\phi}(\alpha|_{M_c}) = \phi(P_c)h_{\mu_c}(T|_{X_c}) \leq h_{\mu}(T).$$

Now letting $c \rightarrow 0$ we conclude that $h_{\phi}(\alpha) = h_{\mu}(T)$. □

It should be remarked that in a special case the above theorem was proved in [GS, Proposition 2.4].

If A is a C^* -algebra and ϕ a state on A , the central measure μ_{ϕ} of ϕ is the measure on the spectrum \hat{A} of A defined by $\mu_{\phi}(F) = \phi(\chi_F)$, where ϕ is regarded as a normal state on A'' , see [P, 4.7.5]. Thus by Theorem 5 and [P, 4.7.6] we have the following:

Corollary 6. *Let (A, ϕ, α) be a C^* -dynamical system with A a separable unital type I C^* -algebra. Then $h_{\phi}(\alpha) = h_{\mu_{\phi}}(\hat{\alpha})$, where $\hat{\alpha}$ is the automorphism of the measure space (\hat{A}, μ_{ϕ}) induced by α .*

Since inner automorphisms act trivially on the center we have:

Corollary 7. *If (M, ϕ, α) is a W^* -dynamical system with M of type I and α an inner automorphism then $h_{\phi}(\alpha) = 0$.*

Note that in the finite case the above corollary also follows from a result of N. Brown [Br, Lemma 2.2].

The next result was shown in [S] when ϕ is a trace.

Corollary 8. *Let R denote the hyperfinite II_1 -factor. Let A be a Cartan subalgebra of R and u a unitary operator in A . If ϕ is a normal state such that u belongs to the centralizer of ϕ then $h_\phi(\text{Ad } u) = 0$.*

Proof. As in [S], it follows from [CFW] that there exists an increasing sequence of full matrix algebras $N_1 \subset N_2 \subset \dots$ with union weakly dense in R such that $A \cong A_n \otimes B_n$, where $A_n = N_n \cap A$ and $B_n = (N'_n \cap R) \cap A$ for all $n \in \mathbb{N}$. Let $M_n = N_n \otimes B_n$. Then M_n is of type I and contains u . Hence $h_\phi(\text{Ad } u|_{M_n}) = 0$. Since $(\cup_n M_n)^\perp = R$, $h_\phi(\text{Ad } u) = 0$ by [SV, Lemma 3.3]. \square

If (A, ϕ, α) and (B, ψ, β) are C^* -dynamical systems we always have

$$h_{\phi \otimes \psi}(\alpha \otimes \beta) \geq h_\phi(\alpha) + h_\psi(\beta),$$

see [SV, Lemma 3.4]. Equality does not always hold, see [NST] or [Sa]. However, we have:

Theorem 9. *Let (A, ϕ, α) and (B, ψ, β) be W^* -dynamical systems. Suppose that A is of type I, and B is injective. Then*

$$h_{\phi \otimes \psi}(\alpha \otimes \beta) = h_\phi(\alpha) + h_\psi(\beta).$$

Proof. We shall rather prove that $h_{\phi \otimes \psi}(\alpha \otimes \beta) = h_\phi(\alpha|_{Z(A)}) + h_\psi(\beta)$. For this it suffices to consider the case when A is abelian; the general case will follow by the same arguments as in the proof of Theorem 5. (Note that the mapping $x \mapsto \text{Tr}(x) - x$ on $\text{Mat}_n(\mathbb{C})$ is not completely positive, but the mapping $x \mapsto \text{Tr}(x) - \frac{1}{n}x$ is by the Pimsner-Popa inequality. Thus replacing M with $M \otimes B$ and Z with $Z \otimes B$ in the proof of Theorem 5 we have to replace the inequality $E(s) \geq cs$ in the proof with $E(s) \geq \frac{c}{n}s$.)

So suppose that A is abelian. It is clear that it suffices to prove that if A_1, \dots, A_n are finite-dimensional subalgebras of A , and B_1, \dots, B_n are finite-dimensional subalgebras of B , then

$$H_{\phi \otimes \psi}(A_1 \otimes B_1, \dots, A_n \otimes B_n) = H_\phi(A_1, \dots, A_n) + H_\psi(B_1, \dots, B_n).$$

We always have the inequality " \geq ", [SV, Lemma 3.4]. To prove the opposite inequality consider a decomposition

$$\phi \otimes \psi = \sum_{i_1, \dots, i_n} \omega_{i_1 \dots i_n}.$$

Let $H_{\{\phi \otimes \psi = \sum \omega_{i_1 \dots i_n}\}}(A_1 \otimes B_1, \dots, A_n \otimes B_n)$ be the entropy of the corresponding abelian model, so

$$\begin{aligned} & H_{\{\phi \otimes \psi = \sum \omega_{i_1 \dots i_n}\}}(A_1 \otimes B_1, \dots, A_n \otimes B_n) \\ &= \sum_{i_1, \dots, i_n} \eta_{\omega_{i_1 \dots i_n}}(1) + \sum_{k=1}^n \sum_i S \left(\phi \otimes \psi|_{A_k \otimes B_k}, \sum_{i_k=i} \omega_{i_1 \dots i_n}|_{A_k \otimes B_k} \right). \end{aligned}$$

Set $C = \bigvee_{k=1}^n A_k$. Let p_1, \dots, p_r be those atoms p of C for which $\phi(p) > 0$. Define positive linear functionals $\psi_{m,i_1\dots i_n}$ on B ,

$$\psi_{m,i_1\dots i_n}(b) = \frac{\omega_{i_1\dots i_n}(p_m \otimes b)}{\phi(p_m)}.$$

Let also ϕ_m be the linear functional on C defined by the equality $\phi_m(a) = \phi(ap_m)$. Then

$$\omega_{i_1\dots i_n} = \sum_{m=1}^r \phi_m \otimes \psi_{m,i_1\dots i_n} \text{ on } C \otimes B,$$

and

$$\psi = \sum_{i_1, \dots, i_n} \psi_{m,i_1\dots i_n} \text{ for } m = 1, \dots, r.$$

Since the supports of the positive functionals ϕ_m are mutually orthogonal minimal projections in C , we have

$$\begin{aligned} & \sum_{k=1}^n \sum_i S \left(\phi \otimes \psi|_{A_k \otimes B_k}, \sum_{i_k=i} \omega_{i_1\dots i_n}|_{A_k \otimes B_k} \right) \\ & \leq \sum_{k=1}^n \sum_i S \left(\phi \otimes \psi|_{C \otimes B_k}, \sum_{i_k=i} \omega_{i_1\dots i_n}|_{C \otimes B_k} \right) \\ & = \sum_{k=1}^n \sum_i S \left(\phi \otimes \psi|_{C \otimes B_k}, \sum_{m=1}^r \phi_m \otimes \left(\sum_{i_k=i} \psi_{m,i_1\dots i_n} \right) |_{C \otimes B_k} \right) \\ & = \sum_{k=1}^n \sum_i \sum_{m=1}^r \phi(p_m) S \left(\psi|_{B_k}, \sum_{i_k=i} \psi_{m,i_1\dots i_n}|_{B_k} \right). \end{aligned}$$

If $a_i \geq 0$ then $\eta \left(\sum_i a_i \right) \leq \sum_i \eta(a_i)$. Hence we have

$$\begin{aligned} & \sum_{i_1, \dots, i_n} \eta \omega_{i_1\dots i_n}(1) \\ & \leq \sum_{m=1}^r \sum_{i_1, \dots, i_n} \eta(\phi_m \otimes \psi_{m,i_1\dots i_n})(1) \\ & = \sum_{m=1}^r \eta\phi(p_m) \sum_{i_1, \dots, i_n} \psi_{m,i_1\dots i_n}(1) + \sum_{m=1}^r \phi(p_m) \sum_{i_1, \dots, i_n} \eta\psi_{m,i_1\dots i_n}(1) \\ & = \sum_{m=1}^r \eta\phi(p_m) + \sum_{m=1}^r \phi(p_m) \sum_{i_1, \dots, i_n} \eta\psi_{m,i_1\dots i_n}(1). \end{aligned}$$

Thus

$$\begin{aligned} & H_{\{\phi \otimes \psi = \sum \omega_{i_1 \dots i_n}\}}(A_1 \otimes B_1, \dots, A_n \otimes B_n) \\ & \leq \sum_{m=1}^r \eta \phi(p_m) + \sum_{m=1}^r \phi(p_m) H_{\{\psi = \sum \psi_{m, i_1 \dots i_n}\}}(B_1, \dots, B_n). \end{aligned}$$

Since $\sum_m \eta \phi(p_m) = H_\phi(C) = H_\phi(A_1, \dots, A_n)$, we conclude that

$$H_{\phi \otimes \psi}(A_1 \otimes B_1, \dots, A_n \otimes B_n) \leq H_\phi(A_1, \dots, A_n) + H_\psi(B_1, \dots, B_n),$$

completing the proof of the Theorem. \square

References

- [Br] N.P. Brown, *A note on topological entropy, embeddings and unitaries in nuclear quasidiagonal C^* -algebras*, Proc. Amer. Math. Soc., **128** (2000), 2603–2609, MR 2000m:46138.
- [CFW] A. Connes, J. Feldman and B. Weiss, *An amenable equivalence relation is generated by a single transformation*, Ergod. Th. & Dynam. Sys., **1** (1981), 431–450, MR 84h:46090, Zbl 491.28018.
- [CNT] A. Connes, H. Narnhofer and W. Thirring, *Dynamical entropy of C^* -algebras and von Neumann algebras*, Commun. Math. Phys., **112** (1987), 691–719, MR 89b:46078, Zbl 637.46073.
- [GN] V.Ya. Golodets and S.V. Neshveyev, *Entropy of automorphisms of II_1 -factors arising from the dynamical systems theory*, J. Funct. Anal., **181**(1) (2001), 14–28, CMP 1 818 110.
- [GS] V.Ya. Golodets and E. Størmer, *Entropy of C^* -dynamical systems defined by bitstreams*, Ergod. Th. & Dynam. Sys., **18** (1998), 859–874, MR 99m:46157, Zbl 932.46057.
- [NST] H. Narnhofer, E. Størmer and W. Thirring, *C^* -dynamical systems for which the tensor product formula for entropy fails*, Ergod. Th. & Dynam. Sys., **15** (1995), 961–968, MR 97g:46088, Zbl 832.46059.
- [OP] M. Ohya and D. Petz, *Quantum Entropy and Its Use*, Springer, 1993, MR 94k:81002, Zbl 891.94008.
- [P] G.K. Pedersen, *C^* -Algebras and their Automorphism Groups*, Academic Press, 1979, MR 81e:46037, Zbl 416.46043.
- [Sa] J.-L. Sauvageot, *Ergodic properties of the action of a matrix in $SL(2, \mathbb{Z})$ on a non-commutative 2-torus*, preprint.
- [ST] J.-L. Sauvageot and J.-P. Thouvenot, *Une nouvelle definition de l'entropie dynamique des systemes non-commutatifs*, Commun. Math. Phys., **145** (1992), 411–423, MR 93d:46113, Zbl 772.46036.
- [S] E. Størmer, *Entropy of some inner automorphisms of the hyperfinite II_1 -factor*, Intern. J. Math., **4** (1993), 319–322, MR 94h:46099, Zbl 804.46075.

- [SV] E. Størmer and D. Voiculescu, *Entropy of Bogoliubov automorphisms of the canonical anticommutation relations*, Commun. Math. Phys., **133** (1990), 521-542, MR 92e:46125, Zbl 743.46051.

Received January 31, 2000. The first author was partially supported by NATO grant SA (PST.CLG.976206)5273. The second author was partially supported by the Norwegian Research Council.

B. VERKIN INSTITUTE FOR LOW TEMPERATURE PHYSICS AND ENGINEERING
NATIONAL ACADEMY OF SCIENCES OF UKRAINE,
47, LENIN AVE.
310164, KHARKOV
UKRAINE

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OSLO
P.O. Box 1053
BLINDERN, 0316 OSLO
NORWAY
E-mail address: erlings@math.uio.no