

*Pacific
Journal of
Mathematics*

FLOW EQUIVALENCE OF SHIFTS OF FINITE TYPE VIA
POSITIVE FACTORIZATIONS

MIKE BOYLE

Volume 204 No. 2

June 2002

FLOW EQUIVALENCE OF SHIFTS OF FINITE TYPE VIA POSITIVE FACTORIZATIONS

MIKE BOYLE

Together with M. Boyle and D. Huang (2000), this paper gives an alternate development of the Huang classification of shifts of finite type up to flow equivalence, and provides additional functorial information, used to analyze the action of the mapping class group of the mapping torus of a shift of finite type on the “isotopy futures” group, which is introduced here. For a shift of finite type σ_A , this group is isomorphic to the Bowen-Franks group $\text{cok}(I - A)$. The action on the isotopy futures group of a subshift is the flow equivalence analogue of the dimension group representation.

1. Introduction.

Shifts of finite type (SFTs) are the fundamental building blocks of symbolic dynamics, with applications to hyperbolic dynamics, ergodic theory, topological dynamics, matrix theory and other areas [Bow, DGS, Ki, LM, Rob, S]. Any SFT is conjugate to an SFT σ_A defined by a matrix A with nonnegative integer entries. A fundamental question about SFTs, when are they flow equivalent, is important also for the study of certain C^* -algebras [C, CK, H2, H3, R]. This question was solved in the irreducible case by Franks [F], extending earlier work of Parry and Sullivan [PS] and Bowen and Franks [BowF], and then in the general case by Huang [H4, H5], following earlier work on more tractable special cases [H1, H3]. Huang [H4, H5] developed complete algebraic invariants (defined in terms of the given matrix A) for flow equivalence of SFTs.

This paper has three main features.

- (1) Taken together with [BH], the paper gives a self-contained alternate development of the Huang classification of SFTs up to flow equivalence. This development separates algebraic and positivity issues, and provides additional functorial information.
- (2) We introduce the isotopy futures group \mathcal{F}_S of the mapping torus Y_S of a subshift S , and when S is an SFT σ_A we construct an isomorphism of \mathcal{F}_S and the Bowen-Franks group $\text{cok}(I - A)$, and analyze the induced action of the mapping class group of Y_S on \mathcal{F}_S .

- (3) We integrate the study of flow equivalence of SFTs into the “positive K -theory” framework for classification problems in symbolic dynamics.

We now discuss these features in more detail.

1. To study reducible SFTs, we work with certain infinite block triangular integral matrices with block rows and columns indexed by a finite poset \mathcal{P} : If $i \not\leq j$ in \mathcal{P} , then the ij block of the matrix must be zero. The elements of \mathcal{P} , and their ordering, correspond to the irreducible components of the SFT, and their asymptotic transitions; the isomorphism class of the poset \mathcal{P} is an invariant of flow equivalence. We say two such matrices B, B' are $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalent if there are matrices U, V satisfying the same zero-subblock conditions, and with all diagonal blocks having determinant 1, such that $UBV = B'$. After fixing a choice of \mathcal{P} , and allowing a permutation of \mathcal{P} , we show that A, A' define flow equivalent SFTs if and only if the matrices $I - A$ and $I - A'$ are $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalent by an equivalence which is “positive on cycle components” (a technical condition which may be removed after reduction to a standard form, see Theorem 3.4). The key to this result is the Factorization Theorem 3.3, which gives necessary and sufficient conditions for an $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence to be a composition of “positive” elementary equivalences (which induce flow equivalences). Complete algebraic invariants for $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence are contained in the joint work [BH] with Danrun Huang. (The proofs in the current paper are very different from those of Huang [H4, H5], but the algebraic sequel [BH] depends completely on the ideas introduced by Huang in [H4, H5].)

In Huang’s development, the proofs involve creating positive matrix models realizing given isomorphisms of an associated “ K -web” of exact sequences of associated groups; the difficult positivity and algebraic issues are intertwined. By interposing the $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence relation between the SFTs and the complicated K -web algebraic invariants, we separate the positivity issues (which we address in this paper) from purely algebraic issues (which are addressed in [BH]). This clarifies the meaning of the invariants and the structure of the problem. It also facilitates the application of algebraic results.

2. The analysis of the induced action on $\mathrm{cok}(I - A)$ uses the Factorization Theorem 3.3 together with purely algebraic results from [BH] on $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence. There is a plausible program (7.15) for extending these ideas to obtain more information.

It seems to be nontrivial to construct a functor which attaches isomorphisms of Bowen-Franks groups to isotopy classes of flow equivalences of SFTs. (For example, we do not know if it is possible to construct a homomorphism from $\check{H}^1(Y_{\sigma_A})$ onto the Bowen-Franks group $\mathrm{cok}(I - A)$ such that the natural action of the mapping class group on $\check{H}^1(Y_{\sigma_A})$ induces an

automorphism of $\text{cok}(I - A)$; and we suspect there is no such homomorphism.) An alternate approach using work of Badoian is discussed at the end of Section 7. Another possible approach would be to extend ideas of Bowen and Franks, who computed $\text{cok}(I - A)$ as a relative homology group in the context of basic sets of Smale flows ([**BowF**, **F**]).

3. In the framework of positive K -theory (a term introduced by Wagoner), some class of matrices A presents some category of dynamical system, and multiplication of $I - A$ by elementary matrices satisfying some positivity condition induces isomorphisms of the system presented by A . This framework was born in [**KRW2**, **KRW3**], where matrices over $t\mathbb{Z}_+[t]$ presented SFTs, and multiplication of $I - A$ by certain elementary matrices over $\mathbb{Z}[t]$ gave a completely new method of constructing topological conjugacies, which allowed the solution of a difficult and important open problem. This framework for SFT's is developed or exploited further in [**BW**, **B1**, **KR1**, **W2**]; in the last reference [**W2**], the K -theory connection is more than a formal analogy and gives new counterexamples to Williams' shift equivalence conjecture. In [**G**], the matrix entries lie in a certain ring of formal power series, and the elementary matrix multiplications induce good finitary isomorphisms of Markov chains. In this paper and in [**Ba1**], the matrices have integer (or zero-one) entries, and the elementary multiplications induce flow equivalences. There is a passage from the topological conjugacy case to the flow equivalence case by "setting t equal to 1" (applying the coinvariants functor), as described in [**B1**]. The positive K -theory approach gives a unified and useful framework for classification problems in symbolic dynamics, and we view this paper as a significant piece of the theory for the case of flow equivalence of SFTs. It is possible that the methods of this paper may be suggestive for the case of topological conjugacy of SFTs.

Some of our results on flow equivalence have alternate proofs based on the work of Leslie Badoian [**Ba1**], who develops for irreducible SFTs a flow equivalence theory analogous to the theory created by Wagoner for topological conjugacies of SFTs. At the end of Section 7, we summarize the main results of [**Ba1**], and discuss those alternate proofs.

Now some words on the structure of the paper. In Section 2 we give some definitions and technical results necessary for the statement of the main results in Section 3. The proof of the Factorization Theorem is carried out in Sections 4-5 and the Appendix. Shifts of finite type and the relation of flow equivalence to the matrix results are addressed in Section 6. The isotopy futures group and connections to flow equivalence are studied in Section 7. For the simple general statement of our Factorization Theorem for matrices, we need preliminary technical arguments to reduce our matrices to a nondegenerate form. These preliminaries are complicated, and we banish them to the Appendix.

The basic approach of this paper, and the Factorization Theorem in the “no cycle components” case under additional technical assumptions since removed, were announced in [B1].

I thank Danrun Huang for many helpful comments, and for a very satisfying collaboration in our sequel paper [BH]. Also, without his earlier work, this paper would not exist.

2. Definitions.

2.1. Poset blocked matrices. For the rest of the paper, we let $\mathcal{P} = \{1, \dots, N\}$ denote a finite poset (partially ordered set). We describe the order with a relation \prec satisfying the following conditions (in which $<$ refers to the usual order on \mathbb{N}) for all i, j, k in \mathcal{P} :

$$\begin{aligned} i \prec j &\implies i < j, \\ i \prec j \prec k &\implies i \prec k. \end{aligned}$$

We write $i \preceq j$ to mean that $i \prec j$ or $i = j$. We can visualize the poset as an acyclic directed graph with vertex set $\{1, \dots, N\}$ and transitions $i \rightarrow j$ iff $i \prec j$.

We say that a matrix (or a block in a matrix) is *square* if its rows and columns are indexed by the same set (which may be finite or countably infinite). Suppose that n_1, \dots, n_N lie in the set $\{1, 2, \dots, \infty\}$. Let $\mathbf{n} = (n_1, \dots, n_N)$. We say a square matrix M is “ \mathbf{n} -blocked” if it splits into blocks M_{ij} , $1 \leq i, j \leq N$, where M_{ij} denotes the intersection of the i th block row and the j th block column, and has size $n_i \times n_j$. (We will also use the notation $M\{i, j\}$ to denote M_{ij} .) Given an \mathbf{n} -blocked matrix M , we let \mathcal{I}_j denote the set of indices for rows/columns through the block M_{jj} .

Definition 2.1. $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ is the set of \mathbf{n} -blocked matrices with entries in \mathbb{Z} satisfying the following conditions:

- (1) For $1 \leq i \leq N$, the block M_{ii} equals the identity matrix in all but finitely many entries.
- (2) For $1 \leq i, j \leq N$ and $i \neq j$, the block M_{ij} is zero in all but finitely many entries.
- (3) If $i \not\preceq j$, then the block M_{ij} is zero.

The matrices in the semiring $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ are block upper triangular and in addition certain blocks above the diagonal must be zero. $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ is closed under addition and (because \prec is transitive) under matrix multiplication.

A nonnegative matrix A is *irreducible* if it is square with all entries non-negative, and for every (i, j) there exists $n > 0$ such that $A^n(i, j) > 0$. (In particular, for us a zero matrix is not irreducible.) A square matrix is *essentially irreducible* if it has a unique principal submatrix which is irreducible and which is contained in no larger irreducible principal submatrix.

Definition 2.2. $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ is the set of \mathbf{n} -blocked nonnegative integral matrices with only finitely many nonzero entries, satisfying the following conditions:

- (1) Each diagonal block M_{ii} is essentially irreducible.
- (2) If $i \not\prec j$, then the block M_{ij} is zero.
- (3) If $i \prec j$, then there is an index i' occurring on a cycle of M_{ii} and an index j' occurring on a cycle of M_{jj} and a positive integer n , such that $A^n(i', j') > 0$.

Definition 2.3. $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ is the set of matrices M in $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ such that $\det(M_{ii}) = 1$ for $1 \leq i \leq N$.

Abbreviations 2.4. $\mathfrak{M}_{\mathcal{P}}(\mathbb{Z})$, $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ and $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ denote the sets $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$, $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ and $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ for which $\mathbf{n} = (n_1, \dots, n_N)$ with every $n_i = \infty$. Whenever any such matrix family appears with no subscript \mathcal{P} , it means that $\mathcal{P} = \{1\}$ (the block structure is trivial).

We say two matrices B, B' in $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ are $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ -equivalent in $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ if there are matrices U, V in $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ such that $UBV = B'$, and we write this as $(U, V): B \rightarrow B'$. We say a matrix is a basic elementary matrix if it equals the identity matrix except in at most one offdiagonal entry. It is not difficult to check that $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ is a group under multiplication which is generated by basic elementary matrices [BH]. Given $\mathbf{n} \leq \mathbf{r}$, we have natural truncation and embedding maps between \mathbf{n} -blocked and \mathbf{r} -blocked matrices,

$$\begin{aligned} \text{tru}_{\mathbf{n}}: \mathfrak{M}_{\mathcal{P}}(\mathbf{r}, \mathbb{Z}) &\rightarrow \mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z}), \\ \iota_{\mathbf{r}}: \mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z}) &\rightarrow \mathfrak{M}_{\mathcal{P}}(\mathbf{r}, \mathbb{Z}). \end{aligned}$$

The truncation map replaces an ij block with its $n_i \times n_j$ upper left corner. The embedding map embeds an ij block as the upper left corner of an ij block. If $i \neq j$, then the image ij block is zero outside this embedded left corner; if $i = j$, it is the identity outside this left corner. We will use A_{∞} to abbreviate $\iota_{\mathbf{n}}(A)$ in the case that every $n_i = \infty$. We will also use $\text{tru}_{\mathbf{n}}$, $\iota_{\mathbf{r}}$ and A_{∞} for matrix families other than $\mathfrak{M}_{\mathcal{P}}$. The only potentially ambiguous point, which should be clear from context, is whether the embedded block corners should be extended as above with $\mathfrak{M}_{\mathcal{P}}$ to match the identity matrix, or should be extended to match the zero matrix (e.g., when the range is $\mathfrak{M}_{\mathcal{P},+}^{\circ}$).

2.2. Positive equivalence. Suppose for some (i, j) that E is a basic elementary matrix with offdiagonal entry $E(i, j) = 1$, $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, and $A(i, j) > 0$. Then we say that each of the equivalences

$$\begin{aligned} (E, I): (I - A) &\rightarrow E(I - A), & (E^{-1}, I): E(I - A) &\rightarrow (I - A), \\ (I, E): (I - A) &\rightarrow (I - A)E, & (I, E^{-1}): (I - A)E &\rightarrow (I - A) \end{aligned}$$

is a *basic positive equivalence* in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$. Note, $E \in \text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$. We say that an $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence is a *positive equivalence* in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ if it is a composition of basic positive equivalences in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$.

To understand the meaning of a basic positive equivalence, suppose $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ with $A(i, j) > 0$ and E is basic elementary matrix with offdiagonal entry $E(i, j) = 1$. We first discuss the case $(E, I): (I - A) \rightarrow E(I - A)$. Define A' by the requirement $E(I - A) = (I - A')$. Then A and A' agree except perhaps in row i , where

$$\begin{aligned} A'(i, k) &= A(i, k) + A(j, k) && \text{if } j \neq k, && \text{and} \\ A'(i, j) &= A(i, j) + A(j, j) - 1. \end{aligned}$$

View A as the adjacency matrix of a directed graph \mathcal{G}_A with edge set \mathcal{E}_A and vertex set given by the $n_1 + \dots + n_N$ indices for the rows/columns of A . (There can be edges joining only finitely many of those vertices.) We can describe a directed graph $\mathcal{G}_{A'}$ which has A' as its adjacency matrix as follows. $\mathcal{G}_{A'}$ has the same vertex set as \mathcal{G}_A . Now pick an edge e which runs from vertex i to vertex j in \mathcal{G}_A (e exists because by assumption $A(i, j) > 0$). The edge set $\mathcal{E}_{A'}$ will be derived from \mathcal{E}_A as follows: Remove e from \mathcal{E}_A ; and then for every vertex k , for every edge f in \mathcal{E}_A from j to k add in a new edge (named $[ef]$) from i to k . It is easy to verify that with this edge set $\mathcal{E}_{A'}$, the directed graph $\mathcal{G}_{A'}$ has adjacency matrix A' .

With this notation, now define a map $\gamma: \mathcal{E}_{A'} \rightarrow (\mathcal{E}_A)^*$ by $\gamma: f \mapsto f$ and $\gamma: [ef] \mapsto ef$. Then γ induces an injective map (also called γ), from the set $\Sigma_{A'}$ of biinfinite paths through $\mathcal{G}_{A'}$ to the set Σ_A of biinfinite paths through \mathcal{G}_A , sending x' to x by the rule

$$\gamma: \dots x'_{-2}x'_{-1}|x'_0x'_1\dots \mapsto \dots \gamma(x'_{-2})\gamma(x'_{-1})|\gamma(x'_0)\gamma(x'_1)\dots$$

(in which the placement of the vertical bar indicates the indexing for x , e.g., $x_0x_1\dots = \gamma(x'_0)\gamma(x'_1)\dots$). Briefly: We get x from x' by replacing each A' edge $[ef]$ with ef .

The injective map $\gamma: \Sigma_{A'} \rightarrow \Sigma_A$ is not surjective precisely because the image will not contain points x for which $x_{-1} = e$ (the image will contain the shifted point $\sigma^{-1}x$ which is defined by $(\sigma^{-1}x)_i = x_{i-1}$). However, although γ is generally not a bijection, it does induce a bijection of orbits (under the shift) in Σ_A and $\Sigma_{A'}$. Also, γ induces a bijection of finite orbits: That is, γ induces a bijection (also called γ) of cycles in \mathcal{G}_A and $\mathcal{G}_{A'}$ (which need not respect the cycle length). If $1 \leq t \leq N$ and c is a cycle for the block A_{tt} , then $\gamma(c)$ is a cycle for A'_{tt} , because if i and j are not indices for the same component then γ is the identity on cycles. Also, if x in Σ_A is backwardly asymptotic (under the shift) to a cycle c and forwardly asymptotic to a cycle \tilde{c} , then $\gamma(x)$ is backwardly asymptotic to $\gamma(c)$ and forwardly asymptotic to $\gamma(\tilde{c})$. It follows that the matrix A' satisfies the conditions of Definition 2.2 and lies in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$.

The discussion of the case $(I, E): (I - A) \rightarrow (I - A)E = (I - A')$ is much the same. Let f be an A -edge from i to j . To form the A' graph from the A graph in this case, delete f , and add a new edge $[ef]$ for each edge e with terminal vertex i . Then define γ as before.

The following lemma, fundamental to the sequel, is implicit in Franks' paper [F].

Lemma 2.5. *Suppose $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, E is a basic elementary matrix in $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ whose nonzero offdiagonal entry is $E(i, j) = 1$, and there is a positive integer k such that $A^k(i, j) > 0$.*

- (1) *If $(E(I - A))(i, j) \leq 0$, then $(E, I): (I - A) \rightarrow E(I - A)$ is a positive equivalence in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$.*
- (2) *If $((I - A)E)(i, j) \leq 0$, then $(I, E): (I - A) \rightarrow (I - A)E$ is a positive equivalence in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$.*

Proof. We will consider the claim for the first equivalence (E, I) ; the other case is similar. By assumption, there is a list $i = i_0, i_1, \dots, i_k = j$ (which we take to be of minimal length, so the indices i_0, i_1, \dots, i_k are distinct) such that for $0 \leq t < k$ we have $A(i_t, i_{t+1}) > 0$. If $k = 1$, then the equivalence is a basic positive equivalence (and we know a basic positive equivalence takes a matrix in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ to a matrix in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$). So suppose $k > 1$. Let F_t be the elementary matrix whose which acts to add row i_t to row i . Let $F = F_{k-1} \dots F_1$. Then we have basic positive equivalences

$$\begin{aligned} (I - A) &\rightarrow F_1(I - A) \rightarrow F_2F_1(I - A) \rightarrow \dots \rightarrow (F_{k-1} \dots F_2F_1)(I - A) \\ &= F(I - A) \rightarrow EF(I - A) \rightarrow (F_{k-1})^{-1}EF(I - A) \\ &\rightarrow \dots \rightarrow (F_1)^{-1} \dots (F_{k-2})^{-1}(F_{k-1})^{-1}EF(I - A) \\ &= F^{-1}EF(I - A) = E(I - A). \end{aligned}$$

□

2.3. Cycle components. The technical discussion of this subsection is only required for the case when the matrix A in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ has a diagonal block whose maximal irreducible submatrix is a permutation matrix.

Lemma 2.6. *Suppose A is an $\mathcal{S} \times \mathcal{S}$ nonnegative integral matrix which has as its unique irreducible submatrix a cyclic permutation matrix. Then the cokernel group $\text{cok}(I - A) = \mathbb{Z}^{\mathcal{S}} / (I - A)\mathbb{Z}^{\mathcal{S}}$ is isomorphic to \mathbb{Z} . Let \mathcal{I} denote the set of indices involved in the cyclic permutation. Then the canonical basis vectors satisfy the following conditions:*

- (1) $[e_i]$ is a generator of $\text{cok}(I - A)$ if $i \in \mathcal{I}$.
- (2) $[e_i] = [e_j]$ if i and j are in \mathcal{I} .
- (3) $[e_i] = 0$ if $i \notin \mathcal{I}$.

Proof. (3) If $i \notin \mathcal{I}$, then for large n , $A^n e_i = 0$, and $e_i = (I - A^n)e_i = (I - A)(I + A + \dots + A^{n-1})e_i$. Then $[e_i] = 0$ in $\text{cok}(I - A)$.

(2) Let π denote the given permutation and suppose i and j are in \mathcal{I} . Then there exists $n > 0$ such that $A^n e_i = e_j$, so in $\text{cok}(I - A)$ we have $[e_i] - [e_j] = [e_i - e_j] = [(I - A^n)e_i] = 0$.

(1) Clearly now, if $i \in \mathcal{I}$, then $[e_i]$ generates $\text{cok}(I - A)$. Also, because $\det(I - A) = 0$, \mathbb{Z} is a subgroup of $\text{cok}(I - A)$. Therefore $\text{cok}(I - A) \cong \mathbb{Z}$. \square

For a matrix A satisfying the hypotheses of the lemma, we make $\text{cok}(I - A)$ an ordered group by declaring its positive set to be the collection of those $[w]$ such that (in the notation of the statement of the lemma) $\sum_{i \in \mathcal{I}} w_i \geq 0$. (This sum does not depend on the representative w of $[w]$.) We say an isomorphism between two such cokernel groups is *positive* if it takes the positive set in the domain to the positive set in the range.

If $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, then for $1 \leq i \leq N$ the diagonal block A_{ii} contains a unique maximal irreducible principal submatrix. If this matrix is a permutation matrix, then we say that i is a *cycle component* of A . We let \mathcal{C}_A denote the set of cycle components of A . For each i in \mathcal{C}_A , we make the cokernel group

$$\text{cok}(I - A)_{ii} = \mathbb{Z}^{n_i} / (I - A)_{ii} \mathbb{Z}^{n_i} \cong \mathbb{Z}$$

an ordered group as described above. For A and A' in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, if (U, V) is an $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence from A to A' , then for $1 \leq i \leq N$ the equivalence (U, V) induces an $\text{SL}(\mathbb{Z})$ equivalence (U_{ii}, V_{ii}) from A_{ii} to A'_{ii} , and this induces an isomorphism from $\text{cok}(I - A)_{ii}$ to $\text{cok}(I - A')_{ii}$ by the rule $[x] \mapsto [U_{ii}x]$. We say that the $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence (U, V) is *positive on cycle components* if this induced isomorphism of the i th component cokernel groups is positive whenever i is a cycle component for both A and A' . For example, if

$$A_{ii} = A'_{ii} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad U_{ii} = V_{ii} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix},$$

then (U, V) is not positive on cycle components.

Proposition 2.7. *Suppose (U, V) is a positive $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence from $(I - A)$ to $(I - A')$ in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$. Then:*

- (1) A and A' have the same cycle components, and
- (2) (U, V) is positive on cycle components.

Proof. It suffices to consider the case $(U, V) = (E, I)$ where E is a basic elementary matrix with offdiagonal entry $E(i_1, j_1) = 1$ such that i_1 and j_1 index rows through A_{ii} and i is a cycle component of A .

(1) It is clear from the earlier discussion on positive equivalence that the i th component of A has a unique cycle iff the i th component of A' has a unique cycle.

(2) For any canonical basis vector e_s , the vector Ee_s is nonnegative because E is nonnegative. It follows that (E, I) must be positive on components. \square

3. Statement of results.

In this section we state the main results (Theorem 3.1 and Theorem 3.3) which do not involve the mapping class group. We also give Theorems 3.4 and 3.5, which clarify computational issues. The definition of flow equivalence is given in Section 6, and all discussion of the mapping class group results is deferred to Section 7.

We need a little more notation. Given \mathcal{P} , we will use the same index set $\mathcal{I}^{\mathcal{P}}$, a disjoint union of countably infinite sets $\mathcal{I}_p^{\mathcal{P}}$, $p \in \mathcal{P}$, for all matrices with $\mathcal{P} \times \mathcal{P}$ blocking into infinite subblocks. Given finite posets $\mathcal{P}, \mathcal{P}'$, let $\text{Iso}[\mathcal{P}, \mathcal{P}']$ be the set of poset isomorphisms from \mathcal{P} to \mathcal{P}' . For each ν in $\text{Iso}[\mathcal{P}, \mathcal{P}']$, fix an infinite permutation matrix $P = P_\nu$ such that

$$P(i, j) = 1 \text{ and } j \in \mathcal{I}_p^{\mathcal{P}} \Rightarrow i \in \mathcal{I}_{\nu(p)}^{\mathcal{P}'}$$

Informally, a block $P\{p, q\}$ is zero if $q \neq \nu(p)$ and is the (infinite) identity matrix if $q = \nu(p)$.

Theorem 3.1 (Classification Theorem). *Suppose A is in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ and A' is in $\mathfrak{M}_{\mathcal{P}',+}^{\circ}(\mathbb{Z})$. The following are equivalent:*

- (1) *The SFTs σ_A and $\sigma_{A'}$ are flow equivalent.*
- (2) *For some $\nu \in \text{Iso}[\mathcal{P}, \mathcal{P}']$, with $P = P_\nu$: there exists a positive $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence from $(I - A)$ to $(I - P^{-1}A'P)$ in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$.*
- (3) *For some $\nu \in \text{Iso}[\mathcal{P}, \mathcal{P}']$, with $P = P_\nu$: A and $P^{-1}A'P$ have the same cycle components, and there exists an $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence from $(I - A)$ to $(I - P^{-1}A'P)$ which is positive on cycle components.*

Remarks 3.2.

- (1) There are only finitely many automorphisms $\nu: \mathcal{P} \rightarrow \mathcal{P}'$, and they are easily computed. So, we can decide (3) in Theorem 3.1 if we can decide it in the case where $P = I$ and $\mathcal{P} = \mathcal{P}'$.
- (2) The content of Theorem 3.1 is contained in [H4, H5]. We will prove the equivalence (1) \iff (2) in Section 6. The implication (2) \implies (3) is trivial. The implication (3) \implies (2) follows from the main contribution of this paper, which is the next theorem.

Theorem 3.3 (Factorization Theorem). *Suppose A and A' are in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$, and $(U, V): (I - A) \rightarrow (I - A')$ is an $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence. The following are equivalent:*

- (1) *$(U, V): (I - A) \rightarrow (I - A')$ is a positive $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$.*

- (2) A and A' have the same cycle components, and (U, V) is positive on cycle components.

Below, given a matrix A in any $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, we let A_{∞} denote its embedding in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$.

Theorem 3.4. *Suppose A and A' are in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, where $\mathbf{n} = (n_1, \dots, n_N)$ and the following hold for $1 \leq i \leq N$:*

- $n_i = 1 \Leftrightarrow i$ is a cycle component of $A \Leftrightarrow i$ is a cycle component of A' ,
- $n_i = 1$ or $n_i = \infty$.

Then the following are equivalent:

- (1) *There exists a positive $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence from $(I - A_{\infty})$ to $(I - A'_{\infty})$ in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$.*
- (2) *$(I - A)$ and $(I - A')$ are $\mathrm{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalent.*

Proof of Theorem 3.4. (2) \implies (1) Suppose (U, V) is the $\mathrm{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence. If $n_i = 1$, then $U_{ii} = V_{ii} = 1$ because $\{U, V\} \subset \mathrm{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$. So, the embeddings of U and V in $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ give an $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence $(I - A_{\infty}) \rightarrow (I - A'_{\infty})$ in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ which is positive on cycle components.

(1) \implies (2) This follows from Lemmas A.3 and A.7. □

The point of Theorem 3.4 is to give a flow equivalence criterion in terms of just $\mathrm{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence (which is characterized in [BH]), without a “positive on cycle components” condition. Given matrices A_1 and A'_1 in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, Lemmas A.1 and A.2 give us positive equivalences, from $I - A_1$ to $I - A$ and from $I - A'_1$ to $I - A'$, such that A_{∞} and A'_{∞} are of the form described in Theorem 3.4.

Theorem 3.5 ([BH]). *Suppose B and B' are matrices in $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ such that for each diagonal block in B or B' , the greatest common divisor of the entries of the block is 1. Suppose $\mathbf{n} \leq \mathbf{r}$, and let ι be the embedding of $\mathfrak{M}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ into $\mathfrak{M}_{\mathcal{P}}(\mathbf{r}, \mathbb{Z})$.*

Then B is $\mathrm{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalent to B' if and only if ιB is $\mathrm{SL}_{\mathcal{P}}(\mathbf{r}, \mathbb{Z})$ equivalent to $\iota B'$.

Theorem 3.5, taken from the Stabilization result in [BH], reduces the problem of $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence of the matrices $(I - A)$ and $(I - A')$ in Theorem 3.4 to an equivalence problem for finite matrices.

4. Factorization: The proof.

To begin, we describe a matrix class in which our positivity considerations will be simplified.

Definition 4.1. Given a subset \mathcal{C} of $\{1, \dots, N\}$, and a vector \mathbf{n} with positive integer entries such that $n_i = 1$ if $i \in \mathcal{C}$, define $\mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$ to be the set of \mathbf{n} -blocked integral matrices M whose blocks M_{ij} satisfy the following conditions:

- $M_{ii} = 0$ if $i \in \mathcal{C}$,
- $M_{ij} = 0$ if $i \neq j$ and $i \not\preceq j$,
- $M_{ij} > 0$ otherwise.

(So, each block of M has all entries zero or all entries greater than zero, $M_{ii} = 0$ when $i \in \mathcal{C}$, and otherwise $M_{ij} > 0$ if and only if $i \preceq j$. If $-M = I - A$, then \mathcal{C} is the set of cycle components of A .)

Definition 4.2. An elementary positive equivalence in $\mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$ is an $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence $(U, V): B \rightarrow B' = UBV$ such that $\{B, B'\} \subset \mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$; one of U, V equals Id ; and the other is a basic elementary matrix. A positive equivalence in $\mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$ is a composition of elementary positive equivalences in $\mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$. For such an equivalence (U, V) , we use notations such as

$$(U, V): B \xrightarrow{+} B' \quad \text{or} \quad B \xrightarrow{+ (U, V)} B' \quad \text{or} \quad B \xrightarrow{+} B'.$$

Observation 4.3. Suppose $B = (A - I)$, $B' = (A' - I)$ and

$$(U, V): B \xrightarrow{+} B'.$$

Then $(U, V): (I - A) \rightarrow (I - A')$ is a positive equivalence in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$.

Outline of the proof. Now we can give an outline of the proof of the Factorization Theorem (3.3), which we break into four steps.

Step 1 of the proof (“block positive reduction”) is to reduce it to proving the following theorem:

Theorem 4.4. Suppose $B = A - I$ and $B' = A' - I$, satisfying

- B and B' are in $\mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$
- $(U, V): B \rightarrow B'$ is an $\text{SL}(\mathbf{n}, \mathbb{Z})$ equivalence
- if $i \notin \mathcal{C}$, then $\dim(\ker(A_{ii})) \geq 2$.

Then

$$(U, V): B \xrightarrow{+} B'.$$

Step 2 (“the positive case”) is to prove Theorem 4.4 in the case B and B' are positive (i.e., $\mathcal{P} = \{1\}$ and $\mathcal{C} = \emptyset$). This step is the heart of the proof and it is carried out in Section 5. This is the only step which uses the condition $\dim(\ker(A_{ii})) \geq 2$.

Step 3 (“the unipotent case”) is to prove Theorem 4.4 in the case that U and V lie in $\mathcal{U}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$, where $\mathcal{U}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ denotes the set of matrices M in $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ such that $M_{ii} = I$ for all i in \mathcal{P} .

Step 4 (“the general case”) is to finish the proof of Theorem 4.4.

Step 1: Block positive reduction. We will accomplish this step by proving the following proposition. Let $\mathcal{C}_A = \mathcal{C}$ denote the set of cycle components of A . For each cycle component i , let $\mathcal{C}_i^{\text{sec}}$ denote the set of indices for rows/columns through A_{ii} such that i does not lie on a cycle, and let $\mathcal{C}_i^{\text{prim}}$ denote the set of indices i for rows/columns through A_{ii} which lie on the unique cycle in A_{ii} . Let $\mathcal{C}^{\text{prim}} = \cup \mathcal{C}_i^{\text{prim}}$ and $\mathcal{C}^{\text{sec}} = \cup \mathcal{C}_i^{\text{sec}}$.

Proposition 4.5. *Suppose $\{A, A'\} \subset \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$; $\mathcal{C}_A = \mathcal{C}_{A'} = \mathcal{C}$; $(U, V) : (I - A) \rightarrow (I - A')$ is an $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence which is positive on cycle components; and $\mathbf{n} = (n_1, \dots, n_N)$ has positive integer entries. Then there is a commuting diagram of $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalences*

$$\begin{array}{ccc} (I - A) & \longrightarrow & (I - \bar{A}) \\ (U, V) \downarrow & & \downarrow (\bar{U}, \bar{V}) \\ (I - A') & \longrightarrow & (I - \bar{A}') \end{array}$$

such that:

- (1) *The horizontal arrows are positive $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalences in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$.*
- (2) *For both \bar{A} and \bar{A}' , the principal submatrix indexed by the complement of \mathcal{C}^{sec} is strictly positive wherever the \mathcal{P} ordering permits a nonzero entry, and in addition the diagonal blocks $(\bar{A} - I)_{tt}$ and $(\bar{A}' - I)_{tt}$ are strictly positive whenever $t \notin \mathcal{C}$.*
- (3) *For both \bar{A} and \bar{A}' , $\mathcal{C}^{\text{prim}}$ is the set of indices ℓ such that for some $i \in \mathcal{C}$, (ℓ, ℓ) indexes the upper left corner of the ii block.*
- (4) *$\bar{A}(i, j) = \bar{A}'(i, j) = 0$ whenever $\{i, j\} \cap \mathcal{C}^{\text{sec}} \neq \emptyset$.*
- (5) *$\bar{U}(i, j) = \bar{V}(i, j) = \delta_{ij}$ whenever $\{i, j\} \cap \mathcal{C}^{\text{sec}} \neq \emptyset$.*

For matrices \bar{A}, \bar{A}' in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, we say an $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence $(\bar{U}, \bar{V}) : (I - \bar{A}) \rightarrow (I - \bar{A}')$ is **nondegenerate** if it satisfies Conditions (2), (3), (4) and (5) of Proposition 4.5. Note Condition (3) implies that $\bar{A}(\ell, \ell) = 1$ if $\ell \in \mathcal{C}^{\text{prim}}$.

Let us see that Proposition 4.5 reduces the proof of (2) \implies (1) in the Factorization Theorem 3.3 to the proof of Theorem 4.4. Given $(I - A)$, $(I - A')$ and (U, V) satisfying (2) in the statement of Theorem 3.3, pick a vector \mathbf{n} with positive integer entries large enough that:

- For all i, j in \mathcal{P} , the ij blocks of $U, V, I - A$ and $I - A'$ agree with $\delta_{ij}I$ outside the upper left $n_i \times n_j$ corner, and
- if $i \notin \mathcal{C}$, then the upper left $n_i \times n_i$ corners of A_{ii} and A'_{ii} have kernels of dimension at least two.

Replace A, A', U and V with their truncations to \mathbf{n} -blocked matrices. Then it suffices to prove that $(U, V): (I - A) \rightarrow (I - A')$ is a positive $\text{SL}(\mathbf{n}, \mathbb{Z})$ equivalence in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$. To do this, first apply Proposition 4.5 to the matrices U, V, A, A' . Then truncate the resulting $\overline{U}, \overline{V}, \overline{A}, \overline{A}'$ by removing all rows and columns indexed by \mathcal{C}^{sec} , and call the resulting matrices U, V, A, A' . To finish the proof of the Factorization Theorem 3.3, it suffices to show $(U, V): (I - A) \rightarrow (I - A')$ is a positive equivalence, and this now follows by an application of Theorem 4.4 and Observation 4.3.

We want Proposition 4.5 in order to have a completely general result about factoring equivalences into positive equivalences, and in order to see the main arguments more clearly in the less technical setting of $\mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$. Because the proof of Proposition 4.5 is tedious (almost entirely on account of technicalities involving cycle components), we relegate the proof of Proposition 4.5 to Appendix A.

Below, we use $\mathcal{U}_{\mathcal{P}}$ to denote $\mathcal{U}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ and we use $\mathfrak{M}_{\mathcal{P}}^{++}$ to denote $\mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$. For i, j in \mathcal{P} and B a matrix with a \mathcal{P} -indexed block structure, we let $B\{i, j\}$ denote the ij block of B .

Step 2: The positive case. This is carried out in Section 5.

Step 3: The unipotent case.

Lemma 4.6. *Suppose U and V are matrices in $\mathcal{U}_{\mathcal{P}}$, B and B' are in $\mathfrak{M}_{\mathcal{P}}^{++}$, and $UBV = B'$. Then*

$$B \xrightarrow[+]{(U,V)} B'.$$

Proof. Write U as a product of matrices in $\mathcal{U}_{\mathcal{P}}$, $U = U_n \cdots U_1$, where for each U_t there is an associated pair (i_t, j_t) , such that the following hold:

- $U_t = I$, except in the block $U_t\{i_t, j_t\}$, and
- if $s \neq t$, then $(i_s, j_s) \neq (i_t, j_t)$.

Note, whenever i_s is an immediate predecessor of j_s in \mathcal{P} and $B\{i_s, i_s\} = 0$, these conditions imply

$$(4.7) \quad (U_s B)\{i_s, j_s\} = B'\{i_s, j_s\}.$$

We claim there are nonnegative matrices Q_1, \dots, Q_n in $\mathcal{U}_{\mathcal{P}}$ such that (with $Q = Q_1 \cdots Q_n$)

$$(4.8) \quad B \xrightarrow[+]{(U_1, Q_1)} \cdot \xrightarrow[+]{(U_2, Q_2)} \dots \xrightarrow[+]{(U_n, Q_n)} U_n \cdots U_1 B Q_1 \cdots Q_n = UBQ.$$

To show (4.8), first we will produce Q_1 such that

$$B \xrightarrow[+]{(U_1, Q_1)} U_1 B Q_1.$$

Denote (i_1, j_1) as (i, j) . Factor U_1 as $U_1 = U_1^- U_1^+$, where U_1^- and U_1^+ equal I outside the $\{i, j\}$ block, $U_1^+ \{i, j\}$ is the nonnegative part of $U_1 \{i, j\}$, and $U_1^- \{i, j\}$ is the nonpositive part of $U_1 \{i, j\}$. Clearly

$$(U_1^+, I) : B \xrightarrow{+} U_1^+ B.$$

For U_1^- there are two cases.

Case I: $B \{i, i\} > 0$. We have $U_1^-(U_1^+ B) = U_1^+ B$ outside blocks $\{i, k\}$ such that $i \prec j \preceq k$. Because $(U_1^+ B) \{i, i\} = B \{i, i\} > 0$, we can pick Q_1 in $\mathcal{U}_{\mathcal{P}}$, with sufficiently large nonnegative entries in such blocks $\{i, k\}$, to put $U_1^-(U_1^+ B)Q_1$ into $\mathfrak{M}_{\mathcal{P}}^{++}$. Then

$$U_1^+ B \xrightarrow{+ (I, Q_1)} U_1^+ B Q_1 \xrightarrow{+ (U_1^-, I)} U_1^- U_1^+ B Q_1 = U_1 B Q_1.$$

Case II: $B \{i, i\} = 0$. Again, $U_1^-(U_1^+ B) = U_1^+ B$ outside blocks $\{i, k\}$ such that $i \prec j \preceq k$. Because $B \{i, j\} > 0$, we can choose Q_1 nonnegative in $\mathcal{U}_{\mathcal{P}}$ such that for all k satisfying $i \prec j \prec k$, we have $U_1^-(U_1^+ B)Q_1 \{i, k\} > 0$. (A positive entry in the block $Q \{j, k\}$ acts here to add a multiple of a column through the $\{i, j\}$ block to a column in the $\{i, k\}$ block.) If there is some h such that $i \prec h \prec j$, then suitable positive entries in $Q_1 \{h, k\}$ will also achieve $U_1^-(U_1^+ B)Q_1 \{i, j\} > 0$. If there is no such h , then i is an immediate predecessor of j in \mathcal{P} , and by appeal to (4.7) we have

$$\begin{aligned} (U_1^-(U_1^+ B)Q_1) \{i, j\} &= (U_1 B) \{i, j\} \\ &= B' \{i, j\} > 0. \end{aligned}$$

Therefore

$$U_1^+ B \xrightarrow{+ (I, Q_1)} U_1^+ B Q_1 \xrightarrow{+ (U_1^-, I)} U_1 B Q_1$$

as required.

Thus in either case we have

$$B \xrightarrow{+ (U_1, Q_1)} U_1 B Q_1 \in \mathfrak{M}_{\mathcal{P}}^{++}.$$

An easy induction on the argument gives (4.8), with

$$B \xrightarrow{+ (U, Q)} U B Q \in \mathfrak{M}_{\mathcal{P}}^{++},$$

with Q a product of nonnegative elementary matrices in $\mathcal{U}_{\mathcal{P}}$. The transposed argument gives a matrix P in $\mathcal{U}_{\mathcal{P}}$ such that P is a product of nonnegative elementary matrices such that

$$B' \xrightarrow{+ (P, V^{-1})} P B' V^{-1} \in \mathfrak{M}_{\mathcal{P}}^{++}.$$

Then

$$B \xrightarrow[+]{(U,Q)} UBQ \xrightarrow[+]{(P,I)} PUBQ = PB'V^{-1}Q \xleftarrow[+]{(I,Q)} PB'V^{-1} \xleftarrow[+]{(P,V^{-1})} B'$$

so

$$(P^{-1}PU, QQ^{-1}V) = (U, V): B \xrightarrow[+]{} B'$$

as required. □

Step 4: The general case.

Lemma 4.9. *Suppose $i \notin \mathcal{C}$, E is a basic elementary matrix in $\mathrm{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$, $E\{j, k\} = (\mathrm{Id})\{j, k\}$ when $(j, k) \neq (i, i)$, $\{B, B'\} \subset \mathfrak{M}_{\mathcal{P}}^{++}(\mathcal{C}, \mathbf{n}, \mathbb{Z})$ and*

$$(E\{i, i\}, \mathrm{Id}): B\{i, i\} \xrightarrow[+]{} B'\{i, i\}.$$

Then there exists V in $\mathcal{U}_{\mathcal{P}}$ such that

$$(E, V): B \xrightarrow[+]{} B'.$$

Similarly, if

$$(\mathrm{Id}, E\{i, i\}): B\{i, i\} \xrightarrow[+]{} B'\{i, i\}$$

then there exists U in $\mathcal{U}_{\mathcal{P}}$ such that

$$(U, E): B \xrightarrow[+]{} B'.$$

Proof. We will consider the equivalence (E, I) , the other case is similar. Let $E(s, t)$ be the nonzero offdiagonal entry of E . If $E(s, t) = 1$, then set $V = \mathrm{Id}$. Now suppose $E(s, t) = -1$, so E acts from the the left to subtract row t from row s . Then possibly there are nonpositive entries in blocks $(EB)\{i, j\}$ where $i \prec j$. To correct for this, pick r an index for a column through the ii block; note that $B(s, r) > B(t, r)$ because $(EB)\{i, i\} > 0$ by assumption; consider a positive integer M ; and let V be the matrix in $\mathcal{U}_{\mathcal{P}}$ which acts from the right to add column r to column q , M times, for every q indexing a column through an ij block for which $i \prec j$. For these q ,

$$(EBV)(s, q) = M(B(s, r) - B(t, r)) + B(s, q) - B(t, q).$$

So, if M is large enough, then this gives

$$B \xrightarrow[+]{(I,V)} BV \xrightarrow[+]{(E,I)} EBV$$

as required. □

Proof of the general case. Now let $(U, V): B \rightarrow B'$ be the $\mathrm{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence, with $\{B, B'\} \subset \mathfrak{M}_{\mathcal{P}}^+(\mathcal{C}, \mathbf{n}, \mathbb{Z})$. By Step 2 (Theorem 5.1), for each $i \in \mathcal{P} \setminus \mathcal{C}$, we have

$$(U\{i, i\}, V\{i, i\}): B\{i, i\} \xrightarrow{+} B'\{i, i\}.$$

So, we may find a string of elementary equivalences, say $(E_1, F_1), \dots, (E_t, F_t)$, with every $E_t\{i, j\} = F_t\{i, j\} = (\mathrm{Id})\{i, j\}$ unless $i = j \notin \mathcal{C}$, which accomplishes the elementary positive equivalence decomposition inside the diagonal blocks. By Lemma 4.9, we may find $(U_1, V_1), \dots, (U_t, V_t)$ with each U_s and V_s in $\mathcal{U}_{\mathcal{P}}$, such that

$$B \xrightarrow{+ (U_1, F_1)} \dots \xrightarrow{+ (E_1, V_1)} \dots \xrightarrow{+ (U_t, F_t)} \dots \xrightarrow{+ (E_t, V_t)} B''.$$

Let $X = E_t U_t \cdots E_2 U_2 E_1 U_1$. Let $Y = F_1 V_1 F_2 V_2 \cdots F_t V_t$. Then for all i in \mathcal{P} , $X\{i, i\} = U\{i, i\}$ and $Y\{i, i\} = V\{i, i\}$, so $UX^{-1} \in \mathcal{U}_{\mathcal{P}}$ and $Y^{-1}V \in \mathcal{U}_{\mathcal{P}}$. It follows from Step 3 (Lemma 4.6) that

$$B'' \xrightarrow{+ (UX^{-1}, Y^{-1}V)} B'.$$

Thus $(U, V): B \rightarrow B'$ is the composition

$$B \xrightarrow{+ (X, Y)} B'' \xrightarrow{+ (UX^{-1}, Y^{-1}V)} B'$$

and this finishes the proof. \square

5. Factorization: The positive case.

In this section, all matrices are $K \times K$, where K is a positive integer and $K > 1$. We let \mathfrak{M}_+ denote the set of $K \times K$ matrices with strictly positive integer entries.

We say an equivalence $(U, V): B \rightarrow B'$ is a positive equivalence through \mathfrak{M}_+ if it can be given as a chain of positive elementary equivalences

$$B = B_0 \rightarrow B_1 \rightarrow B_2 \rightarrow \cdots \rightarrow B_n = B'$$

in which every B_i is in \mathfrak{M}_+ .

The purpose of this section is to prove the following theorem.

Theorem 5.1. *Suppose U and V are in $\mathrm{SL}(K, \mathbb{Z})$, and B and UBV are in \mathfrak{M}_+ . Suppose also that B is $\mathrm{SL}(K, \mathbb{Z})$ equivalent to a diagonal matrix in which at least two entries equal 1.*

Then $(U, V): B \rightarrow UBV$ is a positive equivalence through \mathfrak{M}_+ .

Remark 5.2. The “two entries” technical assumption may be excessive, but is harmless for our applications. Except for the final argument which addresses the possibility that UB is nonpositive, we only use the weaker assumption that B has rank greater than one.

The proof of Theorem 5.1 rests on three lemmas. We begin the preparations.

By a *signed transposition matrix*, we mean a matrix which is the matrix of a transposition, but with one of the off-diagonal 1's replaced by -1. By a *signed permutation matrix* we mean a product of signed transposition matrices. Since $K > 1$, any $K \times K$ permutation matrix with determinant 1 is a signed permutation matrix. A $K \times K$ matrix S is a signed permutation matrix if and only if $\det S = 1$ and the matrix $|S|$ is a permutation matrix (where $|S|(i, j) := |S(i, j)|$).

Lemma 5.3. *Suppose $B \in \mathfrak{M}_+$, E is a basic elementary matrix with non-zero offdiagonal entry $E(i, j)$, and the i th row of EB is not the zero row.*

Then in $SL(K, \mathbb{Z})$ there are a nonnegative matrix Q and a signed permutation matrix S such that $(SE, Q): B \rightarrow SEBQ$ is a positive equivalence through \mathfrak{M}_+ .

Proof. If $E(i, j) = 1$, then let $Q = I = S$. Now suppose $E(i, j) = -1$, so E acts from the left by subtracting row j from row i , and the rows i and j of B are not equal.

Case I: For some k , $B(i, k) > B(j, k)$.

Here we may repeatedly add column k of B to other columns, until we have a matrix B' with $B'(i, m) > B'(j, m)$ for all m . This B' is BQ for some Q which is a product of nonnegative basic elementary matrices. Now $(E, Q): B \rightarrow EBQ$ is the composition of positive equivalences, $(I, Q): B \rightarrow BQ$ followed by $(E, I): BQ \rightarrow EBQ$. Let $S = I$.

Case II: For every k , $B(i, k) \leq B(j, k)$.

Because the rows i and j of B are not equal, after multiplying from the right by a suitable Q we can assume in this Case that $0 < B(i, k) < B(j, k)$ for all k . Now $(I, Q): B \rightarrow BQ$ in \mathfrak{M}_+ , so for notational simplicity from here we may assume $Q = I$.

For concreteness of notation, let $(i, j) = (1, 2)$. For the rest of this Case, for simplicity we will restrict what we write to these two rows, e.g.,

$$E = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} B_1 \\ B_2 \end{pmatrix},$$

where B_1 and B_2 denote the first and second rows of B , and we have $B_1 < B_2$. Let $S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Then

$$(SE)B = \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} B_1 \\ B_2 \end{pmatrix} = \begin{pmatrix} B_2 \\ B_2 - B_1 \end{pmatrix}$$

and the latter matrix is positive. Let $E' = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and let $E'' = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$, then

$$SE = \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix} = E'E''.$$

Now $(E'', I): B \rightarrow E''B$ is a positive equivalence in \mathfrak{M}_+ , since row 2 of B is positive and greater than row 1; and $(E', I): E''B \rightarrow E'E''B$ is also a positive equivalence in \mathfrak{M}_+ . \square

Lemma 5.4. *Suppose B is a $K \times K$ integral matrix of rank at least 2, and U is in $SL(K, \mathbb{Z})$, and no row of B or UB is the zero row. Then U is the product of basic elementary matrices, $U = E_k \cdots E_1$, such that for $1 \leq j \leq k$ the matrix $E_j E_{j-1} \cdots E_1 B$ has no zero row.*

Proof. Without loss of generality, assume $K \geq 3$ and U is not the identity. Let $\mathcal{E}(i)$ denote the set of integral matrices which equal I both on the diagonal and outside of row i . Let \mathcal{U} be the set of factorizations $U = U_n \cdots U_1$ such that for $1 \leq h \leq n$, the matrix U_h is not the identity and there is an index i_h such that $U_h \in \mathcal{E}(i_h)$. Given such a factorization $U = U_n \cdots U_1$, let

$$z = \#\{h : 1 \leq h \leq n \text{ and row } i_h \text{ of } U_h \cdots U_1 B \text{ is the zero row}\}.$$

Step 1. We will produce an element of \mathcal{U} for which $z = 0$.

By induction, it suffices to begin with a factorization $U = U_n \cdots U_1$ from \mathcal{U} for which $z > 0$, and produce another factorization from \mathcal{U} with reduced z . Pick s minimal such that row i_s of $U_s \cdots U_1 B$ is zero, and let t be minimal such that $t > s$ and $i_t = i_s$. (This t exists because row i_s of UB is nonzero.) We will change the factorization by replacing the subword $U_t \cdots U_s$ with a suitable word $U'_t \cdots U'_s$, to be defined recursively.

First pick $j_s \neq i_s$ such that row j_s of $U_{s-1} \cdots U_1 B$ is nonzero ($U_{s-1} \cdots U_1 B$ just denotes B in the case that $s = 1$). Choose F_s an elementary matrix which acts to add a multiple of row j_s to row i_s , such that (for notational simplicity) $F_s^{-1}U_s \neq I$. Define $U'_s = F_s^{-1}U_s \in \mathcal{E}(i_s)$. Now $U_t \cdots U_s = U_t \cdots U_{s+1}F_s U'_s$ and row i_s of $U'_s U_{s-1} \cdots U_1 B$ is not zero.

Now we give the recursive step. Suppose $s < m \leq t$ and we have produced $U_t \cdots F_{m-1}U'_r \cdots U'_s = U_t \cdots U_s$ such that there is a nonzero integer c_{m-1} and an index $j_{m-1} \neq i_s$ such that $F_{m-1}(i_s, j_{m-1}) = c_{m-1}$ and otherwise $F_{m-1} = I$. We will replace $U_m F_{m-1}$ with new terms. There are three cases.

Case 1: $m < t$ and $j_{m-1} \neq i_m$. Set $F_m = F_{m-1}$ and $U'_{r+1} = F_m^{-1}U_m F_m$. For example, if $K = 3$ and $(i_s, i_m, j_{m-1}) = (1, 2, 3)$, then we would have for

some a, b, c that

$$\begin{aligned} U'_{r+1} = F_m^{-1}U_mF_m &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -c & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ a & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ c & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -c & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ a+bc & 1 & b \\ c & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ a+bc & 1 & b \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

Now $U'_{r+1} \in \mathcal{E}(i_m)$ and $F_mU'_{r+1} = U_mF_{m-1}$ and row i_m of $U'_{r+1}U'_r \cdots U'_sU_{s-1} \cdots U_1B$ equals row i_m of $U_m \cdots U_1B$.

Case 2: $m < t$ and $j_{m-1} = i_m$. Choose an index j_m such that $j_m \notin \{i_m, i_s\}$ and row j_m of $U'_r \cdots U'_sU_{s-1} \cdots U_1B$ is not zero. This is possible because rows i_s and j_{m-1} of $U'_r \cdots U'_sU_{s-1} \cdots U_1B$ are linearly dependent (since row i_s of $F_mU'_r \cdots U'_sU_{s-1} \cdots U_1B$ equals row i_s of $U_m \cdots U_1B$ which is the zero row) and $\text{rank}(B) \geq 2$. Pick F_m with $F_m(i_s, j_m) = 1$ and otherwise $F_m = I$. Set $U'_{r+1} = F_m^{-1}F_{m-1}$ and $U'_{r+2} = F_m^{-1}U_mF_m$. Now

- $F_mU'_{r+2}U'_{r+1} = F_m(F_m^{-1}U_mF_m)(F_m^{-1}F_{m-1}) = U_mF_{m-1}$,
- $U'_{r+1} \in \mathcal{E}(i_s)$ and row i_s of $U'_{r+1} \cdots U'_sU_{s-1} \cdots U_1B$ is not zero,
- $U'_{r+2} \in \mathcal{E}(i_m)$ and row i_m of $U'_{r+2} \cdots U'_sU_{s-1} \cdots U_1B$ equals row i_m of $U_m \cdots U_1B$.

Case 3: $m = t$. If $U_tF_{t-1} \neq I$, then set $U'_T = U'_{r+1} = U_tF_{t-1} \in \mathcal{E}(i_s)$: Row i_s is the same in the matrices $U_m \cdots U_1B$ and $U'_T \cdots U'_sU_{s-1} \cdots U_1B$. If $U_tF_{t-1} = I$, then simply delete U_tF_{t-1} , so $U'_T = U'_r$.

The new factorization has z reduced. This concludes Step 1.

Step 2. Suppose we have the factorization from \mathcal{U} with $z = 0$, $U = U_n \cdots U_1$, with $U_h \in \mathcal{E}(i_h)$. For $1 \leq h \leq n$, we will replace U_h with a suitable product of elementary matrices in $\mathcal{E}(i_h)$. The argument will be clear from the case $h = 1$. For notational simplicity, suppose $i_1 = 1$. Write U_1 as a product $U_1 = E_k \cdots E_1$ of basic elementary matrices which agree with I outside row 1. Now, choose a row $m > 1$ of B which is not a rational multiple of row 1 of U_1B (such a row m exists because $\text{rank}(B) > 1$). Let E_0 be the elementary matrix which adds row m to row 1: If $s > 0$, then $(E_0)^sB$ has row 1 nonzero. Choose a nonnegative integer M large enough that for $1 \leq j \leq k$, row 1 of $[E_j \cdots E_1(E_0)^M]B$ is nonzero. Then for $0 \leq s \leq M$,

$$\begin{aligned} [E_0^{-s}][E_k \cdots E_1(E_0)^M]B &= [E_0^{M-s}][E_k \cdots E_1]B \\ &= [E_0^{M-s}]U_1B \end{aligned}$$

and therefore row 1 of $[E_0^{-s}][E_k \cdots E_1(E_0)^M]B$ cannot be zero. Thus the factorization $U_1 = (E_0)^{-M}E_k \cdots E_1(E_0)^M$ has the required properties. \square

Lemma 5.5 (Key Lemma). *Suppose B and B' are in \mathfrak{M}_+ , U and W are in $\text{SL}(K, \mathbb{Z})$, the matrix UB has at least one strictly positive entry, and $UB = B'W$. Then the equivalence $(U, W^{-1}): B \rightarrow B'$ is a positive equivalence through \mathfrak{M}_+ .*

Proof. Step 1: Reduction to the case $UB > 0$.

Consider an entry $(UB)(i, j) > 0$. We can repeatedly add column j to other columns until row i of UB has all entries strictly positive. This corresponds to multiplying from the left by a nonnegative matrix Q in $\text{SL}(K, \mathbb{Z})$, giving $UBQ = B'WQ$. Then we can repeatedly add row i of UBQ to other rows until all entries of UBQ are positive. This corresponds to multiplying from the left by a matrix P in $\text{SL}(K, \mathbb{Z})$, giving

$$(PU)(BQ) = (PB')(WQ) > 0$$

with positive equivalences in \mathfrak{M}_+ given by

$$(I, Q): B \rightarrow BQ, \quad (P, I): B' \rightarrow PB'$$

Therefore, after replacing (U, B, B', W) with (PU, BQ, PB', WQ) , we may assume without loss of generality that $UB > 0$.

Step 2: Reducing the length of an elementary factorization.

By Lemma 5.4, we can write U has a product of basic elementary matrices, $U = E_k \cdots E_1$, such that for $1 \leq j \leq k$, the matrix $B_j = E_j \cdots E_1 B$ has no zero row. By Lemma 5.3, given the pair (E_1, B) , there is a nonnegative Q_1 in $\text{SL}(K, \mathbb{Z})$ and a signed permutation matrix S_1 such that

$$(S_1 E_1, Q_1): B \rightarrow S_1 E_1 B Q_1$$

is a positive equivalence in \mathfrak{M}_+ . We observe that

$$UBQ_1 = S_1^{-1} [S_1 E_k S_1^{-1}] \cdots [S_1 E_2 S_1^{-1}] [S_1 E_1] B Q_1.$$

Now, for $2 \leq j \leq k$, the matrix $S_1 E_j S_1^{-1}$ is again a basic elementary matrix E'_j , and the matrix $E'_j \cdots E'_2 (S_1 E_1 B Q_1)$ has no zero rows.

Again using Lemma 5.3, for the pair $([S_1 E_2 S_1^{-1}], [S_1 E_1 B Q_1])$ choose a signed permutation matrix S_2 and nonnegative Q_2 producing a positive equivalence in \mathfrak{M}_+

$$(S_2 [S_1 E_2 S_1^{-1}], Q_2): S_1 E_1 B Q_1 \rightarrow S_2 [S_1 E_2 S_1^{-1}] S_1 E_1 B Q_1 Q_2$$

so that we get a positive equivalence in \mathfrak{M}_+

$$([S_2 S_1 E_2 S_1^{-1}] [S_1 E_1], Q_1 Q_2): B \rightarrow [S_2 S_1 E_2 S_1^{-1}] [S_1 E_1] B Q_1 Q_2$$

and we observe that

$$UBQ_1 Q_2 = S_1^{-1} S_2^{-1} [S_2 S_1 E_k S_1^{-1} S_2^{-1}] \cdots [S_2 S_1 E_3 S_1^{-1} S_2^{-1}] [S_2 S_1 E_2 S_1^{-1}] [S_1 E_1] B Q_1 Q_2.$$

Continue this, to obtain a signed permutation matrix $S = S_k \cdots S_1$ and nonnegative $Q = Q_1 \cdots Q_k$ such that

$$\begin{aligned} UBQ &= S^{-1}[S_k \cdots S_1 E_k S_1^{-1} \cdots S_{k-1}^{-1}] \cdots [S_2 S_1 E_2 S_1^{-1}][S_1 E_1]BQ \\ &= S^{-1}(SUBQ) \end{aligned}$$

and $(SU, Q): B \rightarrow SUBQ$ is a positive equivalence in \mathfrak{M}_+ .

Step 3: Realizing the permutation.

We continue from Step 2. It remains to show that

$$(S, I): UBQ \rightarrow SUBQ$$

is a positive equivalence in \mathfrak{M}_+ . Since S is a product of signed transposition matrices, it may be described as a permutation matrix in which some rows have been multiplied by -1 . Since UBQ and $SUBQ$ are strictly positive, it must be that S is a permutation matrix. Also, $\det(S) = 1$, so if $S \neq I$ then S is the matrix of a permutation which is a product of 3-cycles. So it is enough to realize the positive equivalence in \mathfrak{M}_+ in the case that S is the matrix of a 3-cycle. For this we write the matrix

$$C = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

as the following product $C_0 C_1 \cdots C_5$:

$$\begin{aligned} &\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &\quad \cdot \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

For $0 \leq i \leq 5$, the matrix $C_i C_{i+1} \cdots C_5$ is nonnegative. Therefore the equivalence $(C, I): B \rightarrow CB$ is a positive equivalence through \mathfrak{M}_+ whenever $B \in \mathfrak{M}_+$. □

We can now complete the proof of Theorem 5.1. It only remains to address the technical point that in the equivalence $(U, V): B \rightarrow B'$, all the entries of UB might be nonpositive. (For example, with K even we could have $(U, V) = (-I, -I)$.)

Proof of Theorem 5.1. By assumption there are X, Y in $SL(K, \mathbb{Z})$ such that $XYB = D$, where D is diagonal and has the block form $D = \begin{pmatrix} I & 0 \\ 0 & F \end{pmatrix}$, where I is 2×2 . For any H in $SL(2, \mathbb{Z})$, the $K \times K$ matrix $G = G_H = \begin{pmatrix} H & 0 \\ 0 & I \end{pmatrix}$ yields a self equivalence $(X^{-1}GX, YG^{-1}Y^{-1}): B \rightarrow B$.

For a matrix Q , we let $Q\{12;*\}$ denote the submatrix consisting of the first two rows. The matrix $(XBY)\{12;*\} = D\{12;*\}$ has rank two, so the matrix $(XB)\{12;*\}$ has rank two, and we may choose $H' \in \text{SL}(2, \mathbb{Z})$ such that the first row r of $H'[(XB)\{12;*\}]$ has both a positive entry and a negative entry. For $M \in \mathbb{N}$, let $H_M = \begin{pmatrix} M & -1 \\ 1 & 0 \end{pmatrix}$, $H = H_M H'$, and $G = G_H$. Let c denote the first column of X^{-1} . Since c is not the zero vector, the $K \times K$ matrix cr has a positive entry and a negative entry.

If M is sufficiently large, then the entries of the two matrices $X^{-1}GXB$ and Mcr will have the same sign wherever the entries of Mcr are nonzero, and $X^{-1}GXB$ will have a positive entry. Then the Key Lemma 5.5 shows that $(X^{-1}GX, YG^{-1}Y^{-1})$ gives a positive equivalence in \mathfrak{M}_+ from B to B .

Similarly, for large enough M the entries of $UX^{-1}GXB$ will agree in sign with the entries of $UMcr$ wherever the entries of the latter matrix are nonzero. Because U is nonsingular, the matrix Ucr is nonzero, and then contains positive and negative entries because r does.

So, using M sufficiently large, we obtain $(U, V): B \rightarrow B'$ as a positive equivalence in \mathfrak{M}_+ , the inverse of $(X^{-1}GX, YG^{-1}Y^{-1})$ followed by $(UX^{-1}GX, YG^{-1}Y^{-1}V)$. \square

6. Flow equivalence.

The purpose of this section is to prove the claims of Theorem 3.1 involving flow equivalence. As sketched in [B1] (see also [Ba1]), the positive K -theory framework is most natural for this. Because a complete development of this connection has not yet appeared, for brevity we will make no direct use of it below.

We begin with some background. For S a selfhomeomorphism of a compact metric space X , the mapping torus Y_S of S is the quotient space $(X \times \mathbb{R}) / \sim$ where $(x, n + t) \sim (S^n x, t)$ if $n \in \mathbb{Z}$. Y_S admits a natural flow,

$$Y_S \times \mathbb{R} \rightarrow Y_S$$

$$([(x, t)], s) \mapsto [(x, s + t)].$$

This flow has the copy $X_0 = \{[(x, 0)]: x \in X\}$ of X as a cross section, and the return map to X_0 under the flow (given by $[(x, 0)] \mapsto [(Sx, 0)]$) is obviously topologically conjugate to S . Let T be another selfhomeomorphism of a compact metric space. Then S and T are *flow equivalent* if and only if there is a homeomorphism $Y_S \rightarrow Y_T$ which takes flow lines onto flow lines and respects the direction of the associated flows. (Equivalently: S and T are conjugate to return maps of cross sections of a common flow.)

For example, consider $S = \sigma_A$, $T = \sigma_{A'}$ and the map γ arising from a basic positive equivalence in Subsection 2.2. It is not difficult to see that γ

is the restriction of a homeomorphism $Y_S \rightarrow Y_T$ which takes flow lines onto flow lines and respects the direction of the associated flows, and therefore σ_A and $\sigma_{A'}$ are flow equivalent.

Now fix A in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ and A' in $\mathfrak{M}_{\mathcal{P}',+}^{\circ}(\mathbb{Z})$. Let F and F' be finite matrices such that $F_{\infty} = A$ and $F'_{\infty} = A'$. Let $\sigma_A = \sigma_F$ and $\sigma_{A'} = \sigma_{F'}$ be the associated SFTs. (So, for example σ_A is the left shift on the path space Σ_A , which is given the natural zero dimensional metrizable topology.) Parry and Sullivan [PS] showed that σ_F and $\sigma_{F'}$ are flow equivalent if and only if σ_F is topologically conjugate to some SFT which after a time change is topologically conjugate to $\sigma_{F'}$. It follows ([PS]) that σ_F and $\sigma_{F'}$ are flow equivalent if and only if F' can be obtained from F by a finite sequence of basic flow moves, which are state splittings and stretchings and their inverses. The inverse of a splitting is called an amalgamation. We will describe the splitting and stretching moves now.

Let B and B' be finite square matrices. B' is obtained from B by an elementary row amalgamation if there exist indices i_1, i_2 and i such that the columns i_1 and i_2 of A' are equal, and A is obtained from A' as follows: Add row i_1 to row i_2 , then remove the row and column indexed by i_1 . The reverse move is that B is obtained from B' by a row splitting. Analogously there are column splittings and amalgamations. By state splittings we mean row splittings and column splittings.

We say B' is obtained from B by a state stretching if for some indices i, j the following hold: $B'(i, j) = 1$, the other entries of row i and column j are zero, and B is the matrix obtained from B' by adding column i to column j and then removing row i and column i .

We are now ready for the proof. Suppose A is in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ and A' is in $\mathfrak{M}_{\mathcal{P}',+}^{\circ}(\mathbb{Z})$. We will show the following are equivalent:

- (1) σ_A and $\sigma_{A'}$ are flow equivalent.
- (2) There exists $\nu \in \text{Iso}[\mathcal{P}, \mathcal{P}']$ such that for $P = P_{\nu}$, there exists a positive $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence from $(I - A)$ to $(I - P^{-1}A'P)$ in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$.

Proof. Given (2), it follows from Lemma 2.5 that there is a chain of basic positive $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalences from $(I - A)$ to $(I - P^{-1}A'P)$. Each basic positive equivalence gives rise to a flow equivalence as discussed above. It follows that (2) implies (1).

Now we assume (1) and will deduce (2). Let F and F' be finite matrices such that $F_{\infty} = A$ and $F'_{\infty} = A'$. After using Lemmas A.1 and A.2 to pass to flow equivalent SFTs, we may assume that for each $i \in \mathcal{P}$, the diagonal blocks F_{ii} and F'_{ii} are strictly positive.

From [PS] we are given a sequence of basic moves through finite matrices, $F = F_0 \rightarrow F_1 \rightarrow \dots \rightarrow F_m = F'$. We may regard \mathcal{P} and \mathcal{P}' as the posets of irreducible components of F and F' respectively, where e.g., $i \preceq j$ in \mathcal{P} when there exists a transition from i to j (by which we mean that there exists a

point in the SFT Σ_A forwardly asymptotic to a cycle from component j and backwardly asymptotic to a cycle from component i). Each move $F_i \rightarrow F_{i+1}$ induces a bijection of irreducible components, respecting transitions, and thus the composition induces a poset isomorphism $\nu: \mathcal{P}(A) \rightarrow \mathcal{P}(A')$. After replacing A' with $P^{-1}A'P$, where $P = P_\nu$, we may assume $\mathcal{P} = \mathcal{P}'$ and $\nu = \text{Id}$.

Next, for $1 \leq i \leq n$, we will associate to F_i a matrix A_i in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ such that (modulo permutations of indices) $\text{tru}(A_i) = F_i$. We must take a little care with the indices, to be able to lift each of the moves $F_i \rightarrow F_{i+1}$ to a (positive) $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence $(I - A_i) \rightarrow (I - A_{i+1})$. Let $\text{Ind}(B)$ denote the set indexing the rows and columns of a square matrix B . For each F_j , we will define an injection $\tau_j: \text{Ind}(F_j) \rightarrow \mathcal{I}^{\mathcal{P}}$, and then define $A_j = \iota(F_j)$ by setting

$$A_j(s, t) = \begin{cases} F(s', t') & \text{if } (s, t) = (\tau_j(s'), \tau_j(t')) \\ = 0 & \text{otherwise.} \end{cases}$$

The maps τ_j will be defined recursively. For $j = 0$, we set $A_0 = A$ and take τ_j to be compatible with the embedding of F as a principal submatrix of A . Now suppose $0 \leq j < n$ and τ_j and A_j are given. The transition $F_j \rightarrow F_{j+1}$ is given by a basic flow move, and under such a move, every element of $\text{Ind}(F_{j+1})$ is naturally related to one or two elements of $\text{Ind}(F_j)$. (An element i of $\text{Ind}(F_{j+1})$ is related to two elements i_1, i_2 of $\text{Ind}(F_j)$ when the move $F_{j+1} \rightarrow F_j$ is a splitting or stretching of the state i into the states i_1, i_2 .) In any case, for each i in $\text{Ind}(F_{j+1})$, fix a related vertex $\text{rel}(i)$ in $\text{Ind}(F_j)$. Then choose any map $\tau_{j+1}: \text{Ind}(F_{j+1}) \rightarrow \mathcal{I}^{\mathcal{P}}$ such that $\tau_j(\text{rel}(i))$ and $\tau_{j+1}(i)$ lie in $\mathcal{I}_p^{\mathcal{P}}$ for the same element p of \mathcal{P} . (When i is related to two indices, this p may depend on the choice for $\text{rel}(i)$.) This defines the matrices $A = A_0, A_1, \dots, A_m$.

Next we will show that each elementary flow move $F_j \rightarrow F_{j+1}$ gives rise to a positive $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence $(I - A_j) \rightarrow (I - A_{j+1})$. Each of the equivalences we give will be accomplished by elementary matrices which must lie in $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ on account of our choices of indices.

First we show how an elementary row splitting gives rise to a positive $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence. The general construction can be understood from the example

$$B = \begin{pmatrix} a & b & 0 \\ c_1 + c_2 & d_1 + d_2 & 0 \\ 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} a & b & b \\ c_1 & d_1 & d_1 \\ c_2 & d_2 & d_2 \end{pmatrix} = B'.$$

Here the positive equivalence $(I - B) \rightarrow (I - B')$ is accomplished as follows:

$$\begin{aligned}
 (I - B) &= \begin{pmatrix} 1 - a & -b & 0 \\ -(c_1 + c_2) & 1 - (d_1 + d_2) & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
 &\rightarrow \begin{pmatrix} 1 - a & -b & 0 \\ -(c_1 + c_2) & 1 - (d_1 + d_2) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -c_2 & -d_2 & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1 - a & -b & 0 \\ -(c_1 + c_2) & 1 - (d_1 + d_2) & 0 \\ -c_2 & -d_2 & 1 \end{pmatrix} \\
 &\rightarrow \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 - a & -b & 0 \\ -(c_1 + c_2) & 1 - (d_1 + d_2) & 0 \\ -c_2 & -d_2 & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1 - a & -b & 0 \\ -c_1 & 1 - d_1 & -1 \\ -c_2 & -d_2 & 1 \end{pmatrix} \\
 &\rightarrow \begin{pmatrix} 1 - a & -b & 0 \\ -c_1 & 1 - d_1 & -1 \\ -c_2 & -d_2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1 - a & -b & -b \\ -c_1 & 1 - d_1 & -d_1 \\ -c_2 & -d_2 & 1 - d_2 \end{pmatrix}.
 \end{aligned}$$

The positive equivalence for a column splitting is constructed similarly.

Next we show that a state stretching gives rise to a positive equivalence. The general construction can be understood from the example

$$B = \begin{pmatrix} 0 & 0 & 0 \\ 0 & a & b \\ 0 & c & d \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 1 & 0 \\ a & 0 & b \\ c & 0 & d \end{pmatrix} = B'.$$

Here the positive equivalence $(I - B) \rightarrow (I - B')$ is accomplished as follows.

$$\begin{aligned}
 (I - B) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 - a & -b \\ 0 & -c & 1 - d \end{pmatrix} \\
 &\rightarrow \begin{pmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ -c & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 - a & -b \\ 0 & -c & 1 - d \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -a & 1 - a & -b \\ -c & -c & 1 - d \end{pmatrix} \\
 &\rightarrow \begin{pmatrix} 1 & 0 & 0 \\ -a & 1 - a & -b \\ -c & -c & 1 - d \end{pmatrix} \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -1 & 0 \\ -a & 1 & -b \\ -c & 0 & 1 - d \end{pmatrix}.
 \end{aligned}$$

At this point we have a positive $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence $(I - A) \rightarrow (I - A'_n)$, where there is a permutation matrix Q such that $Q^{-1}A'_nQ = A'$. Because $A' = F'_\infty$ and F' has all diagonal blocks positive, if $A'(i, j) > 0$ then i lies on an A' cycle and j lies on an A' cycle. Therefore the permutation given by Q can be chosen compatible with the poset isomorphism $\nu = \mathrm{Id}$, and the matrix Q is a block diagonal matrix in $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$. It remains to check that $I - A'_n \rightarrow Q^{-1}A'_nQ$ is accomplished by a positive $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence. Because Q is block diagonal and we can use compositions, it is enough to give the equivalence in the case that Q is the transposition matrix for indices i, j which lie in some \mathcal{I}_p^P . Choose indices α, β in \mathcal{I}_p^P such that A'_n is identically zero in the rows and columns indexed by α and β . Let P be the permutation matrix for the product of transpositions $(i, j)(\alpha, \beta)$. Then P is in $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ and $PA'_nP = Q^{-1}A'_nQ$. This finishes the proof. \square

7. The mapping class group.

In this section the symbols S, T denote subshifts (e.g., T is the restriction of some full shift $\sigma_{[n]}$ to a closed shift-invariant subset, which we also call T). As in Section 6, let Y_S denote the mapping torus of S . We regard Y_S as an oriented space, in the sense that the associated flow gives an orientation to each of its orbits (i.e., to each connected component of Y_S). A *flow equivalence* from a subshift S to a subshift T is an orientation preserving homeomorphism $\varphi: Y_S \rightarrow Y_T$ (where “orientation preserving” means orientation preserving on each orbit). Two such homeomorphisms φ_0, φ_1 are *isotopic* ($\varphi_0 \sim \varphi_1$) if there is a continuous map $t \mapsto \varphi_t, 0 \leq t \leq 1$, which connects them in the metrizable space of homeomorphisms from Y_S to Y_T . Let $\mathrm{Is}(S, T)$ denote the set of isotopy classes of flow equivalences from S to T . We let $\mathrm{Is}(S, S) = \mathrm{Is}(S)$ and call this the mapping class group of the oriented space Y_S .

The isotopy futures group of Y_S .

Given $S, x \in S$ and $n \in \mathbb{Z}$, define

$$r(x, n) = \{[(w, 0)] \in Y_S : w \in S, w_i = x_i \text{ for } i \leq n\}.$$

We call such a set a *ray* in Y_S . We say two sets E, E' in Y_S are *isotopic* ($E \sim E'$) if there is a homeomorphism $\varphi: Y_S \rightarrow Y_S$ such that $\varphi(E) = E'$ and φ is isotopic to the identity. An *isotopy ray* is a set isotopic to a ray. A *beam* is a disjoint union of finitely many rays. An *isotopy beam* is a set isotopic to a beam. Let $\mathcal{B} = \mathcal{B}(S)$ denote the set of isotopy beams of Y_S .

We define $\mathcal{F}(S)$, the *isotopy futures group of S* , to be $\mathbb{Z}\mathcal{B}/K$, where $\mathbb{Z}\mathcal{B}$ is the free abelian group with generating set $\mathcal{B} = \mathcal{B}(S)$, and $K = K(S)$ is

the subgroup of $\mathbb{Z}\mathcal{B}$ generated by all elements of the following forms:

$$(7.1) \quad b - b', \quad \text{if } \{b, b'\} \subset \mathcal{B} \text{ and } b \sim b',$$

$$(7.2) \quad b - \sum_{j=1}^k b_j, \quad \text{if } \{b, b_1, \dots, b_k\} \subset \mathcal{B}, k \in \mathbb{N}, \text{ and } b = \cup_j b_j.$$

For S a subshift, let $\mathcal{P}_n(S)$ denote the partition of S into clopen sets of the form $C(x, n) = \{w \in S : w_i = x_i \text{ if } |i| \leq n\}$.

Lemma 7.3. *Suppose S, T are subshifts and $\varphi : Y_S \rightarrow Y_T$ is an orientation preserving homeomorphism and $b \in \mathcal{B}(S)$. Then $\varphi(b) \in \mathcal{B}(T)$.*

Proof. Exploiting the zero dimensionality of S as in [PS], after postcomposing φ with a suitable map isotopic to the identity we may assume that there is a positive integer M such that for any C in $\mathcal{P}_M(S)$ there is a constant $h = h_C$ and a homeomorphism $f = f_C$ from C to a clopen subset D of T such that $\varphi([(x, 0)]) = [(f(x), h)]$, for all x in C .

Because φ respects disjoint union and pushes $\text{Is}(S)$ forward to $\text{Is}(T)$ (by the rule $[h] \mapsto [\varphi h \varphi^{-1}]$), it suffices to consider the case that b is a ray $r(x, n)$ with $n \geq M$. Let $C' = \{w \in S : w_i = x_i \text{ if } i \leq n\} \subset C \in \mathcal{P}_M(S)$, with $h = h_C$ and $f = f_C$. Choose $k \in \mathbb{N}$ such that for all x in C , the sequence $x(-\infty, n]$ determines $(fx)(-\infty, n - k]$ and the sequence $(fx)(-\infty, n + k]$ determines $x(-\infty, n]$. So, if $w \in f(C')$, then $\{z \in T : z_i = w_i \text{ if } i \leq n + k\} \subset f(C')$. Let \mathcal{W} be the (finite) set of words $\{w[n - k, n + k] : w \in f(C')\}$. Then $\varphi(b) = \varphi(r(x, n)) = \cup_{W \in \mathcal{W}} \{[(z, h)] : z(-\infty, n - k - 1] = (fx)(-\infty, n - k - 1]$ and $z[n - k, n + k] = W\}$, so $\varphi(b)$ is an isotopy beam. \square

The following proposition follows easily from the lemma.

Proposition 7.4. *Suppose S and T are subshifts and $\varphi : Y_S \rightarrow Y_T$ is an orientation preserving homeomorphism. Then the mapping of isotopy beams $b \mapsto \varphi(b)$ induces an isomorphism $\varphi_* : \mathcal{F}(S) \rightarrow \mathcal{F}(T)$.*

Let $\text{Iso}(\mathcal{F}(S), \mathcal{F}(T))$ denote the set of group isomorphisms from $\mathcal{F}(S)$ to $\mathcal{F}(T)$. Let $\text{Aut}(\mathcal{F}(S)) = \text{Iso}(\mathcal{F}(S), \mathcal{F}(S))$. The next proposition is now obvious.

Proposition 7.5. *The rule $\varphi \mapsto \varphi_*$ induces a group homomorphism $\rho : \text{Is}(S) \rightarrow \text{Aut}(\mathcal{F}(S))$.*

Remark 7.6. The construction of \mathcal{F}_S is one of several variations on the dimension group construction introduced by Krieger [Kr1, Kr2]; our construction was influenced also by [LM] and [BFF]. The construction of \mathcal{F}_S is a flow equivalence analogue of Krieger’s construction of a dimension group from a subshift S . The map $\rho : \text{Is}(Y_S) \rightarrow \text{Aut}(\mathcal{F}_S)$ is the analogue for flow equivalence of the dimension representation of the automorphism group of a subshift.

The isomorphism $\beta: \mathcal{F}(\sigma_A) \rightarrow \mathbf{cok}(I - A)$.

Suppose A is a matrix in $\mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$. Let \mathcal{I} denote the index set of the rows and columns of A . Let $\mathbb{Z}^{\mathcal{I}}$ be the group of (infinite) row vectors indexed by \mathcal{I} , with all but finitely many entries zero. For a symbol/edge x_n of σ_A , let $\tau(x_n)$ denote the terminal vertex of the edge x_n (so, $\tau(x_n) \in \mathcal{I}$).

The group $\mathbf{cok}(I - A)$ is the cokernel of the map $\mathbb{Z}^{\mathcal{I}} \rightarrow \mathbb{Z}^{\mathcal{I}}$ given by $v \mapsto v(I - A)$ (i.e., $\mathbf{cok}(I - A) = \mathbb{Z}^{\mathcal{I}}/\text{image}(I - A)$). Given a ray $r = r(x, n)$ with $i = \tau(x_n)$, let e_i be the i th canonical basis vector in $\mathbb{Z}^{\mathcal{I}}$, and define $\beta(r) = [e_i] \in \mathbf{cok}(I - A)$.

First note, given $k \in \mathbb{Z}$ and a ray $r = r(x, n)$, if we set r' equal to $\{[(w, k)]: [(w, 0)] \in r(x, n)\}$, then r' is again a ray,

$$(7.7) \quad r' = r(\sigma^k x, n - k) \quad \text{and} \quad \beta(r') = \beta(r).$$

Here the equality of sets follow from the manipulations

$$\begin{aligned} \{[(w, k)]: [(w, 0)] \in r(x, n)\} &= \{[(\sigma^k w, 0)]: w(-\infty, n] = x(-\infty, n]\} \\ &= \{[(z, 0)]: z(-\infty, n - k] = x(-\infty, n]\} \\ &= \{[(z, 0)]: z(-\infty, n - k] = (\sigma^k x)(-\infty, n - k]\} \end{aligned}$$

and then $\beta(r) = \beta(r')$ because the edges x_n and $(\sigma^k x)_{n-k}$ are equal.

Next, given $x \in \sigma_A$, $n \in \mathbb{Z}$ and $k \in \mathbb{N}$, for each σ_A -word $W = W_1 \cdots W_k$ which can follow x_n , choose a point $y = y_W$ such that $y(-\infty, n] = x(-\infty, n]$ and $y[n + 1, n + k] = W$. Then the equality

$$(7.8) \quad \beta(r(x, n]) = \sum_W \beta(r(y_W, n + k)) \in \mathbf{cok}(I - A)$$

follows for $k = 1$ by direct computation and for $k > 1$ by induction.

Given a beam b which is a disjoint union of finitely many rays $r(x^{(i)}, n^{(i)})$, we now define

$$\beta(b) = \sum_i \beta(r(x^{(i)}, n^{(i)})).$$

(We will use the symbol β for various maps derived from the map β on rays.) To see that this definition is independent of the particular choice of rays, suppose b is also the union of rays $r(w^{(j)}, m^{(j)})$. Choose $M \geq \max_{i,j} \{n^{(i)}, m^{(j)}\}$. Then b is the disjoint union of rays $r(z^{(k)}, M)$, each of the $r(x^{(i)}, n^{(i)})$ and $r(w^{(j)}, m^{(j)})$ is a union of some of the rays $r(z^{(k)}, M)$, and by (7.8) we have

$$\sum_i \beta(r(x^{(i)}, n^{(i)})) = \sum_k \beta(r(z^{(k)}, M)) = \sum_j \beta(r(w^{(j)}, m^{(j)})).$$

Therefore $\beta(b)$ is well-defined.

We will write Y_A for the mapping torus of σ_A .

Lemma 7.9. *If b and b' are beams in Y_A such that $b \sim b'$, then $\beta(b) = \beta(b')$.*

Proof. Without loss of generality, choose $M \in \mathbb{N}$ and a finite set E of Y_A such that b is the disjoint union of rays $r(x, M)$, $x \in E$. Let $\{\varphi_t\}$ be an isotopy such that $\varphi_0 = \text{id}$ and $\varphi_1(b) = b'$. Because $\varphi_1 \sim \text{id}$, there is a continuous function $k(x)$ such that for all $[(x, 0)]$ in b , $\varphi_1: [(x, 0)] \mapsto [(x, k(x))]$. Because $\varphi(b)$ is a beam, the function k is integer valued. Possibly after increasing our choice of M , we may assume that k is constant on each ray $r(x, M)$. By (7.7), φ_1 takes each ray $r = r(x, M)$ onto a ray r' such that $\beta(r) = \beta(r')$, b' is the disjoint union of these rays r' , and

$$\beta(b) = \sum_r \beta(r) = \sum_{r'} \beta(r') = \beta(b').$$

□

An isotopy beam b is isotopic to some beam b' . Define $\beta(b) = \beta(b')$. It follows from the lemma that $\beta(b)$ does not depend on the choice of b' . Likewise we have a well-defined homomorphism of groups

$$(7.10) \quad \begin{aligned} \beta: \mathbb{Z}\mathcal{B} &\rightarrow \text{cok}(I - A), \\ \sum n_i b_i &\mapsto \sum n_i \beta(b_i). \end{aligned}$$

Proposition 7.11. *The kernel of the map β in (7.10) is the subgroup K with generators (7.1, 7.2). So, there is an induced isomorphism of groups*

$$\beta_A: \mathcal{F}(\sigma_A) \rightarrow \text{cok}(I - A).$$

Proof. First we show $K \subset \text{Ker}\beta$ by showing that β vanishes on the generators of K . For (7.1), suppose $b \sim b'$; then $\beta(b - b') = 0$ by Lemma 7.9. For (7.2), suppose b is an isotopy beam and b is the disjoint union of finitely many isotopy beams b_i . Without loss of generality, suppose b is a beam. The b_i are a finite collection of disjoint compact sets, so for sufficiently large m , for any C in $\mathcal{P}_m(S)$ such that $b \cap C \neq \emptyset$, the set $(b \cap C)$ will be contained in one of the b_i . If m is large enough, then $b \cap C$ if nonempty will be a ray. Thus, taking sums over C in \mathcal{P}_m , and for notational convenience defining β to be zero on the empty set, we get

$$\begin{aligned} \beta(b) &= \sum_C \beta(C \cap b) \\ &= \sum_j \sum_C \beta(C \cap b_j) = \sum_j \beta(b_j). \end{aligned}$$

Now we show $\text{ker}\beta \subset K$. Suppose $g = \sum n_j b_j \in \text{ker}\beta$. There exists $M \geq 0$ such that for each j , there are rays $r(x^{(jk)}, M)$ such that $b_j - \sum_k r(x^{(jk)}, M) \in K$. so $g = \sum_{jk} n_j r(x^{(jk)}, M) \pmod{K}$. For any x , $\beta(r(\sigma^M x, 0)) = \beta(r(x, M))$; also, $r(x, 0) - r(x', 0) \in K$ if x_0 and x'_0 have the same terminal vertex i . So, we may choose for each i an element $x^{(i)}$

such that $(x^{(i)})_0$ has terminal vertex i , for each $x^{(jk)}$ replace $x^{(jk)}$ with the appropriate $x^{(i)}$, and after reindexing obtain integers m_i such that

$$g = \sum_i m_i r(x^{(i)}, 0) \pmod{K}.$$

Because $\beta(g) = 0$ and $K \subset \ker \beta$, there is an integral row vector w such that $\sum_i m_i e_i = w(I - A)$, and therefore

$$g = \sum_i w_i \left(\left[r(x^{(i)}, 0) \right] - \sum_j A_{ij} \left[r(x^{(j)}, 0) \right] \right) \pmod{K}.$$

For each i , we have $r(\sigma_A x^{(i)}, -1) - r(x^{(i)}, 0) \in K$, and also

$$r(\sigma_A x^{(i)}, -1) - \sum_j A_{ij} r(x^{(j)}, 0) \in K.$$

It follows that $g = 0 \pmod{K}$. □

In our definition of $\mathcal{F}(S)$ and $\text{cok}(I - A)$ we used sets $r(x, n)$ and row vectors. (So, $\text{cok}(I - A) = \text{rowcok}(I - A)$.) In the same way, using sets

$$r^+(x, n) := \{[(w, 0)] \in Y_S : w \in S \text{ and } w_i = x_i \text{ for } i \geq n\}$$

we may define the pasts group $\mathcal{P}(S)$; and using column vectors, we obtain an isomorphism $\mathcal{P}(\sigma_A) \rightarrow \text{colcok}(I - A)$. For a flow equivalence $\varphi: Y_A \rightarrow Y_B$, the isomorphism $\varphi_*: \mathcal{F}(\sigma_A) \rightarrow \mathcal{F}(\sigma_B)$ given by Proposition 7.4 induces the isomorphism

$$\varphi_*^{\text{row}} := (\beta_B)\varphi_*(\beta_A)^{-1}: \text{rowcok}(I - A) \rightarrow \text{rowcok}(I - B).$$

Likewise, the action of φ on $\mathcal{P}(\sigma_A)$ induces an isomorphism

$$\varphi_*^{\text{col}}: \text{colcok}(I - A) \rightarrow \text{colcok}(I - B).$$

The action of $\text{Is}(\sigma_A)$ on $\text{cok}(I - A)$.

For a flow equivalence $\varphi: Y_A \rightarrow Y_B$, we have group homomorphisms

$$\text{Is}(\sigma_A) \rightarrow \text{Aut}(\text{rowcok}(I - A)) \quad \text{and} \quad \text{Is}(\sigma_A) \rightarrow \text{Aut}(\text{colcok}(I - A))$$

$$\varphi \mapsto \varphi_*^{\text{row}} \qquad \qquad \qquad \varphi \mapsto \varphi_*^{\text{col}}.$$

As described in Subsection 2.2, if (U, V) is a basic positive $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence from $(I - A)$ to $(I - B) = U(I - A)V$, and B plays the role of A' in Subsection 2.2, then there is an associated map γ from σ_A to σ_B , and it is easy to see that this map is the restriction (to the cross section σ_A) of an orientation preserving homeomorphism $Y_A \rightarrow Y_B$. More generally, if (U, V) is the composition of basic positive $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalences (U_i, V_i) , and φ is the corresponding composition of the flow equivalences associated to the (U_i, V_i) , then we will write $\varphi = \varphi_{(U, V)}$. This is an abuse of notation in that we are not claiming that (U, V) determines φ (the map φ may depend on

the particular factorization of (U, V)); we are only indicating that φ arises via some factorization of (U, V) .

Proposition 7.12. *Suppose $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$. Suppose $(U, V): (I - A) \rightarrow (I - B)$ is a positive $\text{SL}_{\mathcal{P}}$ equivalence, and $\varphi_{(U,V)}$ is an associated flow equivalence.*

Then the induced map $\varphi_^{\text{row}}: \text{rowcok}(I - A) \rightarrow \text{rowcok}(I - B)$ is given by the rule $[w] \mapsto [wV]$, and the induced map $\varphi_*^{\text{col}}: \text{colcok}(I - A) \rightarrow \text{colcok}(I - B)$ is given by the rule $[w] \mapsto [Uw]$.*

Proof. We will check the proposition in the case that $(U, V) = (E, I)$ and E is a basic elementary matrix with unique offdiagonal entry $E(i, j) = 1$. The argument for (I, E) is similar and then the proposition follows by composition. For concreteness, suppose $E(1, 2) = 1$ (in other entries $E = I$). Let $\varphi = \varphi_{(E,I)}$ be defined via the map γ and edge e described in Subsection 2.2.

Suppose $x \in \sigma_A$ and x_{-1} has terminal vertex i . Then the edge $(\gamma x)_{-1}$ has terminal vertex i and φ maps $r(x, -1)$ onto $r(\gamma x, -1)$. It follows that the diagram

$$\begin{array}{ccc} \mathcal{F}(\sigma_A) & \xrightarrow{\varphi_*} & \mathcal{F}(\sigma_B) \\ \beta_A \downarrow & & \beta_B \downarrow \\ \text{rowcok}(I - A) & \xrightarrow{\text{Id}} & \text{rowcok}(I - B) \end{array}$$

commutes; that is, $\varphi_*^{\text{row}} = \text{Id}$.

If $x \in \sigma_B$ and x_0 has initial vertex not equal to 2, then φ maps $r^+(x, 0)$ onto $r^+(\gamma x, 0)$. Thus the map $(\varphi_*^{\text{col}})^{-1}$ sends $[e_i]$ in $\text{colcok}(I - A)$ to $[e_i]$ in $\text{colcok}(I - B)$ whenever $i \neq 2$. If the initial vertex of x_0 is 2, then φ^{-1} sends $r^+(x, 0)$ to the set of all points $(w, 0)$ in $r^+(\gamma^{-1}x, 0)$ such that $w_{-1} \neq e$. Consequently, if y is a point in σ_A such that $y_i = (\gamma x)_i$ if $i \geq 0$ and $y_{-1} = e$, then

$$\varphi^{-1}(r^+(x, 0)) = r^+(\gamma^{-1}x, 0) \setminus r^+(y, -1).$$

We also have

$$\begin{aligned} \beta_B^{\text{col}}: \quad r^+(x, 0) &\mapsto [e_2] \in \text{colcok}(I - B), \\ \beta_A^{\text{col}}: \quad r^+(\gamma^{-1}x, 0) &\mapsto [e_2] \in \text{colcok}(I - A), \\ \beta_A^{\text{col}}: \quad r^+(y, -1) &\mapsto [e_1] \in \text{colcok}(I - A). \end{aligned}$$

Therefore $(\varphi_*^{\text{col}})^{-1}: [e_2] \mapsto [e_2] - [e_1]$, hence for all integral column vectors v we have $(\varphi_*^{\text{col}})^{-1}: [v] \mapsto [E^{-1}v]$ as required. \square

Theorem 7.13. *Suppose $A \in \mathfrak{M}_+^{\circ}(\mathbb{Z})$ and the mapping torus of σ_A is not a circle. Then the induced map $\text{Is}(Y_A) \rightarrow \text{Aut}(\text{cok}(I - A))$ is surjective.*

Remark 7.14. Of course, the theorem is true for colcok as well as for rowcok. In the case that the mapping torus of σ_A is a circle (i.e., A has a unique irreducible component, and this component is a permutation matrix), any orientation preserving homeomorphism from Y_A to Y_A is isotopic to the identity, but $\text{cok}(I - A) \cong \mathbb{Z}$ and $\text{Aut}(\text{cok}(I - A)) \cong \mathbb{Z}/2$, so the map $\text{Is}(Y_A) \rightarrow \text{Aut}(\text{cok}(I - A))$ is not surjective. Theorem 7.13 says that apart from this case, every automorphism of the isotopy futures group of an irreducible shift of finite type is induced by a flow equivalence.

Proof of Theorem 7.13. It is proved in [BH] that any automorphism of rowcok($I - A$) or colcok($I - A$) is induced by an $\text{SL}(\mathbb{Z})$ equivalence (by the rules described in the statement of Proposition 7.12). By the Factorization Theorem 3.3, such an equivalence is a positive equivalence. By Proposition 7.12, a flow equivalence associated to this positive equivalence has the desired action on the cokernel group. \square

From the view of symbolic dynamics, Theorem 7.13 stands in contrast to the Kim-Roush-Wagoner result [KRW1] that the dimension representation of a mixing shift of finite type is not in general surjective. (The contrast is meaningful because the invariants are related by “setting t equal to 1” [B1].)

When $A \in \mathfrak{M}_+^{\circ}(\mathbb{Z})$ (i.e., A is essentially irreducible) and σ_A is not a circle, the flow equivalence class of σ_A is given by the $\text{SL}(\mathbb{Z})$ equivalence class of $I - A$, for which $\det(I - A)$ and $\text{cok}(I - A)$ give complete invariants. When \mathcal{P} is nontrivial and $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ (i.e., the SFT σ_A is reducible), the flow equivalence class of A (modulo a permutation of \mathcal{P}) is given by its positive $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence class, and the complete algebraic invariants (introduced by Huang) are more subtle, involving the “ K -web” of the matrix $I - A$, denoted $K(I - A)$. The K -web is a diagram of exact sequences of certain kernel and cokernel groups of submatrices of $I - A$. The K -web invariants are completely analyzed in [BH], which also characterizes the automorphisms of $K(A)$ which can be induced by an $\text{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence. We believe that the type of analysis carried out to describe the action of $\text{Is}(\sigma_A)$ on $\text{cok}(I - A)$ in the irreducible case can be extended to describe the possible actions of $\text{Is}(\sigma_A)$ on the more complicated algebraic structure of the K -web which classifies in the reducible case. Specifically, we expect that the following program can be carried out. Together with [BH], this program would give a complete description of the possible actions of $\text{Is}(\sigma_A)$ on the K -web.

Program 7.15. For A, B in $\mathfrak{M}_{\mathcal{P}}(\mathbb{Z}_+)$, we conjecture the following.

- (1) The K -web data for $I - A$ can be described in terms of isotopy beams of subsystems of Y_A , and the map on isotopy beams by an orientation preserving homeomorphism $\varphi: Y_A \rightarrow Y_B$ induces an isomorphism $K(A) \rightarrow K(B)$.

- (2) For a positive $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence (U, V) from A to B , the isomorphism $K(A) \rightarrow K(B)$ induced by $\varphi_{(U,V)}$ is the natural isomorphism induced by (U, V) as described in [BH].
- (3) If φ is an orientation preserving homeomorphism from Y_A to Y_B , then there is a positive $\mathrm{SL}_{\mathcal{P}}(\mathbb{Z})$ equivalence (U, V) such that φ is isotopic to $\varphi_{(U,V)}$.

The most fundamental of the three steps above is the last one, and a version of this has already been carried out in the irreducible case (i.e., $\mathcal{P} = \{1\}$) by Badoian [Ba1], as we discuss below.

The work of Badoian.

We'll describe some of the work [Ba1] of Leslie Badoian, which gives alternate proofs of some of our results. The work [Ba1] is too extensive for a full summary here; roughly speaking, Badoian carries out for irreducible shifts of finite type a flow equivalence version of the strong shift equivalence theory Wagoner [W1] built on the foundation laid by Williams [Wi].

Badoian builds an infinite oriented CW complex, denoted FK . A zero-cell for FK is an equivalence class of infinite, essentially irreducible, finitely supported zero-one matrices, where two matrices are equivalent iff their unique maximal irreducible principal submatrices are equal. A one-cell $[A] \rightarrow [B]$ corresponds to an elementary equivalence $(I - B) = U(I - A)V$ satisfying certain conditions. Two-cells are also defined, by certain matrix relations. The two main results of [Ba1] are the following:

- *Classification Theorem.* σ_A and σ_B are flow equivalent if and only if A and B lie in the same connected component of FK .
- *Flow Equivalence Theorem.* $\pi_1(FK_A) \cong \mathrm{Is}(\sigma_A)$. (I.e., a path along one-cells gives rise to a flow equivalence, and two paths give rise to isotopic flow equivalences if and only if the paths are homotopic in FK .)

The elementary equivalences of [Ba1] are not the same as our elementary positive equivalences, but Badoian has found short arguments [Ba2] which show directly that that her elementary equivalences and ours generate the same set of flow equivalences up to isotopy. With this fact and some technical remarks, the results of Section 6 for irreducible shifts of finite type follow directly from Badoian's Classification Theorem (which in turn rests on Parry-Sullivan [PS] and Williams [Wi], as does our Section 6).

The Flow Equivalence Theorem gives an alternate route in the irreducible case to the representation $\mathrm{Is}(\sigma_A, \sigma_B) \rightarrow \mathrm{Iso}(\mathrm{cok}(I - A), \mathrm{cok}(I - B))$: We could take the natural definition along an edge (given by the associated flow equivalence), compose along paths of edges, and consult the definition of two-cells in FK to verify that the definition only depends on the homotopy class of the path of edges. All of this is parallel to the development of the dimension representation in Wagoner's strong shift equivalence theory [W1].

We have not relied in proofs on citation of [Ba1], for a few reasons. Although there should be no fundamental problem with extending Badoian’s approach to reducible shifts of finite type, the results in [Ba1] are only for irreducible shifts of finite type. We also wanted self-contained and reasonably brief arguments. (The long work [Ba1] deals with a fundamental difficult problem which we avoid: We do not try to understand when two paths give rise to the same flow equivalence up to isotopy.) Finally, although the CW complex approach has rather spectacularly proved its worth [KR2, KRW1, W1], the Krieger-style construction remains important, and its more earthy definition (by actions on sets) makes sense directly for general subshifts. Matsumoto [Ma] has a far reaching extension of Williams’ theory to general subshifts, and this offers hope for some analogue of Wagoner’s strong shift equivalence theory for general subshifts; but there is no such theory yet.

Appendix A. Reduction to nondegenerate form.

This appendix is devoted to the proof of Proposition 4.5.

We will prove Proposition 4.5 by composition in a larger commuting diagram (to be assembled in three stages):

$$\begin{array}{ccccccc}
 (I - A) & \longrightarrow & (I - \overline{A_1}) & \longrightarrow & (I - \overline{A_2}) & \longrightarrow & (I - \overline{A}) \\
 (U, V) \downarrow & & (U_1, V_1) \downarrow & & (U_2, V_2) \downarrow & & \downarrow (\overline{U}, \overline{V}) \\
 (I - A') & \longrightarrow & (I - \overline{A'_1}) & \longrightarrow & (I - \overline{A'_2}) & \longrightarrow & (I - \overline{A'}) .
 \end{array}$$

The horizontal arrows will be positive equivalences and the vertical equivalences to the right of (U, V) will be defined from them by composition (then the diagram will commute). Stage I will produce the left square with $\overline{A_1}$ and $\overline{A'_1}$ satisfying Conditions (2), (3) and (4) of Proposition 4.5. Stage II will produce the middle square, with $(U_2)_{ii} = (V_2)_{ii} = \text{Id}$ for $i \in \mathcal{C}$, and with $\overline{A_2}$ and $\overline{A'_2}$ still satisfying Conditions (2), (3) and (4) of Proposition 4.5. Stage III will produce the right square to finish the proof. The individual stages will follow from several lemmas.

Lemma A.1. *Suppose $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$. Then there is a positive $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ from $I - A$ to a matrix $I - C$ such that for all $i \in \mathcal{C}_A$ the following hold:*

- (1) *The block C_{ii} has its upper left corner entry equal to 1, and C_{ii} has no other nonzero entry.*
- (2) *Let (ℓ, ℓ) be the entry of C which is the upper left corner of C_{ii} . Then for $j \neq i$, every row of a block C_{ij} other than row ℓ is zero, and every column of a block C_{ji} other than column ℓ is equal to zero.*

Proof. Suppose $i \in \mathcal{C}_A$. Let i_1, \dots, i_k be nonrepeated indices such that $A_{ii}(i_t, i_{t+1}) = 1, 1 \leq t < k$, and $A_{ii}(i_k, i_1) = 1$.

Cycle-shortening construction. Suppose $k > 1$. Let $A = A^{(0)}$. For $1 \leq j < k$, define $A^{(j)}$ by the equation $I - A^{(j)} = E_j(I - A^{(j-1)})$, where E_j denotes the basic elementary matrix which acts to add row i_{k-j+1} to row i_{k-j} . Each $A^{(j)}$ is nonnegative. Then add the columns i_2, \dots, i_k of $A^{(k)}$ to column i_1 of $A^{(k)}$. By Lemma 2.5, each step in this process gives a positive equivalence in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, and in the last matrix A' , the block A'_{ii} has as its unique cycle the 1-cycle (i_1) . Below is an example of the process, with $(i_1, i_2, i_3, i_4) = (1, 2, 3, 4)$, viewed in the principal submatrices on indices 1, 2, 3, 4:

$$\begin{pmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \\ -1 & 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \rightarrow$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Now without loss of generality, we suppose $k = 1$ with $A(i_1, i_1) = 1$. Because (i_1) is the unique A_{ii} cycle, if A_{ii} is nonzero at any entry other than (i_1, i_1) , then it is nonzero at some entry (j, l) such that row l of A is zero or column j of A is zero. In the former case, let E be the elementary matrix which acts from the left to add row l to row j , then $(E, I): (I - A) \rightarrow E(I - A) = (I - A')$ is a positive equivalence in which $A' = A$ except that $A'(j, l) = 0$. The latter case is treated similarly, by adding column j to column l . Iterating, we produce a positive equivalence in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$ from $I - A$ to a matrix $I - A'$ such that $A'(i_1, i_1)$ is the only nonzero entry of A'_{ii} .

Next, given i in $\mathcal{C}_{A'}$ with $A'(i_1, i_1) = 1$, we may for each $j \prec i$ add rows of $I - A'$ through the ii block to rows through the ji block (never adding row i_1) until every column of the block $(I - A')_{ji}$ except column i_1 is zero. We do this for all the cycle components i , for i in decreasing order, so that no block zeroed out for some i is made nonzero by subsequent operations. Then similarly, taking i in \mathcal{C}_A in increasing order, we add columns through the ii block to columns through the ji blocks with $i \prec j$, to end with a matrix C' which satisfies the statement of the lemma (with C' in place of C), except that the distinguished indices i_1 might not be the corner indices ℓ .

So, suppose i is a cycle component for which $\ell \neq i_1$. We apply four basic positive equivalences to give $(I - C') \rightarrow (I - C'')$, as viewed below

in principal submatrices on indices $\{i_1, \ell, k\}$ (where k is any index not in $\{i_1, \ell\}$). (In the very special case that C' is 2×2 , there can be no third index k and these principal submatrices should be restricted to indices $\{i_1, \ell\}$.) For concreteness we use $(\ell, i_1, k) = (1, 2, 3)$:

$$\begin{aligned} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & x \\ 0 & y & z \end{pmatrix} &\rightarrow \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & x \\ 0 & y & z \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -1 & 0 & x \\ 0 & y & z \end{pmatrix} \\ &\rightarrow \begin{pmatrix} 1 & 0 & 0 \\ -1 & 0 & x \\ 0 & y & z \end{pmatrix} \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & x \\ 0 & y & z \end{pmatrix} \\ &\rightarrow \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & x \\ 0 & y & z \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 1 & x \\ y & y & z \end{pmatrix} \\ &\rightarrow \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 & 0 \\ 0 & 1 & x \\ y & y & z \end{pmatrix} = \begin{pmatrix} 0 & 0 & x \\ 0 & 1 & x \\ y & y & z \end{pmatrix}. \end{aligned}$$

If above for any k we have $x \neq 0$, then $x < 0$ and the (i_1, k) entry lies in an ij block with $i < j$; then $y = 0$ and it is a positive equivalence to add column i_1 of C'' $|x|$ times to column k . Doing this as needed, and dealing similarly with nonzero entries y using rows in place of columns, we produce another version of C'' which enjoys the additional property that $i_1 = \ell$ for the cycle component i . Then we repeat until $i_1 = \ell$ for every cycle component i . The resulting matrix C satisfies the statement of the lemma. \square

Lemma A.2. *Suppose $A \in \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbf{n}, \mathbb{Z})$, and $\mathbf{n} = (n_1, \dots, n_N)$ is a vector with positive integer entries. Then there is a positive $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$ from $(I - A)$ to a matrix with Properties (2), (3) and (4) of Proposition 4.5.*

Proof of Lemma A.2. We will describe a sequence of row and column operations (corresponding, by repeated tacit appeal to Lemma 2.5, to positive $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalences in $I - \mathfrak{M}_{\mathcal{P},+}^{\circ}(\mathbb{Z})$) which put the matrix $I - A$ into the required form. To simplify notation, rather than renaming $I - A$ after an equivalence, we will discuss changing properties of $I - A$. We begin with a matrix A with the properties stated (for C) in Lemma A.1, i.e., A satisfies Properties (3) and (4) of Proposition 4.5.

Our first goal will be, given $t \in \mathcal{P}$ which is not a cycle component, to arrange that the block $(I - A)_{tt}$ be strictly negative. Recall \mathcal{I}_t denotes the index set for rows/columns of A_{tt} . Let \mathcal{S} denote the index set for the unique maximal irreducible submatrix of A_{tt} , let \mathcal{S}' denote the complement of \mathcal{S} in \mathcal{I}_t , and e.g., let $A\{\mathcal{S}\}$ denote the principal submatrix of A on index set \mathcal{S} . We will arrange (in order) the following properties (after each stage keeping the

properties achieved at earlier stages, and not changing entries in any block A_{ss} with $s \neq t$, and not losing Properties (3) and (4) of Proposition 4.5).

- (1) $\exists i \in \mathcal{S}$ such that $(I - A)(i, i) \leq 0$.
- (2) $\{i, j\} \cap \mathcal{S}' \neq \emptyset \implies (I - A)(i, j) = \delta_{ij}$.
- (3) If $\mathcal{S}' \neq \emptyset$, then $|\mathcal{S}'| > 1$.
- (4) $\mathcal{S}' = \emptyset$.
- (5) The block $(I - A)_{tt}$ is strictly negative.

(1) If necessary achieve this with the initial row operations of the cycle-shortening construction of the Lemma A.1.

(2) First suppose this condition does not hold for some $\{i, j\} \subset \mathcal{I}_t$. Then pick some $i \in \mathcal{S}'$ and $j \in \mathcal{I}_t$ such that $j \neq i$ and one of the following hold:

- $(I - A)(i, j) \neq 0$ and column i of A_{tt} is zero, or
- $(I - A)(j, i) \neq 0$ and row i of A_{tt} is zero.

In the former case, add column i of $(I - A)$ to other columns j where $(I - A)(i, j) < 0$, until $(I - A)(i, j) = \delta_{ij}$ for all $j \in \mathcal{I}_t$. In the latter case, similarly use row additions to achieve $(I - A)(j, i) = \delta_{ij}$ for all $j \in \mathcal{I}_t$. This procedure reduces the cardinality of the set of entries in $(I - A)_{tt}$ at which Condition (2) fails, and it may be repeated until Condition (2) holds for $\{i, j\} \subset \mathcal{I}_t$. We then add rows and columns indexed by \mathcal{S}' to others as needed until (2) holds in general.

(3) Suppose (for concreteness) that $\mathcal{S} = \{1\}$ and $2 \in \mathcal{S}'$. Then we must have $A(1, 1) = k > 1$ (since t is not a cycle component). Now, subtract row 2 of $(I - A)$ from row 1; then subtract column 2 from column 1. The effect of these moves is to enlarge $\mathcal{S} = \{1\}$ to $\mathcal{S} = \{1, 2\}$. The moves are summarized below in principal submatrices on indices $\{1, 2, 3\}$, where 3 is an arbitrary additional index:

$$\begin{pmatrix} 1 - k & 0 & w \\ 0 & 1 & 0 \\ x & 0 & z \end{pmatrix} \rightarrow \begin{pmatrix} 1 - k & -1 & w \\ 0 & 1 & 0 \\ x & 0 & z \end{pmatrix} \rightarrow \begin{pmatrix} -(k - 2) & -1 & w \\ -1 & 1 & 0 \\ x & 0 & z \end{pmatrix}.$$

(4) Suppose $\mathcal{S}' \neq \emptyset$. By (1) and (3), we may pick i_1, j_1 in \mathcal{S} such that $i_1 \neq j_1$, $(I - A)(i_1, i_1) \leq 0$, and $(I - A)(i_1, j_1) \leq -1$. Add row i_1 of $(I - A)$ to row j_1 , $(|\mathcal{S}'| + 1)$ times, producing $(I - A)(j_1, j_1) \leq -|\mathcal{S}'|$. For each j in \mathcal{S}' , subtract row j of $(I - A)$ from row j_1 . Then subtract each \mathcal{S}' column from column j_1 . This produces A with $\mathcal{S}' = \emptyset$.

(5) With i_1, j_1 as in (4): Add row i_1 to row j_1 (now $(I - A)(j_1, j_1) < 0$); for each i in \mathcal{S} with $i \neq j_1$, add column j_1 to column i (now row j_1 of $(I - A)$ is negative); and for each i in \mathcal{S} with $i \neq j_1$, add row j_1 to row i . We now have $(I - A)_{tt}$ strictly negative as required.

After applying a positive equivalence, then, we may assume that $(I - A)_{ii} < 0$ for every noncycle component i . Consequently, if $i < j$, and i or j is not a cycle component, then for large n the block $(A^n)_{ij}$ is strictly positive.

We can then get a positive equivalence to $(I - A)$ whose block $(I - A)_{ij}$ is strictly negative, by adding columns through i to columns through j (if $i \notin \mathcal{C}_A$) or by adding rows through j to rows through i (if $j \notin \mathcal{C}_A$). Similarly, for every noncycle component j and cycle component i , with $\mathcal{C}_i^{\text{prim}} = \{\ell\}$, add a j -row to row ℓ if $i \prec j$, and add a j -column to column ℓ if $j \prec i$.

Note, if $i \prec j$ and $\{i, j\} \subset \mathcal{C}$, with say $\mathcal{C}_i^{\text{prim}} = \{\ell_i\}$ and $\mathcal{C}_j^{\text{prim}} = \{\ell_j\}$, then $A(\ell_i, \ell_j) > 0$, because the block A_{ij} is not the zero block (because $A \in \mathfrak{M}_{\mathcal{P},+}^o(\mathcal{C}, \mathbf{n}, \mathbb{Z})$) and the only possible nonzero entry is $A(\ell_i, \ell_j) > 0$. Finally, whenever $(I - A)_{ij} < 0$ with $\{i, j\} \subset \mathcal{C}_A$ and $i \prec k \prec j$ for some k in \mathcal{P} , pick k such that $i \prec k \prec j$, and add columns of $I - A$ through component k to columns through component j . The resulting matrix satisfies the statement of the lemma. \square

Lemmas A.1 and A.2 finish the proof for Stage I. We now shift our focus to the form of the equivalence (U, V) . The next lemma gives the proof for Stage II.

Lemma A.3. *Suppose $(U, V): (I - A) \rightarrow (I - A')$ is an $\text{SL}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalence which is positive on cycle components, and A, A' satisfy Conditions (2), (3) and (4) of Proposition 4.5. Then there is a commuting diagram*

$$\begin{array}{ccc} (I - A) & \longrightarrow & (I - \widetilde{A}) \\ (U, V) \downarrow & & \downarrow (\widetilde{U}, \widetilde{V}) \\ (I - A') & \longrightarrow & (I - \widetilde{A}') \end{array}$$

in which the horizontal arrows are positive equivalences; \widetilde{A} and \widetilde{A}' still satisfy Conditions (2), (3) and (4); and for each $i \in \mathcal{C}$, $\widetilde{U}_{ii} = \widetilde{V}_{ii} = \text{Id}$.

Proof. Suppose i is a cycle component for which $n_i > 1$ (otherwise there is nothing to prove). Then $(I - A)_{ii} = (I - A')_{ii} = Q$, where $Q = \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix}$, in which I is $(n_i - 1) \times (n_i - 1)$. Considering blocks of $U_{ii}Q = QV_{ii}^{-1}$, we see U_{ii} and V_{ii} have the corresponding block forms $U_{ii} = \begin{pmatrix} a & 0 \\ x & Z \end{pmatrix}$ and $V_{ii} = \begin{pmatrix} b & y \\ 0 & Z^{-1} \end{pmatrix}$. The positive on cycle components assumption implies $a = 1$. Then $\det(U) = 1$ implies $\det(Z) = 1$. Then $\det(Z^{-1}) = 1 = \det V$ implies $b = 1$. So we have

$$(A.4) \quad U_{ii} = \begin{pmatrix} 1 & 0 \\ x & Z \end{pmatrix} \quad \text{and} \quad V_{ii} = \begin{pmatrix} 1 & y \\ 0 & Z^{-1} \end{pmatrix}$$

for some Z in $\text{SL}(n_i - 1, \mathbb{Z})$.

Now suppose E is a basic elementary matrix with offdiagonal entry $E(j, k) = 1$, where j, k index rows of the ii block other than the first row. Then

$$(A.5) \quad (I - A) \xrightarrow{(I, E^{-1})} \cdot \xrightarrow{(E, I)} (I - A)$$

gives a factorization of $(E, E^{-1}): (I - A) \rightarrow (I - A)$ into basic positive equivalences. For example, if rows 1,2,3 run through Q and $Q(1, 1) = 0$, then

in the principal submatrix on indices 1,2,3 we could have $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$,

and (A.5) would become

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{(I, E^{-1})} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{(E, I)} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Now we can factor (U, V) as

$$(A.6) \quad (I - A) \xrightarrow{(I, E^{-1})} \cdot \xrightarrow{(E, I)} (I - A) \xrightarrow{(UE^{-1}, EV)} (I - A').$$

Because Z is a composition of elementary matrices, and Conditions (2), (3) and (4) are not disturbed by this move, we can repeat this move to obtain a positive equivalence $(G, G^{-1}): (I - A) \rightarrow (I - A)$ such that the (U, V) equals (G, G^{-1}) followed by (UG^{-1}, GV) where $(UG^{-1})_{ii}$ and $(GV)_{ii}$ have the forms (A.4) with $Z = I$. After doing this as needed for every cycle component i , we can assume for each $i \in \mathcal{C}$ with $n_i > 1$ that we have the forms $U_{ii} = \begin{pmatrix} 1 & 0 \\ x^{(i)} & I \end{pmatrix}$ and $V_{ii} = \begin{pmatrix} 1 & y^{(i)} \\ 0 & I \end{pmatrix}$.

Let D and D' be the block diagonal matrices equal to Id except in cycle component diagonal blocks, where $D_{ii} = U_{ii}$ and $D'_{ii} = V_{ii}^{-1}$. We will produce matrices P, Q in $\mathcal{U}_P(\mathbf{n}, \mathbb{Z})$ such that $(D, Q): (I - A) \rightarrow D(I - A)Q$ and $(P, D'): (I - A') \rightarrow P(I - A')D'$ are positive equivalences, and the matrices $D(I - A)Q$ and $P(I - A')D'$ satisfy Conditions (2), (3) and (4). Then the lemma will follow by defining (\tilde{U}, \tilde{V}) by requiring the following diagram to commute:

$$\begin{array}{ccc} (I - A) & \xrightarrow{(D, Q)} & D(I - A)Q \\ (U, V) \downarrow & & \downarrow (\tilde{U}, \tilde{V}) \\ (I - A') & \xrightarrow{(P, D')} & P(I - A')D' \end{array}$$

We will prove the first claim, for (D, Q) ; the proof of the second claim is similar. Let $i_1 < i_2 < \dots < i_k$ be the elements of \mathcal{C} . (Recall, $i \prec j \implies i < j$.) To begin, let $i = i_k$ and let $1, 2, \dots, m$ index the rows through U_{ii} . For $2 \leq j \leq m$, let R_j be the elementary matrix which acts from the right

to subtract column j from column 1. Let $R = R_2 \cdots R_m$ and let $\mathbf{1}$ denote a vector with every entry equal to 1, then

$$\begin{aligned} ((I - A)R)_{ii} &= \begin{pmatrix} 0 & 0 \\ -\mathbf{1} & I \end{pmatrix}, \quad \text{and} \\ ((I - A)R)_{rs} &= (I - A)_{rs} \quad \text{if } rs \neq ii, \end{aligned}$$

and we get a positive equivalence

$$(I - A) \xrightarrow{(I, R_2)} \cdots \xrightarrow{(I, R_m)} (I - A)R.$$

Next, let D_k be a product of elementary matrices, $E = E_n \cdots E_1$, where E_t acts from the left to add ϵ_t ($\epsilon_t = 1$ or $\epsilon_t = -1$) times row 1 to row j_t , and $2 \leq j_t \leq m$. Consider the equivalence $(E_1, I): (I - A)R \rightarrow E_1(I - A)R$. Notice $(E_1(I - A)R)_{ii} = ((I - A)R)_{ii}$. So, this equivalence (E_1, I) is positive unless $(E_1(I - A)R)(j, k) > 0$ for some columns p to the right of the ii block. Let F_1 be the product of basic elementary matrices $F_{1,t}$, $1 \leq t \leq T$ say, which act from the right to subtract column j_1 from such columns p enough times to guarantee (with $F_1 = F_{1,1} \cdots F_{1,T}$) that $(E_1(I - A)RF_1)(j_1, p) < 0$. Then

$$(I - A)R \xrightarrow{(I, F_{1,1})} \cdots \xrightarrow{(I, F_{1,T})} \cdot \xrightarrow{(E_1, I)} E_1(I - A)RF_1$$

gives a positive equivalence $(E_1, F_1): (I - A)R \rightarrow E_1(I - A)RF_1$. Recursively, for $1 \leq t < m$, apply this procedure, to produce F_{t+1} giving a positive equivalence

$$\begin{aligned} E_t \cdots E_1(I - A)RF_1 \cdots F_t &\xrightarrow{(I, F_{t+1})} \cdot \xrightarrow{(E_{t+1}, I)} E_{t+1} \cdots \\ &E_1(I - A)RF_1 \cdots F_{t+1}. \end{aligned}$$

Let $Q_k = F_1 \cdots F_m$: then we have a positive equivalence

$$(I - A) \xrightarrow{(D_k, RQ_k)} D_k(I - A)RQ_k \xrightarrow{(I, R^{-1})} D_k(I - A)RQ_k R^{-1}.$$

Because $RQ_k = Q_k R$, altogether we get

$$(I - A) \xrightarrow{(D_k, Q_k)} D_k(I - A)RQ_k.$$

Notice, $Q_k \in \mathcal{U}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$. Moreover, if $j \in \mathcal{P}$ and $j < i_k$, then for any t the tj blocks of $(I - A)$ and $(D_k(I - A))_k Q_k$ are equal.

Next, for the cycle components i_{k-1}, \dots, i_1 (in that order) we repeat the procedure used above for (D_k, Q_k) to produce pairs $(D_{k-1}, Q_{k-1}), \dots, (D_1, Q_1)$ with $D = D_k D_{k-1} \cdots D_1$ and $Q_{(-)} := Q_k Q_{k-1} \cdots Q_1$ giving a positive equivalence

$$(I - A) \xrightarrow{(D_k, Q_k)} \cdot \xrightarrow{(D_{k-1}, Q_{k-1})} \cdots \xrightarrow{(D_1, Q_1)} D(I - A)Q_{(-)}.$$

To see that the (D_i, Q_i) define positive equivalences, note that for $i_s \neq i_k$, the column-subtracting moves we use to prepare the entries in a block $i_s j$ to the right of the $i_s i_s$ block do not change the sign of entries outside the

$i_s j$ block (because we are subtracting columns through the $i_s i_s$ block with diagonal entry 1, and these columns have no other nonzero entry at this stage, because the earlier subtractions of columns through block $i_r i_r$ with $r \geq s$ do not affect the i_s block column).

For every cycle component i , the ii block of the matrix $D(I - A)Q_{(-)}$ equals Id . Suppose there exists (r, s) such that $r \in \mathcal{C}^{\text{sec}}$ and $(D(I - A)Q_{(-)})(r, s) < 0$; then choose such an (r, s) with r minimal, and add column r to column s . Because the (r, s) entry cannot lie in a diagonal block, this elementary positive equivalence is implemented by multiplication from the right by a matrix in $\mathcal{U}_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$. Repeat this move until a matrix is produced in which the (r, s) entry is zero whenever $r \in \mathcal{C}^{\text{sec}}$. Let the corresponding positive equivalence be denoted $(I, Q_{(+)}) : D(I - A)Q_{(-)} \rightarrow D(I - A)Q_{(-)}Q_{(+)}$. The proof is finished by setting $Q = Q_{(-)}Q_{(+)}$. \square

The next lemma gives the last ingredient, Stage III, for the proof of Proposition 4.5.

Lemma A.7. *Suppose U, V, A, A' satisfy the assumptions of Lemma A.3 and in addition assume that $U_{ii} = V_{ii} = \text{Id}$ for every $i \in \mathcal{C}$. Then there is a commuting diagram of $SL_{\mathcal{P}}(\mathbf{n}, \mathbb{Z})$ equivalences*

$$\begin{array}{ccc} (I - A) & \longrightarrow & (I - \bar{A}) \\ (U, V) \downarrow & & \downarrow (\bar{U}, \bar{V}) \\ (I - A') & \longrightarrow & (I - \bar{A}') \end{array}$$

satisfying the conclusion of Proposition 4.5. (Moreover, $\bar{A} = A$ and $\bar{A}' = A'$.)

Proof. We will build a suitable commuting diagram

$$\begin{array}{ccccc} (I - A) & \xrightarrow{(E^{-1}, H)} & (I - A) & \xrightarrow{(\bar{H}, \bar{E}^{-1})} & (I - A) \\ (U, V) \downarrow & & (U_3, V_3) \downarrow & & \downarrow (\bar{U}, \bar{V}) \\ (I - A') & \xrightarrow{(I, I)} & (I - A') & \xrightarrow{(I, I)} & (I - A') \end{array}$$

and then use $(\bar{H}E^{-1}, H\bar{E}^{-1})$ and (I, I) for the upper and lower horizontal arrows in the diagram required for the lemma. First we work on the left half of the diagram. We will choose E, H, U_3 satisfying:

- (i) $U_3(i, j) = \delta_{ij}, \quad \forall i \in \mathcal{C}^{\text{sec}},$
- (ii) $(E^{-1}, H) : (I - A) \rightarrow (I - A)$ is a positive equivalence, and
- (iii) $H^{-1}(i, j) = \delta_{ij}, \quad \forall i \notin \mathcal{C}^{\text{sec}}.$

Recall, \mathcal{I}_s denotes the set of indices for rows/columns through A_{ss} . To choose E , let the entries (i, j) for which $i \in \mathcal{C}^{\text{sec}}$ and $U(i, j) \neq \delta_{ij}$ be listed as $(i_1, j_1), \dots, (i_n, j_n)$, where $i_k \in \mathcal{I}_{s(k)}$ and $s(1) \preceq s(2) \preceq \dots \preceq s(n)$.

(So, $j_k \in \mathcal{I}_{t(k)}$ with $s(k) \prec t(k)$ since by assumption $U_{s(k)s(k)} = \text{Id.}$) Let $\mu_k = U(i_k, j_k)$. Define matrices E_k , $1 \leq k \leq n$, by $E_k(i_k, j_k) = -\mu_k$ and otherwise $E_k(i, j) = \delta_{ij}$. Then $(UE_k)(i_k, j_k) = 0$. Define $E = E_1 E_2 \cdots E_n$. Then by our ordering $s(1) \preceq s(2) \preceq \cdots \preceq s(n)$, we have $(UE)(i, j) = \delta_{ij}$ for $i \in \mathcal{C}^{\text{sec}}$. Let $U_3 = UE$, now (i) holds, and $U = (UE)E^{-1} = U_3 E^{-1}$ as required for the diagram to commute. Also $E(i, j) = \delta_{ij}$ if $i \notin \mathcal{C}^{\text{sec}}$, so $E^{-1}(i, j) = \delta_{ij}$ if $i \notin \mathcal{C}^{\text{sec}}$.

Next for $1 \leq k \leq n$, we will define H_k such that $(E_k^{-1}, H_k): (I - A) \rightarrow (I - A)$ is a positive equivalence and $H_k(i, j) = \delta_{ij}$ when $i \notin \mathcal{C}^{\text{sec}}$. Then we will set $(E^{-1}, H) = (E_n^{-1} \cdots E_1^{-1}, H_1 \cdots H_n)$, so that $(E^{-1}, H): (I - A) \rightarrow (I - A)$ is the composition of positive equivalences and satisfies (ii). To prepare for the definition of H_k , given k pick M a positive integer greater than the absolute value of any entry in row i_k of $E_k^{-1}(I - A)$, and define a matrix F_k as follows: $F_k(i_k, j) = -M$ if $i_k \neq j$ and $(E_k^{-1}(I - A))(i_k, j) \neq 0$, and $F_k(i, j) = \delta_{ij}$ otherwise. Define a matrix G_k by setting $G_k(i_k, j) = -(E_k^{-1}(I - A)F_k)(i_k, j)$ and $G_k(i, j) = \delta_{ij}$ otherwise. Then we have the positive equivalence

$$(I - A) \xrightarrow{(I, F_k)} \cdot \xrightarrow{(E_k^{-1}, I)} \cdot \xrightarrow{(I, G_k)} (I - A).$$

Let $H_k = F_k G_k$. Note $H_k(i, j) = \delta_{ij}$ if $i \notin \mathcal{C}^{\text{sec}}$, so $H(i, j) = \delta_{ij}$ if $i \notin \mathcal{C}^{\text{sec}}$, and therefore also $H^{-1}(i, j) = \delta_{ij}$ if $i \notin \mathcal{C}^{\text{sec}}$. We now have E, H, U_3 satisfying (i)-(iii).

To get the right half of the commuting diagram, we apply to the equivalence (U_3, V_3) the transpose of the procedure above to get matrices $\overline{E}, \overline{H}, \overline{V}, \overline{U}$ satisfying:

- (i) $\overline{V}(i, j) = \delta_{ij}, \quad \forall j \in \mathcal{C}^{\text{sec}},$
- (ii) $(\overline{H}, \overline{E}^{-1}): (I - A) \rightarrow (I - A)$ is a positive equivalence, and
- (iii) $\overline{H}^{-1}(i, j) = \delta_{ij}, \quad \forall j \notin \mathcal{C}^{\text{sec}},$

where \overline{U} and \overline{V} are defined by $\overline{U} = U_3 \overline{H}^{-1}$ and $\overline{V} = \overline{E} V_3$. Using (i) and the forms of $(I - A)$ and $(I - A')$, we get for every $j \in \mathcal{C}^{\text{sec}}$ and every i that

$$\overline{U}(i, j) = (\overline{U}(I - A))(i, j) = (\overline{U}(I - A)\overline{V})(i, j) = (I - A')(i, j) = \delta_{ij} = \overline{V}(i, j).$$

Now suppose $i \in \mathcal{C}^{\text{sec}}$. We claim that $\overline{U}(i, j) = \delta_{ij}$. Suppose not. Pick $j \neq i$ such that $\overline{U}(i, j) \neq 0$. Because $\overline{U} = U_3 \overline{H}^{-1}$, it follows from (i) that $\overline{U}(i, j) = \overline{H}^{-1}(i, j)$, and then from (iii) that $j \in \mathcal{C}^{\text{sec}}$. This is a contradiction.

Finally, for $i \in \mathcal{C}^{\text{sec}}$ we obtain

$$\overline{V}(i, j) = ((I - A)\overline{V})(i, j) = (\overline{U}(I - A)\overline{V})(i, j) = (I - A')(i, j) = \delta_{ij}.$$

This finishes the proof. □

References

- [Ba1] L. Badoian, *Flow equivalences of shifts of finite type and flow K-theory*, Ph.D. thesis, University of California, Berkeley, 1998.
- [Ba2] ———, *personal communication*, 2000.
- [Bow] R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Springer Lecture Notes in Math., **470** (1975), Springer-Verlag, MR 56 #1364, Zbl 0308.28010.
- [BowF] R. Bowen and J. Franks, *Homology for zero-dimensional basic sets*, Annals of Math., **106** (1977), 73-92, MR 56 #16692, Zbl 0375.58018.
- [B1] M. Boyle, *Positive K-theory and symbolic dynamics*, in ‘Dynamics and Randomness’ (editors A. Maass, S. Martinez and J. San Martin), Kluwer, 2002, 31-52.
- [B2] ———, *Talk at Mt. Holyoke conference ‘Classification problems in C^* -algebras and dynamics,’* 1996.
- [BFF] M. Boyle, D. Fiebig and U. Fiebig, *A dimension group for local homeomorphisms and endomorphisms of one-sided shifts of finite type*, J. Reine Angew. Math., **487** (1997), 27-59, MR 98i:54020, Zbl 0877.58038.
- [BHa] M. Boyle and D. Handelman, *Orbit equivalence, flow equivalence and ordered cohomology*, Israel J. Math., **95** (1996), 169-210, MR 98a:46082, Zbl 0871.58071.
- [BH] M. Boyle and D. Huang, *Poset block equivalence of integral matrices*, Trans. Amer. Math. Soc., to appear.
- [BW] M. Boyle and J. B. Wagoner, in preparation.
- [C] J. Cuntz, *A class of C^* -algebras and topological Markov chains II: Reducible chains and the Ext-functor for C^* -algebras*, Inventiones Math., **63** (1981), 25-40, MR 82f:46073b, Zbl 0461.46047.
- [CK] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Inventiones Math., **56** (1980), 251-268, MR 82f:46073a, Zbl 0434.46045.
- [DGS] M. Denker, C. Grillenberger and K. Sigmund, *Ergodic theory on compact spaces*, Springer Lecture Notes in Math., **527** (1976), MR 56 #15879, Zbl 0328.28008.
- [F] J. Franks, *Flow equivalence of subshifts of finite type*, Ergod. Th. & Dynam. Sys., **4** (1984), 53-66, MR 86j:58078, Zbl 0555.54026.
- [G] R. Gomez, *Finitary isomorphisms of Markov chains via positive K-theory*, Ph.D. thesis, University of Maryland, College Park, 2000.
- [H1] D. Huang, *Flow equivalence of reducible shifts of finite type*, Ergod. Th. & Dynam. Sys., **14** (1994), 695-720, MR 95k:46110, Zbl 0819.46051.
- [H2] ———, *The classification of two-component Cuntz-Krieger algebras*, Proc. Amer. Math. Soc., **124**(2) (1996), 505-512, MR 96d:46078, Zbl 0846.46040.
- [H3] ———, *Flow equivalence of reducible shifts of finite type and Cuntz-Krieger algebra*, J. Reine. Angew. Math., **462** (1995), 185-217, MR 96m:46123, Zbl 0820.46065.
- [H4] ———, *Automorphisms of Bowen-Franks groups for shifts of finite type*, Ergod. Th. & Dynam. Sys., **21**(4) (2001), 1113-1137, CMP 1 849 604.

- [H5] ———, *The K-web invariant and flow equivalence of reducible shifts of finite type*, in preparation.
- [KR1] K.H. Kim and F.W. Roush, *Free Z_p actions on subshifts*, Pure Math. Appl., **8(2-4)** (1997), 293-322, MR 99f:58066, Zbl 0910.58012.
- [KR2] ———, *The Williams conjecture is false for irreducible subshifts*, Annals of Math., **149(2)** (1999), 545-558, MR 2001b:37012.
- [KRW1] K.H. Kim, F.W. Roush and J.B. Wagoner, *Automorphisms of the dimension group and gyration numbers of automorphisms of a shift*, J. Amer. Math. Soc., **5** (1992), 191-212, MR 93h:54026, Zbl 0749.54012.
- [KRW2] ———, *Characterization of inert actions on periodic points*, Part I, Forum Math., **12** (2000), 565-602, MR 2001g:37009.
- [KRW3] ———, *Characterization of inert actions on periodic points*, Part II, Forum Math., **12(6)** (2000), 671-712, MR 2001j:37024.
- [Ki] B. Kitchens, *Symbolic Dynamics. One-Sided, Two-Sided and Countable State Markov Shifts*, Springer-Verlag, 1998, MR 98k:58079, Zbl 0892.58020.
- [Kr1] W. Krieger, *On a dimension for a class of homeomorphism groups*, Math. Ann., **252** (1980), 239-250, MR 82b:46083, Zbl 0472.54028.
- [Kr2] ———, *On dimension functions and topological Markov chains*, Inventiones Math., **56** (1980), 239-250, MR 81m:28018, Zbl 0431.54024.
- [LM] D. Lind and B. Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge University Press, 1995, MR 97a:58050.
- [Ma] K. Matsumoto, *Presentations of subshifts and their topological conjugacy invariants*, Documenta Math., **4** (1999), 285-340, MR 2000h:37013, Zbl 0926.37002.
- [Ne] M. Newman, *Integral Matrices*, Academic Press, New York, 1972, MR 49 #5038, Zbl 0254.15009.
- [PS] W. Parry and D. Sullivan, *A topological invariant for flows on one-dimensional spaces*, Topology, **14** (1975), 297-299, MR 53 #9179, Zbl 0314.54045.
- [PT] W. Parry and S. Tuncel, *Classification problems in ergodic theory*, LMS Lecture Note Series, **67**, Cambridge Press, Cambridge, 1982, MR 84g:28024, Zbl 0487.28014.
- [Rob] C. Robinson, *Dynamical systems. Stability, symbolic dynamics, and chaos*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1995, MR 97e:58064, Zbl 0914.58021.
- [R] M. Rørdam, *Classification of Cuntz-Krieger algebras*, K-theory, **9** (1995), 31-58, MR 96k:46103, Zbl 0826.46064.
- [Ros] J. Rosenberg, *Algebraic K-theory and its applications*, Graduate Texts in Mathematics, **147**, Springer-Verlag, 1994, MR 95e:19001, Zbl 0801.19001.
- [S] S. Smale, *Differentiable dynamical systems*, Bull. Amer. Math. Soc., **73** (1967), 747-817, MR 37 #3598, Zbl 0202.55202.
- [W1] J.B. Wagoner, *Strong shift equivalence theory and the shift equivalence problem*, Bull. Amer. Math. Soc. (N.S.), **36(3)** (1999), 271-296, MR 2001e:54061, Zbl 0927.19001.
- [W2] ———, *Strong shift equivalence and K_2 of the dual numbers*, J. Reine Angew. Math., **521** (2000), 119-160, MR 2001i:19001, Zbl 0969.37003.

[Wi] R.F. Williams, *Classification of subshifts of finite type*, Annals of Math., **98** (1973), 120-153; *Erratum*, Annals of Math., **99** (1974), 380-381, MR 48 #9769, Zbl 0282.58008.

Received September 20, 2000 and revised December 1, 2000. The research of the author was supported by NSF Grant DMS9706852.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MARYLAND
COLLEGE PARK, MD 20742-4015
E-mail address: mmb@math.umd.edu

