

*Pacific
Journal of
Mathematics*

GENERALIZED FOCK SPACES AND WEYL
COMMUTATION RELATIONS FOR THE DUNKL KERNEL

FETHI SOLTANI

Volume 214 No. 2

April 2004

GENERALIZED FOCK SPACES AND WEYL COMMUTATION RELATIONS FOR THE DUNKL KERNEL

FETHI SOLTANI

In this paper we study a class of generalized Fock spaces associated with the Dunkl operators. Next we introduce the commutator relations between the Dunkl operators and multiplication operators which lead to a generalized class of Weyl commutation relations for the Dunkl kernel.

1. Introduction.

Fock space (called also Segal-Bargmann space [5]) is the Hilbert space of entire functions of \mathbb{C}^d with inner product given by

$$(f, g) := \frac{1}{\pi^d} \int_{\mathbb{C}^d} f(z) \overline{g(z)} e^{-|z|^2} dx dy, \quad z = x + iy,$$

where

$$|z|^2 = \sum_{i=1}^d x_i^2 + y_i^2, \quad dx dy = \prod_{i=1}^d dx_i dy_i.$$

This space which associated with Fock's [12] realization of the creation and annihilation operators of Bose particles is studied by Bargmann [4]. Next, the ordinary Fock space \mathcal{A} is the subject of many works ([5, 7] and [8]).

In 2001, M. Sifi and F. Soltani [21] introduced a Hilbert space \mathcal{A}_γ of entire functions where the inner product is weighted by a generalized Gaussian distribution. On \mathcal{A}_γ the Dunkl operator on the real line:

$$T_\gamma(f)(z) := \frac{d}{dz} f(z) + \frac{2\gamma}{z} \left[\frac{f(z) - f(-z)}{2} \right], \quad \gamma > 0,$$

and the multiplication by z are adjoints and satisfy the commutation rule

$$[T_\gamma, z] = I + 2\gamma B, \quad \text{where } Bf(x) = f(-x).$$

This commutator rule leads to a generalized class of Weyl commutation relations for the Dunkl kernel in the one dimensional.

In this paper we consider the differential-difference operators T_j , $j = 1, \dots, d$, on \mathbb{R}^d introduced by C.F. Dunkl in [9] and called Dunkl operators in the literature. These operators are very important in pure Mathematics and in Physics. They provide a useful tool in the study of special functions associated with root systems [10]. They are closely related to certain

representations of degenerated affine Hecke algebras ([6] and [16]). Moreover the commutative algebra generated by these operators has been used in the study of certain exactly solvable models of quantum Mechanics, namely the Calogero-Sutherland-Moser models, which deal with systems of identical particles in the one dimensional space ([2, 3] and [14]).

The Dunkl kernel $E_k(x, y)$ is the unique solution of the initial problem

$$T_j^x u(x, y) = y_j u(x, y); \quad u(0, y) = 1; \quad j = 1, \dots, d,$$

see [10, 17] and [18]. This kernel has a unique holomorphic extension to $\mathbb{C}^d \times \mathbb{C}^d$. Furthermore, the Dunkl kernel $E_k(z, w); z, w \in \mathbb{C}^d$ can be expanded in a power series in the form

$$E_k(z, w) = \sum_{\nu \in \mathbb{N}^d} \varphi_\nu(z) \varphi_\nu(w),$$

with some homogeneous orthonormal basis $\{\varphi_\nu\}_{\nu \in \mathbb{N}^d}$ of polynomials ([17] and [19]).

We introduce in this paper the generalized Fock space \mathcal{A}_k associated with the Dunkl operators. This is a Hilbert space of functions f on \mathbb{C}^d which can be written $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z)$ with

$$\|f\|_k^2 = (f, f)_k := \sum_{\nu \in \mathbb{N}^d} |a_\nu|^2 < \infty.$$

If $f, g \in \mathcal{A}_k$, having series expansions $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z)$ and $g(z) = \sum_{\nu \in \mathbb{N}^d} b_\nu \varphi_\nu(z)$. Then the inner product is given by the generalized spherical harmonics

$$(f, g)_k = \left(f(T) \tilde{g} \right) (0),$$

where $f(T) = f(T_1, \dots, T_d)$ and $\tilde{g}(z) = \sum_{\nu \in \mathbb{N}^d} \bar{b}_\nu \varphi_\nu(z)$.

The generalized Fock space \mathcal{A}_k , has also a reproducing kernel \mathcal{K} given by $\mathcal{K}(z, w) = E_k(z, \bar{w}); z, w \in \mathbb{C}^d$. If $f \in \mathcal{A}_k$, then we have

$$f(w) = (f, E_k(\cdot, \bar{w}))_k, \quad w \in \mathbb{C}^d.$$

Thus the Dunkl kernel serves as the generalized Dirac delta function in \mathcal{A}_k .

The associated operators for the generalized Fock space \mathcal{A}_k are T_j and the multiplication operator by z_j . They are adjoints in \mathcal{A}_k and satisfy a commutation rule:

$$[T_i, z_j] = \delta_{i,j} I + \sum_{\alpha \in R_+} k(\alpha) \alpha_i \alpha_j B_\alpha; \quad i, j = 1, \dots, d,$$

where B_α a reflection operator, $k(\alpha)$ a multiplicity function and R_+ is a positive root system.

These commutators rule lead to a generalized class of Weyl commutation relations for the Dunkl kernel.

These relations are studied in the classical case ($k = 0$) in [13].

Throughout this paper we shall use the standard multi-index notations. This for multi-indices $\nu, s \in \mathbb{N}^d$, we write $|\nu| = \sum_{i=1}^d \nu_i$, $\nu! = \prod_{i=1}^d \nu_i!$, $\binom{\nu}{s} = \prod_{i=1}^d \binom{\nu_i}{s_i}$ as well as $z^\nu = \prod_{i=1}^d z_i^{\nu_i}$, $D^\nu = \prod_{i=1}^d D_i^{\nu_i}$, for $z = (z_1, \dots, z_d) \in \mathbb{C}^d$ and any family $D = (D_1, \dots, D_d)$ of commuting operators. Finally, we will need the partial ordering \leq on \mathbb{N}^d , which is defined by $s \leq \nu \Leftrightarrow s_i \leq \nu_i, i = 1, \dots, d$.

2. Preliminaries.

In this section we collect some notations and results on Dunkl operators and Dunkl kernel that will be important later on. General references here are [9, 17, 18, 19] and [20].

We consider \mathbb{R}^d with the Euclidean scalar $\langle \cdot, \cdot \rangle$ and $|x| = \sqrt{\langle x, x \rangle}$. On \mathbb{C}^d , $|\cdot|$ denotes also the standard Hermitian norm, while $\langle z, w \rangle = \sum_{j=1}^d z_j w_j$ and $\ell(z) = \langle z, z \rangle$.

For $\alpha \in \mathbb{R}^d \setminus \{0\}$, let σ_α be the reflection in the hyperplane $H_\alpha \subset \mathbb{R}^d$ orthogonal to α ,

$$\sigma_\alpha x := x - \frac{2\langle \alpha, x \rangle}{|\alpha|^2} \alpha.$$

A finite set $R \subset \mathbb{R}^d \setminus \{0\}$ is called a root system if $R \cap \mathbb{R}\alpha = \{-\alpha, \alpha\}$ and $\sigma_\alpha R = R$ for all $\alpha \in R$. We assume that it is normalized by $|\alpha|^2 = 2$ for all $\alpha \in R$. For a given root system R the reflections $\sigma_\alpha, \alpha \in R$ generated a finite group $G \subset O(d)$, the reflection group associated with R . All reflections in G correspond to suitable pairs of roots. For a given $\beta \in H = \mathbb{R}^d \setminus \bigcup_{\alpha \in R} H_\alpha$, we fix the positive subsystem $R_+ = \{\alpha \in R / \langle \alpha, \beta \rangle > 0\}$, then for each $\alpha \in R$ either $\alpha \in R_+$ or $-\alpha \in R_+$. The connected components of H are called the Weyl chambers of G .

A function $k : R \rightarrow \mathbb{C}$ on a root system R is called a multiplicity function if it is invariant under the action of the associated reflection group G . If one regards k as a function on the corresponding reflections, this means that k is constant on the conjugacy classes of reflections in G . For abbreviation, we introduce the index

$$\gamma = \gamma(k) := \sum_{\alpha \in R_+} k(\alpha).$$

Moreover, let w_k denotes the weight function:

$$w_k(x) := \prod_{\alpha \in R_+} |\langle \alpha, x \rangle|^{2k(\alpha)}, \quad x \in \mathbb{R}^d,$$

which is G -invariant and homogeneous of degree 2γ .

For $d = 1$ and $G = \mathbb{Z}_2$, the multiplicity function k is a simple parameter denoted $\gamma > 0$ and

$$w_k(x) = |x|^{2\gamma}, \quad x \in \mathbb{R}.$$

The Dunkl operators T_j ; $j = 1, \dots, d$, on \mathbb{R}^d associated with the finite reflection group G and multiplicity function k are given for a function f of class C^1 on \mathbb{R}^d , by

$$T_j f(x) := \frac{\partial}{\partial x_j} f(x) + \sum_{\alpha \in R_+} k(\alpha) \alpha_j \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle}.$$

In the case $k = 0$, the T_j ; $j = 1 \dots, d$, reduce to the corresponding partial derivatives. In this paper we will assume throughout that $k \geq 0$.

For $y \in \mathbb{R}^d$, the initial problem

$$\begin{cases} T_j^x u(x, y) = y_j u(x, y); & j = 1, \dots, d, \\ u(0, y) = 1, \end{cases}$$

admits a unique analytic solution on \mathbb{R}^d , which will be denoted $E_k(x, y)$ and called the Dunkl kernel ([17, 18, 19] and [20]). This kernel has a unique holomorphic extension to $\mathbb{C}^d \times \mathbb{C}^d$.

Examples.

- 1) If $k = 0$, then $E_k(z, w) = e^{\langle z, w \rangle}$ for $z, w \in \mathbb{C}^d$. (Recall that $\langle \cdot, \cdot \rangle$ was defined to be bilinear on $\mathbb{C}^d \times \mathbb{C}^d$.)
- 2) If $d = 1$ and $G = \mathbb{Z}_2$, the Dunkl kernel is given by

$$E_\gamma(z, w) = \mathfrak{S}_{\gamma-\frac{1}{2}}(zw) + \frac{zw}{2\gamma+1} \mathfrak{S}_{\gamma+\frac{1}{2}}(zw),$$

where

$$\mathfrak{S}_{\gamma-\frac{1}{2}}(zw) = \Gamma\left(\gamma + \frac{1}{2}\right) \sum_{n=0}^{\infty} \frac{1}{n! \Gamma(n + \gamma + \frac{1}{2})} \left(\frac{zw}{2}\right)^{2n},$$

is the modified Bessel function of order $\gamma - \frac{1}{2}$ [21].

Let $\mathcal{P} = \mathbb{C}[\mathbb{R}^d]$ denotes the \mathbb{C} - Algebra of polynomial functions on \mathbb{R}^d and \mathcal{P}_n , $n \in \mathbb{N}$, the subspace of homogeneous polynomials of degree n . In the context of generalized spherical harmonics, C.F. Dunkl in [9] introduced on \mathcal{P} the following bilinear form:

$$(1) \quad (p, q)_k := (p(T)q)(0); \quad p, q \in \mathcal{P}.$$

Here $p(T)$ is the operator derived from $p(x)$ by replacing x_i by T_i . A useful collection of its properties can be found in [9] and [17]. We recall that $(\cdot, \cdot)_k$ is symmetric, positive-definite and $(p, q)_k = 0$, for $p \in \mathcal{P}_n$, $q \in \mathcal{P}_m$ with $n \neq m$. Moreover, for all $j = 1, \dots, d$ and $p, q \in \mathcal{P}$,

$$(x_j p, q)_k = (p, T_j q)_k.$$

Let $\{\varphi_\nu\}_{\nu \in \mathbb{N}^d}$ be an orthonormal basis of \mathcal{P} with respect to the scalar product $(\cdot, \cdot)_k$ such that $\varphi_\nu \in \mathcal{P}_{|\nu|}$ and the coefficients of the φ_ν are real. As $\mathcal{P} = \bigoplus_{n \in \mathbb{N}} \mathcal{P}_n$ and $\mathcal{P}_n \perp \mathcal{P}_m$ for $n \neq m$, the φ_ν with $|\nu| = n$ can for example be constructed by Gram-Schmidt orthogonalization within \mathcal{P}_n from an arbitrary ordered real-coefficients basis of \mathcal{P}_n . If $k = 0$ the canonical choice of the homogeneous orthonormal basis φ_ν is just $\varphi_\nu(x) = \frac{x^\nu}{\sqrt{\nu!}}$.

As in the classical case, M. Rösler obtained in [17, p. 524] the following connection of the basis φ_ν with the Dunkl kernel:

$$(2) \quad E_k(z, w) = \sum_{\nu \in \mathbb{N}^d} \varphi_\nu(z)\varphi_\nu(w); \quad z, w \in \mathbb{C}^d,$$

where the convergence is normal on $\mathbb{C}^d \times \mathbb{C}^d$.

Example. If $d = 1$ and $G = \mathbb{Z}_2$ every homogeneous orthonormal basis is of the form

$$(3) \quad \varphi_n(z) = \frac{z^n}{\sqrt{b_n(\gamma)}}, \quad b_n(\gamma) = \frac{2^n([n/2])!}{\Gamma(\gamma + \frac{1}{2})} \Gamma\left(\left[\frac{n+1}{2}\right] + \gamma + \frac{1}{2}\right).$$

Here $[n/2]$ is the integer part of $n/2$.

From (2), the Dunkl kernel E_k possesses the following properties ([17, 19] and [20]): For all $z, w \in \mathbb{C}^d$ and $\lambda \in \mathbb{C}$,

$$(4) \quad E_k(z, w) = E_k(w, z), \quad E_k(\lambda z, w) = E_k(z, \lambda w),$$

$$(5) \quad \overline{E_k(z, w)} = E_k(\bar{z}, \bar{w}), \quad E_k(z, \bar{z}) = \sum_{\nu \in \mathbb{N}^d} |\varphi_\nu(z)|^2,$$

$$(6) \quad |E_k(z, w)| \leq e^{|z||w|}.$$

In [18], M. Rösler establish the Bochner-type representation of the Dunkl kernel

$$(7) \quad E_k(x, z) = \int_{\mathbb{R}^d} e^{\langle \xi, z \rangle} d\mu_x(\xi); \quad x \in \mathbb{R}^d, \quad z \in \mathbb{C}^d,$$

where μ_x is a probability measure on \mathbb{R}^d with support in $\{\xi \in \mathbb{R}^d / |\xi| \leq |x|\}$.

The Dunkl kernel E_k is analytic on $\mathbb{C}^d \times \mathbb{C}^d$. Therefore, there exist unique analytic functions m_ν , $\nu \in \mathbb{N}^d$, on \mathbb{C}^d with

$$(8) \quad E_k(z, w) = \sum_{\nu \in \mathbb{N}^d} \frac{m_\nu(z)}{\nu!} w^\nu; \quad z, w \in \mathbb{C}^d.$$

The restriction of m_ν to \mathbb{R}^d are called the ν -th moment functions ([18, 19] and [20]). It is given explicitly by

$$m_\nu(x) = \int_{\mathbb{R}^d} \xi^\nu d\mu_x(\xi), \quad x \in \mathbb{R}^d,$$

where μ_x is the measure given by (7).

The functions m_ν are homogeneous polynomials of degree $|\nu|$. Among the applications of these moments, we mention the construction of martingales from Dunkl-type Markov processes [19].

3. Fock spaces for the Dunkl kernel.

In this section we define and study the generalized Fock space for the Dunkl kernel in d -dimensions.

Definition 1. The generalized Fock space \mathcal{A}_k associated with the Dunkl operators is the space of holomorphic functions f on \mathbb{C}^d which can be written $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z)$ with

$$\|f\|_k^2 := \sum_{\nu \in \mathbb{N}^d} |a_\nu|^2 < \infty.$$

Hence the inner product in \mathcal{A}_k is given for $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z) \in \mathcal{A}_k$ and $g(z) = \sum_{\nu \in \mathbb{N}^d} b_\nu \varphi_\nu(z) \in \mathcal{A}_k$, by

$$(9) \quad (f, g)_k := \sum_{\nu \in \mathbb{N}^d} a_\nu \bar{b}_\nu.$$

Remark. If $k = 0$, \mathcal{A}_0 is the ordinary Fock space \mathcal{A} [4].

Proposition 1.

- i) If $f, g \in \mathcal{A}_k$ with $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z)$ and $g(z) = \sum_{\nu \in \mathbb{N}^d} b_\nu \varphi_\nu(z)$, we have

$$(f, g)_k = (f(T)\tilde{g})(0),$$

where $\tilde{g}(z) = \sum_{\nu \in \mathbb{N}^d} \bar{b}_\nu \varphi_\nu(z)$.

- ii) If $f \in \mathcal{A}_k$ with $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z)$, we have

$$|f(z)| \leq e^{|z|^2/2} \|f\|_k.$$

Proof. i) From [17, p. 529], we have

$$(\varphi_\nu(T)\varphi_s)(0) = \delta_{\nu,s},$$

where $\delta_{\nu,s}$ is the Kronecker symbol.

Thus

$$(f, g)_k = \sum_{\nu, s \in \mathbb{N}^d} a_\nu \bar{b}_s (\varphi_\nu(T)\varphi_s)(0).$$

Using the continuity of the inner product, we obtain the result.

- ii) Using Cauchy-Schwarz's inequality, then

$$|f(z)|^2 \leq \left[\sum_{\nu \in \mathbb{N}^d} |a_\nu|^2 \right] \left[\sum_{\nu \in \mathbb{N}^d} |\varphi_\nu(z)|^2 \right] = \|f\|_k^2 E_k(z, \bar{z}).$$

Thus

$$|f(z)| \leq [E_k(z, \bar{z})]^{1/2} \|f\|_k.$$

The result follows from the inequality (6). □

From Proposition 1 ii), the map $f \rightarrow f(z)$, $z \in \mathbb{C}^d$, is a continuous linear functional on \mathcal{A}_k . Thus from Riesz theorem [1], \mathcal{A}_k has a reproducing kernel.

Proposition 2. *The function \mathcal{K} given for $w, z \in \mathbb{C}^d$, by*

$$\mathcal{K}(z, w) = E_k(z, \bar{w}),$$

is a reproducing kernel for the generalized Fock spaces \mathcal{A}_k , that is:

- i) *For every $w \in \mathbb{C}^d$, the function $z \rightarrow \mathcal{K}(z, w)$ belongs to \mathcal{A}_k .*
- ii) *The reproducing property: For every $w \in \mathbb{C}^d$ and $f \in \mathcal{A}_k$, we have*

$$(f, \mathcal{K}(\cdot, w))_k = f(w).$$

Proof. i) Using (5) and (6), we deduce for $w \in \mathbb{C}^d$,

$$\|E_k(\cdot, \bar{w})\|_k^2 = \sum_{\nu \in \mathbb{N}^d} |\varphi_\nu(\bar{w})|^2 = E_k(w, \bar{w}) \leq e^{|w|^2},$$

which proves i).

ii) If $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z) \in \mathcal{A}_k$, it follows from (9) that

$$(f, E_k(\cdot, \bar{w}))_k = \sum_{\nu \in \mathbb{N}^d} a_\nu \overline{\varphi_\nu(\bar{w})} = f(w).$$

□

Corollary 1.

- i) *The set $\{E_k(\cdot, \bar{w}), w \in \mathbb{C}^d\}$ is complete in \mathcal{A}_k .*
- ii) *For all $z, w \in \mathbb{C}^d$, we have*

$$E_k(\bar{z}, w) = (E_k(\cdot, \bar{z}), E_k(\cdot, \bar{w}))_k.$$

iii) *Let $m \in \mathbb{N} \setminus \{0\}$ and $z_1, z_2, \dots, z_m \in \mathbb{C}^d$, with $z_i \neq z_j$, then*

$$\det [E_k(\bar{z}_i, z_j)]_{i,j=1}^m > 0.$$

Notation. We denote by $L^2(\mu_k)$ the Hilbert space of measurable functions on \mathbb{R}^d , for which

$$\|f\|_{2,k} := \left[\int_{\mathbb{R}^d} |f(x)|^2 d\mu_k(x) \right]^{1/2} < \infty.$$

Here μ_k is the measure defined on \mathbb{R}^d , by

$$d\mu_k(x) := c_k w_k(x) dx, \quad \text{with } c_k = \left(\int_{\mathbb{R}^d} e^{-|x|^2} d\mu_k(x) \right)^{-1},$$

is the Mehta-type constant.

In the next part of this section we establish the unitary equivalence of $L^2(\mu_k)$ and \mathcal{A}_k . First we recall some properties of the generalized Hermite functions ([17] and [19]):

Definition 2. The generalized Hermite polynomials $\{H_\nu\}_{\nu \in \mathbb{N}^d}$ associated with the basis $\{\varphi_\nu\}_{\nu \in \mathbb{N}^d}$ on \mathbb{C}^d , are given by

$$H_\nu(z) := 2^{|\nu|} e^{-\Delta_k/4} \varphi_\nu(z) = 2^{|\nu|} \sum_{n=0}^{[\nu/2]} \frac{(-1)^n}{2^{2n} n!} \Delta_k^n \varphi_\nu(z),$$

where $\Delta_k = \sum_{i=1}^d T_i^2$ is the Dunkl Laplacian [17].

Moreover, we define the generalized Hermite functions on \mathbb{C}^d , by

$$h_\nu(z) := 2^{-|\nu|/2} e^{-\ell(z)/2} H_\nu(z).$$

Examples.

1) If $k = 0$, we obtain

$$H_\nu(x) = \frac{2^{|\nu|}}{\sqrt{\nu!}} \prod_{i=1}^d e^{-\frac{1}{4} \frac{\partial^2}{\partial x_i^2}} (x_i^{\nu_i}) = \frac{1}{\sqrt{\nu!}} \prod_{i=1}^d \widehat{H}_{\nu_i}(x_i), \quad x \in \mathbb{R}^d,$$

where

$$\widehat{H}_{x_i} = (-1)^{\nu_i} e^{x_i^2} \frac{\partial^{\nu_i}}{\partial x_i^{\nu_i}} (e^{-x_i^2}).$$

2) If $d = 1$ and $G = \mathbb{Z}_2$, we obtain

$$H_n(z) = \sum_{i=0}^{[n/2]} \frac{(-1)^i}{i! b_{n-2i}(\gamma)} (2x)^{n-2i}, \quad x \in \mathbb{R},$$

where $b_n(\gamma)$ are the constants given by (3).

The following lemma is shown in [17, p. 525-529]:

Lemma 1.

- i) The set $\{h_\nu\}_{\nu \in \mathbb{N}^d}$ is an orthonormal basis of $L^2(\mu_k)$.
- ii) For all $z, w \in \mathbb{C}^d$, there is a generating function for the generalized Hermite polynomials,

$$e^{-\ell(w)} E_k(2z, w) = \sum_{\nu \in \mathbb{N}^d} h_\nu(z) \varphi_\nu(w).$$

Notation. We denote by U_k the kernel given for $z, w \in \mathbb{C}^d$, by

$$(10) \quad U_k(z, w) := e^{-(\ell(z)+\ell(w))/2} E_k(\sqrt{2}z, w).$$

Lemma 2. For $w, z \in \mathbb{C}^d$, we have

$$U_k(z, w) = \sum_{\nu \in \mathbb{N}^d} h_\nu(z) \varphi_\nu(w).$$

Proof. From Definition 2, we have

$$\sum_{\nu \in \mathbb{N}^d} h_\nu(z) \varphi_\nu(w) = e^{-\ell(z)/2} \sum_{\nu \in \mathbb{N}^d} 2^{-|\nu|/2} H_\nu(z) \varphi_\nu(w).$$

As φ_ν is homogeneous of degree $|\nu|$, then

$$\varphi_\nu\left(\frac{w}{\sqrt{2}}\right) = 2^{-|\nu|/2} \varphi_\nu(w).$$

Thus

$$\sum_{\nu \in \mathbb{N}^d} h_\nu(z) \varphi_\nu(w) = e^{-\ell(z)/2} \sum_{\nu \in \mathbb{N}^d} H_\nu(z) \varphi_\nu\left(\frac{w}{\sqrt{2}}\right).$$

Applying Lemma 1 ii) and (4), we obtain

$$\sum_{\nu \in \mathbb{N}^d} h_\nu(z) \varphi_\nu(w) = e^{-(\ell(z)+\ell(w))/2} E_k\left(2z, \frac{w}{\sqrt{2}}\right) = U_k(z, w).$$

□

Lemma 3.

i) For all $z, w \in \mathbb{C}^d$, we have

$$E_k(z, w) = \int_{\mathbb{R}^d} U_k(z, x) U_k(w, x) d\mu_k(x).$$

ii) For all $z \in \mathbb{C}^d$, the function $x \rightarrow U_k(z, x)$ belongs to $L^2(\mu_k)$, and we have

$$\|U_k(z, \cdot)\|_{2,k}^2 = E_k(z, \bar{z}).$$

iii) For all $x \in \mathbb{R}^d$, the function $z \rightarrow U_k(z, x)$ belongs to \mathcal{A}_k , and we have

$$\|U_k(\cdot, x)\|_k^2 = e^{-3|x|^2} E_k(2x, x).$$

Proof. i) We put

$$I = \int_{\mathbb{R}^d} U_k(z, x) U_k(w, x) d\mu_k(x).$$

From (10), we have

$$I = e^{-(\ell(z)+\ell(w))/2} \int_{\mathbb{R}^d} e^{-|x|^2} E_k(\sqrt{2}z, x) E_k(\sqrt{2}w, x) d\mu_k(x).$$

So from [17, p. 523] and (4), we get

$$\int_{\mathbb{R}^d} e^{-|x|^2} E_k(\sqrt{2}z, x) E_k(\sqrt{2}w, x) d\mu_k(x) = e^{(\ell(z)+\ell(w))/2} E_k(z, w),$$

which proves i).

ii) This assertion follows from i) and (5).

iii) For $z \in \mathbb{C}^d$, we put

$$\phi(z) := e^{-(\ell(z)+\ell(\bar{z}))/2}.$$

Let $x \in \mathbb{R}^d$, then from Proposition 2 ii), (10) and (4), we have

$$\|U_k(\cdot, x)\|_k^2 = e^{-|x|^2} (\phi(\cdot)E_k(\cdot, \sqrt{2x}), E_k(\cdot, \sqrt{2x}))_k = e^{-3|x|^2} E_k(2x, x).$$

□

Definition 3. The chaotic transform \mathcal{C}_k (also called S -transform in the stochastic calculus [15]) is the transformation defined on $L^2(\mu_k)$, by

$$\mathcal{C}_k(f)(z) := \int_{\mathbb{R}^d} U_k(z, x)f(x)d\mu_k(x), \quad z \in \mathbb{C}^d.$$

Remark. The basis elements of $L^2(\mu_k)$ and \mathcal{A}_k are called chaos. In the following theorem we shall prove that the transformation \mathcal{C}_k maps the chaos of $L^2(\mu_k)$ to these of \mathcal{A}_k .

Theorem 1. *The chaotic transform \mathcal{C}_k is a unitary mapping of $L^2(\mu_k)$ on \mathcal{A}_k . Moreover, the basis elements are related by*

$$\mathcal{C}_k(h_\nu) = \varphi_\nu.$$

Proof. It follows directly from Lemma 1 i) and Lemma 2, that for $\nu \in \mathbb{N}^d$,

$$\mathcal{C}(h_\nu)(z) = \int_{\mathbb{R}^d} U_k(z, x)h_\nu(x)d\mu_k(x) = \varphi_\nu(z), \quad z \in \mathbb{C}^d.$$

Consequently \mathcal{C}_k maps the subspace generated by the family $\{h_\nu\}_{\nu \in \mathbb{N}^d}$ into the polynomials in \mathcal{A}_k . Thus \mathcal{C}_k maps a dense set in $L^2(\mu_k)$ into a dense set in \mathcal{A}_k . Further, if $f \in L^2(\mu_k)$, then $f(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu h_\nu(x)$. For $\nu \in \mathbb{N}^d$, let $f_N(x) = \sum_{j=0}^N \sum_{|\nu|=j} a_\nu h_\nu(x)$, $x \in \mathbb{R}$. Then

$$\mathcal{C}_k(f_N)(z) = \sum_{j=0}^N \sum_{|\nu|=j} a_\nu \varphi_\nu(z); \quad \lim_{N \rightarrow \infty} \|f - f_N\|_{2,k} = 0.$$

On the other hand, from Hölder’s inequality and Lemma 3 ii), we have

$$|\mathcal{C}_k(f - f_N)(z)| \leq [E_k(z, \bar{z})]^{1/2} \|f - f_N\|_{2,k}.$$

Thus we obtain

$$\mathcal{C}_k(f)(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z).$$

Hence

$$\|\mathcal{C}_k(f)\|_k^2 = \sum_{\nu \in \mathbb{N}^d} |a_\nu|^2 = \|f\|_{2,k}^2.$$

It follows that \mathcal{C}_k is a unitary transformation from $L^2(\mu_k)$ into \mathcal{A}_k .

Clearly, if $g(z) = \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z) \in \mathcal{A}_k$, we have

$$(11) \quad \mathcal{C}_k^{-1}(g)(x) = \sum_{\nu \in \mathbb{N}^d} a_\nu h_\nu(x), \quad x \in \mathbb{R}^d.$$

Which completes the proof. □

Proposition 3. *If $g \in \mathcal{A}_k$, we have*

$$\mathcal{C}_k^{-1}(g)(x) = (g, U_k(\cdot, x))_k, \quad x \in \mathbb{R}^d.$$

Proof. Let $g \in \mathcal{A}_k$. We put for $x \in \mathbb{R}^d$,

$$\Psi_k(g)(x) = (g, U_k(\cdot, x))_k.$$

Using Lemma 2, Lemma 3 iii) and the same method as in the proof of Theorem 1 we obtain

$$\Psi_k(g)(x) = \sum_{\nu \in \mathbb{N}^d} a_\nu h_\nu(x) = \mathcal{C}_k^{-1}(g)(x), \quad x \in \mathbb{R}^d.$$

□

4. Commutators and Weyl relations for the Dunkl kernel.

We define the multiplication operators Q_i ; $i = 1, \dots, d$ on \mathcal{A}_k by

$$Q_i f(z) := z_i f(z), \quad z \in \mathbb{C}^d.$$

We denote also by T_i ; $i = 1, \dots, d$ the operators defined on \mathcal{A}_k .

Let

$$\begin{aligned} \mathcal{D}(Q_i) &= \{f \in \mathcal{A}_k / Q_i(f) \in \mathcal{A}_k\}, \\ \mathcal{D}(T_i) &= \{f \in \mathcal{A}_k / T_i f \in \mathcal{A}_k\} \end{aligned}$$

denote the domains of Q_i and T_i respectively.

We denote by $[\cdot, \cdot]$ the commutator product ($[A, B] = AB - BA$). As in [11], we have the following relations:

Lemma 4. *The operators Q_i and T_i , $i = 1, \dots, d$ satisfy on \mathcal{A}_k the commutation relations:*

$$(12) \quad [T_i, T_j] = [Q_i, Q_j] = 0; \quad i, j = 1, \dots, d,$$

$$(13) \quad [T_i, Q_j] = \delta_{ij} I + \sum_{\alpha \in R_+} k(\alpha) \alpha_i \alpha_j B_\alpha; \quad i, j = 1, \dots, d,$$

where I the identity operator and B_α is the reflection operator ($B_\alpha^2 = I$) given by

$$(14) \quad B_\alpha f(z) = f(\sigma_\alpha z).$$

Proof. Using the fact that $\sigma_\alpha^2 = id$ and $\langle \alpha, \sigma_\alpha z \rangle = -\langle \alpha, z \rangle$, we obtain

$$T_i T_j f(z) = T_i \left(\frac{\partial}{\partial z_j} f \right) (z) + \sum_{\alpha \in R_+} k(\alpha) \alpha_j \frac{\partial}{\partial z_i} \left(\frac{f(z) - f(\sigma_\alpha z)}{\langle \alpha, z \rangle} \right).$$

Since

$$\frac{\partial}{\partial z_i} (f(\sigma_\alpha z)) = \frac{\partial}{\partial z_i} f(\sigma_\alpha z) - \sum_{\ell=1}^d \alpha_i \alpha_\ell \frac{\partial}{\partial z_\ell} f(\sigma_\alpha z),$$

we have

$$T_i T_j f(z) = -\frac{\partial^2}{\partial z_j \partial z_i} f(z) + T_i \left(\frac{\partial}{\partial z_j} f \right) (z) + T_j \left(\frac{\partial}{\partial z_i} f \right) (z) - \sum_{\alpha \in R_+} k(\alpha) \alpha_i \alpha_j \left[\frac{f(z) - f(\sigma_\alpha z)}{(\langle \alpha, z \rangle)^2} - \sum_{\ell=1}^d \alpha_\ell \frac{\frac{\partial}{\partial z_\ell} f(\sigma_\alpha z)}{\langle \alpha, z \rangle} \right].$$

Thus

$$[T_i, T_j]f(z) = 0.$$

The other relations are evident. □

Proposition 4. *Let*

$$\begin{aligned} f(z) &= \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z) \in \mathcal{D}(Q_i) && \text{and} \\ g(z) &= \sum_{\nu \in \mathbb{N}^d} a_\nu \varphi_\nu(z) \in \mathcal{D}(T_i), && \text{then} \\ (Q_i f, g)_k &= (f, T_i g)_k. \end{aligned}$$

Proof. Applying Proposition 1 i), we get

$$(Q_i f, g)_k = (Q_i f(T)\tilde{g})(0) = \sum_{\nu, s \in \mathbb{N}^d} a_\nu \bar{b}_s T_i \varphi_\nu(T) \varphi_s(0).$$

Then from (12) we obtain

$$(Q_i f, g)_k = \sum_{\nu, s \in \mathbb{N}^d} a_\nu \bar{b}_s \varphi_\nu(T) T_i \varphi_s(0) = (f, T_i g)_k. \quad \square$$

Lemma 5. *If $f \in \mathcal{A}_k$, then $B_\alpha f \in \mathcal{A}_k$, and we have*

$$\|Q_i f\|_k^2 = \|T_i f\|_k^2 + \|f\|_k^2 + \sum_{\alpha \in R_+} k(\alpha) \alpha_i^2 (f, B_\alpha f)_k,$$

where B_α is the operator given by (14).

Proof. Let $f \in \mathcal{A}_k$. Applying the chaotic transform, in view of Theorem 1, it suffices to show that $\mathcal{C}_k^{-1}(B_\alpha f) \in L^2(\mu_k)$. From (11), we have

$$\mathcal{C}_k^{-1}(B_\alpha f)(x) = \mathcal{C}_k^{-1}(f)(\sigma_\alpha x), \quad x \in \mathbb{R}^d.$$

Putting $u = \sigma_\alpha x$, we get

$$d\mu_k(x) = |J_\alpha| d\mu_k(u) \quad \text{where} \quad J_\alpha = \det \left[\delta_{i,j} - \alpha_i \alpha_j \right]_{i,j=1}^d.$$

Since $J_\alpha = -1$, we obtain

$$\|\mathcal{C}_k^{-1}(B_\alpha f)\|_{2,k}^2 = \int_{\mathbb{R}^d} |\mathcal{C}_k^{-1}(f)(u)|^2 d\mu_k(u).$$

Which proves that $B_\alpha f \in \mathcal{A}_k$.

On the other hand, from Proposition 4, we deduce

$$\|Q_i f\|_k^2 = (f, T_i Q_i f)_k.$$

But from (13), we have

$$T_i Q_i f = Q_i T_i f + f + \sum_{\alpha \in R_+} k(\alpha) \alpha_i^2 B_\alpha f.$$

Thus

$$\|Q_i f\|_k^2 = (f, Q_i T_i f)_k + \|f\|_k^2 + \sum_{\alpha \in R_+} k(\alpha) \alpha_i^2 (f, B_\alpha f)_k.$$

Using another time Proposition 4, we obtain the result. □

Proposition 5. *The operators Q_i and T_i are closed densely defined operators on \mathcal{A}_k , and we have*

$$\mathcal{D}(Q_i) = \mathcal{D}(T_i); \quad Q_i^* = T_i; \quad T_i^* = Q_i,$$

where Q_i^* and T_i^* are the adjoints operators of Q_i and T_i , respectively.

Proof. These results follow from [4, Theorem 1.2], Lemma 5 and Proposition 4 by using the same method as [21, Proposition 6]. □

Lemma 6. *For $\nu \in \mathbb{N}^d \setminus \{0\}$, we have the following relations:*

i)

$$\begin{aligned} [T^\nu, Q_j] &= \nu_j T_1^{\nu_1} \dots T_{i-1}^{\nu_{i-1}} T_i^{\nu_i-1} T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d} \\ &+ B_\alpha \sum_{i=1}^d \sum_{\ell=0}^{\nu_i-1} \sum_{\alpha \in R_+} k(\alpha) \alpha_i \alpha_j H_1^{\nu_1} \dots H_{i-1}^{\nu_{i-1}} H_i^\ell T_i^{\nu_i-\ell-1} T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d}, \end{aligned}$$

where $H_i; i = 1, \dots, d$, are given by the differential-difference operators

$$H_i = -T_i + 2 \frac{\partial}{\partial x_i} - \sum_{\ell=1}^d \alpha_i \alpha_\ell \frac{\partial}{\partial x_\ell}.$$

ii)

$$\begin{aligned} [T_j, Q^\nu] &= \nu_j Q_1^{\nu_1} \dots Q_{i-1}^{\nu_{i-1}} Q_i^{\nu_i-1} Q_{i+1}^{\nu_{i+1}} \dots Q_d^{\nu_d} \\ &+ B_\alpha \sum_{i=1}^d \sum_{\ell=0}^{\nu_i-1} \sum_{\alpha \in R_+} k(\alpha) \alpha_i \alpha_j Z_1^{\nu_1} \dots Z_{i-1}^{\nu_{i-1}} Z_i^\ell Q_i^{\nu_i-\ell-1} Q_{i+1}^{\nu_{i+1}} \dots Q_d^{\nu_d}, \end{aligned}$$

where $Z_i; i = 1, \dots, d$, are the multiplication operators given by

$$Z_i = Q_i - \sum_{\ell=1}^d \alpha_i \alpha_\ell Q_\ell.$$

Proof. From (13), we have

$$\begin{aligned}
 [T_i^{\nu_i}, Q_j] &= \sum_{\ell=0}^{\nu_i-1} T_i^\ell [T_i, Q_j] T_i^{\nu_i-\ell-1} \\
 &= \nu_i \delta_{i,j} T_i^{\nu_i-1} + \sum_{\ell=0}^{\nu_i-1} k(\alpha) \alpha_i \alpha_j T_i^\ell B_\alpha T_i^{\nu_i-\ell-1}.
 \end{aligned}$$

From this equality, we get

$$\begin{aligned}
 [T^\nu, Q_j] &= \sum_{i=1}^d T_1^{\nu_1} \dots T_{i-1}^{\nu_{i-1}} [T_i^{\nu_i}, Q_j] T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d} \\
 &= \nu_j T_1^{\nu_1} \dots T_{i-1}^{\nu_{i-1}} T_i^{\nu_i-1} T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d} \\
 &\quad + \sum_{i=1}^d \sum_{\ell=0}^{\nu_i-1} \sum_{\alpha \in R_+} k(\alpha) \alpha_i \alpha_j T_1^{\nu_1} \dots T_{i-1}^{\nu_{i-1}} T_i^\ell B_\alpha T_i^{\nu_i-\ell-1} T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d}.
 \end{aligned}$$

But

$$T_i^{\nu_i} B_\alpha = B_\alpha H_i^{\nu_i},$$

where

$$H_i = -T_i + 2 \frac{\partial}{\partial x_i} - \sum_{\ell=1}^d \alpha_i \alpha_\ell \frac{\partial}{\partial x_\ell}.$$

Thus we obtain Assertion i). And similarly we get ii). □

Notation. For $x \in \mathbb{R}^d$ and $z \in \mathbb{C}^d$, we denote by

$$I_k(z, x) := \frac{E_k(z, x) - E_k(z, \sigma_\alpha x)}{\langle \alpha, x \rangle}.$$

From [17, p. 533], we can write the function $I_k(z, x)$ in the form

$$(15) \quad I_k(z, x) = \langle \nabla_x E_k(z, x), \alpha \rangle + \frac{1}{2} \langle \alpha, x \rangle \alpha^t D_x^2 E_k(z, \xi) \alpha,$$

with some ξ on the line segment between x and $\sigma_\alpha x$.

(Here ∇ and $D^2 f(x)$ denote the usual gradient and Hessian of f in x .)

Lemma 7. For $a, b \in \mathbb{C}^d$, we have the following commutation relations:

i) $[E_k(a, T), Q_j] = a_j E_k(a, T) - R_{k,j}(a, T)$, where

$$\begin{aligned}
 R_{k,j}(a, T) &= \sum_{\alpha \in R_+} k(\alpha) \alpha_j I_k(a, T) \\
 &\quad - B_\alpha \sum_{\substack{\nu \in \mathbb{N}^d \\ \alpha \in R_+}} \sum_{i=1}^d \sum_{\ell=0}^{\nu_i-1} k(\alpha) \alpha_i \alpha_j \frac{m_\nu(a)}{\nu!} H_1^{\nu_1} \dots H_{i-1}^{\nu_{i-1}} H_i^\ell T_i^{\nu_i-\ell-1} T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d}.
 \end{aligned}$$

ii) $[T_j, E_k(b, Q)] = b_j E_k(b, Q) - S_{k,j}(b, Q)$, where

$$S_{k,j}(b, Q) = \sum_{\alpha \in R_+} k(\alpha) \alpha_j I_k(b, Q) - B_\alpha \sum_{\substack{\nu \in \mathbb{N}^d \\ \alpha \in R_+}} \sum_{i=1}^d \sum_{\ell=0}^{\nu_i-1} k(\alpha) \alpha_i \alpha_j \frac{m_\nu(b)}{\nu!} Z_1^{\nu_1} \dots Z_{i-1}^{\nu_{i-1}} Z_i^\ell Q_i^{\nu_i-\ell-1} Q_{i+1}^{\nu_{i+1}} \dots Q_d^{\nu_d}.$$

Proof. Using (8) and Lemma 6 i), we obtain

$$\begin{aligned} & [E_k(a, T), Q_j] \\ &= \sum_{\nu \in \mathbb{N}^d} \frac{m_\nu(a)}{\nu!} [T^\nu, Q_j] \\ &= \sum_{\nu \in \mathbb{N}^d} \frac{m_\nu(a)}{\nu!} \nu_j T_1^{\nu_1} \dots T_{i-1}^{\nu_{i-1}} T_i^{\nu_i-1} T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d} \\ &+ B_\alpha \sum_{\nu \in \mathbb{N}^d} \sum_{i=1}^d \sum_{\ell=0}^{\nu_i-1} \sum_{\alpha \in R_+} k(\alpha) \alpha_i \alpha_j \frac{m_\nu(a)}{\nu!} H_1^{\nu_1} \dots H_{i-1}^{\nu_{i-1}} H_i^\ell T_i^{\nu_i-\ell-1} T_{i+1}^{\nu_{i+1}} \dots T_d^{\nu_d}. \end{aligned}$$

Applying the relation

$$\frac{\partial}{\partial w_j} E_k(z, w) = z_j E_k(z, w) - \sum_{\alpha \in R_+} k(\alpha) \alpha_j I_k(z, w); \quad z, w \in \mathbb{C}^d,$$

we obtain

$$[E_k(a, T), Q_j] = a_j E_k(a, T) - R_{k,j}(a, T).$$

This proves i). Similarly, we can prove ii). □

Remark. If $d = 1$ and $G = \mathbb{Z}_2$ [21], we have

$$\begin{aligned} R_\gamma(a, T_\gamma) &= \frac{2\gamma}{2\gamma + 1} a(I - B) \mathfrak{S}_{\gamma+\frac{1}{2}}(aT_\gamma), \\ S_\gamma(b, Q) &= \frac{2\gamma}{2\gamma + 1} b(I - B) \mathfrak{S}_{\gamma+\frac{1}{2}}(bQ), \end{aligned}$$

where $Bf(x) = f(-x)$.

Since $E_k(a, 0) = 1$, the Dunkl kernel $E_k(a, z)$; $a, z \in \mathbb{C}^d$, is a unit in the integral domain formal power series over \mathbb{C}^d . We define

$$E_k^{-1}(a, z) := \sum_{\nu \in \mathbb{N}^d} \frac{t_\nu(a)}{\nu!} z^\nu.$$

Writing

$$E_k(a, z) E_k^{-1}(a, z) = E_k^{-1}(a, z) E_k(a, z) = 1,$$

we obtain

$$t_0(a) = 1, \quad \sum_{\nu \in \mathbb{N}^d} \left\{ \sum_{s \leq \nu} \binom{\nu}{s} m_{\nu-s}(a) t_s(a) \right\} \frac{z^\nu}{\nu!} = 1.$$

Thus $\{t_\nu(a)\}_{\nu \in \mathbb{N}^d}$ is a sequence of moment functions in a determined by

$$t_0(a) = 1, \quad t_\nu(a) = - \sum_{s \leq \nu-1} \binom{\nu}{s} m_{\nu-s}(a) t_s(a).$$

The function $E_k^{-1}(a, z)$ occurs in the generalized Weyl commutation relations for the Dunkl kernel.

Theorem 2. *Let $a, b \in \mathbb{C}^d$, then:*

- i) $E_k(b, Q)E_k(a, T) = E_k(a, T)E_k(b, P_a)$, $P_a = (P_{a,1}, \dots, P_{a,d})$, where

$$P_{a,j} = Q_j - a_j I + E_k^{-1}(a, T)R_{k,j}(a, T).$$

- ii) $E_k(a, T)E_k(b, Q) = E_k(b, Q)E_k(a, L_b)$, $L_b = (L_{b,1}, \dots, L_{b,d})$, where

$$L_{b,j} = T_j + b_j I - E_k^{-1}(b, Q)S_{k,j}(b, Q).$$

- iii) $E_k(a, Q)E_k(b, Q) = E_k(a \# b, Q)$, $E_k(a, T)E_k(b, T) = E_k(a \# b, T)$, where $a \# b$ is the convolution of a and b given by

$$m_\nu(a \# b) = \sum_{s \leq \nu} \binom{\nu}{s} m_s(a) m_{\nu-s}(b).$$

Proof. We shall prove i), ii) follows in the same way. For $j = 1, 2, \dots, d$, we have

$$E_k^{-1}(a, T)Q_j E_k(a, T) = E_k^{-1}(a, T) \left\{ E_k(a, T)Q_j - [E_k(a, T), Q_j] \right\}.$$

Using Lemma 7 i), we obtain

$$E_k^{-1}(a, T)Q_j E_k(a, T) = Q_j - a_j I + E_k^{-1}(a, T)R_{k,j}(a, T).$$

Thus implies that for $\nu \in \mathbb{N}^d$:

$$E_k^{-1}(a, T)Q^\nu E_k(a, T) = P_a^\nu, \quad P_a = (P_{a,1}, \dots, P_{a,d}),$$

where

$$P_{a,j} = Q_j - a_j I + E_k^{-1}(a, T)R_{k,j}(a, T).$$

Multiplying by $\frac{m_\nu(b)}{\nu!}$ and summing, we get

$$E_k^{-1}(a, T)E_k(b, Q)E_k(a, T) = E_k(b, P_a).$$

Then i) follows upon multiplication by $E_k(a, T)$.

iii) It suffices to prove the first relation.

Using (8) and (12), we can write

$$\begin{aligned}
 E_k(a, Q)E_k(b, Q) &= \sum_{\nu, s \in \mathbb{N}^d} \frac{m_\nu(a)m_s(b)}{\nu! s!} Q^{\nu+s} \\
 &= \sum_{\nu \in \mathbb{N}^d} \left\{ \sum_{s \leq \nu} \binom{\nu}{s} m_s(a)m_{\nu-s}(b) \right\} \frac{Q^\nu}{\nu!} \\
 &= \sum_{\nu \in \mathbb{N}^d} \frac{m_\nu(a\#b)}{\nu!} Q^\nu.
 \end{aligned}$$

Thus we obtain

$$E_k(a, Q)E_k(b, Q) = E_k(a\#b, Q).$$

□

Remarks. 1) In the classical case ($k = 0$) [13, p. 223], the Weyl commutation relations are given by

$$\begin{aligned}
 e^{\langle a, P \rangle} e^{\langle b, Q \rangle} &= e^{\langle a, b \rangle} e^{\langle b, Q \rangle} e^{\langle a, P \rangle}, \\
 e^{\langle a, P \rangle} e^{\langle b, P \rangle} &= e^{\langle a+b, P \rangle}, \\
 e^{\langle a, Q \rangle} e^{\langle b, Q \rangle} &= e^{\langle a+b, Q \rangle},
 \end{aligned}$$

where $P = (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_d})$ and $Q = (Q_1, \dots, Q_d)$.

2) If $d = 1$ and $G = \mathbb{Z}_2$ [21], the Weyl commutation relations are given by

$$\begin{aligned}
 E_\gamma(bQ)E_\gamma(aT_\gamma) &= E_\gamma(aT_\gamma)E_\gamma(bP_a); \\
 E_\gamma(aT_\gamma)E_\gamma(bQ) &= E_\gamma(bQ)E_\gamma(aL_b),
 \end{aligned}$$

where

$$P_a = Q - aI + \frac{2\gamma}{2\gamma + 1} aE_\gamma^{-1}(aT_\gamma)(I - B)\mathfrak{S}_{\alpha+1}(aT_\gamma),$$

and

$$L_b = T_\gamma + bI - \frac{2\gamma}{2\gamma + 1} bE_\gamma^{-1}(bQ)(I - B)\mathfrak{S}_{\alpha+1}(bQ).$$

Acknowledgments. It is our pleasure to thank Professor Dr. M. Rösler and Professor M. Voit for their interesting discussions.

References

- [1] N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc., **68** (1950), 337-404, MR 0051437, Zbl 0037.20701.
- [2] T.H. Baker and P.J. Forrester, *The Calogero-Sutherland model and generalized classical polynomials*, Comm. Math. Phys., **188** (1997), 175-216, MR 1471336, Zbl 0903.33010.
- [3] ———, *The Calogero-Sutherland model and polynomials with prescribed symmetry*, Nucl. Phys., **B 492** (1997), 682-716, MR 1456121, Zbl 0986.33500.
- [4] V. Bargmann, *On a Hilbert space of analytic functions and an associated integral transform*, Comm. Pure Appl. Math., **14** (1961), 187-214, MR 0157250, Zbl 0107.09102.
- [5] C.A. Berger and L.A. Coburn, *Toeplitz operators on the Segal-Bargmann space*, Trans. Amer. Math. Soc., **301** (1987), 813-829, MR 0882716, Zbl 0625.47019.
- [6] I. Cherednik, *A unification of the Knizhnik-Zamolodchikov equation and Dunkl operators via affine Hecke algebras*, Invent. Math., **106** (1991), 411-432, MR 1128220, Zbl 0742.20019.
- [7] F.M. Cholewinski, *Generalized Fock spaces and associated operators*, SIAM, J. Math. Anal., **15** (1984), 177-202, MR 0728694, Zbl 0596.46017.
- [8] L.A. Coburn and J. Xia, *Toeplitz algebras and Rieffel deformations*, Comm. Math. Phys., **168** (1995), 23-38, MR 1324389, Zbl 0834.47021.
- [9] C.F. Dunkl, *Differential-difference operators associated to reflection groups*, Trans. Amer. Math. Soc., **311** (1989), 167-183, MR 0951883, Zbl 0652.33004.
- [10] ———, *Integral kernels with reflection group invariance*, Canad. J. Math., **43** (1991), 1213-1227, MR 1145585, Zbl 0827.33010.
- [11] P. Etingof and V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*, Invent. Math., **147** (2002), 243-348, MR 1881922.
- [12] V. Fock, *Verallgemeinerung und Lösung der Diracschen statistischen Gleichung*, Z. Phys., **49** (1928), 339-357.
- [13] T. Hida, *Brownian motion*, Applications of Mathematics, **11**, Springer-Verlag, New York-Berlin, 1980, MR 0562914, Zbl 0432.60002.
- [14] L. Lapointe and L. Vinet, *Exact operator solution of the Calogero-Sutherland model*, Comm. Math. Phys., **178** (1996), 425-452, MR 1389912, Zbl 0859.35103.
- [15] N. Obata, *White noise calculus and Fock space*, Lecture Notes in Mathematics, **1577**, Springer-Verlag, Berlin, 1994, MR 1301775, Zbl 0814.60058.
- [16] E.M. Opdam, *Harmonic analysis for certain representations of graded Hecke Algebras*, Acta Math., **175** (1995), 75-121, MR 1353018, Zbl 0836.43017.
- [17] M. Rösler, *Generalized Hermite polynomials and the heat equation for Dunkl operators*, Comm. Math. Phys., **192** (1998), 519-542, MR 1620515, Zbl 0908.33005.
- [18] ———, *Positivity of Dunkl's intertwining operator*, Duke Math. J., **98** (1999), 445-463, MR 1695797, Zbl 0947.33013.
- [19] M. Rösler and M. Voit, *Markov processes related with Dunkl operators*, Adv. Appl. Math., **21** (1998), 575-643, MR 1652182, Zbl 0919.60072.
- [20] ———, *Biorthogonal polynomials associated with reflection groups and a formula of Macdonald*, J. Comp. Appl. Math., **99** (1998), 337-351, MR 1662706, Zbl 0928.33012.

- [21] M. Sifi and F. Soltani, *Generalized Fock spaces and Weyl relations for the Dunkl kernel on the real line*, J. Math. Anal. Appl., **270** (2002), 92-106, MR 1911753, Zbl 1012.46033.

Received July 25, 2002 and revised June 26, 2003.

DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCES OF TUNIS
UNIVERSITY OF TUNIS-EL MANAR
1060 TUNIS
TUNISIA
E-mail address: Fethi.Soltani@fst.rnu.tn

