

*Pacific
Journal of
Mathematics*

TRANSITION DENSITY ESTIMATES FOR STABLE
PROCESSES ON SYMMETRIC SPACES

PIOTR GRACZYK

Volume 217 No. 1

November 2004

TRANSITION DENSITY ESTIMATES FOR STABLE PROCESSES ON SYMMETRIC SPACES

PIOTR GRACZYK

We present estimates of the transition densities for stable processes on Riemannian symmetric spaces of noncompact type. We show that these processes have a weak scaling property and we address in this way a question of Gettoor about the stability properties of pseudostable measures on symmetric spaces.

1. Introduction

Stable Lévy processes on a group, where stability is meant with respect to group automorphisms, can only exist on nilpotent groups (see [Ku], [App1], [App2]). In [Ge], using a subordination procedure, Gettoor defined stable processes with respect to Brownian motion on hyperbolic spaces of noncompact type. He asked whether stability properties of such a process and its semigroup can be found. In this paper we answer this question positively (see Theorem 4.3 and Remark 1 thereafter), proving a *weak scaling* property of the transition densities of the stable process in the sense of Gettoor. We also solve in this way, in the case of symmetric spaces, Open Problem 4 from [App2]. The main results of our paper are given in Theorem 4.3 and Corollaries 5.3 and 5.6.

2. Preliminaries

Let G denote a noncompact semisimple Lie group, K a maximal compact subgroup, and $X = G/K$ the associated Riemannian symmetric space with nonpositive curvature. We adopt the notation and conventions from [AJ]. In particular, if \mathfrak{a} is the Cartan space and $\lambda \in \mathfrak{a}$, then we denote by ϕ_λ the spherical functions on X .

It is well-known that the heat kernel on $X = G/K$ is given by

$$h_t(x) = C \int_{\mathfrak{a}} \frac{d\lambda}{|c(\lambda)|^2} e^{-t(|\lambda|^2 + |\rho|^2)} \phi_\lambda(x)$$

where $C = C(X)$ is a constant, $c(\lambda)$ is the c -function appearing in the inverse spherical Fourier transform formula and $\rho = \frac{1}{2} \sum_{\alpha > 0} m_\alpha \alpha$ is the half-sum of the positive roots with multiplicities m_α .

We set $n = \dim X$, $m = \sum_{\alpha > 0} m_\alpha$, and we denote by Σ^{++} the set of positive indivisible roots and by $\mathfrak{a}^+ \subset \mathfrak{a}$ the positive Weyl chamber. We have global estimates:

Theorem 2.1 ([AJ], Theorem 3.7). *Let $k > 0$. Then*

$$(1) \quad h_t(\exp H) \asymp t^{-\frac{n}{2}}(1+t)^{\frac{m}{2}-|\Sigma^{++}|} \left(\prod_{\alpha \in \Sigma^{++}} (1 + \langle \alpha, H \rangle) \right) e^{-|\rho|^2 t - \langle \rho, H \rangle - \frac{|H|^2}{4t}},$$

provided $|H| < k(1+t)$, $H \in \overline{\mathfrak{a}^+}$.

Observe that $(\prod_{\alpha \in \Sigma^{++}} (1 + \langle \alpha, H \rangle)) e^{-\langle \rho, H \rangle} \asymp \phi_0(\exp H)$ when $H \in \overline{\mathfrak{a}^+}$. Thus for any $x \in X$ we have

$$(2) \quad h_t(x) \asymp t^{-\frac{n}{2}}(1+t)^{\frac{m}{2}-|\Sigma^{++}|} \phi_0(x) e^{-|\rho|^2 t - \frac{|x|^2}{4t}},$$

where $|x| = d(x_0, x)$, the Riemannian distance between $x \in X$ and $x_0 = eK$.

From the probabilistic point of view, the $h_t(x)$ are the densities of the Brownian motion on $X = G/K$. If (X_t) is the corresponding diffusion on G with stationary independent (left) increments and the distribution of X_0 given by the Haar measure on K then its transition function is

$$P_t(x, y) = h_t(x^{-1}y), \quad x, y \in G.$$

Convention: by c without subscripts we denote a positive constant that may vary from term to term, but otherwise depends only on the underlying space and α (see below).

3. Stable semigroups

In the sequel we understand the objects under discussion (processes, transition probabilities, etc.) equivalently on the symmetric space $X = G/K$ or on the group G , without changing notation.

Let $\alpha \in (0, 2)$. The α -stable process on a symmetric space X with transition densities $P_t(x, y) = p_t(x^{-1}y)$ was defined by Gettoor ([Ge]) by means of a subordination procedure. In particular,

$$(3) \quad p_t(x) = \int_0^\infty h_u(x) \eta_t(u) du,$$

where η_t is the density of the $\alpha/2$ -stable subordinator (cf. also [Be]).

In [AJ] estimates for p_t when $\alpha = 1$ were given. In [Gruet] estimates for $p_t(x)$ when $t \rightarrow \infty$ were given in the rank-one case. In this paper we obtain estimates for $p_t(x)$ with respect to both variables x, t for any $\alpha \in (0, 2)$ on all Riemannian symmetric spaces of noncompact type.

It is well-known that the symmetric stable densities on Euclidean spaces cannot be written explicitly, except for $\alpha = 1$. We now recall the exact estimates of the densities $\eta_t(u)$, which will be crucial for our estimates of $p_t(x)$.

By Theorem 37.1 of [D],

$$(4) \quad \lim_{u \rightarrow \infty} \eta_1(u) u^{1+\alpha/2} = \frac{\alpha}{2\Gamma(1 - \frac{\alpha}{2})}.$$

This, together with the boundedness of $\eta_1(\cdot)$ and the scaling property

$$(5) \quad \eta_t(u) = t^{-2/\alpha} \eta_1(t^{-2/\alpha} u), \quad t, u > 0,$$

gives

$$(6) \quad \eta_t(u) \leq ctu^{-1-\alpha/2}, \quad t, u > 0,$$

$$(7) \quad \eta_t(u) \geq \tilde{c}tu^{-1-\alpha/2}, \quad t > 0, u > t^{2/\alpha}.$$

Moreover, by [H] and (5) we know that

$$(8) \quad \eta_t(u) \asymp t^{\frac{1}{2-\alpha}} u^{-\frac{4-\alpha}{4-2\alpha}} e^{-c_1 t^{\frac{2}{2-\alpha}} u^{-\frac{\alpha}{2-\alpha}}}, \quad t^{-2/\alpha} u \in (0, 1),$$

where $c_1 = c_1(\alpha) = \frac{2-\alpha}{2} \left(\frac{\alpha}{2}\right)^{\frac{\alpha}{2-\alpha}}$. Observe that for $u > t^{2/\alpha}$ we have

$$\exp(-c_1 t^{\frac{2}{2-\alpha}} u^{-\frac{\alpha}{2-\alpha}}) \geq \exp(-c_1),$$

so that (6) and (7) give

$$(9) \quad \eta_t(u) \asymp tu^{-1-\alpha/2} e^{-c_1 t^{\frac{2}{2-\alpha}} u^{-\frac{\alpha}{2-\alpha}}}, \quad t^{-2/\alpha} u > 1,$$

which of course simplifies to $\eta_t(u) \asymp tu^{-1-\alpha/2}$, but we want to make the estimates (8) and (9) as similar as possible.

Consider now the case of an n -dimensional Riemannian symmetric space $X = G/K$ when G is a complex Lie group. We have then

$$h_u(x) = \phi_0(x) (4\pi u)^{-n/2} e^{-\frac{|x|^2}{4u} - |\rho|^2 u}$$

and $h_u(\exp H)$ is a probability density with respect to the polar coordinate Jacobian $J(H) dH = \text{vol}(K/M) \prod_{\alpha \in \Sigma^+} \sinh^{m_\alpha} \alpha(H) dH$. Thanks to this explicit formula for h_u we are able to compute the 1-stable density in the complex case, in terms of the modified Bessel function of the third kind (or the MacDonald function) $K_\nu(x)$.

Proposition 3.1. *If G is a complex Lie group and $\alpha = 1$ then*

$$p_t(x) = 2 \left(\frac{2\pi}{|\rho|} \right)^{-\frac{n+1}{2}} t \phi_0(x) (|x|^2 + t^2)^{-\frac{n+1}{4}} K_{\frac{n+1}{2}} (|\rho| \sqrt{|x|^2 + t^2}).$$

Proof. For $\alpha = 1$ we have $\eta_t(u) = \frac{t}{2\sqrt{\pi}} u^{-3/2} \exp(-\frac{t^2}{4u})$. By using the subordination formula and an integral representation for MacDonald function [GR], p. 907, (8.432.6),

$$K_\nu(z) = \frac{1}{2} \left(\frac{z}{2} \right)^\nu \int_0^\infty e^{-t - \frac{z^2}{4t}} t^{-\nu-1} dt,$$

we get the desired assertion. \square

4. Estimates of $p_t(x)$

By (8) and (9) we have

$$(10) \quad p_t(x) \asymp t^{\frac{1}{2-\alpha}} \int_0^{t^{2/\alpha}} h_u(x) u^{-\frac{4-\alpha}{4-2\alpha}} e^{-c_1 t^{\frac{2}{2-\alpha}} u^{-\frac{\alpha}{2-\alpha}}} du \\ + t \int_{t^{2/\alpha}}^{\infty} h_u(x) u^{-1-\alpha/2} e^{-c_1 t^{\frac{2}{2-\alpha}} u^{-\frac{\alpha}{2-\alpha}}} du.$$

As in [AJ] we will see that the main contribution in (10) will come from the interval where

$$u \sim u_0 = u_0(x, t)$$

with u_0 minimizing the function

$$(11) \quad f(u) = \frac{|x|^2}{4u} + |\rho|^2 u + c_1 t^{\frac{2}{2-\alpha}} u^{-\frac{\alpha}{2-\alpha}}.$$

The exponent equal to $-f$ is then maximal in the exponential appearing in (10), if we replace $h_u(x)$ by its estimate given in Theorem 2.1. More precisely, when $|x| < k(1+u)$ and $u > \delta > 0$ with δ fixed, we have

$$(12) \quad h_u(x) \eta_t(u) \asymp \phi_0(x) G(t, u) e^{-\frac{|x|^2}{4u} - |\rho|^2 u - c_1 t^{\frac{2}{2-\alpha}} u^{-\frac{\alpha}{2-\alpha}}},$$

where

$$G(t, u) = \begin{cases} t^{\frac{1}{2-\alpha}} u^{-r(\alpha)}, & u \leq t^{2/\alpha}, \\ t u^{-s(\alpha)}, & u \geq t^{2/\alpha}, \end{cases}$$

with $r(\alpha) = \frac{l}{2} + |\Sigma^{++}| + \frac{4-\alpha}{4-2\alpha}$ and $s(\alpha) = \frac{l}{2} + |\Sigma^{++}| + 1 + \frac{\alpha}{2}$, where $l = \dim \mathfrak{a}$ is the rank of X (we have used $l = n - m$).

Since

$$f'(u) = -\frac{|x|^2}{4u^2} + |\rho|^2 - \frac{c_1 \alpha}{2-\alpha} \frac{t^{\frac{2}{2-\alpha}}}{u^{\frac{2}{2-\alpha}}},$$

we get

$$(13) \quad |\rho|^2 u_0^2 = \frac{|x|^2}{4} + \frac{c_1 \alpha}{2-\alpha} t^{\frac{2}{2-\alpha}} u_0^{\frac{2(1-\alpha)}{2-\alpha}},$$

or, in more convenient form,

$$(14) \quad |\rho|^2 = \frac{1}{4} \left(\frac{|x|}{u_0} \right)^2 + c_2 \left(\frac{t}{u_0} \right)^{\frac{2}{2-\alpha}},$$

with $c_2 = c_1 \alpha / (2 - \alpha)$.

Remark. Equation (13) can be solved explicitly as a (bi)quadratic equation for $\frac{2(1-\alpha)}{2-\alpha} = -2, 0, 1, 2, 4$, corresponding respectively to $\alpha = \frac{3}{2}, 1, 0, \emptyset, 3$, so only for $\alpha = \frac{3}{2}$ or 1. In the case $\alpha = 1$ we obtain $|\rho|^2 u_0^2 = \frac{|x|^2}{4} + c_1 t^2 =$

$\frac{|x|^2+t^2}{4}$ and $u_0 = \sqrt{|x|^2+t^2}/2|\rho|$ as in [AJ]. In the case $\alpha = \frac{3}{2}$ we obtain a biquadratic equation and

$$u_0 = \frac{1}{|\rho|\sqrt{2}} \left(\frac{|x|^2}{4} + \sqrt{\frac{|x|^4}{16} + 4c_2t^4|\rho|^2} \right)^{\frac{1}{2}}.$$

Equation (13) may be solved explicitly for some other particular values of α (see the following lemma). For a generic $\alpha \in (0, 2)$ the function u_0 is given only implicitly by (13).

Lemma 4.1. *For $\alpha = 4/3$ and $t > c_3|x|$ with $c_3 = \frac{1}{\sqrt{3}(4c_2|\rho|)^{1/3}}$, we have*

$$u_0 = \frac{1}{\sqrt[3]{2|\rho|^2}} \left(\left(c_2t^3 - \sqrt{c_2^2t^6 - \frac{|x|^6}{432|\rho|^2}} \right)^{\frac{1}{3}} + \left(c_2t^3 + \sqrt{c_2^2t^6 - \frac{|x|^6}{432|\rho|^2}} \right)^{\frac{1}{3}} \right).$$

Proof. If $\alpha = \frac{4}{3}$ then (13) becomes by rearrangement an equation of degree 3 with respect to u_0 . It can be solved explicitly by Cardano formulas. However, a reasonable expression occurs only when *the determinant* is positive. This is ensured by the second condition in the hypothesis. \square

Lemma 4.2. *For each x and t there exists exactly one solution $u_0(x, t)$ of (13) and (14). We have $u_0 \asymp |x| + t$. The functions $(H, t) \rightarrow u_0(\exp H, t)$ and $(H, t) \rightarrow f(u_0(\exp H, t))$ are homogeneous of degree 1.*

Proof. From (14) we have $(|x|/u_0)^2 < 4|\rho|^2$ and $(t/u_0)^{2/(2-\alpha)} < c_2^{-1}|\rho|^2$, so that $u_0 > |x|/(2|\rho|)$ and $u_0 > tc_2^{(2-\alpha)/2}|\rho|^{-(2-\alpha)}$, which gives the lower bound. On the other hand, for any constant $A > 1$ we have

$$\frac{1}{4} \left(\frac{|x|}{A(|x|+t)} \right)^2 + c_2 \left(\frac{t}{A(|x|+t)} \right)^{\frac{2}{2-\alpha}} \leq \frac{1}{4A^2} + c_2 \left(\frac{1}{A} \right)^{\frac{2}{2-\alpha}},$$

which is less than $|\rho|^2$ for A sufficiently large. Since the right-hand side of (14) is decreasing in u_0 , it follows that $u_0 \leq A(|x| + t)$. By a similar argument the solution of (14) exists and is unique.

Write $u_0(H, t) = u_0(\exp H, t)$. If $u_0 = u_0(H, t)$ is a solution of (14) then for $a > 0$ the solution of (14) for aH and at is au_0 , so $au_0 = u_0(aH, at)$. Thus $u_0(H, t)$ is 1-homogeneous. The homogeneity of $f(u_0)$ is now evident by (11). \square

We now give the exact bounds of the α -stable density $p_t(x)$ in terms of u_0 .

Theorem 4.3.

$$p_t(x) \asymp \begin{cases} t(t^{1/\alpha} + |x|)^{-n-\alpha} & \text{if } t + |x| \leq 1, \\ \phi_0(x)t^{\frac{1}{2-\alpha}}(|x| + t)^{-r(\alpha)+\frac{1}{2}}e^{-f(u_0)} & \text{if } t + |x| \geq 1 \text{ and } u_0 \leq t^{2/\alpha}, \\ \phi_0(x)t(|x| + t)^{-s(\alpha)+\frac{1}{2}}e^{-f(u_0)} & \text{if } t + |x| \geq 1 \text{ and } u_0 \geq t^{2/\alpha}, \end{cases}$$

with $r(\alpha) = \frac{l}{2} + |\Sigma^{++}| + \frac{4-\alpha}{4-2\alpha}$, $s(\alpha) = \frac{l}{2} + |\Sigma^{++}| + 1 + \frac{\alpha}{2}$, $u_0 = u_0(x, t)$ given implicitly by (13) and the function f defined in (11).

Remark 1. Theorem 4.3 gives a weak scaling property for stable densities on symmetric spaces (the possibility of recovering, at least asymptotically, the density $p_t(x)$ if $p_1(x)$ is known).

More precisely, when $t + |x| \leq 1$, then $p_t(\exp H) \asymp t^{-n/\alpha} p_1(\exp(t^{-1/\alpha} H))$ (in the Euclidean case we have equality).

When $t + |x| \geq 1$, it is not the function $p_t(x)$ itself but the function $\phi_0(x)^{-1} p_t(x)$ that has a weak scaling property (the role of the function ϕ_0 may be explained as the influence of the non-Euclidean structure of X). It is clear that the factors $t^{\frac{1}{2-\alpha}} (|H| + t)^{-r(\alpha)+\frac{1}{2}}$ and $t (|H| + t)^{-s(\alpha)+\frac{1}{2}}$ of the bounds of the function $\phi_0(x)^{-1} p_t(\exp H)$ may be obtained from the factors $(|H| + 1)^{-r(\alpha)+\frac{1}{2}}$ and $(|H| + 1)^{-s(\alpha)+\frac{1}{2}}$ of the bounds of $\phi_0(x)^{-1} p_1(\exp H)$ by transformations of the form $H \mapsto t^\beta p_1(\exp(t^{-1} H))$, with a convenient value of β . The same is true for the function $f(u_0)$ since, by the 1-homogeneity of $f(u_0)$, we have $tf(u_0(\exp(t^{-1} H), 1)) = f(u_0(\exp H, t))$.

On structures different from \mathbb{R} stable measures defined by subordination may preserve such a weaker scaling property (cf. [BSS] for fractals). We answer in this way a question raised by Gettoor in [Ge] about looking for “stability” properties of the densities p_t .

Remark 2. After the proof of the theorem we will give a simple criterion concerning the conditions $u_0 \geq t^{2/\alpha}$ and $u_0 \leq t^{2/\alpha}$. Note that for $u_0 \asymp t^{2/\alpha}$ the second and the third estimates coincide.

Proof of Theorem 4.3. The estimates for small t and $|x|$ are the same as on \mathbb{R}^n and may be found in [BSS] (cf. also [Ben]). From Lemma 4.2 it follows that $u_0(x, t) \rightarrow \infty$ when $\max(x, t) \rightarrow \infty$. Following [AJ], Theorem 4.3.1, we will look for the estimates of $p_t(x)$ in terms of u_0 , when $u_0 \rightarrow \infty$. We split the integral (3) for $p_t(x)$ into

$$(15) \quad \int_0^{\kappa^{-1}u_0} + \int_{\kappa^{-1}u_0}^\infty = I_1 + I_2$$

for a constant $\kappa > 1$. We will see below that the first integral is essentially smaller than the second one, so $p_t(x) \asymp \int_{\kappa^{-1}u_0}^\infty h_u(x) \eta_t(u) du$. Now we may use the estimate for $h_u(x)$ from the Theorem 2.1. Write $J_2 = \phi_0(x)^{-1} I_2$ and in J_2 apply the change of variables $u = u_0(x, t)v$. Set

$$P = P(x, t) = \frac{|x|^2}{4u_0} / |\rho|^2 u_0 \quad \text{and} \quad Q = Q(x, t) = \frac{c_1 \alpha}{2 - \alpha} t^{\frac{2}{2-\alpha}} u_0^{\frac{-\alpha}{2-\alpha}} / |\rho|^2 u_0.$$

We have $0 < P, Q < 1$ and $P + Q \equiv 1$ (the functions P and Q measure in a sense the proportion of importance of variables x and t respectively in the

function $u_0(x, t)$). We get

$$(16) \quad f(u_0 v) = |\rho|^2 u_0 \left(v + P \frac{1}{v} + Q \frac{2 - \alpha}{\alpha} v^{-\frac{\alpha}{2 - \alpha}} \right).$$

For any values of P and Q , the point v where the function

$$g(v, x, t) = v + P \frac{1}{v} + Q \frac{2 - \alpha}{\alpha} v^{-\frac{\alpha}{2 - \alpha}}$$

attains its minimum must satisfy the equation $P \frac{1}{v^2} + Q v^{-\frac{2}{2 - \alpha}} = 1$, which admits the only solution $v = 1$. The functions u_0 , P and Q depend on x and t . Nevertheless, due to the uniqueness property just mentioned and the fact that $\frac{\partial^2 g}{\partial v^2} \asymp 1$ for all x and t , the proof of the Laplace method (see [O], pp. 80–82)) may be adapted to the present situation. The moderate price to pay is obtaining bounds instead of asymptotics in u_0 . Consequently we get

$$(17) \quad J_2 \asymp c t^{\frac{1}{2 - \alpha}} u_0^{-r(\alpha) + \frac{1}{2}} e^{-f(u_0)}, \quad u_0 \rightarrow \infty, \quad u_0 \leq t^{2/\alpha},$$

$$(18) \quad J_2 \asymp c t u_0^{-s(\alpha) + \frac{1}{2}} e^{-f(u_0)}, \quad u_0 \rightarrow \infty, \quad u_0 \geq t^{2/\alpha}.$$

To complete the proof we need to justify the claim that the first integral in (15), which we denote by I_1 , is essentially smaller than the second one. We use the global upper estimate for the heat kernel ([An], [AJ, (3.3)]), and proceed as with J_2 above, applying the Laplace method. It follows that $I_1/I_2 \rightarrow 0$ when $u_0 \rightarrow \infty$. \square

Lemma 4.4. *There exist positive constants K, M such that if $t \leq K|x|^{\alpha/2}$ then $u_0 \geq t^{2/\alpha}$ and if $t \geq M|x|^{\alpha/2}$ then $u_0 \leq t^{2/\alpha}$. If $t \asymp |x|^{\alpha/2}$ then $u_0 \asymp t^{2/\alpha}$.*

Proof. The first two assertions follow easily from Lemma 4.2. If $t \asymp |x|^{\alpha/2}$ then $|x| \asymp t^{2/\alpha}$ and, consequently, $u_0 \asymp |x| + t \asymp t^{2/\alpha} + t$. Since u_0 and t are supposed to be large ($t \rightarrow \infty$), we have $u_0 \asymp t^{2/\alpha}$. \square

5. Properties of u_0

Note that for general $\alpha \in (0, 2)$ we have no explicit formula for u_0 and $f(u_0)$. In this section we study their properties in greater depth.

Proposition 5.1. *$(H, t) \mapsto u_0 = u_0(\exp H, t)$ is a norm on $\mathfrak{a} \times \mathbb{R}^+$.*

Proof. Again set $u_0(H, t) = u_0(\exp H, t)$. We already know that $u_0(H, t)$ is 1-homogeneous. Set $u_0 = u_0(H, t)$, $u_1 = u_0(H_1, t_1)$, $u_2 = u_0(H + H_1, t + t_1)$, $a = 1/(4|\rho|)^2$ and $b = c_2/|\rho|^2$. For $p = \frac{u_0}{u_0 + u_1} \in (0, 1)$ and $q = 1 - p$ we have

$$(19) \quad \frac{u_0}{p} = \frac{u_1}{q}.$$

Then we obtain

$$ap \left(\frac{|H|}{u_0} \right)^2 + bp \left(\frac{t}{u_0} \right)^{\frac{2}{2-\alpha}} = p, \quad \text{and} \quad aq \left(\frac{|H_1|}{u_1} \right)^2 + bq \left(\frac{t_1}{u_1} \right)^{\frac{2}{2-\alpha}} = q.$$

Summing these equations, using the fact that functions $x \rightarrow x^2$ and $t \rightarrow t^{2/(2-\alpha)}$ are convex, and applying (19), we get $u_2 \leq u_0/p = u_1/q$ and $u_2 \leq u_0 + u_1$. This shows that u_0 is a norm. \square

Proposition 5.2. u_0 extends to a norm on $\mathfrak{a} \times \mathbb{R}$. $f(u_0)$ is a norm on $\mathfrak{a} \times \mathbb{R}^+$ and extends to a norm on $\mathfrak{a} \times \mathbb{R}$.

Proof. Since u_0 is homogeneous, it is determined by its unit sphere $S = \{(H, t) : u_0(H, t) = 1\}$. Since $c_2^{(2-\alpha)/2} = \frac{\alpha}{2}$, S can be described by

$$t = \frac{2}{\alpha} \left(|\rho|^2 - \frac{|H|^2}{4} \right)^{\frac{2-\alpha}{2}}.$$

One may deduce from this formula the convexity of the unit ball $B_0 \subset \mathbb{R}^2$ of the 1-homogeneous function

$$U_0 : U_0(y, t) = u_0(|H|, |t|)$$

with any $H \in \mathfrak{a}$ such that $|y| = |H|$, $y \in \mathbb{R}$. Thus U_0 is a norm on \mathbb{R}^2 . The convexity of B_0 implies the convexity of

$$B = \{(H, t) \in \mathfrak{a} \times \mathbb{R} : (|H|, t) \in B_0\} = \{(H, t) \in \mathfrak{a} \times \mathbb{R} : u_0(H, |t|) = 1\},$$

which is the unit ball of u_0 extended naturally to $\mathfrak{a} \times \mathbb{R}$. This gives the first statement.

For the second one, we solve $f(u_0) = 1$ using (11) and the equality

$$(20) \quad \left(\frac{t}{u_0} \right)^{\frac{2}{2-\alpha}} = \frac{1}{c_2} \left(|\rho|^2 - \frac{1}{4} \left(\frac{|H|}{u_0} \right)^2 \right).$$

We obtain a quadratic equation for u_0 ,

$$(21) \quad \frac{2}{\alpha} |\rho|^2 u_0^2 - u_0 + \frac{\alpha-1}{2\alpha} |H|^2 = 0,$$

and it follows that

$$(22) \quad u_0 = \varphi(H) = \frac{\alpha}{4|\rho|^2} (1 + P(H)), \quad |H| \in \left(0, \frac{1}{|\rho|} \right),$$

where

$$P(H) = \sqrt{1 - \frac{4(\alpha-1)}{\alpha^2} |\rho|^2 |H|^2}.$$

Indeed, let $\varphi_m(H)$ be the alternative formal solution to (21) (this is possible at least for $\alpha > 1$). Then positivity of the right-hand side of (20) implies that $\frac{|H|}{2\varphi_m(H)} \leq |\rho|$ and, consequently, $P(H) \leq 1 - \frac{2|H||\rho|}{\alpha}$. This is possible only for $|H| \leq \frac{\alpha}{2|\rho|}$. Solving this we obtain $|H| \geq \frac{1}{|\rho|}$. The contradiction

shows that (21) admits at most one solution. An analogous argument for φ instead of φ_m establishes the range for H . Note that for $|H| \in (0, 1/|\rho|)$ the determinant of (21) is positive for every $\alpha \in (0, 2)$. Thus, we arrive at (22). Putting $u_0 = \varphi(H)$ into (20) we get

$$t = \psi(H) = \frac{2}{\alpha} \varphi(H) \left(|\rho|^2 - \frac{|H|^2}{4\varphi(H)^2} \right)^{1-\frac{\alpha}{2}}.$$

We have $f(u_0) = 1$ iff $u_0 = \varphi(H)$ and $t = \psi(H)$. To finish the proof it is enough to show that the graph of $y \mapsto \psi(y)$ is concave on \mathbb{R}^+ (i.e., $\psi'' \leq 0$, where we understand $y = |H|$ and ψ as a function of $|H|$ on \mathbb{R}^+) and the symmetry with respect to the t -axis does not affect the concavity (e.g., ψ is decreasing). We obtain

$$\psi''(y) = - \left(\frac{\alpha}{2} \right)^{\frac{\alpha}{2}} \frac{2r^{2-\alpha}(1 + P(y))^\alpha}{s(y)^{1+\frac{\alpha}{2}} P(y)}$$

with the obvious meaning of $P(y)$ and

$$s(y) = \alpha - 2|\rho|^2|y|^2 + \sqrt{\alpha^2 - 4(\alpha - 1)|\rho|^2|y|^2} > 0,$$

since for $y < 1/|\rho|$ we have

$$s(y) \geq \alpha - 2 + \sqrt{\alpha^2 - 4(\alpha - 1)} = \alpha - 2 + |\alpha - 2| = 0.$$

These formulas can be quickly checked using Maple or Mathematica. Thus $\psi''(y) \leq 0$. It is easy to verify that $\psi'(0) = 0$, so $\psi'(y) \leq 0$ as required. \square

Corollary 5.3. *When $t + |x| \geq 1$,*

$$p_t(x) \asymp \begin{cases} \phi_0(x) t^{\frac{1}{2-\alpha}} (|x| + t)^{-r(\alpha)+\frac{1}{2}} e^{\sqrt{|x|^2+t^2}} q_\alpha & \text{if } u_0 \leq t^{2/\alpha}, \\ \phi_0(x) t (|x| + t)^{-s(\alpha)+\frac{1}{2}} e^{\sqrt{|x|^2+t^2}} q_\alpha & \text{if } u_0 \geq t^{2/\alpha}, \end{cases}$$

where the function q_α satisfies

$$q_\alpha(k \exp Hk', t) = q_\alpha \left(\frac{(H, t)}{\|(H, t)\|_2} \right), \quad k, k' \in K,$$

and is a continuous, bounded and bounded away from zero function on the Euclidean unit sphere in $\mathfrak{a} \times \mathbb{R}$.

Proof. Set $q_\alpha(\exp H, t) = f(u_0(\exp H, t))/\sqrt{|H|^2 + t^2}$. By Proposition 5.3 the function $(H, t) \rightarrow q_\alpha(\exp H, t)$ is homogeneous of degree 0. \square

Remark. When $\alpha = 1$ the function q_α is constant. The bounds from the Corollary 5.3 are an extension of the bounds obtained in [AJ] for $\alpha = 1$.

Lemma 4.2 gives enough information to replace u_0 in an estimate when it is a multiplicative factor. To deal with the exponent $e^{-f(u_0)}$ in the estimates of $p_t(x)$ more delicate (additive) properties are required. Namely,

$$\exp(f(x)) \asymp \exp(g(x)) \iff |f - g| \text{ is bounded.}$$

This motivates the following definition: we write

$$f(x) \stackrel{e}{\asymp} g(x)$$

if there exists $M \in \mathbb{R}$ such that $|f(x) - g(x)| \leq M$ for all x .

Proposition 5.4. *If $t \leq |x|^{\alpha/2}$ then*

$$(23) \quad \exp(-f(u_0)) \asymp \exp(-|\rho||x|).$$

In particular, this holds for fixed $t > 0$, as $x \rightarrow \infty$ (that is, for all sufficiently large $x > x_0(t)$). Moreover, if $t = |x|^s$ with $s > \alpha/2$ then (23) does not hold.

Proof. By our assumption, $t^{\frac{2}{2-\alpha}} \leq |x|^{\frac{\alpha}{2-\alpha}}$ and

$$(24) \quad t^{\frac{2}{2-\alpha}} u_0^{-\frac{\alpha}{2-\alpha}} \leq \left(\frac{|x|}{u_0} \right)^{\frac{\alpha}{2-\alpha}}.$$

The right-hand side of this inequality is bounded since $u_0 \asymp |x| + t$. Thus, Equation (14) multiplied by u_0 gives

$$(25) \quad f(u_0) = \frac{|x|^2}{4u_0} + |\rho|^2 u_0 + c_1 t^{\frac{2}{2-\alpha}} u_0^{-\frac{\alpha}{2-\alpha}} \stackrel{e}{\asymp} 2|\rho|^2 u_0.$$

Using (13) and (24) we deduce that $u_0 \stackrel{e}{\asymp} \frac{|x|}{2|\rho|}$, which together with (25) gives the first assertion.

By the definition of $f(u_0)$ for $t = |x|^s$ and by (13) we obtain

$$(26) \quad f(u_0) - |\rho||x| = (c_1 + c_2) \left(\frac{|x|}{u_0} \right)^{\frac{\alpha}{2-\alpha}} |x|^{\frac{2s-\alpha}{2-\alpha}} - |x| \left(|\rho| - \frac{|x|}{2u_0} \right).$$

By (14)

$$(27) \quad \left(|\rho| - \frac{|x|}{2u_0} \right) \left(|\rho| + \frac{|x|}{2u_0} \right) = c_2 \left(\frac{t}{u_0} \right)^{\frac{2}{2-\alpha}}$$

so, transforming (26) with $t = |x|^s$ we get

$$(28) \quad f(u_0) - |\rho||x| = |x|^{\frac{2s-\alpha}{2-\alpha}} \left((c_1 + c_2) \left(\frac{|x|}{u_0} \right)^{\frac{\alpha}{2-\alpha}} - c_2 \left(\frac{|x|}{u_0} \right)^{\frac{2}{2-\alpha}} \left(|\rho| + \frac{|x|}{2u_0} \right)^{-1} \right).$$

Note that for $s \leq \alpha/2$ this is bounded (for $s < \alpha/2$ it even tends to 0), since $u_0 \asymp |x| + t$. (This gives once again the first assertion.) From (14) with $|x| = t^{1/s}$ we deduce that $\frac{t}{u_0} \rightarrow 0$. Observe that it is enough to show the second assertion for $s \in (\alpha/2, 1)$.

Now, from (14) it follows that

$$(29) \quad \frac{|x|}{2u_0} \rightarrow |\rho|.$$

Together with (28) it implies that for $s > \alpha/2$ we have $f(u_0) - |\rho||x| \rightarrow \infty$, which completes the proof. \square

Proposition 5.5. *If $|x| \leq \sqrt{t}$ then $\exp(-f(u_0)) \asymp \exp(-|\rho|^\alpha t)$. This is true, in particular, when x is fixed and $t \rightarrow \infty$ (that is, for all sufficiently large $t > t_0(x)$). If $|x| = t^s$ with $s > \frac{1}{2}$, it is no longer true.*

Proof. Multiplying (14) by u_0 and using

$$(30) \quad \frac{|x|^2}{u_0} \leq \frac{t}{u_0},$$

which is bounded since $u_0 \asymp |x| + t$, we get

$$(31) \quad f(u_0) \stackrel{e}{\asymp} |\rho|^2 u_0 + \frac{2-\alpha}{\alpha} |\rho|^2 u_0 = \frac{2}{\alpha} |\rho|^2 u_0.$$

We claim that

$$(32) \quad \frac{2}{\alpha} |\rho|^2 u_0 \stackrel{e}{\asymp} |\rho|^\alpha t.$$

We will use the following simple consequence of the Mean Value Theorem: for $0 < a < b$ and any γ we have

$$(33) \quad b^\gamma - a^\gamma = \gamma K^{\gamma-1} (b - a),$$

for some $K \in (a, b)$. From (14) multiplied by $u_0^{2/(2-\alpha)}$ we have

$$|\rho|^2 u_0^{\frac{2}{2-\alpha}} - c_2 t^{\frac{2}{2-\alpha}} = \frac{|x|^2}{4} u_0^{-2+\frac{2}{2-\alpha}}.$$

From this and (33) applied to $a = c_2 t^{\frac{2}{2-\alpha}}$, $b = |\rho|^2 u_0^{\frac{2}{2-\alpha}}$ and $\gamma = \frac{2-\alpha}{2}$ we get

$$|\rho|^{2-\alpha} u_0 - c_2^{\frac{2-\alpha}{2}} t = \frac{2-\alpha}{2} K^{-\frac{\alpha}{2}} \frac{|x|^2}{4} u_0^{-2+\frac{2}{2-\alpha}},$$

with some $K \in (a, b)$. Consequently, as $K > a$,

$$|\rho|^{2-\alpha} u_0 - c_2^{\frac{2-\alpha}{2}} t \leq c \left(\frac{u_0}{t} \right)^{\frac{\alpha}{2-\alpha}} \frac{|x|^2}{u_0},$$

which is bounded by virtue of (30) and the fact that $|x|^2 \leq t$ yields $u_0 \asymp |x| + t \asymp t$. It follows that

$$(34) \quad |\rho|^{2-\alpha} u_0 \stackrel{e}{\asymp} c_2^{\frac{2-\alpha}{2}} t.$$

To get our claim, multiply (34) by $\frac{2|\rho|^\alpha}{\alpha}$ and use the fact that $\frac{2}{\alpha} c_2^{\frac{2-\alpha}{2}} = 1$. The first assertion now follows from (31) and (32).

In order to prove the second assertion, we show in a way similar to the proof of the second assertion of the Proposition 5.4 (but using the Mean Value Theorem as above, for $\gamma = \alpha/2$) that the formula

$$(35) \quad f(u_0) - |\rho|^\alpha t = \frac{1}{2} t^{2s-1} \left(\frac{t}{u_0} - \frac{\alpha}{4} \left(\frac{t}{u_0} \right)^2 K^{\frac{\alpha-2}{2}} \right)$$

holds when $|x| = t^s$, with

$$(36) \quad c_2 \left(\frac{t}{u_0} \right)^{\frac{2}{2-\alpha}} < K < |\rho|^2.$$

The case $s > 1$ being evident by (14), we see that for $s < 1$, again using (14), we have $\frac{|x|}{u_0} \rightarrow 0$ and

$$\frac{t}{u_0} \rightarrow c_2^{\frac{\alpha-2}{2}} |\rho|^{2-\alpha}$$

when $|x| = t^s$.

Taking limits in (36) we get $K \rightarrow |\rho|^2$, so $K^{\frac{\alpha-2}{2}} \rightarrow |\rho|^{\alpha-2}$. Thus the expression between the brackets in (35) does not tend to 0. It follows that for $s > 1/2$ the difference $f(u_0) - |\rho|^\alpha t$ is unbounded when $t \rightarrow \infty$ and $|x| = t^s$. \square

Corollary 5.6. *For any constants $K, M > 0$ we have*

$$p_t(x) \asymp \begin{cases} \phi_0(x) t^{\frac{1}{2-\alpha}} (|x| + t)^{-r(\alpha) + \frac{1}{2}} e^{-|\rho|^\alpha t} & \text{if } t + |x| \geq 1 \text{ and } |x| \leq K t^{1/2}, \\ \phi_0(x) t (|x| + t)^{-s(\alpha) + \frac{1}{2}} e^{-|\rho||x|} & \text{if } t + |x| \geq 1 \text{ and } |x| \geq M t^{2/\alpha}. \end{cases}$$

On the other hand, in a region $K t^{1/2} < |x| < M t^{2/\alpha}$ one cannot give a simpler estimate of $e^{-f(u_0)}$ and, consequently, a simpler estimate of $p_t(x)$ than the one given in the Theorem 4.3.

Proof. Only the last assertion needs to be justified. Suppose that $g(|x|, t)$ is a 1-homogeneous function such that $e^{-f(u_0)} \asymp e^{-g}$ on

$$R = \{(|x|, t) : K t^{1/2} < |x| < M t^{2/\alpha}\}.$$

It implies that $f(u_0) \stackrel{c}{\asymp} g$.

Observe that the region R contains half-lines of the form $|x| = at$, where $t > t_0(a)$, $a > 0$. The function $f(u_0) - g$ is homogeneous of degree 1 and bounded on R , so $f(u_0) = g$ on R .

Suppose that the function $f(u_0)$ is explicitly determined. Writing (11) in the form

$$f(u_0) = \frac{|x|^2}{4u_0} + |\rho|^2 u_0 + c_1 u_0 \left(\frac{t}{u_0} \right)^{\frac{2}{2-\alpha}}$$

and replacing $(\frac{t}{u_0})^{\frac{2}{2-\alpha}}$ by

$$c_2^{-1} \left(|\rho|^2 - \frac{1}{4} \left(\frac{|x|}{u_0} \right)^2 \right)$$

we get according to (14) a quadratic equation with respect to u_0 that could be solved explicitly. Except for some special values of $\alpha = 1, \frac{3}{2}, \dots$, this is not possible. \square

Acknowledgements. We thank J.-P. Anker, K. Bogdan and T. Byczkowski for discussions and helpful comments on the subject of the paper. The second named author acknowledges the hospitality of Université d'Angers during a post-doctoral fellowship granted by the region Pays de la Loire.

References

- [An] J.-P. Anker, *Sharp estimates for some functions of the Laplacian on noncompact symmetric spaces*, Duke Math. J., **65** (1992), 257–297, MR 1150587 (93b:43007), Zbl 0764.43005.
- [AJ] J.-P. Anker and L. Ji, *Heat kernel and Green function estimates on noncompact symmetric spaces*, Geom. Funct. Anal., **9(6)** (1999), 1035–1091, MR 1736928 (2001b:58038), Zbl 0942.43005.
- [App1] D. Applebaum, *Lévy processes in stochastic differential geometry*, Lévy processes, 111–137, Birkhäuser, Boston, MA, 2001, MR 1833695 (2002d:60038), Zbl 0984.60056.
- [App2] D. Applebaum, *On the subordination of spherically symmetric Lévy processes in Lie groups*, Int. Math. J., **1(2)** (2002), 185–194, MR 1828652 (2002f:60011), Zbl 0984.60018.
- [Ben] A. Bendikov, *Asymptotic formulas for symmetric stable semigroups*, Exposition. Math., **12** (1994), 381–384, MR 1297844 (95j:60029), Zbl 0810.60070.
- [Be] J. Bertoin, *Lévy Processes*, Cambridge Tracts in Mathematics, **121**, Cambridge University Press, Cambridge, 1996, MR 1406564 (98e:60117), Zbl 0861.60003.
- [BG] R.M. Blumenthal and R.K. Gettoor, *Markov Processes and Potential Theory*, Pure and Applied Mathematics, **29**, Academic Press, New York, 1968, MR 0264757 (41 #9348), Zbl 0169.49204.
- [BSS] K. Bogdan, A. Stós and P. Sztonyk, *Harnack inequality for symmetric stable processes on fractals*, C.R. Acad. Sci. Paris, Ser. I, **335(1)** (2002), 59–63, MR 1920996 (2003k:60100), Zbl 1019.60078.
- [D] G. Doetsch, *Introduction to the Theory and Application of the Laplace Transformation*, Springer, Berlin-Heidelberg-New York, 1974, MR 0344810 (49 #9549), Zbl 0278.44001.
- [Ge] R.K. Gettoor, *Infinitely divisible probabilities on the hyperbolic plane*, Pacific J. Math., **11** (1961), 1287–1308, MR 0133858 (24 #A3682), Zbl 0124.34502.
- [GR] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products*, 6th edition, Academic Press, London, 1965, MR 0197789 (33 #5952), Zbl 0981.65001.

- [Gruet] J-C. Gruet, *Jacobi radial stable processes*, Ann. Math. Blaise Pascal, **5** (1998), 39–48, MR 1671691 (99m:60015), Zbl 0926.60020.
- [H] J. Hawkes, *A lower Lipschitz condition for the stable subordinator*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, **17** (1971), 23–32, MR 0282413 (43 #8125), Zbl 0193.45002.
- [Ku] H. Kunita, *Stable Lévy processes on nilpotent Lie groups*, in ‘Stochastic analysis on infinite-dimensional spaces’ (Baton Rouge, LA, 1994), 167–182, Pitman Res. Notes Math. Ser., **310**, Longman Sci. Tech., Harlow, 1994, MR 1415667 (97k:60019), Zbl 0814.60003.
- [O] F.W.J. Olver, *Asymptotics and Special Functions*, Computer Science and Applied Mathematics, Academic Press, New York-London, 1974, MR 0435697 (55 #8655), Zbl 0303.41035.

Received December 3, 2002 and revised January 14, 2004. Both authors supported by KBN grant 2 P03A 041 22 and RTN Harmonic Analysis and Related Problems contract HPRN-CT-2001-00273-HARP.

DÉPARTEMENT DE MATHÉMATIQUES
UNIVERSITÉ D’ANGERS
FRANCE
E-mail address: graczyk@univ-angers.fr

INSTITUTE OF MATHEMATICS
WROCLAW UNIVERSITY OF TECHNOLOGY
WROCLAW
POLAND
E-mail address: stos@im.pwr.wroc.pl

LABORATOIRE DE MATHÉMATIQUES UMR 6620
UNIVERSITÉ BLAISE PASCAL
FRANCE