

*Pacific  
Journal of  
Mathematics*

**PEAK-INTERPOLATING CURVES FOR  $A(\Omega)$  FOR  
FINITE-TYPE DOMAINS IN  $\mathbb{C}^2$**

GAUTAM BHARALI

Volume 218 No. 2

February 2005



## PEAK-INTERPOLATING CURVES FOR $A(\Omega)$ FOR FINITE-TYPE DOMAINS IN $\mathbb{C}^2$

GAUTAM BHARALI

Let  $\Omega$  be a bounded, weakly pseudoconvex domain in  $\mathbb{C}^2$ , having smooth boundary.  $A(\Omega)$  is the algebra of all functions holomorphic in  $\Omega$  and continuous up to the boundary. A smooth curve  $C \subset \partial\Omega$  is said to be complex-tangential if  $T_p(C)$  lies in the maximal complex subspace of  $T_p(\partial\Omega)$  for each  $p \in C$ . We show that if  $C$  is complex-tangential and  $\partial\Omega$  is of constant type along  $C$ , then every compact subset of  $C$  is a peak-interpolation set for  $A(\Omega)$ . Furthermore, we show that if  $\partial\Omega$  is real-analytic and  $C$  is an arbitrary real-analytic, complex-tangential curve in  $\partial\Omega$ , compact subsets of  $C$  are peak-interpolation sets for  $A(\Omega)$ .

### 1. Statement of the main result

Let  $\Omega$  be a bounded domain in  $\mathbb{C}^n$ , and let  $A(\Omega)$  be the algebra of functions continuous on  $\bar{\Omega}$  and holomorphic in  $\Omega$ . Recall that a compact subset  $K \subset \partial\Omega$  is called a *peak-interpolation set* for  $A(\Omega)$  if given any  $f \in \mathcal{C}(K)$ ,  $f \not\equiv 0$ , there exists a function  $F \in A(\Omega)$  such that  $F|_K = f$  and  $|F(\zeta)| < \sup_K |f|$  for every  $\zeta \in \bar{\Omega} \setminus K$ .

We are interested in determining when a smooth submanifold  $M \subset \partial\Omega$  is a peak-interpolation set for  $A(\Omega)$ . When  $\Omega$  is a strictly pseudoconvex domain having  $\mathcal{C}^2$  boundary and  $M$  is of class  $\mathcal{C}^2$ , the situation is very well understood; see [Henkin and Tumanov 1976; Nagel 1976; Rudin 1978]. In the strictly pseudoconvex setting,  $M$  is a peak-interpolation set for  $A(\Omega)$  if and only if  $M$  is *complex-tangential*, meaning that  $T_p(M) \subset H_p(\partial\Omega)$  for all  $p \in M$ . (Here and in what follows, for any submanifold  $M \subseteq \partial\Omega$ ,  $T_p(M)$  will denote the real tangent space to  $M$  at the point  $p \in M$ , while  $H_p(\partial\Omega)$  will denote the maximal complex subspace of  $T_p(\partial\Omega)$ .)

Very little is known, however, when  $\Omega$  is a weakly pseudoconvex of finite type. (There are several notions of type for domains in  $\mathbb{C}^n$ ,  $n \geq 2$ , but they all coincide for pseudoconvex domains in  $\mathbb{C}^2$ . See Section 2 below.) In view of a result by Henkin and Tumanov [1976] or a similar result by Nagel and Rudin [1978], it is still necessary for  $M$  to be complex-tangential. It was recently shown [Bharali 2004] that for bounded (weakly) convex domains  $\Omega \subset \mathbb{C}^n$  with real-analytic boundaries,

---

MSC2000: primary 32A38, 32T25; secondary 32C25, 32D99.

Keywords: complex-tangential, finite type domain, interpolation set, pseudoconvex domain.

complex-tangential submanifolds  $M \subset \partial\Omega$  are peak-interpolation sets. However, showing even that any smooth compact complex-tangential arc in  $\partial\Omega$  is a peak-interpolation set for  $A(\Omega)$ , for a general smoothly bounded weakly pseudoconvex domain of finite type, is a difficult problem. This is because doing so would necessarily imply that every point in  $\partial\Omega$  is a peak point for  $A(\Omega)$ . Whether or not this is true for general pseudoconvex domains of finite type is an extremely difficult open question in the theory of functions in several complex variables, but this fact is certainly known for smoothly bounded finite type domains in  $\mathbb{C}^2$  [Bedford and Fornæss 1978; Fornæss and McNeal 1994; Fornæss and Sibony 1989], and we will use it in one of our results below. In this paper we show, among other things, that when  $\Omega$  is a bounded domain in  $\mathbb{C}^2$ ,  $\partial\Omega$  is real-analytic and  $C \subset \partial\Omega$  is a real-analytic curve, it suffices that  $C$  be complex-tangential for every compact subset of  $C$  to be a peak-interpolation set for  $A(\Omega)$ .

More precisely, our main result is:

**Theorem 1.1.** *Let  $\Omega$  be a bounded pseudoconvex domain in  $\mathbb{C}^2$  having smooth boundary, and let  $C \subset \partial\Omega$  be a smooth curve.*

- (i) *Let  $\partial\Omega$  be of class  $\mathcal{C}^\infty$  and  $\Omega$  be of finite type. If  $C$  is complex-tangential, and if  $\partial\Omega$  is of constant type along  $C$ , then each compact subset of  $C$  is a peak-interpolation set for  $A(\Omega)$ .*
- (ii) *Let  $\Omega$  have real-analytic boundary and let  $C \subset \partial\Omega$  be a real-analytic complex-tangential curve. Then each compact subset of  $C$  is a peak interpolation set for  $A(\Omega)$ .*

In (ii) above, we do *not* assume that  $\partial\Omega$  is of constant type along  $C$ .

## 2. Some notation and introductory remarks

We begin by defining the notion of type.

**Definition 2.1.** Let  $\Omega \subset \mathbb{C}^2$  be a bounded domain having a smooth boundary. Let  $p \in \partial\Omega$ . The *type* of  $p$ , denoted by  $\tau(p)$ , is the maximum order of contact that the germ of a 1-dimensional complex variety through  $p$  can have with  $\partial\Omega$  at  $p$ . The point  $p$  is said to be of finite type if  $\tau(p) < \infty$ . The domain  $\Omega$  is said to be of *finite type* if there is an  $N \in \mathbb{N}$  such that  $\tau(p) \leq N$  for each  $p \in \partial\Omega$ .

**Remark 2.2.** Let  $\Omega \subset \mathbb{C}^2$  be a smoothly bounded pseudoconvex domain. Suppose  $p \in \partial\Omega$  has type  $\tau(p) = N$  and there are local holomorphic coordinates  $(U; \zeta_1, \zeta_2)$ , near  $p$ , relative to which  $p = 0$  and relative to which  $U \cap \partial\Omega$  is defined by

$$(2-1) \quad \rho(\zeta) = A(\zeta_1) + O(v_2^2, |\zeta_1||v_2|) - u_2,$$

where  $\zeta_k := u_k + iv_k$ , and  $A(\zeta_1) = O(|\zeta_1|^2)$ . Then:

- (1)  $N$  is the leading order in  $\zeta_1$  of  $A$ .
- (2)  $N$  is an even number, because  $\Omega$  is pseudoconvex.

These are consequences of a computation on smoothly bounded pseudoconvex domains in  $\mathbb{C}^2$  of finite type at  $p \in \partial\Omega$ , given in [Fornæss and Stensønes 1987, Lecture 28]. Examining this calculation, we can infer that:

- (3) Suppose  $\Phi = (\phi_1, \phi_2) : (U, p) \rightarrow (\mathbb{C}^2, 0)$  is a smooth change of coordinate such that  $\bar{\partial}\phi_1$  and  $\bar{\partial}\phi_2$  vanish to infinite order at  $p$ , and such that  $(U \cap \partial\Omega)$  (with respect to these new coordinates) has a defining function of the form (2–1), where we have written  $\zeta_j = \phi_j(z_1, z_2)$  for  $j = 1, 2$ . Then conclusions (1) and (2) above continue to hold.

We now present some notation. For a  $\mathcal{C}^2$  function  $\phi$  defined in some open set in  $\mathbb{C}^n$ , we set

$$\begin{aligned} \partial_j \phi &= \frac{\partial \phi}{\partial z_j}, & \bar{\partial}_j \phi &= \frac{\partial \phi}{\partial \bar{z}_j}, \\ \partial_{jk}^2 \phi &= \frac{\partial^2 \phi}{\partial z_j \partial z_k}, & \bar{\partial}_{jk}^2 \phi &= \frac{\partial^2 \phi}{\partial \bar{z}_j \partial \bar{z}_k}. \end{aligned}$$

If  $F$  is a smooth function defined in a neighborhood of  $0 \in \mathbb{R}^N$ , we define (borrowing our notation from [Bloom 1978a])

$$\begin{aligned} \text{In}(F) &:= \text{the leading homogeneous polynomial} \\ &\quad \text{in the Taylor expansion of } F \text{ around } 0, \\ \text{ord}(F) &:= \text{the degree of } \text{In}(F). \end{aligned}$$

In what follows,  $B(p; r)$  will denote the open Euclidean ball in  $\mathbb{C}^2$  centered at  $p \in \mathbb{C}^2$  and having radius  $r$ , while  $D(a; r)$  will denote the open disc in  $\mathbb{C}$  centered at  $a \in \mathbb{C}$  and having radius  $r$ . Several parameters occur in our analysis and the independence of the quantitative estimates in the results below from these parameters will be of some concern. We will express such estimates via the notation  $X \lesssim Y$  — meaning that there is a constant  $C > 0$ , independent of all parameters, such that  $X \leq CY$ .

A standard approach [Henkin and Tumanov 1976; Rudin 1978] to proving that  $C \subset \partial\Omega$ , with  $C, \partial\Omega$  smooth, is a peak-interpolation set makes use of Bishop’s theorem:

**Theorem** [Bishop 1962]. *Let  $\Omega$  be a bounded domain in  $\mathbb{C}^n$ . A compact subset  $K \subset \partial\Omega$  is a peak-interpolation set for  $A(\Omega)$  if and only if  $|\mu|(K) = 0$  for every annihilating measure  $\mu \perp A(\Omega)$ .*

In this theorem, an *annihilating measure* is a regular, complex Borel measure on  $\bar{\Omega}$  which, viewed as a bounded linear functional on  $\mathcal{C}(\bar{\Omega})$ , annihilates  $A(\Omega)$ . A variation of the aforementioned approach—needed in the proof of our main theorem— involves showing that if for any  $p \in C$  there is a small neighborhood  $V_p \ni p$  such that for each bump function  $\chi \in \mathcal{C}_c^\infty(V_p; [0, 1])$  with  $\text{int}(\chi^{-1}\{1\}) \cap C$  being an open arc in  $C$ , there is a sequence of functions  $\{h_k\}_{k \in \mathbb{N}}$  such that

- (i)  $\{h_k\}_{k \in \mathbb{N}} \subset A(\Omega)$  and is uniformly bounded on  $\bar{\Omega}$ ;
- (ii)  $\lim_{k \rightarrow \infty} h_k(z) = 0$  for all  $z \in \bar{\Omega} \setminus (C \cap V_p)$ ;
- (iii)  $\lim_{k \rightarrow \infty} h_k(z) = \chi(z)$  for all  $z \in C \cap V_p$ .

We explain in the next section why Theorem 1.1(i) follows from the existence of such a  $\{h_k\}_{k \in \mathbb{N}}$ .

The key step in our proof is to show that if  $C$  is as described in Theorem 1.1(i), then for each  $p \in C$  we can find a small neighborhood  $V_p \ni p$  such that for any  $U \Subset V_p$  for which  $C \cap U$  is an arc, there is a smooth function  $G$  in  $V_p$  that is almost holomorphic with respect to  $C \cap V_p$  and peaks on  $C \cap \bar{U}$ . Further, one requires that this almost holomorphic peak function must approach the value 1 at a controlled rate. We show that

$$(2-2) \quad |G(z)| \leq 1 - C \text{dist}[z, C \cap V_p]^{2M} \quad \text{for all } z \in \bar{\Omega} \cap V_p.$$

Here  $2M$  represents the type of  $\partial\Omega$  along  $C$ . The above result is strongly reminiscent of [Noell 1985, Lemma 2.1]. In that lemma, if  $C$ —where  $C$  is *not necessarily* complex-tangential, but  $\partial\Omega$  is of type  $2M$  along  $C$ —has the property that at each  $p \in C$  there is a holomorphic function, smooth up to  $\partial\Omega$ , that peaks on a small closed sub-arc of  $C$  passing through  $p$ , then we can find a holomorphic peak function, smooth up to  $\partial\Omega$ , that satisfies the estimate (2-2). In our situation we do not, of course, have holomorphic functions that peak locally along  $C$ . However, we can use some of Noell's ideas (which in turn rely on an estimate from [Bloom 1978a]) and exploit the complex-tangency of  $C$  to construct an *almost-holomorphic* local peak function that satisfies good estimates. This construction is presented in Section 4.

We complete the proof of Theorem 1.1 in Section 5. Part (i) of the theorem will follow from the construction of the family  $\{h_k\}_{k \in \mathbb{N}}$  described above. Each  $h_k$  is, near  $C$ , a holomorphic correction of the  $k$ -th power of  $G$  ( $G$  as introduced above). This correction is achieved by solving an appropriate  $\bar{\partial}$ -equation in  $\bar{\Omega}$ , and the estimate (2-2) is used to show that  $h_k$  satisfies the three properties listed above. Theorem 1.1(ii) will follow from the fact that in the real-analytic setting  $\partial\Omega$  is of constant type along  $C$  *except* for a discrete set of points in  $C$ . Using part (i) of the

theorem and the fact that each point in this discrete set is a peak point for  $A(\Omega)$ , we deduce part (ii).

### 3. A technical lemma

In this section, we present an abstract lemma that is instrumental to the proof of our main theorem.

**Definition 3.1.** Given an open set  $V \subset \mathbb{R}^N$ , a *bump function*  $f$  in  $V$  is a function belonging to  $\mathcal{C}_c^\infty(V; [0, 1])$  such that  $\text{int}(f^{-1}\{1\}) \neq \emptyset$ .

**Lemma 3.2.** Let  $\Omega$  be a bounded domain in  $\mathbb{C}^2$  having smooth boundary and let  $C$  be a smooth curve in  $\partial\Omega$ . Assume that for each  $p \in C$ , there exists a small neighborhood  $V_p$  of  $p$  such that for each bump function  $\chi \in \mathcal{C}_c^\infty(V_p; [0, 1])$  for which  $\text{int}(\chi^{-1}\{1\}) \cap C$  is an arc, we can find a sequence of functions  $\{h_k\}_{k \in \mathbb{N}} \subset A(\Omega)$  (depending on  $\chi$ ) satisfying

- (i)  $\{h_k\}_{k \in \mathbb{N}} \subset A(\Omega)$  is uniformly bounded on  $\bar{\Omega}$ ;
- (ii)  $\lim_{k \rightarrow \infty} h_k(z) = 0$  for all  $z \in \bar{\Omega} \setminus (C \cap V_p)$ ;
- (iii)  $\lim_{k \rightarrow \infty} h_k(z) = \chi(z)$  for all  $z \in C \cap V_p$ .

Then  $C$  is a countable union of peak-interpolation sets for  $A(\Omega)$ .

**Remark 3.3.** A form of this lemma is true if  $\Omega$  is a bounded domain in  $\mathbb{C}^n$  and  $C$  is replaced by  $M \subset \partial\Omega$ , where  $M$  is a smooth submanifold of  $\partial\Omega \cap U$ ,  $U$  being an open subset of  $\mathbb{C}^n$ . However, to be able to derive the conclusion of the lemma in this new setting with  $\dim_{\mathbb{R}}(M) > 1$ , one would have to produce, for every bump function  $\chi \in \mathcal{C}_c^\infty(V_p; [0, 1])$  (not merely those for which  $\text{int}(\chi^{-1}\{1\}) \cap M$  is nice), an  $h \in A(\Omega)$  such that  $\{h^k : k \in \mathbb{N}\}$  would satisfy conditions (i)–(iii) above. Being able to find such an  $h$  could be rather difficult if  $\dim_{\mathbb{R}}(M) > 1$ , because  $\text{int}(\chi^{-1}\{1\}) \cap M$  could be structurally quite complicated in this situation. We add that if  $\partial\Omega$  is *strictly pseudoconvex*, a less exacting form of the above lemma — see, for instance, [Henkin and Tumanov 1976, Lemma 6] — suffices to infer peak-interpolation in higher dimensions.

*Proof of Lemma 3.2.* Fix  $p \in C$ . We may assume that  $C \cap V_p$  is an arc in  $C$ . Let  $K$  be any compact subset of  $C \cap V_p$  and let  $\mu$  be any annihilating measure. Then

$$K = (C \cap V_p) \setminus \bigsqcup_{k \in \mathbb{N}} \mathcal{A}_k,$$

where each  $\mathcal{A}_k$  is an open sub-arc of  $C \cap V_p$ . If we could show that  $\mu(\mathcal{A}_k) = 0$  for each  $k$  and that  $\mu(C \cap V_p) = 0$ , we could conclude by the additivity of  $\mu$  that  $\mu(K) = 0$ .

Let  $\mathcal{C} \subset C \cap V_p$  be any closed sub-arc of  $C$ . Let  $\{\mathcal{D}_\nu\}_{\nu \in \mathbb{N}}$  be a shrinking family of compact subsets of  $\mathbb{C}^2$  such that

- (a)  $\mathfrak{D}_{v+1} \subset \text{int}(\mathfrak{D}_v)$ ,
- (b)  $\bigcap_{v \in \mathbb{N}} \mathfrak{D}_v = \mathcal{C}$ ,
- (c)  $\mathfrak{D}_v \subseteq V_p$ ,
- (d)  $C \cap \mathfrak{D}_v$  is an arc.

Let  $\chi_v \in \mathcal{C}_c^\infty(V_p; [0, 1])$  be a bump function with

$$\chi_v|_{\mathfrak{D}_{v+1}} \equiv 1 \quad \text{and} \quad \text{supp } \chi_v \subseteq \mathfrak{D}_v.$$

Finally, define  $\{h_{k,v}\}_{k \in \mathbb{N}}$  to be the sequence of functions corresponding to  $\chi_v$  given by the hypothesis of this lemma.

Choose any  $\mu \perp A(\Omega)$ . By the bounded convergence theorem,

$$0 = \lim_{k \rightarrow \infty} \int_{\bar{\Omega}} h_{k,v} d\mu = \int_{C \cap V_p} \chi_v d\mu.$$

Another passage to the limit yields  $\mu(\mathcal{C}) = 0$ , and this is true for any  $\mu \perp A(\Omega)$ . As  $\mu$  is a regular measure, this shows that  $\mu(\mathcal{A}) = 0$  for any *open* sub-arc  $\mathcal{A} \subset C \cap V_p$ ; in particular  $\mu(C \cap V_p) = 0$ . Let  $\mathcal{V}_p$  be any neighborhood of  $p$  such that  $\mathcal{V}_p \Subset V_p$ . In view of our remarks in the first paragraph of this proof we have just shown that  $|\mu|(C \cap \overline{\mathcal{V}_p}) = 0$  for any  $\mu \perp A(\Omega)$ . By Bishop’s theorem,  $C \cap \overline{\mathcal{V}_p}$  is a peak-interpolation set for  $A(\Omega)$ . Letting  $p$  vary over a countable dense subset of  $C$ , we have the desired result. □

#### 4. Constructing an almost holomorphic function that peaks locally on $C$

Let  $p \in \partial\Omega$ . In this section, we will study  $\partial\Omega$  near  $p$  with respect to a convenient system of local coordinates that are almost holomorphic with respect to  $C$  (near  $p$ ), where  $\Omega$  and  $C$  are as in Theorem 1.1(i). The following lemma asserts the existence of local coordinates having the desired properties:

**Lemma 4.1.** *Let  $\Omega$  be a bounded domain in  $\mathbb{C}^2$  having smooth boundary and let  $C \subset \partial\Omega$  be a complex-tangential curve. Let  $p \in C$ . There is a neighborhood  $\omega \ni p$  and a  $\mathcal{C}^\infty$ -diffeomorphism  $\Phi : (\omega, p) \rightarrow (\mathbb{C}^2, 0)$  which is almost holomorphic with respect to  $(C \cap \omega)$  and such that, writing  $(\zeta_1, \zeta_2) := \Phi(z_1, z_2)$ , we have:*

- (1)  $\Phi(C \cap \omega) \subset \{(\zeta_1, \zeta_2) : \text{im}(\zeta_1) = \zeta_2 = 0\}$ .
- (2)  $\Phi(\partial\Omega \cap \omega)$  is defined by a defining function of the form

$$\rho(\zeta) = A(\zeta_1) + B(\zeta_1)v_2 + R(\zeta_1, v_2) - u_2,$$

where  $\zeta_k = u_k + i v_k$  for  $k = 1, 2$ ,  $A(\zeta_1) = O(|\zeta_1|^2)$ ,  $R(\zeta_1, v_2) = O(|v_2|^2)$ , and

$$A(u_1) = B(u_1) = 0 \quad \text{and} \quad \nabla A(u_1) = \nabla B(u_1) = 0 \quad \text{for all } u_1 \text{ near } 0.$$

*Proof.* Without loss of generality, we may let  $p$  be the origin, and assume that, near  $p$ ,  $\partial\Omega$  is defined by

$$r(z_1, z_2) = h(z_1, \operatorname{im} z_2) - \operatorname{Re} z_2,$$

where  $h(0) = 0$  and  $\nabla h(0) = 0$ .

Let  $\omega$  be a neighborhood of  $p = 0$  and let  $\mathcal{M} \subset \partial\Omega$  be the smooth 2-manifold of  $\omega$  formed by the integral curves to the vector-field  $-\mathbb{J}(\nabla r)$  passing through  $(C \cap \omega)$ .  $\mathcal{M}$  is totally real. Let

$$\gamma = (\gamma_1, \gamma_2) : (B(0; \varepsilon), (u_1, u_2) = 0) \rightarrow ((\mathcal{M} \cap \omega), p = 0)$$

parametrize  $\mathcal{M}$  near  $p = 0$  in such a way that, for each  $c$ ,  $\operatorname{Image}(\gamma|_{\{u_2=c\}})$  is an integral curve to the unit section of  $T(\mathcal{M}) \cap H(\partial\Omega)|_{\mathcal{M}}$ , with

$$\operatorname{Image}(\gamma|_{\{u_2=0\}}) = C \cap \omega \quad \text{and} \quad \frac{\partial \gamma(0, 0)}{\partial u_2} = -\mathbb{J}(\nabla r)(0, 0).$$

Shrinking  $\omega$  if necessary, we construct a diffeomorphism  $\Phi : (\omega, p = 0) \rightarrow (\mathbb{C}^2, 0)$  of class  $\mathcal{C}^\infty$  that is almost holomorphic with respect to  $(\mathcal{M} \cap \omega)$ , by defining

$$\Phi^{-1}(\zeta_1, \zeta_2) = (\Gamma_1(\zeta_1, -i\zeta_2), \Gamma_2(\zeta_1, -i\zeta_2)) := \eta(\zeta_1, \zeta_2),$$

where  $\zeta_k := u_k + i v_k$  for  $k = 1, 2$  and  $\Gamma_k$  is an almost holomorphic extension of  $\gamma_k$  for  $k = 1, 2$ . By construction,

$$(4-1) \quad \begin{aligned} \Phi(\mathcal{M} \cap \omega) &\subset \{(\zeta_1, \zeta_2) : v_1 = u_2 = 0\}, \\ \Phi(C \cap \omega) &\subset \{(\zeta_1, \zeta_2) : v_1 = \zeta_2 = 0\}. \end{aligned}$$

Now,  $\Phi(\partial\Omega \cap \omega)$  is defined by

$$\rho(\zeta_1, \zeta_2) = r \circ \Phi^{-1}(\zeta_1, \zeta_2).$$

We expand  $\rho$  around the origin in a Taylor series. We make use of the fact that  $\Gamma_k$  are almost holomorphic with respect to  $\{(\zeta_1, \zeta_2) \mid v_1 = v_2 = 0\}$  to get

$$\rho(\zeta) = 2 \operatorname{Re} \left( \sum_{j=1}^2 \frac{\partial r}{\partial z_j}(\eta(0, 0)) \left( \frac{\partial \Gamma_j}{\partial \zeta_1}(0, 0) \zeta_1 + (-i) \frac{\partial \Gamma_j}{\partial \zeta_2}(0, 0) \zeta_2 \right) \right) + \mathcal{O}(|\zeta|^2).$$

Using the fact that

$$\frac{\partial \Gamma_j}{\partial \zeta_k}(0, 0) = \frac{\partial \gamma_j}{\partial u_k}(0, 0) \quad \text{for } j, k = 1, 2,$$

we get

$$\begin{aligned} \rho(\zeta) &= 2 \operatorname{Re} \left( \sum_{j=1}^2 \frac{\partial r}{\partial z_j}(\gamma(0, 0)) \frac{\partial \gamma_j}{\partial u_1}(0, 0) \zeta_1 \right. \\ &\quad \left. + (-i) \sum_{j=1}^2 \frac{\partial r}{\partial z_j}(\gamma(0, 0)) \frac{\partial \gamma_j}{\partial u_2}(0, 0) \zeta_2 \right) + O(|\zeta|^2) \\ &= 2 \operatorname{Re} \left( (-i) \sum_{j=1}^2 \frac{\partial r}{\partial z_j}(\gamma(0, 0)) \frac{\partial \gamma_j}{\partial u_2}(0, 0) \zeta_2 \right) + O(|\zeta|^2) \\ &= -u_2 + O(|\zeta|^2). \end{aligned}$$

The second equality follows from the complex-tangency of  $\operatorname{Image}(\gamma(\cdot, 0))$ , which implies

$$\sum_{j=1}^2 \frac{\partial r}{\partial z_j}(\gamma(u_1, 0)) \frac{\partial \gamma_j}{\partial u_1}(u_1, 0) = 0 \quad \text{for all } u_1 \in (-\varepsilon, \varepsilon),$$

and the last equality follows from the normalization condition on  $\partial\gamma(0, 0)/\partial u_2$ . We see that the only term in the expansion above that has first order in either  $\zeta_1$  or  $\zeta_2$  is  $-u_2$ . Hence, the hypersurface  $\Phi(\partial\Omega \cap \omega)$  is tangent at 0 to the hyperplane  $\mathbf{H} := \{(\zeta_1, \zeta_2) \in \mathbb{C}^2 \mid u_2 = 0\}$ . Thus, we can find, near  $0 \in \mathbb{C}^2$ , a defining function — and for convenience of notation, we will continue to call it  $\rho$  — having the form

$$(4-2) \quad \rho(\zeta) = A(\zeta_1) + B(\zeta_1)v_2 + R(\zeta_1, v_2) - u_2,$$

where  $A(\zeta_1) = O(|\zeta_1|^2)$  and  $R(\zeta_1, v_2) = O(|v_2|^2)$ . Then, since  $\Phi(C \cap \omega)$  is contained in  $\Phi(\partial\Omega \cap \omega)$ , setting  $v_1 = \zeta_2 = 0$  in (4-2), we get

$$(4-3) \quad A(u_1) = 0 \quad \text{for all } (u_1, 0) \in \Phi(C \cap \omega).$$

And since  $\Phi(\mathcal{M} \cap \omega) \subset \Phi(\partial\Omega \cap \omega)$ , setting  $v_1 = u_2 = 0$  in (4-2), we see that  $B(u_1)v_2 + O(|v_2|^2) = 0$  for all  $(u_1, v_2)$  belonging to a small neighborhood 0. Thus

$$(4-4) \quad B(u_1) = 0 \quad \text{for all } (u_1, 0) \in \Phi(C \cap \omega).$$

By construction,  $(\nabla\rho)(u_1, v_2)$  is a normal vector to  $\Phi(\mathcal{M} \cap \omega)$  for all  $(u_1, v_2) \in \Phi(\mathcal{M} \cap \omega)$ . This implies that  $T_{(u_1, v_2)}(\Phi(\partial\Omega \cap \omega)) = \mathbf{H}$  for all  $(u_1, v_2) \in \Phi(\mathcal{M} \cap \omega)$ . Computing  $(\nabla\rho)(u_1, v_2)$ , we see that  $\nabla A(u_1) + \nabla B(u_1)v_2 = 0$  for all  $(u_1, v_2)$  in a neighborhood of 0. Thus

$$(4-5) \quad \nabla A(u_1) = \nabla B(u_1) = 0 \quad \text{for all } (u_1, 0) \in \Phi(C \cap \omega).$$

By (4-3), (4-4) and (4-5), we have the desired result. □

We now state the key lemma of this paper. It concerns the construction of an almost holomorphic peak function of the type discussed in Section 2.

**Proposition 4.2.** *Let  $\Omega$  be a bounded pseudoconvex domain in  $\mathbb{C}^2$  of finite type, and let  $\partial\Omega$  be of class  $\mathcal{C}^\infty$ . Let  $C \subset \partial\Omega$  be a complex-tangential curve of class  $\mathcal{C}^\infty$ , and let  $\partial\Omega$  be of constant type  $2M$  along  $C$ . Let  $p \in C$ . There exists a neighborhood  $V \equiv V(p)$  of  $p$  and a uniform constant  $C > 0$ , and for any open set  $U \Subset V$  such that  $C \cap U$  is an arc, there is a neighborhood  $V_1 \equiv V(p, U)$  of  $p$  satisfying  $C \cap V_1 = C \cap V$  and a function  $G \in \mathcal{C}^\infty(V_1)$  —  $G$  depending on  $p$  and  $U$  — that satisfies*

- (1)  $G^{-1}\{1\} = C \cap \bar{U}$ ;
- (2)  $\bar{\partial}G$  vanishes to infinite order on  $V \cap C$ ;
- (3)  $|G(z)| \leq 1 - C \operatorname{dist}[z, C \cap V]^{2M}$  for each  $z \in \bar{\Omega} \cap V_1$ .

*Proof.* Let  $\omega \ni p$  and  $\Phi : (\omega, p) \rightarrow (\mathbb{C}^2, 0)$  be the change of coordinate described in Lemma 4.1. Let  $\Phi(\partial\Omega \cap \omega)$  be defined by

$$(4-6) \quad \rho(\zeta_1, \zeta_2) = A(\zeta_1) + B(\zeta_1)v_2 + R(\zeta_1, v_2) - u_2.$$

Consider a point  $(x_0, 0) \in \Phi(C \cap \omega)$  and let

$$(4-7) \quad \varrho_{x_0}(\zeta_1^*, \zeta_2) = \mathcal{A}_{x_0}(\zeta_1^*) + \mathcal{B}_{x_0}(\zeta_1^*)v_2 + \mathcal{R}_{x_0}(\zeta_1^*, v_2) - u_2$$

represent the expansion of  $\rho$  in (4-6) around  $(x_0, 0)$ , where  $\zeta_1^* := \zeta_1 - x_0$ .

**Claim 1.** Shrinking  $\omega$  if necessary, there is a  $c > 0$  such that

$$(4-8) \quad A(u_1 + iv_1) \geq cv_1^{2M}, \quad \text{for all } \zeta_1 \text{ such that } \zeta \in \Phi(\omega).$$

As  $A(x_0) = B(x_0) = 0$  and  $\nabla A(x_0) = \nabla B(x_0) = 0$  for each  $(x_0, 0) \in \Phi(\partial\Omega \cap \omega)$ , the right-hand side of (4-7) represents a defining function of the form (2-1). By Remark 2.2(3), the function  $\mathcal{A}_{x_0}$  in (4-7) must vanish to order  $2M$  at 0, whereby the function  $A$  in (4-6) must vanish precisely to order  $2M$  at each  $(u_1, 0) \in \Phi(\partial\Omega \cap \omega)$ . Now write

$$(4-9) \quad A(u_1 + iv_1) = a_J(u_1)v_1^J + O(|v_1|^{J+1}),$$

where  $J$  is the least positive integer  $k$  such that  $a_k \neq 0$  near  $u_1 = 0$ . By our remarks above, it is clear that  $J \leq 2M$ . But, if  $J < 2M$ , then if  $\tilde{u}_1$  is such that  $a_J(\tilde{u}_1) \neq 0$ , then  $A$  vanishes to order  $< 2M$  at  $u_1 + iv_1 = \tilde{u}_1$ , which contradicts our remarks above. Thus,  $J = 2M$  in (4-9) and

$$A(u_1 + iv_1) = a_{2M}(u_1)v_1^{2M} + O(|v_1|^{2M+1}).$$

and  $a_{2M}(0) \neq 0$ . Now recall that  $\Phi$  is almost-holomorphic with respect to  $(\mathcal{M} \cap \omega)$ . If, in fact,  $(u_1 + iv_1, u_2 + iv_2)$  were *holomorphic* coordinates, the pseudoconvexity

of  $\Omega$  would have implied that

$$\alpha : (u_1, v_1) \mapsto a_{2M}(u_1)v_1^{2M} \text{ is subharmonic,}$$

$$\Delta\alpha(u_1, v_1) > 0 \text{ off } \{v_1 = 0\}, \text{ and } (u_1, v_1) \text{ close to } 0.$$

This would have implied that  $a_{2M}(u_1) > 0$  for  $u_1$  close to 0 (the second statement above follows from an obvious calculation). In our present situation, the coordinates  $(u_1 + iv_1, u_2 + iv_2)$  differ from holomorphic ones by terms vanishing to arbitrarily high order along  $(C \cap \omega)$ . From the last two facts, we can conclude, after shrinking  $\omega$  if necessary, that

$$a_{2M}(u_1) > 0 \text{ for all } (u_1, 0) \in \Phi(\partial\Omega \cap \omega).$$

From this final fact, we deduce (4–8). Hence the claim.

**Claim 2.** We can find  $\omega_1 \Subset \omega$  and a uniform constant  $T > 0$  such that

$$(4-10) \quad B(\zeta_1)^2 \leq TA(\zeta_1) \text{ for all } \zeta \in \Phi(\bar{\Omega} \cap \omega_1).$$

To see this, we use a procedure originating in [Bloom 1978a, Section 3]. Write  $q = (x_0, 0) \in \Phi(C \cap \omega)$ . The positivity of the Levi form for  $\partial\Omega$  on the complex tangent vectors implies that, *were*  $(u_1 + iv_1, u_2 + iv_2)$  *holomorphic coordinates*, there would be a  $\delta > 0$  such that the function  $\mathfrak{L}$  induced by the Levi form

$$\mathfrak{L} : D(x_0; \delta) \times (-\delta, \delta) \rightarrow \mathbb{R}$$

defined by

$$\mathfrak{L} = |\partial_{\bar{2}}\rho|^2 \partial_{11}^2\rho + |\partial_{\bar{1}}\rho|^2 \partial_{22}^2\rho - 2\text{Re}(\partial_1\rho \partial_{\bar{2}}\rho \partial_{12}^2\rho)$$

would be nonnegative (notice that  $\mathfrak{L}$  is independent of  $u_2$ ). In our present situation, however,  $\mathfrak{L}(u_1, v_2) \geq 0$  for all  $(u_1, v_2) \in \Phi(\mathcal{M} \cap \omega)$ .

Write

$$\mathfrak{L}(\zeta_1, v_2) = \mathfrak{L}^{(0)}(\zeta_1) + v_2\mathfrak{L}^{(1)}(\zeta_1) + v_2^2\mathfrak{L}^{(2)}(\zeta_1) + O(|v_2|^3).$$

It has been shown in [Bloom 1978a] that if  $\text{ord } B < \text{ord } A$ , then

$$(4-11) \quad \begin{aligned} \text{In}(\mathfrak{L}^{(0)}) &= \frac{1}{4}\text{In}(\partial_{11}^2 A), & \text{ord } \mathfrak{L}^{(0)} &= \text{ord } A - 2, \\ \text{In}(\mathfrak{L}^{(1)}) &= \frac{1}{4}\text{In}(\partial_{11}^2 B), & \text{ord } \mathfrak{L}^{(1)} &= \text{ord } B - 2. \end{aligned}$$

If already  $2 \text{ord } B \geq \text{ord } A$ , then (4–10) would follow trivially. Thus, *assume* that  $2 \text{ord } B < \text{ord } A$ . Write  $r = \text{ord } B$ . We have

$$\frac{1}{\lambda^{2r-2}}\mathfrak{L}(\lambda(u_1 + i0), \lambda^r v_2) \geq 0$$

for all  $(u_1, v_2) \in (x_0 - \delta, x_0 + \delta) \times (-\delta, \delta)$  and  $\lambda \in \mathbb{R}_+$ . But from (4–11) and our assumption, we get

$$(4-12) \quad \lim_{\lambda \rightarrow 0^+} \frac{1}{\lambda^{2r-2}} \mathfrak{L}(\lambda u_1, \lambda^r v_2) = \frac{v_2}{4} \text{In}(\partial_{1\bar{1}}^2 B)(\zeta_1).$$

Write

$$B(u_1 + i v_1) = b_J(u_1) v_1^J + O(|v_1|^{J+1}),$$

where  $J$  is the least positive integer  $k$  such that  $b_k \neq 0$  near  $u_1 = 0$ . By Lemma 4.1(2),  $J \geq 2$ , whence  $\text{In}(B)$  is nonharmonic near 0. So, as  $v_2$  occurs linearly in the right-hand side of (4–12), it is impossible that

$$\frac{1}{4} v_2 \text{In}(\partial_{1\bar{1}}^2 B)(u_1) \geq 0, \quad \text{for all } (u_1, v_2) \in (x_0 - \delta, x_0 + \delta) \times (-\delta, \delta).$$

This results in a contradiction. So  $2 \text{ord } B \geq \text{ord } A$ , which, in conjunction with the positivity of  $A$ , namely (4–8), yields (4–10).

Finally, define  $H : \Phi(\bar{\Omega} \cap \omega_1) \rightarrow \mathbb{C}$  by

$$H(\zeta) = \zeta_2 - \alpha \zeta_2^2,$$

for  $\alpha > 0$  chosen appropriately large. We choose  $\alpha$  as follows: Observe that

$$\begin{aligned} & \frac{1}{2} A(\zeta_1) + B(\zeta_1) v_2 + R(\zeta_1, v_2) + \frac{1}{6} \alpha v_2^2 \\ &= \left( \frac{T}{\sqrt{2}} v_2 + \frac{B(\zeta_1)}{\sqrt{2}T} \right)^2 + \frac{1}{2T} (TA(\zeta_1) - B(\zeta_1)^2) - \frac{T^2}{2} v_2^2 + R(\zeta_1, v_2) + \frac{\alpha}{6} v_2^2. \end{aligned}$$

The first two terms of the right-hand side are positive, in view of (4–10). So we shrink  $\omega_1$  appropriately and choose  $\alpha > 0$  so large that

$$(4-13) \quad \frac{1}{2} A(\zeta_1) + B(\zeta_1) v_2 + R(\zeta_1, v_2) + \frac{1}{6} \alpha v_2^2 \geq 0, \quad \text{for all } \zeta \in \Phi(\bar{\Omega} \cap \omega_1).$$

Now consider:

*Case (i):*  $u_2 \geq 0$ . Let  $\varepsilon_1 > 0$  be so small that  $B(p; \varepsilon_1) \subset \omega_1$  and

$$(u_2 - \alpha u_2^2) \geq \frac{1}{2} u_2 \quad \text{for } \zeta \in \Phi(\bar{\Omega} \cap B(p; \varepsilon_1)).$$

Then, for all such  $\zeta$ , we have

$$\begin{aligned} (4-14) \quad \text{Re } H(\zeta) &= (u_2 - \alpha u_2^2) + \alpha v_2^2 \\ &\geq \frac{1}{2} u_2 + \alpha v_2^2 = \frac{1}{4} u_2 + \frac{1}{2} \alpha v_2^2 + \frac{1}{4} (u_2 + 2\alpha v_2^2) \\ &\geq \frac{1}{4} u_2 + \frac{1}{2} \alpha v_2^2 + \frac{1}{4} ((A(\zeta_1) + B(\zeta_1) v_2 + R(\zeta_1, v_2)) + 2\alpha v_2^2) \\ &= \frac{1}{4} u_2 + \frac{1}{8} A(\zeta_1) + \frac{1}{2} \alpha v_2^2 + \frac{1}{4} (\frac{1}{2} A(\zeta_1) + B(\zeta_1) v_2 + R(\zeta_1, v_2) + 2\alpha v_2^2) \\ &\gtrsim u_2^2 + v_2^2 + A(\zeta_1), \quad \text{using (4-13)}. \end{aligned}$$

Case (ii):  $u_2 < 0$ . Let  $\varepsilon_2 > 0$  be so small that  $B(p; \varepsilon_2) \subset \omega_1$  and that  $(u_2 - \alpha u_2^2) \geq 2u_2$  for  $\zeta \in \Phi(\bar{\Omega} \cap B(p; \varepsilon_2))$ . Then, for all such  $\zeta$ , we have (arguing exactly as before)

(4-15)

$$\begin{aligned} \operatorname{Re} H(\zeta) &\geq -u_2 + \frac{1}{2}\alpha v_2^2 + 3(u_2 + \frac{1}{6}\alpha v_2^2) \\ &\geq -u_2 + \frac{3}{2}A(\zeta_1) + \frac{1}{2}\alpha v_2^2 + 3(\frac{1}{2}A(\zeta_1) + B(\zeta_1)v_2 + R(\zeta_1, v_2) + \frac{1}{6}\alpha v_2^2) \\ &\gtrsim u_2^2 + v_2^2 + A(\zeta_1), \quad \text{using (4-13)}. \end{aligned}$$

Now let  $\varepsilon_0 = \min(\varepsilon_1, \varepsilon_2)$ . From (4-8), (4-14) and (4-15) we see that there is a uniform constant  $\kappa > 0$  such that

$$\begin{aligned} (4-16) \quad \operatorname{Re} H(\zeta) &\geq \kappa(u_2^2 + v_2^2 + v_1^{2M}) \\ &\geq \kappa \operatorname{dist}[\zeta, \Phi(C \cap B(p; \varepsilon_0))]^{2M} \quad \text{for } \zeta \in \Phi(\bar{\Omega} \cap B(p; \varepsilon_0)). \end{aligned}$$

Write  $\Phi(C \cap U) = (a, b)$ , and without loss of generality, assume that  $a < 0 < b$ . Define the function  $\phi$  by

$$\phi(u_1) = \begin{cases} \exp(1/(u_1 - a)) & \text{if } u_1 < a, \\ 0 & \text{if } a \leq u_1 \leq b, \\ \exp(-1/(u_1 - b)) & \text{if } u_1 > b. \end{cases}$$

Let  $r > 0$  such that  $B(0; r) \supset \Phi(B(p; \varepsilon_0))$ , and let  $R(\sigma)$  be the rectangle

$$R(\sigma) = \{(u_1 + i v_1) \in \mathbb{C} \mid |u_1| < r, |v_1| < \sigma\}.$$

By an argument given in [Noell 1985, Lemma 2.1], there exists a smooth almost holomorphic extension  $\tilde{\phi}$  of  $\phi$  and a  $\sigma > 0$  small enough that

$$(4-17) \quad \operatorname{Re}(\tilde{\phi}(u_1 + i v_1)) \geq -\frac{1}{2}\kappa v_1^{2M}, \quad u_1 + i v_1 \in R(\sigma).$$

We set

$$V_1(p, U) = B(p; \varepsilon_0) \cap \Phi^{-1}(\operatorname{Image} \Phi \cap (R(\sigma) \times \mathbb{C})).$$

From (4-16) and (4-17), we infer that the function  $G(z) = (1 - \tilde{\phi}) \circ \Phi(z) - H \circ \Phi(z)$  satisfies (1)-(3). □

### 5. The proof of Theorem 1.1

Statement (i). Let  $C$  be as in Theorem 1.1(i), and fix  $p \in C$ . Let  $V(p)$  be a neighborhood of  $p$  as given by Proposition 4.2. We will use Lemma 3.2 to provide a proof. Take  $V_p$ , in the notation of that lemma, to be  $V(p)$ . In the notation of Lemma 3.2, let  $\chi \in \mathcal{C}_c^\infty(V_p; [0, 1])$  be a bump function such that  $\operatorname{int}(\chi^{-1}\{1\}) \cap C$

is an arc. Write  $U = \text{int}(\chi^{-1}\{1\})$ . Now set  $V_1 = V_1(p, U)$  and let  $G \in \mathcal{C}^\infty(V_1)$  be as given by Proposition 4.2.

Define

$$G_k(z) = \begin{cases} G(z)^k \chi(z) & \text{if } z \in \bar{\Omega} \cap V_1, \\ 0 & \text{if } z \in \bar{\Omega} \setminus V_1. \end{cases}$$

Also define

$$(5-1) \quad f_k(z) = \bar{\partial} G_k(z) = k G(z)^{k-1} \bar{\partial} G(z) \chi(z) + G(z)^k \bar{\partial} \chi(z).$$

For a  $(0, 1)$  form  $\phi(z) = \phi_1(z)d\bar{z}_1 + \phi_2(z)d\bar{z}_2$  defined on  $\bar{\Omega}$ , define

$$\|\phi\|_{\bar{\Omega}} := \max \{ \sup_{\bar{\Omega}} |\phi_1(z)|, \sup_{\bar{\Omega}} |\phi_2(z)| \}.$$

By construction,

$$(5-2) \quad \|G^k \bar{\partial} \chi\|_{\bar{\Omega}} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Notice that  $\bar{\partial} G$  vanishes to infinite order wherever  $G(z) = 1$ . Thus, for  $j = 1, 2$ ,

$$(5-3) \quad |k G(z)^{k-1} \partial_j G(z) \chi(z)| \lesssim k(1 - C \text{dist}[z, C \cap V_p]^{2M})^{k-1} |\partial_j G(z)| \rightarrow 0$$

uniformly as  $k \rightarrow \infty$ .

From (5-2) and (5-3),

$$(5-4) \quad \|f_k\|_{\bar{\Omega}} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Now consider on  $\Omega$  the  $\bar{\partial}$ -equations

$$\bar{\partial} u_k = f_k.$$

We need Lipschitz estimates for the solution of the  $\bar{\partial}$ -equation on pseudoconvex domains in  $\mathbb{C}^2$  of finite type. Such estimates may be found in several places in the literature; for instance, in the results of Chang, Nagel and Stein [Chang et al. 1992], which imply that

$$(5-5) \quad \|u_k\|_{\bar{\Omega}} \leq \|u_k\|_{\Lambda^{1/N}(\bar{\Omega})} \leq C^* \|f_k\|_{\bar{\Omega}},$$

where  $N$  is a positive integer such that  $\tau(p) \leq N$  for each  $p \in \partial\Omega$ ,  $\Lambda^{1/N}(\bar{\Omega})$  is the class of complex-valued Lipschitz functions on  $\bar{\Omega}$  of order  $1/N$ , and  $C^* > 0$  is a constant depending only on  $\Omega$ . From (5-4) and (5-5) we see that  $\|u_k\|_{\bar{\Omega}} \rightarrow 0$ , whence, defining

$$h_k(z) = G_k(z) - u_k(z) \quad \text{for all } z \in \bar{\Omega},$$

we have a sequence of  $A(\Omega)$  functions with

$$\lim_{k \rightarrow \infty} h_k(z) = \lim_{k \rightarrow \infty} G_k(z) = \begin{cases} \chi(z) & \text{if } z \in C \cap V_p, \\ 0 & \text{if } z \in \bar{\Omega} \setminus (C \cap V_p). \end{cases}$$

Notice that, by construction, the sequence  $\{h_k\}_{k \in \mathbb{N}}$  is uniformly bounded. The sequence  $\{h_k\}_{k \in \mathbb{N}} \subset A(\Omega)$  satisfies hypotheses (i)–(iii) in Lemma 3.2 for the bump function  $\chi \in \mathcal{C}_c^\infty(V_p; [0, 1])$  such that  $\text{int}(\chi^{-1}\{1\}) \cap C$  is an arc. Thus we conclude, using Lemma 3.2, that any compact subset of  $C$  is a peak-interpolation set for  $A(\Omega)$ .

Statement (ii). In the present situation,  $\Omega$  is a bounded domain having a real-analytic boundary and  $C$  is a real-analytic complex-tangential curve. Let  $B$  be an open ball in  $\mathbb{C}^2$  and let  $\gamma : (-2\varepsilon, 2\varepsilon) \rightarrow C$  be an injective real-analytic parametrization of  $C$  locally such that  $\text{Image}(\gamma|_{[-\varepsilon, \varepsilon]}) = (C \cap \bar{B})$ . Let  $p \in (C \cap \bar{B})$  be such that

$$\tau(p) = \min_{q \in C \cap \bar{B}} \tau(q).$$

Write  $\tau(p) = 2M$ .

Recall that

$$H_p \otimes \mathbb{C}(\partial\Omega) = H_p^{1,0}(\partial\Omega) \oplus H_p^{0,1}(\partial\Omega),$$

where  $H \otimes \mathbb{C}(\partial\Omega)$  is the complexification of  $H(\partial\Omega)$ , and that  $H_p^{1,0}(\partial\Omega)$  and  $H_p^{0,1}(\partial\Omega)$  are the eigenspaces of the complex-structure map  $\mathbb{J}$  corresponding to  $+i$  and  $-i$  respectively. Without loss of generality, we may assume that there is an open set  $U \supset \bar{B}$  and a real-analytic section  $L$  of  $H^{1,0}(\partial\Omega)|_U$  such that  $L(q)$  spans  $H_q^{1,0}(\partial\Omega)$  and

$$L(q) \in \{v \in H_q^{1,0}(\partial\Omega) : \|v\| = 1\}$$

for each  $q \in (\partial\Omega \cap U)$ . Now consider the real-analytic function  $\mathfrak{L} : S^1 \times I \rightarrow \mathbb{R}$  defined by

$$\mathfrak{L}(\zeta, t) = \sum_{\substack{j+k=2M \\ 1 \leq j < 2M}} L^{j-1} \bar{L}^{k-1} \langle [L, \bar{L}], \partial\rho \rangle (\gamma(t)) \zeta^j \bar{\zeta}^k,$$

where  $I$  is an open interval around  $[-\varepsilon, \varepsilon]$ ,  $S^1$  is the unit circle in  $\mathbb{C}$  and  $\rho$  is a defining function of  $\partial\Omega$ . Let  $t_0$  be such that  $\gamma(t_0) = p$ . By [Bloom 1978b, Theorem 3.3],  $\tau(p) = 2M$  implies that there exists a  $\zeta_0 \in S^1$  such that  $\mathfrak{L}(\zeta_0, t_0) \neq 0$ . Then, by the real-analyticity of  $\mathfrak{L}$ , we conclude that

$$\{t \in [-\varepsilon, \varepsilon] : \mathfrak{L}(\zeta_0, t) = 0\} \text{ is a finite set } \mathfrak{S} \subset [-\varepsilon, \varepsilon].$$

Write  $\mathfrak{S} = \{t_1, \dots, t_N\}$ . Again by [Bloom 1978b, Theorem 3.3],  $\partial\Omega$  is of constant type  $2M$  in each connected component of  $(C \cap \bar{B}) \setminus \{\gamma(t_1), \dots, \gamma(t_N)\}$ . Therefore, by Theorem 1.1(i),

$(C \cap \bar{B}) \setminus \{\gamma(t_1), \dots, \gamma(t_N)\}$  is a countable union of peak-interpolation sets.

Recall that  $\Omega$  is a bounded domain with real-analytic boundary. By [Bedford and Fornæss 1978], therefore, every point of  $\partial\Omega$  is a peak point for  $A(\Omega)$ . So, each  $\gamma(t_j)$ , for  $j = 1, \dots, N$ , is a peak point for  $A(\Omega)$ . This, together with the fact that  $(C \cap \bar{B}) \setminus \{\gamma(t_1), \dots, \gamma(t_N)\}$  is a countable union of peak-interpolation sets, implies that  $C$  is a countable union of peak-interpolation sets for  $A(\Omega)$ , and that each compact subset of  $C$  is a peak-interpolation set for  $A(\Omega)$ .

### Acknowledgement

The author wishes to thank Alexander Nagel for several useful discussions during the course of this work.

### References

- [Bedford and Fornæss 1978] E. Bedford and J. E. Fornæss, “A construction of peak functions on weakly pseudoconvex domains”, *Ann. of Math. (2)* **107**:3 (1978), 555–568. MR 58 #11520 Zbl 0392.32004
- [Bharali 2004] G. Bharali, “On peak-interpolation manifolds for  $A(\Omega)$  for convex domains in  $\mathbb{C}^n$ ”, *Trans. Amer. Math. Soc.* **356**:12 (2004), 4811–4827. MR 2084399 Zbl 02097164
- [Bishop 1962] E. Bishop, “A general Rudin–Carleson theorem”, *Proc. Amer. Math. Soc.* **13** (1962), 140–143. MR 24 #A3293 Zbl 0101.08807
- [Bloom 1978a] T. Bloom, “ $\mathcal{C}^\infty$  peak functions for pseudoconvex domains of strict type”, *Duke Math. J.* **45**:1 (1978), 133–147. MR 58 #6353 Zbl 0376.32014
- [Bloom 1978b] T. Bloom, “Remarks on type conditions for real hypersurfaces in  $\mathbb{C}^n$ ”, pp. 14–24 in *Several complex variables* (Cortona, 1976/1977), Scuola Norm. Sup. Pisa, 1978. MR 84f:32022 Zbl 0421.32022
- [Chang et al. 1992] D.-C. Chang, A. Nagel, and E. M. Stein, “Estimates for the  $\bar{\partial}$ -Neumann problem in pseudoconvex domains of finite type in  $\mathbb{C}^2$ ”, *Acta Math.* **169**:3 (1992), 153–228. MR 93k:32025 Zbl 0821.32011
- [Fornæss and McNeal 1994] J. E. Fornæss and J. D. McNeal, “A construction of peak functions on some finite type domains”, *Amer. J. Math.* **116**:3 (1994), 737–755. MR 95j:32023 Zbl 0809.32005
- [Fornæss and Sibony 1989] J. E. Fornæss and N. Sibony, “Construction of P.S.H. functions on weakly pseudoconvex domains”, *Duke Math. J.* **58**:3 (1989), 633–655. MR 90m:32034 ZBL 0679.32017
- [Fornæss and Stensønes 1987] J. E. Fornæss and B. Stensønes, *Lectures on counterexamples in several complex variables*, Mathematical Notes **33**, Princeton University Press, Princeton, NJ, 1987. MR 88f:32001 Zbl 0626.32001
- [Henkin and Tumanov 1976] A. E. Tumanov and G. M. Henkin, “Interpolation submanifolds of pseudoconvex manifolds”, pp. 74–86 in *Математическое программирование и смежные вопросы: Выпуклое программирование* (Drogobych, 1974), edited by B. S. Mitjagin and B. P. Titarenko, Central. Èkonom.-Mat. Inst. Akad. Nauk SSSR, Moscow, 1976. In Russian; translated in *Translations Amer. Math. Soc.*, **115** (1980), 59–69. MR 58 #22665 Zbl 0455.32009
- [Nagel 1976] A. Nagel, “Smooth zero sets and interpolation sets for some algebras of holomorphic functions on strictly pseudoconvex domains”, *Duke Math. J.* **43**:2 (1976), 323–348. MR 56 #670 Zbl 0343.32016

- [Nagel and Rudin 1978] A. Nagel and W. Rudin, “Local boundary behavior of bounded holomorphic functions”, *Canad. J. Math.* **30**:3 (1978), 583–592. MR 58 #6315 Zbl 0427.32006
- [Noell 1985] A. V. Noell, “Interpolation in weakly pseudoconvex domains in  $\mathbb{C}^2$ ”, *Math. Ann.* **270**:3 (1985), 339–348. MR 86j:32043 Zbl 0534.32003
- [Rudin 1978] W. Rudin, “Peak-interpolation sets of class  $C^1$ ”, *Pacific J. Math.* **75**:1 (1978), 267–279. MR 58 #6346 Zbl 0383.32007

Received December 2, 2002. Revised April 20, 2004.

GAUTAM BHARALI  
MATHEMATICS DEPARTMENT  
THE UNIVERSITY OF MICHIGAN  
525 EAST UNIVERSITY AVENUE  
ANN ARBOR, MI 48109  
bharali@umich.edu