

*Pacific
Journal of
Mathematics*

**TRANSVERSAL HOLOMORPHIC SECTIONS AND
LOCALIZATION OF ANALYTIC TORSIONS**

HUITAO FENG AND XIAONAN MA

Volume 219 No. 2

April 2005

TRANSVERSAL HOLOMORPHIC SECTIONS AND LOCALIZATION OF ANALYTIC TORSIONS

HUITAO FENG AND XIAONAN MA

We prove a Bott-type residue formula twisted by $\wedge(\mathbb{V}^*)$ with a holomorphic vector bundle \mathbb{V} , and relate certain analytic torsions on the total manifold to the analytic torsions on the zero set of a holomorphic section of \mathbb{V} .

Introduction

Beasley and Witten [2003], studying half-linear models, have described a compactification on any Calabi–Yau threefold Y that is a complete intersection in a compact toric variety X . In particular, a remarkable cancellation involving the instanton effect [Beasley and Witten 2003, (1.3)], involving certain determinants of the $\bar{\partial}$ -operator, was derived directly from a residue theorem. One would like to understand its implications in mathematics, for example in Gromov–Witten theory. Bershadsky, Cecotti, Ooguri and Vafa [Bershadsky et al. 1993; 1994] predicted that the analytic torsion of Ray–Singer will play a role regarding the genus-1 Gromov–Witten invariant. Thus we naturally try to understand the results about analytic torsion first.

As an application of [Bismut and Lebeau 1991] and the localization formula (1–3) in this paper, we were able to relate certain analytic torsions on the total manifold with the zero set of a holomorphic transversal section of \mathbb{V} , generalizing [Bismut 2004, Theorem 6.6] and [Zhang n.d.] with $\mathbb{V} = TX$ therein. We expect our formula will be useful for understanding [Beasley and Witten 2003, (1.3)] from a mathematical point of view.

This paper is organized as follows. In Section 1 we prove a Bott-type residue formula. In Section 2 we get a localization formula for Quillen metrics. In Section 3 we get a localization formula for analytic torsions under extra conditions. In Section 4, for the reader’s convenience, we write down six intermediate results, corresponding to [Bismut and Lebeau 1991, Theorems 6.4–6.9].

MSC2000: 58J52, 32L10, 58J20, 32C35, 57R20.

Keywords: analytic torsion, characteristic classes, characteristic numbers, residue formula.
Feng was partially supported by the NNSF of China (10271059).

1. A Bott-type residue formula

In this section, along the lines of [Bismut 1986, §1], we give a Bott-type residue formula (1–3) by assuming that the holomorphic section is transversal; compare to [Beasley and Witten 2003, (2.32), (2.34)].

Let X be a compact complex manifold with $\dim X = n$ and let \mathbb{V} be a holomorphic vector bundle on X with $\dim \mathbb{V} = l$. We assume that the line bundles $\det TX$ and $\det \mathbb{V}$ are holomorphically isomorphic. We fix a holomorphic isomorphism $\phi : \det \mathbb{V}^* \simeq \det T^*X$, which is clearly unique up to a constant. Thus ϕ defines a map from the \mathbb{Z}_2 -graded tensor product $\wedge(\overline{T^*X}) \widehat{\otimes} \wedge(\mathbb{V}^*)$ to $\wedge(\overline{T^*X}) \widehat{\otimes} \wedge^{\max}(T^*X) \subset \wedge(T_{\mathbb{R}}^*X) \otimes_{\mathbb{R}} \mathbb{C}$. We can define the integral of an element α of $\Omega(X, \wedge(\mathbb{V}^*))$, the set of smooth sections of $\wedge(\overline{T^*X}) \widehat{\otimes} \wedge(\mathbb{V}^*)$ on X , by

$$\int_X \alpha = \int_X \phi(\alpha).$$

Let v be a holomorphic section of \mathbb{V} on X . Assume that v vanishes on a complex manifold $Y \subset X$. Then $\nabla v|_Y : TX|_Y \rightarrow \mathbb{V}|_Y$ mapping U to $\nabla_U v$ does not depend on the choice of a connection ∇ on \mathbb{V} , and $\nabla_U v|_Y = 0$ for $U \in TY$. Let N be the normal bundle to Y in X . Assume also that $\nabla v|_Y : N \rightarrow \mathbb{V}|_Y$ is injective, and there is a holomorphic vector subbundle \mathbb{V}_1 on Y such that

$$(1-1) \quad \mathbb{V}|_Y = \mathbb{V}_1 \oplus \text{Im } \nabla v|_Y.$$

Let $P^{\mathbb{V}_1}$ and $P^{\text{Im } \nabla v}$ be the natural projections from \mathbb{V} onto \mathbb{V}_1 and $\text{Im } \nabla v|_Y$.

Let $i(v)$ be the standard contraction operator acting on $\wedge(\mathbb{V}^*)$. A natural question, posed in [Beasley and Witten 2003, §2], is how to express $\int_X \alpha$ using the local data near the zero set Y of v for a $(\bar{\partial}^X + i(v))$ -closed form α , that is, a form satisfying $(\bar{\partial}^X + i(v))\alpha = 0$.

First we recall an idea due to Bismut [Bismut 1986]; see also [Zhang 1990].

Proposition 1.1. *Let $\alpha \in \Omega(X, \wedge(\mathbb{V}^*))$ be a $(\bar{\partial}^X + i(v))$ -closed form. Then*

$$\int_X \alpha = \int_X e^{-\bar{\partial}^X + i(v))\omega/t} \alpha \quad \text{for any } \omega \in \Omega(X, \wedge(\mathbb{V}^*)) \text{ and } t > 0.$$

Proof. For any $\omega \in \Omega(X, \wedge(\mathbb{V}^*))$

$$(1-2) \quad \int_X \bar{\partial}^X \omega = \int_X \phi(\bar{\partial}^X \omega) = \int_X \bar{\partial}^X \phi(\omega) = \int_X d\phi(\omega) = 0.$$

From $(\bar{\partial}^X + i(v))^2 = 0$ and $(\bar{\partial}^X + i(v))\alpha = 0$, we have

$$\frac{\partial}{\partial s} \int_X e^{-s(\bar{\partial}^X + i(v))\omega} \alpha = - \int_X (\bar{\partial}^X + i(v))(\omega e^{-s(\bar{\partial}^X + i(v))\omega} \alpha) = 0,$$

and the desired equality follows. □

Recall that $\nabla v|_Y : N \rightarrow \text{Im } \nabla v|_Y$ is an isomorphism that induces isomorphisms of holomorphic line bundles $\phi_N = (\det \nabla v|_Y)^* : \det(\text{Im } \nabla v|_Y)^* \rightarrow \det N^*$ and $\phi_Y = \phi|_Y / ((\det \nabla v|_Y)^*) : \det \mathbb{V}_1^* \rightarrow \det T^*Y$. These two isomorphisms make the integral \int_N along the normal bundle N and \int_Y well defined.

Let $h^\mathbb{V}$ be a Hermitian metric on \mathbb{V} such that \mathbb{V}_1 and $\text{Im } \nabla v|_Y$ are orthogonal on Y . Let g_1^N be a Hermitian metric on N such that $\nabla.v|_Y : N \rightarrow \text{Im } \nabla v|_Y$ is an isometry. Let $R^\mathbb{V}$ be the curvature of the holomorphic Hermitian connection $\nabla^\mathbb{V}$ on $(\mathbb{V}, h^\mathbb{V})$. Let $j : Y \rightarrow X$ be the natural embedding, and $\{Y_j\}_j$ the connected components of Y . On Y , define

$$R_v^\mathbb{V} = -(\nabla.v)^{-1} P^{\text{Im } \nabla v} R^\mathbb{V}(\cdot, j_*\cdot) P^{\mathbb{V}_1} \in \overline{T^*Y} \widehat{\otimes} \mathbb{V}_1^* \otimes \text{End } N.$$

$R_v^\mathbb{V}$ is well defined since $P^{\text{Im } \nabla v} R^\mathbb{V}(j_*\cdot, j_*\cdot) P^{\mathbb{V}_1} = 0$. Thus, for $U \in TY$, $W \in \mathbb{V}_1$, $u_1, u_2 \in N$,

$$\langle R_v^\mathbb{V}(\bar{U}, W)u_1, u_2 \rangle_{g_1^N} = -\langle R^\mathbb{V}(u_1, \bar{U})W, \nabla_{u_2} v \rangle = \langle W, R^\mathbb{V}(\bar{u}_1, U)\nabla_{u_2} v \rangle.$$

Certainly $\det_N((1 + R_v^\mathbb{V})/2\pi i)$ is $\bar{\partial}^Y$ -closed.

The following result verifies a formula of Beasley and Whitney [2003, (2.32), (2.34)] and generalizes corresponding results in [Zhang 1990], [Liu 1995] and [Bott 1967].

Theorem 1.2. *For any $(\bar{\partial}^X + i(v))$ -closed form $\alpha \in \Omega(X, \wedge(\mathbb{V}^*))$,*

$$(1-3) \quad \int_X \alpha = \sum_j \int_{Y_j} \frac{(-1)^{(l-n)(n-\dim Y_j)} \alpha}{\det_N((1 + R_v^\mathbb{V})/(-2\pi i))}.$$

Proof. Set

$$S = \langle \cdot, v \rangle_{h^\mathbb{V}} \in C^\infty(X, \mathbb{V}^*).$$

By Proposition 1.1, for any $t \in]0, +\infty[$,

$$(1-4) \quad \int_X \alpha = \int_X e^{-\frac{1}{2t}(\bar{\partial}^X + i(v))S} \alpha = \int_X e^{-\frac{1}{2t}(\bar{\partial}^X S + |v|^2)} \alpha.$$

Thus, as $t \rightarrow 0$, the integral $\int_X \alpha$ is asymptotically equal to $\int_{\mathfrak{U}} e^{-\frac{1}{2t}(\bar{\partial}^X S + |v|^2)} \alpha$ for any neighborhood \mathfrak{U} of Y .

Take $y \in Y$. Since Y is a complex submanifold, we can find holomorphic coordinates $\{z_i\}_{i=1}^n$ of a neighborhood U of y such that y corresponds to 0 and $\{(\partial/\partial z_i)(0)\}_{i=m+1}^n$ is an orthonormal basis of (N, g_1^N) , and, moreover,

$$U \cap Y = \{p \in U, z_{m+1}(p) = \dots = z_n(p) = 0\}.$$

Let $\{\mu_k\}_{k=1}^{l'}$ and $\{\mu_k\}_{k=l'+1}^l$ be holomorphic frames for \mathbb{V}_1 and $\text{Im } \nabla v|_Y$ on $U \cap Y$, with

$$\nabla_{\partial/\partial z_k(0)}^\mathbb{V} v = \mu_k(0) \quad \text{for } l' + 1 \leq k \leq l,$$

and for $z' = (z_1, \dots, z_m)$, $z'' = (z_{m+1}, \dots, z_n)$, $z = (z', z'')$, define $\mu_k(z)$ by parallel transport of $\mu_k(z', 0)$ with respect to $\nabla^{\mathbb{V}}$ along the curve $u \mapsto (z', uz'')$. Identify \mathbb{V}_z with $\mathbb{V}_{(z',0)}$ by identifying $\mu_k(z)$ with $\mu_k(z', 0)$. Denote by $W_y(\varepsilon)$ the ε -neighborhood of y in the normal space N . Then

$$(1-5) \quad \int_{Y \cap U} \int_{W_y(\varepsilon)} e^{-\frac{1}{2t}(\bar{\partial}^X S + |v|^2)} \alpha \\ = \int_{Y \cap U} \int_{z \in W_y(\varepsilon/\sqrt{t})} e^{-\frac{1}{2t}(|v(\sqrt{t}z)|^2 + (\bar{\partial}^X S)(\sqrt{t}z))} t^{n-m} \alpha(y, \sqrt{t}z).$$

Define $z = \sum_j z_j (\partial/\partial z_j)$ and $\bar{z} = \sum_j \bar{z}_j (\partial/\partial \bar{z}_j)$. The tautological vector field is $Z = z + \bar{z}$. Then, for $z \in N_y$,

$$\frac{1}{2t} |v(\sqrt{t}z)|^2 = \frac{1}{2} |\nabla_z^{\mathbb{V}} v|^2 + O(\sqrt{t}) = \frac{1}{2} |z|^2 + O(\sqrt{t})$$

and

$$\bar{\partial}^X S = \sum_{k=1}^l \langle \mu_k, \nabla^{\mathbb{V}} v \rangle \mu^k.$$

From now on, set $z = (0, z'')$ and $Z = z + \bar{z}$. Since $\nabla_Z^{\mathbb{V}} \mu_k(0) = 0$, we know that

$$(1-6) \quad \frac{1}{2t} \bar{\partial}^X S(\sqrt{t}z) \\ = \frac{1}{2t} \sum_{k=1}^l \langle \mu_k, \nabla^{\mathbb{V}} v \rangle (\sqrt{t}z) \mu^k(0) \\ = \frac{1}{2t} \sum_{k=1}^l \left(\langle \mu_k, \nabla^{\mathbb{V}} v \rangle(0) + \sqrt{t} \langle \mu_k, \nabla_Z^{\mathbb{V}} \nabla^{\mathbb{V}} v \rangle(0) \right. \\ \left. + \frac{t}{2} (\langle \nabla_Z^{\mathbb{V}} \nabla_Z^{\mathbb{V}} \mu_k, \nabla^{\mathbb{V}} v \rangle + \langle \mu_k, \nabla_Z^{\mathbb{V}} \nabla_Z^{\mathbb{V}} \nabla^{\mathbb{V}} v \rangle)(0) + O(t^{3/2}) \right) \mu^k(0).$$

Because of the factor t^{n-m} in (1-5), it should be clear that in the limit, only those monomials in the vertical form

$$d\bar{z}_{m+1} \wedge \dots \wedge d\bar{z}_n \widehat{\otimes} \mu^{l'+1} \wedge \dots \wedge \mu^l$$

whose weight is exactly t^{m-n} should be kept. Now,

$$\nabla_Z^{\mathbb{V}} \nabla_{\partial/\partial z_j}^{\mathbb{V}} v = R^{\mathbb{V}} \left(Z, \frac{\partial}{\partial z_j} \right) v + \nabla_{\partial/\partial z_j}^{\mathbb{V}} \nabla_Z^{\mathbb{V}} v - 1_{[m,n]}(j) \nabla_{\partial/\partial z_j}^{\mathbb{V}} v, \\ \nabla_{\bar{z}}^{\mathbb{V}} \nabla_{\partial/\partial z_j}^{\mathbb{V}} v(0) = R^{\mathbb{V}} \left(\bar{z}, \frac{\partial}{\partial z_j} \right) v + \nabla_{\partial/\partial z_j}^{\mathbb{V}} \nabla_{\bar{z}}^{\mathbb{V}} v = 0,$$

where $1_{[m,n]}$ is the characteristic function of the interval $[m, n]$. Note that $\nabla^{\vee} = \nabla^{\vee 1} \oplus \nabla^{\text{Im} \nabla v}$ on Y and that

$$\langle \mu_k, \nabla_z^{\vee} \nabla_{\partial/\partial z_j}^{\vee} v \rangle(0) = 0 \quad \text{for } 1 \leq j \leq m, 1 \leq k \leq l'.$$

It follows that in the expression

$$\frac{1}{2\sqrt{t}} \langle \mu_k, \nabla_Z^{\vee} \nabla^{\vee} v \rangle(0) \mu^k(0)$$

a nonzero contribution can only appear in the term

$$(1-7) \quad \frac{1}{2\sqrt{t}} \left(\sum_{j=1}^m \sum_{k=l'+1}^l + \sum_{j=m+1}^n \sum_{k=1}^{l'} \right) \langle \mu_k, \nabla_z^{\vee} \nabla_{\partial/\partial z_j}^{\vee} v \rangle(0) d\bar{z}^j \otimes \mu^k(0).$$

Similarly, in the last term of (1-6), the only term with a nonzero contribution is

$$\frac{1}{4} \sum_{j=1}^m \sum_{k=1}^{l'} \left(\langle \nabla_Z^{\vee} \nabla_Z^{\vee} \mu_k, \nabla_{\partial/\partial z_j}^{\vee} v \rangle(0) + \langle \mu_k, \nabla_Z^{\vee} \nabla_Z^{\vee} \nabla_{\partial/\partial z_j}^{\vee} v \rangle(0) \right) d\bar{z}^j \otimes \mu^k(0).$$

But for $1 \leq j \leq m$, both $\nabla_{\partial/\partial z_j}^{\vee} v(0)$ and $\nabla_{\partial/\partial z_j}^{\vee} \nabla_{\bar{z}}^{\vee} \nabla_z^{\vee} v(0) = \nabla_{\partial/\partial z_j}^{\vee} (R^{\vee}(\bar{z}, z)v)(0)$ vanish, since $v = 0$ on Y . Thus, for $1 \leq j \leq m$,

$$\nabla_Z^{\vee} \nabla_Z^{\vee} \nabla_{\partial/\partial z_j}^{\vee} v(0) = 2R^{\vee} \left(\bar{z}, \frac{\partial}{\partial z_j} \right) \nabla_z^{\vee} v(0) + \nabla_{\partial/\partial z_j}^{\vee} \nabla_z^{\vee} \nabla_z^{\vee} v(0).$$

By the preceding discussion, as $t \rightarrow 0$, in (1-5), we should replace $\frac{1}{2t} \bar{\partial}^X S(y, \sqrt{t}z)$ by the 2-form

$$\begin{aligned} & \frac{1}{2} \sum_{k=1}^l \langle \mu_k, \nabla^{\vee} v \rangle(0) \mu^k(0) + \sqrt{t} \times \text{expression (1-7)} \\ & + \frac{1}{2} \sum_{j=1}^m \sum_{k=1}^{l'} \left\langle \mu_k, R^{\vee} \left(\bar{z}, \frac{\partial}{\partial z_j} \right) \nabla_z^{\vee} v + \nabla_{\partial/\partial z_j}^{\vee} \nabla_z^{\vee} \nabla_z^{\vee} v \right\rangle(0) d\bar{z}^j \otimes \mu^k(0). \end{aligned}$$

Set $\beta_Y = d\bar{z}_1 \cdots d\bar{z}_m \wedge \mu^1(0) \cdots \mu^{l'}(0)$, $\beta_N = d\bar{z}_{m+1} \cdots d\bar{z}_n \wedge \mu^{l'+1}(0) \cdots \mu^l(0)$, $\phi(\mu^1(0) \cdots \mu^l(0)) = f dz_1 \cdots dz_n$. Then

$$\phi_Y(\mu^1(0) \cdots \mu^{l'}(0)) \phi_N(\mu^{l'+1}(0) \cdots \mu^l(0)) = f dz_1 \cdots dz_n.$$

Thus

$$\begin{aligned} \phi(\beta_Y \wedge \beta_N) &= (-1)^{l(n-m)} f d\bar{z}_1 \cdots d\bar{z}_n \wedge dz_1 \cdots dz_n \\ &= (-1)^{(l-m)(n-m)} \phi_Y(\beta_Y) \phi_N(\beta_N). \end{aligned}$$

Now, observing that $\int_{\mathbb{C}} \bar{z}^i e^{-|z|^2} dz d\bar{z} = 0$ for $i > 0$ and that $\nabla^{\vee} v : (N, g_1^N) \rightarrow (\text{Im} \nabla v, h^{\text{Im} \nabla v})$ is an isometry and $l - l' = n - m$, we find that the limit of (1-4)

as $t \rightarrow 0$ is the sum over j of

$$(1-8) \quad \int_{Y_j} (-1)^{(l-n)(n-m)} j^* \alpha \int_N \exp \left(-\frac{1}{2} \sum_{k=1}^l \langle \mu_k, \nabla^{\mathbb{V}} v \rangle(0) \mu^k(0) \right. \\ \left. -\frac{1}{2} \langle \cdot, P^{\mathbb{V}_1} R^{\mathbb{V}}(\bar{z}, j_* \cdot) \nabla_z^{\mathbb{V}} v \rangle(0) - \frac{1}{2} |\nabla_z^{\mathbb{V}} v|^2 \right).$$

The second integrand in this expression can be rewritten as

$$\exp \left(-\frac{1}{2} \sum_{i=1}^{n-m} d\bar{z}_{m+i} \wedge \mu^{l'+i}(0) + \frac{1}{2} \langle R^{\mathbb{V}}(z, j_* \cdot) P^{\mathbb{V}_1 \cdot}, \nabla_z^{\mathbb{V}} v \rangle(0) - \frac{1}{2} |z|^2 \right) \\ = \exp \left(\frac{1}{2} \langle (\nabla^{\mathbb{V}} v)^{-1} R^{\mathbb{V}}(z, j_* \cdot) P^{\mathbb{V}_1 \cdot}, z \rangle - \frac{1}{2} |z|^2 \right) \left(\frac{1}{2} \right)^{l-l'} dz_{m+1} d\bar{z}_{m+1} \cdots dz_n d\bar{z}_n.$$

Thus the expression in (1-8) is equal to

$$\int_{Y_j} \frac{(-1)^{(l-n)(n-m)} \alpha}{\det_N((1 + R_v^{\mathbb{V}})/(-2\pi i))},$$

which leads to (1-3). □

2. Localization of Quillen metrics via a transversal section

Let X be a compact complex manifold of dimension n . Let \mathbb{V} and ξ be holomorphic vector bundles on X with $\dim \mathbb{V} = m$, and let v be a holomorphic section of \mathbb{V} . Assume that v vanishes on a complex manifold $Y \subset X$ and satisfies (1-1). Then we have a complex of holomorphic vector bundles on X ,

$$(2-1) \quad 0 \rightarrow \wedge^m(\mathbb{V}^*) \xrightarrow{i(v)} \wedge^{m-1}(\mathbb{V}^*) \xrightarrow{i(v)} \cdots \xrightarrow{i(v)} \wedge^1(\mathbb{V}^*) \xrightarrow{i(v)} \wedge^0(\mathbb{V}^*) \rightarrow 0.$$

Let $(\Omega(X, \wedge(\mathbb{V}^*) \otimes \xi), \bar{\partial}^X)$ be the Dolbeault complex associated to the holomorphic vector bundle $\wedge(\mathbb{V}^*) \otimes \xi$. Let $\mathcal{H}_v(X, \wedge(\mathbb{V}^*) \otimes \xi)$ be the hypercohomologies of the bicomplex $(\Omega(X, \wedge(\mathbb{V}^*) \otimes \xi), \bar{\partial}^X, i(v))$. Let $j : Y \rightarrow X$ be the obvious embedding. Now the pullback map j^* induces naturally a map of complexes

$$(2-2) \quad j^* : (\Omega(X, \wedge(\mathbb{V}^*) \otimes \xi), \bar{\partial}^X + i(v)) \rightarrow (\Omega(Y, \wedge(\mathbb{V}_1^*) \otimes \xi), \bar{\partial}^Y).$$

Theorem 2.1. *The map j^* is a quasi-isomorphism of complexes. In particular, j^* induces an isomorphism*

$$(2-3) \quad \mathcal{H}_v(X, \wedge(\mathbb{V}^*) \otimes \xi) \simeq H(Y, \wedge(\mathbb{V}_1^*) \otimes \xi).$$

Proof. In [Feng 2003] there is an analytic proof of this theorem when $\mathbb{V} = TX$. There we used the twisted vector bundle $\wedge(T^*X)$ and here $\wedge(\mathbb{V}^*)$ takes its place; the proof works just the same. For an algebraic proof, we can modify the proof of [Bismut 2004, Theorem 5.1]. □

Let N^X, N_H^X be the number operators on $\wedge(T^*X), \wedge(\mathbb{V}^*)$ corresponding to multiplication by p on $\wedge^p(T^*X), \wedge^p(\mathbb{V}^*)$; do the same replacing X by Y and \mathbb{V}^* by \mathbb{V}_1^* . Then $N^X - N_H^X$ and $N^Y - N_H^Y$ define \mathbb{Z} -gradings on $\Omega(X, \wedge(\mathbb{V}^*) \otimes \xi)$ and $\Omega(Y, \wedge(\mathbb{V}_1^*) \otimes \xi)$, which in turn induce \mathbb{Z} -gradings on $\mathcal{H}_v(X, \wedge(\mathbb{V}^*) \otimes \xi)$ and $H(Y, \wedge(\mathbb{V}_1^*) \otimes \xi)$, respectively. The isomorphism j^* preserves these \mathbb{Z} -gradings.

From [Bismut and Lebeau 1991, (1.24)], we define the complex lines $\lambda_v(\mathbb{V}^*)$ and $\lambda(\mathbb{V}_1^*)$ by

$$\lambda_v(\mathbb{V}^*) = \bigotimes_{p=-m}^n (\det \mathcal{H}_v^p(X, \wedge(\mathbb{V}^*) \otimes \xi))^{(-1)^{p+1}},$$

$$\lambda(\mathbb{V}_1^*) = \bigotimes_{p=0}^n \bigotimes_{q=0}^m (\det H^p(Y, \wedge^q(\mathbb{V}_1^*) \otimes \xi))^{(-1)^{p+q+1}}.$$

By (2–3), we have a canonical isomorphism of complex lines

$$\lambda_v(\mathbb{V}^*) \simeq \lambda(\mathbb{V}_1^*).$$

Let ρ be the nonzero section of $\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_v(\mathbb{V}^*)$ associated with this canonical isomorphism.

Let g^{TX} be a Kähler metric on TX . We identify N with the bundle orthogonal to TY in $TX|_Y$. Let g^{TY} and g^N be the metrics on TY and N induced by g^{TX} . Let h^ξ be a Hermitian metric on ξ . Let $h^\mathbb{V}$ be a metric on \mathbb{V} such that \mathbb{V}_1 and $\text{Im } \nabla v|_Y$ are orthogonal on Y and $\nabla v|_Y : N \rightarrow \text{Im } \nabla v|_Y$ is an isometry.

Let dv_X be the Riemannian volume form on (X, g^{TX}) . Let $\langle \cdot, \cdot \rangle_0$ be the metric on $\wedge(\overline{T^*X}) \widehat{\otimes} \wedge(\mathbb{V}^*) \otimes \xi$ induced by $g^{TX}, h^\mathbb{V}, h^\xi$. The Hermitian product on $\Omega(X, \wedge(\mathbb{V}^*) \otimes \xi)$ is defined by

$$(2-4) \quad \langle \alpha, \alpha' \rangle = \frac{1}{(2\pi)^n} \int_X \langle \alpha, \alpha' \rangle_0 dv_X \quad \text{for } \alpha, \alpha' \in \Omega(X, \wedge(\mathbb{V}^*) \otimes \xi).$$

Let $\bar{\partial}^{X*}$ and $v^* \wedge = i(v)^*$ be the adjoint of $\bar{\partial}^X$ and $i(v)$ with respect to $\langle \cdot, \cdot \rangle$. Set

$$V = i(v) + i(v)^*, \quad D^X = \bar{\partial}^X + \bar{\partial}^{X*}.$$

By Hodge theory,

$$(2-5) \quad \mathcal{H}_v(X, \wedge(\mathbb{V}^*) \otimes \xi) \simeq \text{Ker}(D^X + V).$$

Denote by P be the operator of orthogonal projection from $\Omega(X, \wedge(\mathbb{V}^*) \otimes \xi)$ onto $\text{ker}(D^X + V)$ and set $P^\perp = 1 - P$. Let $h^{\mathcal{H}_v}$ be the L^2 -metric on $\mathcal{H}_v(X, \wedge(\mathbb{V}^*) \otimes \xi)$ induced by the L^2 -product (2–4) via the isomorphism (2–5). Define in the same way a Hermitian product on $\Omega(Y, \wedge(\mathbb{V}_1^*) \otimes \xi)$ associated to $g^{TY}, h^{\mathbb{V}_1}, h^\xi$. Let $\bar{\partial}^{Y*}$ be the adjoint of $\bar{\partial}^Y$, and $h^{H(Y, \wedge(\mathbb{V}_1^*) \otimes \xi)}$ the corresponding L^2 -metric on

$H(Y, \wedge(\mathbb{V}_1^*) \otimes \xi)$. Set

$$D^Y = \bar{\partial}^Y + \bar{\partial}^{Y*}.$$

Let Q be the orthogonal projection operator from $\Omega(Y, \wedge(\mathbb{V}_1^*) \otimes \xi)$ on $\text{Ker } D^Y$, and $Q^\perp = 1 - Q$. Let $|\cdot|_{\lambda_v(\mathbb{V}^*)}$ and $|\cdot|_{\lambda(\mathbb{V}^*)}$ be the L^2 -metrics on $\lambda_v(\mathbb{V}^*)$ and $\lambda(\mathbb{V}^*)$ induced by $h^{\mathcal{K}_v}$ and $h^{H(Y, \wedge(\mathbb{V}_1^*) \otimes \xi)}$. Following [Bismut and Lebeau 1991, (1.49)], let

$$\theta_v^X(s) = -\text{Tr}_s((N^X - N_H^X)((D^X + V)^2)^{-s} P^\perp).$$

Then $\theta_v^X(s)$ extends to a meromorphic function of $s \in \mathbb{C}$, which is holomorphic at $s = 0$.

The Quillen metric $\|\cdot\|_{\lambda_v(\mathbb{V}^*)}$ on the line $\lambda_v(\mathbb{V}^*)$ is defined by

$$\|\cdot\|_{\lambda_v(\mathbb{V}^*)} = |\cdot|_{\lambda_v(\mathbb{V}^*)} \exp\left(-\frac{1}{2} \frac{\partial \theta_v^X}{\partial s}(0)\right).$$

In the same way, the function

$$\theta^Y(s) = -\text{Tr}_s((N^Y - N_H^Y)(D^{Y,2})^{-s} Q^\perp)$$

extends to a meromorphic function of $s \in \mathbb{C}$, holomorphic at $s = 0$. The Quillen metric $\|\cdot\|_{\lambda(\mathbb{V}_1^*)}$ on the line $\lambda(\mathbb{V}_1^*)$ is defined by

$$\|\cdot\|_{\lambda(\mathbb{V}_1^*)} = |\cdot|_{\lambda(\mathbb{V}_1^*)} \exp\left(-\frac{1}{2} \frac{\partial \theta^Y}{\partial s}(0)\right).$$

Let $\|\cdot\|_{\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_v(\mathbb{V}^*)}$ be the Quillen metric on $\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_v(\mathbb{V}^*)$ induced by $\|\cdot\|_{\lambda_v(\mathbb{V}^*)}$ and $\|\cdot\|_{\lambda(\mathbb{V}_1^*)}$ as in [Bismut and Lebeau 1991, §1e].

The purpose of this section is to give a formula for $\|\rho\|_{\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_v(\mathbb{V}^*)}^2$. Now we introduce some notations.

For a holomorphic Hermitian vector bundle (E, h^E) on X , we denote by $\text{Td}(E)$, $\text{ch}(E)$, $c_{\max}(E)$ the Todd class, Chern character, and top Chern class of E , and by $\text{Td}(E, h^E)$, $\text{ch}(E, h^E)$, $c_{\max}(E, h^E)$ the Chern–Weil representatives of $\text{Td}(E)$, $\text{ch}(E)$, $c_{\max}(E)$ with respect to the holomorphic Hermitian connection ∇^E on (E, h^E) .

Let δ_Y be the current of integration on Y . By [Bismut 1992, Theorem 3.6], a current $\tilde{c}_{\max}(\mathbb{V}, h^\mathbb{V})$ on X is well defined by the holomorphic section v (which induces an embedding $v : X \rightarrow \mathbb{V}$), and this current satisfies

$$(2-6) \quad \frac{\bar{\partial} \partial}{2\pi i} \tilde{c}_{\max}(\mathbb{V}, h^\mathbb{V}) = c_{\max}(\mathbb{V}_1, h^{\mathbb{V}_1}) \delta_Y - c_{\max}(\mathbb{V}, h^\mathbb{V}).$$

Let $\widetilde{\text{Td}}(TY, TX, g^{TX|_Y})$ be the Bott–Chern current on Y associated to the exact sequence

$$(2-7) \quad 0 \rightarrow TY \rightarrow TX|_Y \rightarrow N \rightarrow 0$$

constructed in [Bismut et al. 1988a, §1f], which satisfies

$$\frac{\bar{\partial}\partial}{2\pi i} \widetilde{\text{Td}}(TY, TX, g^{TX|_Y}) = \text{Td}(TX|_Y, g^{TX|_Y}) - \text{Td}(TY, g^{TY}) \text{Td}(N, g^N).$$

Finally, let $R(x)$ be the power series introduced in [Gillet and Soulé 1991], which is such that if $\zeta(s)$ is the Riemann zeta function, then

$$R(x) = \sum_{\substack{n \geq 1 \\ n \text{ odd}}} \left(\sum_{j=1}^n \frac{1}{j} \zeta(-n) + 2 \frac{\partial \zeta}{\partial s}(-n) \right) \frac{x^n}{n!}.$$

We identify R with the corresponding additive genus. We also set

$$\text{ch}(\wedge^*(\mathbb{V}_1^*)) = \sum_i (-1)^i \text{ch}(\wedge^i(\mathbb{V}_1^*)),$$

and denote by $\text{ch}(\wedge^*(\mathbb{V}_1^*), h^{\wedge^*(\mathbb{V}_1^*)})$ its Chern–Weil representative.

Theorem 2.2. *The Quillen metric $\|\rho\|_{\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_V(\mathbb{V}^*)}^2$ is given by the exponential of*

$$\begin{aligned} (2-8) \quad & - \int_X \text{Td}(TX, g^{TX}) \text{Td}^{-1}(\mathbb{V}, h^{\mathbb{V}}) \tilde{c}_{\max}(\mathbb{V}, h^{\mathbb{V}}) \text{ch}(\xi, h^\xi) \\ & + \int_Y \text{Td}^{-1}(N, g^N) \widetilde{\text{Td}}(TY, TX|_Y, g^{TX|_Y}) \text{ch}(\wedge^*(\mathbb{V}_1^*), h^{\wedge^*(\mathbb{V}_1^*)}) \text{ch}(\xi, h^\xi) \\ & - \int_Y \text{Td}(TY) R(N) \text{ch}(\wedge^*(\mathbb{V}_1^*)) \text{ch}(\xi). \end{aligned}$$

Proof. Set

$$(2-9) \quad T(\wedge(\mathbb{V}^*), h^{\wedge(\mathbb{V}^*)}) = \text{Td}^{-1}(\mathbb{V}, h^{\mathbb{V}}) \tilde{c}_{\max}(\mathbb{V}, h^{\mathbb{V}}).$$

By the same argument as in [Bismut et al. 1990, Theorem 3.17], the current

$$T(\wedge(\mathbb{V}^*), h^{\wedge(\mathbb{V}^*)})$$

is exactly the current on X associated to (2-1) (evaluated modulo irrelevant ∂ or $\bar{\partial}$ coboundaries).

Now, from the choice of our metric $h^{\mathbb{V}}$, the analogue of [Bismut and Lebeau 1991, Definition 1.21, assumption (A)] is satisfied for the complex (2-1). Then we verify that as far as local index theoretic computations are concerned, the situation is exactly the same as in [Bismut and Lebeau 1991]. Because of the quasi-isomorphism of Theorem 2.1, there are no “small” eigenvalues of the operator $D + TV$ when $T \rightarrow +\infty$. In Section 3, we write down the intermediate results corresponding to [Bismut and Lebeau 1991, §6c]. Comparing to [Bismut and Lebeau 1991, §§6c–6e], the proof of Theorem 2.2 is complete. \square

Remark 2.3. Assume that Y consists only discrete points; then $l \geq n$ and the last two terms of (2–8) are zero. In this case, if $n = l$, then (2–1) is a resolution of $j_*(\mathbb{C}_Y)$ and Theorem 2.2 is a direct consequence of [Bismut and Lebeau 1991, Theorem 0.1]. By [Bismut 1992, Theorem 3.2, Definition 3.5], $\tilde{c}_{\max}(\mathbb{V}, h^{\mathbb{V}})$ is zero if $l > n + 1$.

3. L^2 metrics on $H_v(X, \wedge(\mathbb{V}^*))$ and localization

We keep the assumptions and notations of Section 2.

Let g^{TX} be a Kähler metric on TX , and let g^{TY}, g^N be the metrics on TY, N induced by g^{TX} . Let $h^{\mathbb{V}}$ be a metric on \mathbb{V} such that \mathbb{V}_1 and $\text{Im } \nabla v|_Y$ are orthogonal on Y and $\nabla v|_Y : (N, g^N) \rightarrow \text{Im } \nabla v|_Y$ is an isometry.

Let $\phi_1 : \det \mathbb{V}_1^* \rightarrow \det T^*Y$ be a nonzero holomorphic section. Let $h_1^{\mathbb{V}}$ be a metric on \mathbb{V} such that on Y , \mathbb{V}_1 and $\text{Im } \nabla v|_Y$ are orthogonal and

$$|\phi|_{\det \mathbb{V} \otimes \det T^*X, 1} = |\phi_1|_{\det \mathbb{V}_1 \otimes \det T^*Y, 1} = 1,$$

where $|\cdot|_{\det \mathbb{V} \otimes \det T^*X, 1}$ and $|\cdot|_{\det \mathbb{V}_1 \otimes \det T^*Y, 1}$ are the norms on the holomorphic line bundles $\det \mathbb{V} \otimes \det T^*X$ and $\det \mathbb{V}_1 \otimes \det T^*Y$ induced by $h_1^{\mathbb{V}}$ and g^{TX} .

We will add a subscript 1 to denote the objects induced by $h_1^{\mathbb{V}}$. For

$$\beta \in \wedge^p(\overline{T^*X}) \widehat{\otimes} \wedge^q(\mathbb{V}^*),$$

we define $*_{\mathbb{V}, 1}\beta \in \wedge^{n-p}(\overline{T^*X}) \widehat{\otimes} \wedge^{l-q}(\mathbb{V}^*)$ by

$$\langle \alpha, \beta \rangle_1 \phi^{-1}(dv_X) = \alpha \wedge *_{\mathbb{V}, 1}\beta.$$

It's useful to write down a local expression for $*_{\mathbb{V}, 1}$. if $\{w^i\}_{i=1}^n$ and $\{\mu^i\}_{i=1}^l$ are orthonormal bases of T^*X and $(\mathbb{V}^*, h_1^{\mathbb{V}})$, then

$$dv_X = (-1)^{n(n+1)/2} (\sqrt{-1})^n \bar{w}^1 \wedge \dots \wedge \bar{w}^n \widehat{\otimes} w^1 \wedge \dots \wedge w^n$$

and $\phi^{-1}(w^1 \wedge \dots \wedge w^n) = f \mu^1 \wedge \dots \wedge \mu^l$ with $|f| = 1$. If

$$\beta = \bar{w}^1 \wedge \dots \wedge \bar{w}^p \widehat{\otimes} \mu^1 \wedge \dots \wedge \mu^q,$$

then

$$*_{\mathbb{V}, 1}\beta = (-1)^{(n-p)q+n(n+1)/2} (\sqrt{-1})^n f \bar{w}^{p+1} \wedge \dots \wedge \bar{w}^n \widehat{\otimes} \mu^{q+1} \wedge \dots \wedge \mu^l.$$

Thus $*_{\mathbb{V}, 1}*_{\mathbb{V}, 1}\beta = (-1)^{(p+q)(n+l+1)}\beta$, for any $\beta \in \wedge^p(\overline{T^*X}) \widehat{\otimes} \wedge^q(\mathbb{V}^*)$. Combining this with (1–2), we find that

$$\bar{\partial}^{X*}\beta = (-1)^{p+q+1} *_{\mathbb{V}, 1}^{-1} \bar{\partial}^X *_{\mathbb{V}, 1}\beta, \quad (i(v))^*\beta = (-1)^{p+q+1} *_{\mathbb{V}, 1}^{-1} i(v) *_{\mathbb{V}, 1}\beta.$$

Thus the antilinear map $*_{\mathbb{V}, 1}$ is an isometry from $(\mathcal{H}_v(X, \wedge(\mathbb{V}^*)), h_1^{\mathcal{H}_v})$ to itself.

The bilinear form

$$(3-1) \quad \alpha, \beta \in \mathcal{H}_v(X, \wedge(\mathbb{V}^*)) \mapsto \frac{1}{(2\pi)^n} \int_X \alpha \wedge \beta$$

is nondegenerate; indeed, $\alpha \in \mathcal{H}_v(X, \wedge(\mathbb{V}^*))$ implies $*_{\mathbb{V},1}\alpha \in \mathcal{H}_v(X, \wedge(\mathbb{V}^*))$, so $\alpha \neq 0$ implies

$$\int_X \alpha \wedge *_{\mathbb{V},1}\alpha > 0.$$

Thus the metric $|\cdot|_{\lambda_v(\mathbb{V}^*),1}$ on $\lambda_v(\mathbb{V}^*)$ only depends on the nondegenerate bilinear form (3-1) on $\mathcal{H}_v(X, \wedge(\mathbb{V}^*))$, which is metric-independent.

Recall the definition of $\det \nabla v|_Y$ from Section 1. Now,

$$\frac{\phi|_Y / ((\det \nabla v|_Y)^*)}{\phi_1}$$

is a holomorphic function on Y . Since Y is compact, this function is locally constant. Then we have the following extension of [Bismut 2004, Theorem 5.7].

Theorem 3.1.

$$(3-2) \quad \log(|\rho|_{\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_v(\mathbb{V}^*),1})^2 = \int_Y \text{Td}(TY) \text{ch}(\wedge(\mathbb{V}_1^*)) \log \left| \frac{\phi|_Y / ((\det \nabla v|_Y)^*)}{\phi_1} \right|.$$

Proof. We use ϕ_1 to define the integral $\int_Y \gamma$ for $\gamma \in H(Y, \wedge(\mathbb{V}_1^*))$. Since

$$|\phi_1|_{\det \mathbb{V}_1 \otimes \det T^*Y,1} = 1,$$

following the same considerations as above, we find that the antilinear operator $*_{\mathbb{V},1}$ maps $H(Y, \wedge(\mathbb{V}_1^*))$ into itself isometrically. Therefore, to evaluate the left-hand side of (3-2), we only need to compare the bilinear forms (3-1) with

$$a, b \in H(Y, \wedge(\mathbb{V}_1^*)) \mapsto \frac{1}{(2\pi)^m} \int_Y a \wedge b.$$

Let $A_v \in \text{End}^{\text{even}} H(Y, \wedge(\mathbb{V}_1^*))$ be given by

$$(3-3) \quad a \mapsto \frac{(-1)^{(l-n)(n-m)} a}{(2\pi)^{n-m} \det_N((1 + R_v^{\mathbb{V}})/(-2\pi i))} \frac{\phi|_Y / ((\det \nabla v|_Y)^*)}{\phi_1}.$$

Set

$$\det A_v = \frac{\det A_v|_{H^{\text{even}}(Y, \wedge(\mathbb{V}_1^*))}}{\det A_v|_{H^{\text{odd}}(Y, \wedge(\mathbb{V}_1^*))}};$$

then

$$(|\rho|_{\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_v(\mathbb{V}^*),1})^2 = |\det A_v|.$$

Now, A_v is a degree-increasing operator in $H(Y, \wedge(\mathbb{V}_1^*))$. Therefore it acts like a triangular matrix whose diagonal part is just multiplication by the locally constant

function $\frac{\phi|_Y/((\det \nabla v|_Y)^*)}{\phi_1}$. Using (3–3), we get

$$\det A_v = \left(\frac{\phi|_Y/((\det \nabla v|_Y)^*)}{\phi_1} \right)^{\chi(Y, \wedge(\mathbb{V}_1^*))}.$$

But $\chi(Y, \wedge(\mathbb{V}_1^*)) = \int_Y \text{Td}(TY) \text{ch}(\wedge(\mathbb{V}_1^*))$; thus we get (3–2). □

Let g_1^N be the metric on N such that $\nabla v|_Y : (N, g_1^N) \rightarrow (\text{Im}(\nabla v), h_1^{\text{Im}(\nabla v)})$ is an isometry. Let $\widetilde{\text{Td}}^{-1}(N, g^N, g_1^N)$ be the Bott–Chern class constructed in [Bismut et al. 1988a, §1f] such that

$$\frac{\bar{\partial}\partial}{2\pi i} \widetilde{\text{Td}}^{-1}(N, g^N, g_1^N) = \text{Td}^{-1}(N, g_1^N) - \text{Td}^{-1}(N, g^N).$$

Finally, we can compute the analytic torsion on the total manifold via the zero set of a transversal section v .

Theorem 3.2. *If $h_1^{\mathbb{V}_1} = h^{\mathbb{V}_1}$ on Y , then*

$$\begin{aligned} (3-4) \quad & -\frac{\partial\theta_{v,1}^X}{\partial s}(0) + \frac{\partial\theta^Y}{\partial s}(0) = -\int_X \text{Td}(TX, g^{TX}) \text{Td}^{-1}(\mathbb{V}, h_1^{\mathbb{V}}) \tilde{c}_{\max}(\mathbb{V}, h_1^{\mathbb{V}}) \\ & + \int_Y \left(\text{Td}^{-1}(N, g^N) \widetilde{\text{Td}}(TY, TX|_Y, g^{TX|_Y}) \right. \\ & \quad \left. + \text{Td}(TX, g^{TX}) \widetilde{\text{Td}}^{-1}(N, g^N, g_1^N) \right) \text{ch}(\wedge^*(\mathbb{V}_1^*), h^{\wedge^*(\mathbb{V}_1^*)}) \\ & - \int_Y \text{Td}(TY) \text{ch}(\wedge^*(\mathbb{V}_1^*)) \left(R(N) + \log \left| \frac{\phi|_Y/((\det \nabla v|_Y)^*)}{\phi_1} \right| \right). \end{aligned}$$

Proof. Since $h_1^{\mathbb{V}_1} = h^{\mathbb{V}_1}$, we have $|\cdot|_{\lambda(\mathbb{V}_1^*)} = |\cdot|_{\lambda(\mathbb{V}_1^*),1}$ and $\|\cdot\|_{\lambda(\mathbb{V}_1^*)} = \|\cdot\|_{\lambda(\mathbb{V}_1^*),1}$. Let $\tilde{\text{ch}}(\wedge(\mathbb{V}^*), h_1^{\wedge(\mathbb{V}^*)}, h^{\wedge(\mathbb{V}^*)})$ be the Bott–Chern class constructed in [Bismut et al. 1988a, §1f], so that

$$\frac{\bar{\partial}\partial}{2\pi i} \tilde{\text{ch}}(\wedge(\mathbb{V}^*), h_1^{\wedge(\mathbb{V}^*)}, h^{\wedge(\mathbb{V}^*)}) = \text{ch}(\wedge(\mathbb{V}^*), h^{\wedge(\mathbb{V}^*)}) - \text{ch}(\wedge(\mathbb{V}^*), h_1^{\wedge(\mathbb{V}^*)}).$$

Then by the anomaly formula [Bismut et al. 1988b, Theorem 1.23],

$$\log \left(\frac{\|\cdot\|_{\lambda_v(\mathbb{V}^*)}^2}{\|\cdot\|_{\lambda_v(\mathbb{V}^*),1}^2} \right) = \int_X \text{Td}(TX, g^{TX}) \tilde{\text{ch}}(\wedge(\mathbb{V}^*), h_1^{\wedge(\mathbb{V}^*)}, h^{\wedge(\mathbb{V}^*)}).$$

By [Bismut et al. 1990, Theorem 2.5],

$$\begin{aligned} (3-5) \quad & T(\wedge(\mathbb{V}^*), h^{\wedge(\mathbb{V}^*)}) - T(\wedge(\mathbb{V}^*), h_1^{\wedge(\mathbb{V}^*)}) \\ & = \text{ch}(\wedge^*(\mathbb{V}_1^*), h^{\wedge^*(\mathbb{V}_1^*)}) \widetilde{\text{Td}}^{-1}(N, g_1^N, g^N) \delta_Y - \tilde{\text{ch}}(\wedge(\mathbb{V}^*), h_1^{\wedge(\mathbb{V}^*)}, h^{\wedge(\mathbb{V}^*)}). \end{aligned}$$

By (2–9), Theorems 2.2 and 3.1, and the preceding equations, the proof of Theorem 3.2 is complete. □

Remark 3.3. If Y consists only of discrete points and $n = l$, then $\phi_1 = \text{Id}$. In this case let $g^{\det N}$ and $g_1^{\det N}$ be the metrics on $\det N = \det TX$ induced by g^N and g_1^N . By Remark 2.3 and Theorem 3.2,

$$-\frac{\partial \theta_{v,1}^X}{\partial s}(0) = - \int_X \text{Td}(TX, g^{TX}) \text{Td}^{-1}(\mathbb{V}, h_1^{\mathbb{V}}) \tilde{c}_{\max}(\mathbb{V}, h_1^{\mathbb{V}}) + \sum_{p \in Y} \left(\frac{1}{2} \log(g^{\det N} / g_1^{\det N}) - \log |\phi / (\det \nabla v|_Y)^*| \right).$$

Remark 3.4. If $\mathbb{V} = TX$ and v is a holomorphic Killing vector field, (3–4) is a special case of [Bismut 1992, Theorems 6.2 and 7.7]. In this case, $h_1^{\mathbb{V}} = g^{TX}$, and on Y , we have a holomorphic and orthogonal splitting $TX|_Y = TY \oplus N$. Thus $\widetilde{\text{Td}}(TY, TX|_Y, g^{TX|_Y}) = 0$. To compute $\widetilde{\text{Td}}^{-1}(N, g^N, g_1^N)$, note that $g_1^N = g^N((\nabla v)\cdot, (\nabla v)\cdot)$, as $A = (\nabla v)^*(\nabla v)$ is positive and self-adjoint; thus $(A)^s$ is well defined for $s \in [0, 1]$. Taking $g_s^N = g^N((A)^s\cdot, \cdot)$, we obtain by [Bismut et al. 1988a, Theorem 1.30]

$$\widetilde{\text{Td}}^{-1}(N, g^N, g_1^N) = \int_0^1 \langle (\text{Td}^{-1})'(N, g_s^N), \log A \rangle ds.$$

But ∇v is holomorphic, so the curvature R_s^N associated to the holomorphic connection on (N, g_s^N) is $R_s^N = R^N$ for $s \in [0, 1]$. Thus

$$(3-6) \quad \widetilde{\text{Td}}^{-1}(N, g^N, g_1^N) = \langle (\text{Td}^{-1})'(N, g^N), \log A \rangle.$$

Now

$$(3-7) \quad \text{Td}(TX, g^{TX}) T(\wedge(T^*X), h^{\wedge(T^*X)}) = \tilde{c}_{\max}(TX, g^{TX})$$

is an $(n-1, n-1)$ -form on X .

In this case, we get easily the special case of [Bismut 2004, Theorem 4.15] directly from [Ray and Singer 1973] by using Poincaré duality:

$$(3-8) \quad \frac{\partial \theta^Y}{\partial s}(0) = 0.$$

From (3–4), (3–6), (3–7), and the vanishing of the constant terms of $R(N)$ and $\frac{\text{Td}'}{\text{Td}}(N, g^N) - \frac{1}{2}$, we get

$$(3-9) \quad -\frac{\partial \theta_{v,1}^X}{\partial s}(0) = \int_Y c_{\max}(TY) \left(R(N) - \left\langle \frac{\text{Td}'}{\text{Td}}(N, g^N) - \frac{1}{2}, \log A \right\rangle \right) = 0.$$

4. Appendix: six intermediate results

In this section, to help readers understand how to obtain Theorem 2.2, we write down the corresponding intermediate results from [Bismut and Lebeau 1991, Theorems 6.4-6.9].

Let $\nabla^{\wedge(\mathbb{V}^*)}$ be the connection on $\wedge(\mathbb{V}^*)$ induced by $\nabla^{\mathbb{V}^*}$. Set $C_u = \nabla^{\wedge(\mathbb{V}^*)} + \sqrt{u}V$. Let $\mathcal{B}_{T^2}^2$ and $\text{Tr}_s(N_H^Y \exp(-\mathcal{B}_{T^2}^2))$ be the operator and the generalized trace associated to the complex (2-7) as in [Bismut and Lebeau 1991, §5]. Let Φ be the homomorphism from $\wedge^{\text{even}}(T_{\mathbb{R}}^*X)$ into itself which to $\alpha \in \wedge^{2p}(T_{\mathbb{R}}^*X)$ associates $(2\pi i)^{-p}\alpha$.

Theorem 4.1. *For any $u_0 > 0$, there exists $C > 0$ such that for $u \geq u_0, T \geq 1$,*

$$\begin{aligned} \left| \text{Tr}_s(N_H^X e^{-u(D^X+TV)^2}) - \text{Tr}_s\left(\left(\frac{1}{2} \dim N + N_H^Y\right)e^{-uD^{Y,2}}\right) \right| &\leq \frac{C}{\sqrt{T}}, \\ \left| \text{Tr}_s((N^X - N_H^X)e^{-u(D^X+TV)^2}) - \text{Tr}_s((N^Y - N_H^Y)e^{-uD^{Y,2}}) \right| &\leq \frac{C}{\sqrt{T}}. \end{aligned}$$

Theorem 4.2. *Let \tilde{P}_T be the orthogonal projection operator from $\Omega(X, \wedge(\mathbb{V}^*) \otimes \xi)$ to $\text{Ker}(D^X + TV)$. There exist $c > 0$ and $C > 0$ such that, for any $u \geq 1$ and $T \geq 1$,*

$$\left| \text{Tr}_s((N^X - N_H^X)e^{-u(D^X+TV)^2}) - \text{Tr}_s((N^X - N_H^X)\tilde{P}_T) \right| \leq ce^{-Cu},$$

Theorem 4.3. *There exist $C > 0$ and $\gamma \in]0, 1]$ such that, for any $u \in]0, 1]$ and $0 \leq T \leq 1/u$,*

$$\left| \text{Tr}_s(N_H^X e^{-u(D^X+TV)^2}) - \int_X \text{Td}(TX, g^{TX}) \Phi \text{Tr}_s(N_H^X e^{-C^2/T^2}) \right| \leq C(u(1+T))^\gamma.$$

There exists a constant $C' > 0$ such that for $u \in]0, 1]$ and $0 \leq T \leq 1$,

$$\left| \text{Tr}_s(N_H^X e^{-u(D^X+TV)^2}) - \text{Tr}_s(N_H^X e^{-uD^X}) \right| \leq C'T.$$

Theorem 4.4. *For any $T > 0$,*

$$\lim_{u \rightarrow 0} \text{Tr}_s(N_H^X e^{-u(D^X+(T/u)V)^2}) = \int_Y \Phi \text{Tr}_s(N_H^Y e^{-\mathcal{B}_{T^2}^2}) \text{ch}(\wedge(\mathbb{V}_1^*), h^{\wedge(\mathbb{V}_1^*)}) \text{ch}(\xi, h^\xi).$$

Theorem 4.5. *There exist $C > 0$ and $\delta \in]0, 1]$ such that, for any $u \in]0, 1]$ and $T \geq 1$,*

$$\left| \text{Tr}_s(N_H^X e^{-u(D^X+(T/u)V)^2}) - \text{Tr}_s\left(\left(\frac{1}{2} \dim N + N_H^Y\right)e^{-uD^{Y,2}}\right) \right| \leq \frac{C}{T^\delta}.$$

Let $|\cdot|_{\lambda_v(\mathbb{V}^*), T}^2$ be the L^2 -metric on $\lambda_v(\mathbb{V}^*)$ induced by $g^{TX}, T^2h^{\mathbb{V}}$ as in (2-5).

Theorem 4.6. As $T \rightarrow +\infty$,

$$\log \left(\frac{|\cdot|_{\lambda_v(\mathbb{V}^*), T}^2}{|\cdot|_{\lambda_v(\mathbb{V}^*)}^2} \right) = -\log |\rho|_{\lambda(\mathbb{V}_1^*)^{-1} \otimes \lambda_v(\mathbb{V}^*)}^2 + \text{Tr}_s((\dim N + 2N_H^Y)Q) \log T + O\left(\frac{1}{T}\right).$$

Acknowledgements

Feng thanks Jean-Pierre Bourguignon and the IHES, where part of this research was performed, for their hospitality. Thanks also to K. Liu for drawing our attention to [Beasley and Witten 2003].

References

- [Beasley and Witten 2003] C. Beasley and E. Witten, “Residues and world-sheet instantons”, 2003. hep-th/0304115
- [Bershadsky et al. 1993] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, “Holomorphic anomalies in topological field theories”, *Nuclear Phys. B* **405**:2-3 (1993), 279–304. MR 94j:81254 Zbl 1039.81550
- [Bershadsky et al. 1994] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, “Kodaira–Spencer theory of gravity and exact results for quantum string amplitudes”, *Comm. Math. Phys.* **165**:2 (1994), 311–427. MR 95f:32029 Zbl 0815.53082
- [Bismut 1986] J.-M. Bismut, “Localization formulas, superconnections, and the index theorem for families”, *Comm. Math. Phys.* **103**:1 (1986), 127–166. MR 87f:58147 Zbl 0602.58042
- [Bismut 1992] J.-M. Bismut, “Bott–Chern currents, excess normal bundles and the Chern character”, *Geom. Funct. Anal.* **2**:3 (1992), 285–340. MR 94a:58206 Zbl 0776.32007
- [Bismut 2004] J.-M. Bismut, “Holomorphic and de Rham torsion”, *Compos. Math.* **140**:5 (2004), 1302–1356. MR 2081158 Zbl 02110378
- [Bismut and Lebeau 1991] J.-M. Bismut and G. Lebeau, “Complex immersions and Quillen metrics”, *Inst. Hautes Études Sci. Publ. Math.* **74** (1991), 1–297. MR 94a:58205 Zbl 0784.32010
- [Bismut et al. 1988a] J.-M. Bismut, H. Gillet, and C. Soulé, “Analytic torsion and holomorphic determinant bundles, I: Bott–Chern forms and analytic torsion”, *Comm. Math. Phys.* **115**:1 (1988), 49–78. MR 89g:58192a Zbl 0651.32017
- [Bismut et al. 1988b] J.-M. Bismut, H. Gillet, and C. Soulé, “Analytic torsion and holomorphic determinant bundles, III: Quillen metrics on holomorphic determinants”, *Comm. Math. Phys.* **115**:2 (1988), 301–351. MR 89g:58192c Zbl 0651.32017
- [Bismut et al. 1990] J.-M. Bismut, H. Gillet, and C. Soulé, “Complex immersions and Arakelov geometry”, pp. 249–331 in *The Grothendieck Festschrift*, vol. I, Progr. Math. **86**, Birkhäuser, Boston, 1990. MR 92a:14019 Zbl 0744.14015
- [Bott 1967] R. Bott, “A residue formula for holomorphic vector-fields”, *J. Differential Geometry* **1** (1967), 311–330. MR 38 #730 Zbl 0179.28801
- [Feng 2003] H. Feng, “Holomorphic equivariant cohomology via a transversal holomorphic vector field”, *Internat. J. Math.* **14**:5 (2003), 499–514. MR 2004j:32022 Zbl 1050.32013
- [Gillet and Soulé 1991] H. Gillet and C. Soulé, “Analytic torsion and the arithmetic Todd genus”, *Topology* **30**:1 (1991), 21–54. MR 92d:14015 Zbl 0787.14005

- [Liu 1995] K. Liu, “Holomorphic equivariant cohomology”, *Math. Ann.* **303**:1 (1995), 125–148. MR 97f:32041 Zbl 0835.14006
- [Ray and Singer 1973] D. B. Ray and I. M. Singer, “Analytic torsion for complex manifolds”, *Ann. of Math. (2)* **98** (1973), 154–177. MR 52 #4344 Zbl 0267.32014
- [Zhang 1990] W. Zhang, “A remark on a residue formula of Bott”, *Acta Math. Sinica (N.S.)* **6**:4 (1990), 306–314. MR 91j:58153 Zbl 0738.32007
- [Zhang n.d.] W. Zhang, “Equivariant Dolbeault complex and total Quillen metrics”, preprint.

Received July 25, 2003. Revised February 13, 2004.

HUITAO FENG
COLLEGE OF MATHEMATICAL SCIENCES
NANKAI UNIVERSITY
300071, TIANJIN
CHINA
fht@nankai.edu.cn

XIAONAN MA
CENTRE DE MATHÉMATIQUES
UMR 7640 DU CNRS
ÉCOLE POLYTECHNIQUE
91128 PALAISEAU CEDEX
FRANCE
ma@math.polytechnique.fr