

*Pacific
Journal of
Mathematics*

**UPPER BOUNDS FOR THE SPECTRAL RADIUS
OF THE $n \times n$ HILBERT MATRIX**

PETER OTTE

UPPER BOUNDS FOR THE SPECTRAL RADIUS OF THE $n \times n$ HILBERT MATRIX

PETER OTTE

We derive upper bounds for the spectral radius of the $n \times n$ Hilbert matrix. The key idea is to write the Hilbert matrix as integral operator with positive kernel function and then to use a Wielandt-type min-max principle for the spectral radius. Choosing special trial functions yields a new bound that improves the best bound known heretofore.

1. Introduction

The spectral asymptotics of the Hilbert matrix has attracted a lot of interest concerning both the lowest and the largest eigenvalue. Here we shall focus on the spectral radius ρ_n of the $n \times n$ Hilbert matrix for which we shall prove, particularly, the bound

$$(1) \quad \rho_n \leq 2w_n \arcsin \frac{1}{w_n} \quad \text{with} \quad w_n := 2 \left(\frac{(n!)^2}{(2n)!} \right)^{1/2n}, \quad n \in \mathbb{N}.$$

This improves, at least for large values of n , Cassels' bound, given in (5) below, which is the best hitherto known. Numerical computations suggest that (1) is actually better for all n except $n = 1, 2$.

We base the proof of (1) upon relating the Hilbert matrix to an integral operator H_n whose spectral radius can be expressed by a min-max principle for operators having positive kernel functions:

$$(2) \quad \rho_n = \inf_{\varphi \in M} \sup_{0 < x < 1} \frac{(H_n \varphi)(x)}{\varphi(x)}$$

where M is some set of appropriate trial functions. For the sake of completeness we shall prove (2) without recourse to the general theory. In the matrix case the above min-max principle is due to Wielandt [1950] and related to the enclosure result of Collatz [1942]. It has been generalized in many directions; see [Friedland 1990; Marek 1966; Schaefer 1984], for example.

MSC2000: 15A42, 15A60, 47G10.

Keywords: Hilbert matrix, Hilbert inequality, spectral radius, Wielandt min-max principle, integral operator.

To derive estimates we pick $\varphi(x) := (1-x)^\gamma$ in (2), with $-1 < \gamma < 0$. We restrict ourselves to the case $\gamma = -\frac{1}{2}$, for which the calculations are manageable, and obtain (1).

Hilbert was the first one to investigate spectral properties of the matrix named after him. In his lectures he showed his double series theorem stating that ρ_n stays finite as $n \rightarrow \infty$; this was first published by Weyl [1908] (see also [Wiener 1910]). The concrete inequality

$$(3) \quad \rho_n \leq \pi$$

is due to Schur [1911]. This is the optimal constant that does not depend on the dimension n . However, if we do want the bound to depend on n it is possible to strengthen (3). Frazer [1946] obtained

$$(4) \quad \rho_n \leq n \sin \frac{\pi}{n} \quad \text{for } n \geq 2,$$

by refining a method of Fejér and Riesz [1921], which they used to prove what is now called the Fejér–Riesz inequality for analytic functions. Equation (4) was later rediscovered by Hsiang [1957] and Yahya [1965], and was eventually improved by Cassels [1948] to

$$(5) \quad \rho_n \leq 2 \arctan \sqrt{2n}.$$

Finally, it might be instructive to look at the asymptotic expansion of ρ_n . The first asymptotic result was obtained by Taussky [1949] by computing the quadratic form with special trial vectors having components $c_k := 1/\sqrt{k}$; it was

$$\rho_n = \pi + O\left(\frac{1}{\ln n}\right).$$

The exact asymptotic behaviour

$$\rho_n = \pi - \frac{\pi^5}{2 \ln^2 n} + O\left(\frac{\ln \ln n}{\ln^3 n}\right)$$

was determined by de Bruijn and Wilf [1962], who compared the matrix operator with an integral operator whose spectral asymptotics can be derived from general results of Widom [1958] (see also [Widom 1961]).

2. Estimates for the spectral radius

We start by relating the Hilbert matrix

$$(6) \quad A_n := \left(\frac{1}{j+k+1} \right)_{j,k=0,\dots,n-1}$$

to the integral operator $H_n : C[0, 1] \rightarrow C[0, 1]$ having the kernel function

$$(7) \quad K_n(xy) := \sum_{j=0}^{n-1} (xy)^j = \frac{1 - (xy)^n}{1 - xy}.$$

For $n = \infty$ this operator was used by Magnus [1950] to study the spectrum of the infinite Hilbert matrix. We let H_n act on $C[0, 1]$ because we want to have sufficiently many trial functions at hand. As we hoped, H_n has (almost) the same spectrum as A_n . In particular, they have the same spectral radius, henceforth denoted by ρ_n .

Lemma 1. *Let $C[0, 1]$ be equipped with the usual maximum norm. Then $H_n : C[0, 1] \rightarrow C[0, 1]$ is a bounded linear operator. The respective spectra of the Hilbert matrix A_n and the integral operator H_n are the same apart from 0. Their common spectral radius ρ_n can be expressed by*

$$(8) \quad \rho_n = \inf_{\varphi \in M} \sup_{0 < x < 1} \frac{(H_n \varphi)(x)}{\varphi(x)}, \text{ where } M := \{\varphi \in L^1[0, 1] \mid \varphi > 0, \frac{1}{\varphi} \in C[0, 1]\}.$$

Proof. It is clear from the definition and (7) that H_n is linear and bounded. Also (7) shows that H_n has n -dimensional range spanned by the monomials x^k , for $k = 0, \dots, n-1$, which implies that the spectrum of H_n consists only of eigenvalues. To each $c \in \mathbb{C}^n$ we associate $\varphi_c \in C[0, 1]$ in the natural way:

$$(9) \quad c = (c_0, \dots, c_{n-1}) \in \mathbb{C}^n \longleftrightarrow \varphi_c(x) = \sum_{j=0}^{n-1} c_j x^j.$$

The statement on the spectra then follows from

$$\begin{aligned} (H_n \varphi_c)(x) &= \int_0^1 \sum_{j=0}^{n-1} (xy)^j \sum_{k=0}^{n-1} c_k y^k dy \\ &= \sum_{j,k=0}^{n-1} c_k x^j \int_0^1 y^{j+k} dy = \sum_{j=0}^{n-1} x^j \sum_{k=0}^{n-1} \frac{1}{j+k+1} c_k. \end{aligned}$$

Note that H_n must have a kernel and A_n does not.

To prove Formula (8) we recall from the Perron–Frobenius Theorem that, since A_n has positive entries, ρ_n is an eigenvalue of A_n and hence of H_n . Let v be the corresponding eigenfunction. Writing down the eigenvalue equation for v and dividing by $\varphi \in M$ yields

$$\rho_n \frac{v(x)}{\varphi(x)} = \int_0^1 K_n(xy) \frac{\varphi(y)}{\varphi(x)} \frac{v(y)}{\varphi(y)} dy.$$

This shows that $v/\varphi \in C[0, 1]$ is an eigenfunction of the operator $H_{n,\varphi}$ with kernel

$$K_{n,\varphi}(xy) := K_n(xy) \frac{\varphi(y)}{\varphi(x)},$$

whence $\rho_n \leq \rho(H_{n,\varphi})$, the spectral radius of $H_{n,\varphi}$. Since $\rho(H_{n,\varphi}) \leq \|H_{n,\varphi}\|_\infty$ we conclude that

$$\rho_n \leq \|H_{n,\varphi}\|_\infty = \sup_{0 < x < 1} \int_0^1 K_{n,\varphi}(xy) dy$$

where we have used $\varphi(x) > 0$, $K_n(xy) \geq 0$, and thus $K_{n,\varphi}(xy) \geq 0$. To show equality in (8) we once again invoke the Perron–Frobenius Theorem, according to which the eigenvector of A_n belonging to ρ_n can be chosen to have positive components, whence we can, via (9), likewise choose the eigenfunction $v > 0$. In particular, $v \in M$. \square

We use Lemma 1 to estimate the spectral radius from above by cleverly choosing trial functions in (8):

$$(10) \quad r_n(x) := \frac{(H_n \varphi)(x)}{\varphi(x)} = \frac{1}{\varphi(x)} \int_0^1 \frac{1 - (xy)^n}{1 - xy} \varphi(y) dy \quad \text{for } n \in \mathbb{N}.$$

To get an idea of what the φ 's should look like we cast r_n into a form more amenable to further investigation. The crucial point is to evaluate the integral

$$J_n(x) := \int_0^1 \frac{y^n}{1 - xy} \varphi(y) dy.$$

We start by differentiating with respect to x :

$$(11) \quad J'_n(x) = \int_0^1 \frac{y^{n+1}}{(1 - xy)^2} \varphi(y) dy = \frac{1}{x} \int_0^1 \frac{y^n}{(1 - xy)^2} \varphi(y) dy - \frac{1}{x} J_n(x).$$

The explicitly written integral on the right can also be produced by integration by parts, which we perform in such a way that $\varphi(1)$ is omitted because our trial functions will have a singularity at $x = 1$:

$$\begin{aligned} J_n(x) &= \left[(y - 1) \frac{y^n}{1 - xy} \varphi(y) \right]_0^1 \\ &\quad - \int_0^1 (y - 1) \left(\left(\frac{ny^{n-1}}{1 - xy} + \frac{xy^n}{(1 - xy)^2} \right) \varphi(y) + \frac{y^n}{1 - xy} \varphi'(y) \right) dy \\ &= \delta_n \varphi(0) + n J_{n-1}(x) - n J_n(x) + (x - 1) \int_0^1 \frac{y^n}{(1 - xy)^2} \varphi(y) dy \\ &\quad + J_n(x) + \tilde{J}_n(x), \end{aligned}$$

with $\delta_n := \delta_{n,0}$ the Kronecker delta and

$$\tilde{J}_n(x) := \int_0^1 \frac{y^n}{1-xy} (1-y)\varphi'(y) dy.$$

Hence we can eliminate the integral in question from (11):

$$(12) \quad J'_n(x) = \frac{1}{x(1-x)} (\delta_n \varphi(0) + nJ_{n-1}(x) - nJ_n(x) + \tilde{J}_n(x)) - \frac{1}{x} J_n(x).$$

To eliminate the annoying J_{n-1} we observe that

$$J_n(x) = \int_0^1 \frac{y^n}{1-xy} \varphi(y) dy = \frac{1}{x} J_{n-1}(x) - \frac{1}{x} \int_0^1 y^{n-1} \varphi(y) dy,$$

and therewith (12) becomes

$$(13) \quad J'_n(x) = \frac{\delta_n}{x(1-x)} \varphi(0) - \frac{n+1}{x} J_n(x) + \frac{\kappa_n}{x(1-x)} + \frac{1}{x(1-x)} \tilde{J}_n(x)$$

where we have put

$$(14) \quad \kappa_0 := 0, \quad \kappa_n := n \int_0^1 y^{n-1} \varphi(y) dy \quad \text{for } n \in \mathbb{N}.$$

We are going to express

$$(15) \quad \Phi_n(x) := \frac{x^n}{\varphi(x)} J_n(x)$$

by dint of (13) through a differential equation:

$$\begin{aligned} \Phi'_n(x) &= -\frac{\varphi'(x)}{\varphi^2(x)} x^n J_n(x) + \frac{nx^{n-1}}{\varphi(x)} J_n(x) + \frac{x^n}{\varphi(x)} J'_n(x) \\ &= -\left(\frac{\varphi'(x)}{\varphi(x)} + \frac{1}{x}\right) \Phi_n(x) + \frac{x^{n-1}}{(1-x)\varphi(x)} \tilde{J}_n(x) + \frac{x^{n-1}}{(1-x)\varphi(x)} (\delta_n \varphi(0) + \kappa_n). \end{aligned}$$

At this point we fix our trial function φ in such a way that

$$(16) \quad (1-x)\varphi'(x) = -\gamma\varphi(x),$$

that is, $\varphi(x) = (1-x)^\gamma$ with some $\gamma \in \mathbb{R}$ to be specified later, whereby \tilde{J}_n becomes a multiple of J_n , and we arrive at a differential equation for Φ_n :

$$(17) \quad \Phi'_n(x) = -(\gamma+1)\frac{1}{x}\Phi_n(x) + \frac{x^{n-1}}{(1-x)^{1+\gamma}} (\delta_n + \kappa_n).$$

This is equivalent to

$$(x^{1+\gamma}\Phi_n(x))' = \frac{x^{n+\gamma}}{(1-x)^{1+\gamma}} (\delta_n + \kappa_n),$$

which we can solve immediately for Φ_n :

$$(18) \quad \Phi_n(x) = \frac{\delta_n + \kappa_n}{x^{1+\gamma}} \int_0^x \frac{\xi^{n+\gamma}}{(1-\xi)^{1+\gamma}} d\xi \quad \text{for } n \in \mathbb{N}_0.$$

In particular, we can now see that γ must satisfy $-1 < \gamma < 0$ in order to yield well defined integrals and to have $\varphi \in M$ in (8). We summarize our calculations.

Theorem 2. *The spectral radius ρ_n of the $n \times n$ Hilbert matrix A_n as in (6) can be estimated by*

$$(19) \quad \rho_n \leq \inf_{0 < \alpha < 1} \sup_{0 < x < 1} \frac{1}{x^{1-\alpha}} \int_0^x \frac{1 - \kappa_n \xi^n}{\xi^\alpha (1-\xi)^{1-\alpha}} d\xi,$$

where

$$(20) \quad \kappa_n = \frac{n!}{(n-\alpha)(n-1-\alpha) \cdots (1-\alpha)} \quad \text{for } n \in \mathbb{N}.$$

Proof. Put $\alpha := -\gamma$ and use in turn the min-max principle (8) and the definitions of r_n and Φ_n as in (10) and (15), with φ being chosen according to (16) to obtain

$$\rho_n \leq \inf_{0 < \alpha < 1} \sup_{0 < x < 1} r_n(x) = \inf_{0 < \alpha < 1} \sup_{0 < x < 1} (\Phi_0(x) - \Phi_n(x)).$$

Then (19) follows directly from the representation (18) of Φ_n .

For φ as in (16) the integral in (14) is Euler's beta function. Hence,

$$\kappa_n = nB(n, 1-\alpha) = \frac{n\Gamma(n)\Gamma(1-\alpha)}{\Gamma(n+1-\alpha)}$$

wherefrom we deduce (20). □

The optimal way to derive bounds on ρ_n would be to determine the maximum of the function r_n exactly. Unfortunately, this turns out to be rather complicated, and we content ourselves with narrowing the region where the maximum must lie.

Corollary 3. *The spectral radius ρ_n of the $n \times n$ Hilbert matrix A_n can be estimated by*

$$(21) \quad \rho_n \leq \inf_{0 < \alpha < 1} \kappa_n^{(1-\alpha)/n} \int_0^{1/\kappa_n^{1/n}} \frac{1}{\xi^\alpha (1-\xi)^{1-\alpha}} d\xi,$$

which in the case $\alpha = \frac{1}{2}$ specializes to

$$(22) \quad \rho_n \leq 2w_n \arcsin \frac{1}{w_n} \quad \text{with } w_n := \kappa_n^{1/2n} = 2 \left(\frac{(n!)^2}{(2n)!} \right)^{1/2n}.$$

Proof. When $1 - \kappa_n \xi^n \leq 0$ the function r_n is decreasing, whence the maximum must lie in the interval $[0, x_0]$ for x_0 the unique zero of the integrand in (19):

$$1 - \kappa_n x_0^n = 0, \quad \text{i.e.,} \quad x_0 = 1/\kappa_n^{1/n}.$$

We conclude

$$\sup_{0 < x < 1} r_n(x) = \sup_{0 < x \leq x_0} r_n(x) \leq \sup_{0 < x \leq x_0} \Phi_0(x) = \Phi_0(x_0)$$

because the $\Phi_n(x)$ are nonnegative and Φ_0 increases.

For $\alpha = \frac{1}{2}$ the w_n are easily obtained from (22) and (20) and the integral in (21) can be evaluated by the change of variables $\xi = s^2$. □

Finally, we shall check that our estimate (22) is indeed better than (5). Using some familiar formulae for arctan and arcsin we obtain

$$\begin{aligned} \arctan \sqrt{2n} - w_n \arcsin \frac{1}{w_n} &= \arctan \sqrt{2n} - \arcsin \frac{1}{w_n} - (w_n - 1) \arcsin \frac{1}{w_n} \\ &\geq \arctan \sqrt{2n} - \arctan \frac{1}{\sqrt{w_n^2 - 1}} - \frac{\pi}{2}(w_n - 1) \\ &= \arctan \frac{\sqrt{2n(w_n^2 - 1)} - 1}{\sqrt{w_n^2 - 1} + \sqrt{2n}} - \frac{\pi}{2}(w_n - 1). \end{aligned}$$

Now the asymptotics of the middle binomial coefficient yields

$$w_n \sim 2 \left(\frac{\sqrt{\pi n}}{4^n} \right)^{1/2n} \sim n^{1/4n} \quad \text{as } n \rightarrow \infty,$$

which implies immediately $\lim_{n \rightarrow \infty} w_n = 1$, and further

$$n(w_n - 1) \sim n(n^{1/4n} - 1) = n(e^{(\ln n)/4n} - 1) \sim \frac{1}{4} \ln n \quad \text{as } n \rightarrow \infty.$$

Therefore, for large n ,

$$\frac{\sqrt{2n(w_n^2 - 1)} - 1}{\sqrt{w_n^2 - 1} + \sqrt{2n}} \geq \frac{1}{4} \frac{\sqrt{2n(w_n^2 - 1)}}{\sqrt{2n}} \geq \frac{1}{4} \sqrt{w_n - 1}.$$

Noting $\arctan x \geq cx$ for small x with some constant $c > 0$ and using the monotonicity of arctan we conclude

$$\arctan \sqrt{2n} - w_n \arcsin \frac{1}{w_n} \geq \frac{c}{4} \sqrt{w_n - 1} - \frac{\pi}{2}(w_n - 1) > 0$$

for large values of n . With some care it should be possible to show the statement for smaller values of n , too.

3. Remarks

We suggest some topics that might be worth further investigation.

(1) In order to derive from [Theorem 2](#) a bound that can be computed more or less explicitly we did not determine in [\(19\)](#) the maximum of the function r_n exactly. Thus, the first possibility to strengthen [\(22\)](#) is to study the maximum of r_n .

(2) Also for computational reasons we fixed the exponent $\alpha = \frac{1}{2}$ in [\(21\)](#). However, numerical computations suggest that $\alpha = \frac{1}{2}$ is generally not the optimal choice and that other values of α give much more accurate estimates. According to a theorem of Čebyšev the integral in [\(21\)](#) can be evaluated for any $0 < \alpha < 1$ in closed form by means of elementary functions. It is not clear whether these elementary functions allow for an efficient minimizing procedure.

(3) A vaguer idea is to pick other trial functions than $(1 - x)^{-\alpha}$. Our method will work as long as we arrive at a differential equation for Φ_n as in [\(17\)](#).

(4) Since Wielandt's min-max principle is accompanied by a max-min principle, one can also think of deriving lower bounds for the spectral radius in which case; however, completely different trial functions are needed.

Acknowledgments

I thank Robert Denk and Hubert Kalf for stimulating discussions and their steady interest in this subject.

References

- [de Bruijn and Wilf 1962] N. G. de Bruijn and H. S. Wilf, "On Hilbert's inequality in n dimensions", *Bull. Amer. Math. Soc.* **68** (1962), 70–73. [MR 31 #291](#) [Zbl 0105.04201](#)
- [Cassels 1948] J. W. S. Cassels, "An elementary proof of some inequalities", *J. London Math. Soc.* **23** (1948), 285–290. [MR 10,434g](#) [Zbl 0034.18501](#)
- [Collatz 1942] L. Collatz, "Einschließungssatz für die charakteristischen Zahlen von Matrizen", *Math. Z.* **48** (1942), 221–226. [MR 5,30d](#) [Zbl 0027.00604](#)
- [Fejér and Riesz 1921] L. Fejér and F. Riesz, "Über einige funktionentheoretische Ungleichungen", *Math. Z.* **11** (1921), 305–314. [JFM 48.0327.01](#)
- [Frazer 1946] H. Frazer, "Note on Hilbert's inequality", *J. London Math. Soc.* **21** (1946), 7–9. [MR 8,259h](#) [Zbl 0060.14903](#)
- [Friedland 1990] S. Friedland, "Characterizations of the spectral radius of positive operators", *Linear Algebra Appl.* **134** (1990), 93–105. [MR 92d:47054](#) [Zbl 0707.15005](#)
- [Hsiang 1957] F. C. Hsiang, "An inequality for finite sequences", *Math. Scand.* **5** (1957), 12–14. [MR 19,955a](#) [Zbl 0078.24701](#)
- [Magnus 1950] W. Magnus, "On the spectrum of Hilbert's matrix", *Amer. J. Math.* **72** (1950), 699–704. [MR 12,836e](#) [Zbl 0041.23805](#)

- [Marek 1966] I. Marek, “Spektrale Eigenschaften der \mathcal{K} -positiven Operatoren und Einschließungssätze für den Spektralradius”, *Czechoslovak Math. J.* **16** (91) (1966), 493–517. [MR 36 #711](#) [Zbl 0152.33701](#)
- [Schaefer 1984] H. H. Schaefer, “A minimax theorem for irreducible compact operators in L^p -spaces”, *Israel J. Math.* **48**:2-3 (1984), 196–204. [MR 86g:47042](#) [Zbl 0576.47020](#)
- [Schur 1911] I. Schur, “Bemerkungen zur Theorie der beschränkten Bilinearformen mit unendlich vielen Veränderlichen”, *J. Reine Angew. Math.* **140** (1911), 1–28. [JFM 42.0367.01](#)
- [Taussky 1949] O. Taussky, “A remark concerning the characteristic roots of the finite segments of the Hilbert matrix”, *Quart. J. Math., Oxford Ser.* **20** (1949), 80–83. [MR 11,16a](#) [Zbl 0036.01302](#)
- [Weyl 1908] H. Weyl, *Singuläre Integralgleichungen mit besonderer Berücksichtigung des Fourier-schen Integraltheorems*, Inauguraldissertation, Universität Göttingen, 1908.
- [Widom 1958] H. Widom, “On the eigenvalues of certain Hermitian operators”, *Trans. Amer. Math. Soc.* **88** (1958), 491–522. [MR 20 #4782](#) [Zbl 0101.09202](#)
- [Widom 1961] H. Widom, “Extreme eigenvalues of translation kernels”, *Trans. Amer. Math. Soc.* **100** (1961), 252–262. [MR 25 #2420](#) [Zbl 0197.10903](#)
- [Wielandt 1950] H. Wielandt, “Unzerlegbare, nicht negative Matrizen”, *Math. Z.* **52** (1950), 642–648. [MR 11,710g](#) [Zbl 0035.29101](#)
- [Wiener 1910] F. Wiener, “Elementarer Beweis eines Reihensatzes von Herrn Hilbert”, *Math. Ann.* **68** (1910), 361–366. [JFM 41.0391.04](#)
- [Yahya 1965] Q. A. M. M. Yahya, “On the generalisation of Hilbert’s inequality”, *Amer. Math. Monthly* **72** (1965), 518–520. [MR 32 #4235](#) [Zbl 0125.30101](#)

Received October 20, 2003. Revised March 19, 2004.

PETER OTTE
RUHR-UNIVERSITÄT BOCHUM
FAKULTÄT FÜR MATHEMATIK
UNIVERSITÄTSSTRASSE 150
D-44780 BOCHUM
GERMANY
peter.otte@ruhr-uni-bochum.de