

*Pacific
Journal of
Mathematics*

**A NEW BOUND FOR FINITE FIELD BESICOVITCH SETS
IN FOUR DIMENSIONS**

TERENCE TAO

Volume 222 No. 2

December 2005

A NEW BOUND FOR FINITE FIELD BESICOVITCH SETS IN FOUR DIMENSIONS

TERENCE TAO

Let F be a finite field with characteristic greater than two. A Besicovitch set in F^4 is a set $P \subseteq F^4$ containing a line in every direction. The Kakeya conjecture asserts that P and F^4 have roughly the same size, in the sense that $|P|/|F|^4$ exceeds $C_\varepsilon|F|^{-\varepsilon}$ for $\varepsilon > 0$ arbitrarily small, where C_ε does not depend on P or F . Wolff showed that $|P|$ exceeds a universal constant times $|F|^3$. Here we improve his exponent to $3 + \frac{1}{16} - \varepsilon$ for $\varepsilon > 0$ arbitrarily small. On the other hand, we show that Wolff's bound of $|F|^3$ is sharp if we relax the assumption that the lines point in different directions. One new feature in the argument is the use of some basic algebraic geometry.

1. Introduction

Let F be a finite field with characteristic greater than 2. For any $n \geq 2$, we define a *Besicovitch set* in F^n to be a set $P \subseteq F^n$ containing a line in every direction (every equivalence class under parallelism). The *finite field Kakeya conjecture* (see [Wolff 1998b], for example) asserts that $|P| \geq C_\varepsilon|F|^{n-\varepsilon}$ for any $\varepsilon > 0$, where $|P|$ denotes the cardinality of P and the quantities C_ε are independent of $|F|$. This conjecture is the finite field analogue of the Euclidean Kakeya set conjecture, which is related to several other problems in harmonic analysis; see [Wolff 1998b; Mockenhaupt and Tao 2004] for further discussion on this. Basically, one can view the finite field Kakeya problem as a simplified model for the more interesting Euclidean Kakeya problem, where technical difficulties involving small separations, small angles, and multiple scales complicate the task (as discussed briefly in Section 9).

Informally, the Kakeya conjecture asserts that lines pointing in different directions in F^n cannot have substantial overlap. This conjecture has been proved in two dimensions but remains open in higher dimensions. In [Wolff 1998b] (see also [Wolff 1995; Mockenhaupt and Tao 2004]) it was shown that $|P| \gtrsim |F|^{(n+2)/2}$,

MSC2000: 42B25, 05C35.

Keywords: Besicovitch sets, affine spaces over finite fields, Kakeya conjecture, reguli.

This work was inspired by the April 2002 Instructional Conference on Combinatorial aspects of Mathematical Analysis at University of Edinburgh. The author is a Clay Prize Fellow and is supported by the Packard Foundation.

where $A \gtrsim B$ means that $A \geq C^{-1}B$ for some universal constant C . In fact, more was proved:

Definition 1.1. A family L of lines in F^n is said to obey the *Wolff axiom* if for every $2 \leq k \leq n - 1$, every k -dimensional affine subspace $V \subset F^n$ contains at most $O(|F|^{k-1})$ lines in L . (Here we view the field F as being quite large, and the family L as depending on F . The implied constant in the $O(\cdot)$ notation may depend on n and k but is uniform in F . Also Recall that an affine subspace is a translate of a vector subspace of F^n .)

Theorem 1.2 [Wolff 1995; 1998b]. *If L is a family of $O(|F|^{n-1})$ lines obeying the Wolff axiom, and $P \subseteq F^n$ contains all the lines in L , then $|P| \gtrsim |F|^{(n+2)/2}$.*

In fact one only needs to use the Wolff axiom for $k = 2$. From this theorem and the observation that any family of lines that point in different directions automatically obeys the Wolff axiom, we immediately see that Besicovitch sets have cardinality $\gtrsim |F|^{(n+2)/2}$.

In [Mockenhaupt and Tao 2004] (see also [Katz et al. 2000]) it was observed that the statement of Theorem 1.2 is sharp in three dimensions, in the sense that there exist finite fields F and collections of lines L in F^3 obeying the Wolff axiom and a collection P of points containing all the lines in L , such that $|P| \sim |F|^{5/2}$ (where $A \sim B$ means that $A \gtrsim B$ and $B \gtrsim A$). Indeed, if F contains a subfield G of index 2, with the accompanying involution $z \mapsto \bar{z}$ on F , one can take P to be the *Heisenberg group*

$$P := \{(z_1, z_2, z_3) \in F^3 : \text{Im}(z_3) = \text{Im}(z_1\bar{z}_2)\},$$

where $\text{Im}(z) := (z - \bar{z})/2$. (It is an interesting question whether an example similar to this can be obtained if F does not contain a subfield of index 2.)

Our first observation is that Theorem 1.2 is also sharp in four dimensions:

Proposition 1.3. *Let $\langle \cdot, \cdot \rangle : F^4 \times F^4 \rightarrow F$ be a nondegenerate symmetric quadratic form on F^4 . Let P be the “unit sphere”*

$$(1-1) \quad P := \{x \in F^4 : \langle x, x \rangle = 1\}$$

and let L be the set of all lines of the form $\{x + tv : t \in F\}$, where $x \in F^4$, $v \in F^4 \setminus \{0\}$ are such that $\langle x, x \rangle = 1$, $\langle v, x \rangle = 0$, and $\langle v, v \rangle = 0$. Then L has cardinality $|L| \sim |F|^3$ and obeys the Wolff axiom, while P has cardinality $|P| \sim |F|^3$ and contains all the lines in L .

We prove this in Section 3. A similar counterexample can be created in \mathbb{R}^4 as long as one chooses the form $\langle \cdot, \cdot \rangle$ to be indefinite. The proposition does not contradict the Kakeya conjecture because the lines L do not all point in different directions (despite obeying the Wolff axiom). Nevertheless, it seems of interest

to extend this example (and the Heisenberg group) to higher dimensions, though perhaps the bound of $|F|^{(n+2)/2}$ in Theorem 1.2 need not be sharp for large n .

This example illustrates two things. Firstly, in order to progress toward the Kakeya conjecture in low dimensions one must make better use of the hypothesis that the lines in L point in different directions; merely assuming the Wolff axiom will not by itself suffice. (In high dimensions — say $n \geq 9$ — there are other, more “arithmetic” arguments available to improve upon Theorem 1.2. See [Bourgain 1999; Katz and Tao 1999; 2002b; Rogers 2001; Mockenhaupt and Tao 2004].)

Secondly, the algebraic geometry of quadric surfaces may be relevant to the Kakeya problem.¹ In the three-dimensional case $n = 3$, quadric surfaces are essentially the same thing as *reguli*, those ruled surfaces consisting of all the lines that intersect three fixed lines in general position. In particular, we have the “three-line lemma”, which asserts that given three mutually skew lines in F^3 , there are at most $O(|F|)$ lines in different directions that intersect all three.

Reguli have already come up in the work of Schlag [1998], who used the three-line lemma to give a new proof of Bourgain’s estimate [1991]

$$(1-2) \quad |P| \gtrsim |F|^{7/3}$$

in three dimensions. While it is true that this bound has since been superseded by the estimate in Theorem 1.2, we shall need to follow [Schlag 1998] and make use of reguli and the three-line lemma in what follows. We are indebted to Nets Katz for pointing out the usefulness of reguli in the low-dimensional Kakeya problem. Indeed, our work here was inspired by similar work in three dimensions by Nets Katz (currently in preparation).

The main result of this paper is the following improved bound on the cardinality of Besicovitch sets in four dimensions. We use $A \lesssim B$ to denote the estimate $A \leq C_\varepsilon |F|^\varepsilon B$ for any $\varepsilon > 0$, where C_ε is a quantity depending only on ε .

Theorem 1.4. *If P is a Besicovitch set in $|F|^4$, then $|P| \gtrsim |F|^{3+\frac{1}{16}}$.*

One can probably improve the \gtrsim to a \gtrsim by going through the argument in this paper more carefully, but we will not do so here in order to simplify the exposition.

The paper is organized as follows. After setting out our incidence geometry notation in Section 2, we prove Proposition 1.3 in Section 3. We then review some basic algebraic geometry in Section 4, culminating in a “three-regulus lemma” in F^4 , which will be the analogue of the three-line lemma in F^3 . In Section 5 we review some combinatorial preliminaries, before starting the proof of Theorem 1.4, which occupies the next three sections. The first step is to use a standard

¹There seems to be a parallel phenomenon in recent work on Szemerédi’s theorem on arithmetic progressions, in that while arithmetic progressions are rather “linear” quantities, they give rise rather naturally to other “quadratic” objects which then need to be studied. See [Gowers 1998].

“iterated popularity” argument (as in [Christ 1998], for example), together with a rudimentary version of the “plate number” argument in [Wolff 1998a], in order to refine the Besicovitch set to a uniform, nondegenerate collection of points and lines. After a sufficient number of refinements, we can construct a large number of reguli incident to many lines in the Besicovitch set, and eventually get about $|F|^3$ lines incident to three distinct reguli (if $|P|$ is too close to $|F|^3$); this will contradict the three-regulus lemma mentioned earlier.

2. Incidence notation

We now set some notation for the finite field geometry of the affine space F^4 . A *line* in F^4 is a set of the form $l = \{x + tv : t \in F\}$ where $x, v \in F^4$ and v is nonzero. Two lines are *parallel* if they are translates of each other but not identical; a set of lines is said to *point in different directions* if no two lines in the set are parallel or identical.

A *2-plane* in F^4 is a set of the form $\pi = \{x + t_1v_1 + t_2v_2 : t_1, t_2 \in F\}$ where $x, v_1, v_2 \in F^4$ and v_1, v_2 are linearly independent. Two lines are *coplanar* if they lie in the same 2-plane; observe that coplanar lines must either be identical, parallel, or intersect in a point. A pair of lines are *skew* if they are not coplanar.

A *3-space* in F^4 is a set of the form $\lambda = \{x + t_1v_1 + t_2v_2 + t_3v_3 : t_1, t_2, t_3 \in F\}$ where $x, v_1, v_2, v_3 \in F^4$ and v_1, v_2, v_3 are linearly independent. Observe that any pair of skew lines lies in a unique 3-space. Two 3-spaces are *parallel* if they are disjoint, and one is the translate of the other.

We shall use the symbol p to refer to points, l to lines, π to 2-planes, and λ to 3-spaces. We use the symbol P to refer to sets of points, L to sets of lines, Π to sets of 2-planes, and Λ to sets of 3-spaces. We use $\text{Gr}(F^4, 1)$ to denote the space of all lines, $\text{Gr}(F^4, 2)$ to denote the space of all 2-planes, and $\text{Gr}(F^4, 3)$ to denote the space of all 3-spaces. (Note that these are the *affine* Grassmannians, in that the spaces do not need to contain the origin).

3. The counterexample

It is likely that Proposition 1.3 follows from the standard theory of Fano varieties of quadric surfaces, but we will just give an elementary argument.

Proof of Proposition 1.3. Let P and L be as in the proposition. It is clear from the construction that the lines in L lie in P . Now we verify the cardinality bounds. We begin with a standard lemma on the number of ways of representing a field element as a quadratic form.

Lemma 3.1. *Let $\langle \cdot, \cdot \rangle : F^n \times F^n \rightarrow F$ be a symmetric bilinear form on F^n with rank at least 1, and let $Q(x) := \langle x, x \rangle$ be the associated quadratic form. Then*

$$(3-1) \quad \{(x_1, \dots, x_n) \in F^n : Q(x_1, \dots, x_n) = x\} \lesssim |F|^{n-1}$$

for all $x \in F$.

If we know that $\langle \cdot, \cdot \rangle$ has rank at least 3, we can improve (3-1) to

$$(3-2) \quad \{(x_1, \dots, x_n) \in F^n : Q(x_1, \dots, x_n) = x\} \sim |F|^{n-1}$$

for all $x \in F$, if $|F|$ is sufficiently large.

Proof. By placing the quadratic form Q in normal form (recalling that $\text{char} F \neq 2$) we may assume that

$$Q(x_1, \dots, x_n) = \alpha_1 x_1^2 + \dots + \alpha_k x_k^2,$$

where k is the rank of Q and $\alpha_1, \dots, \alpha_k$ are nonzero elements of F . We may assume that $k = n$ since the general case $n \geq k$ follows by adding $n - k$ dummy variables. In particular $\alpha_j \neq 0$ for $j = 1, \dots, n$.

The bound (3-1) is now clear, since if we fix x_1, \dots, x_{n-1} and x then there are at most 2 choices for x_n . Now let us assume $n \geq 3$, and prove (3-2).

We use Gauss sums. We fix a nonprincipal character e of F , i.e. a multiplicative function $e : F \rightarrow S^1$ that is not identically 1. For instance, if $F = \mathbb{Z}/p\mathbb{Z}$ for some prime p , one can take $e(x) := \exp(2\pi i x/p)$.

For any $y \in F$, let $S(y)$ be the Gauss sum $S(y) := \sum_{x \in F} e(yx^2)$. As is well known (see [Mockenhaupt and Tao 2004], for example), $S(0)$ is equal to $|F|$, while $|S(y)| = |F|^{1/2}$ for all nonzero values of y .

Fix $x \in F$. By expanding the Kronecker delta as a Fourier series, we see that the number of solutions to (3-1) can be written as

$$\begin{aligned} \sum_{x_1, \dots, x_n \in F} \delta(\alpha_1 x_1^2 + \dots + \alpha_n x_n^2 - x) &= \frac{1}{|F|} \sum_{y \in F} \sum_{x_1, \dots, x_n \in F} e((\alpha_1 x_1^2 + \dots + \alpha_n x_n^2 - x)y) \\ &= \frac{1}{|F|} \sum_{y \in F} e(-xy) \prod_{i=1}^n S(\alpha_i y) \\ &= |F|^{n-1} + \frac{1}{|F|} \sum_{y \in F \setminus \{0\}} e(-xy) \prod_{i=1}^n S(\alpha_i y) \\ &= |F|^{n-1} + \frac{1}{|F|} \sum_{y \in F \setminus \{0\}} O(|F|^{n/2}) \\ &= |F|^{n-1} + O(|F|^{n/2}) \end{aligned}$$

as desired, since $n \geq 3$. □

From the lemma we see that $|P| \sim |F|^3$, as desired. Now we count the lines in L .

The lemma yields $\sim |F|^3$ choices of null direction $\{v \in F^4 \setminus 0 : \langle v, v \rangle = 0\}$. For each such v , the space $v^\perp := \{x \in F^4 : \langle x, v \rangle = 0\}$ is 3-dimensional (since Q is nondegenerate). Furthermore, since Q is nondegenerate on F^4 and v is a null vector, we see that Q must also be nondegenerate on v^\perp . Restricting Q to v^\perp (which is of course isomorphic to F^3) we see from Lemma 3.1 that there are $\sim |F|^2$ choices for x . Thus there are $\sim |F|^5$ possible pairs (x, v) that generate a line in L . However, each line in L is generated by $\sim |F|^2$ such pairs (x, v) , so we have $|L| \sim |F|^3$ as desired.

It remains to verify the Wolff axiom. First pick a 3-space λ and consider the lines in L which go through λ .

Pick an arbitrary point x_0 in λ , so that $\lambda - x_0$ is a three-dimensional subspace of F^4 . By Lemma 3.1, the number of null vectors $\{v \in \lambda - x_0 : \langle v, v \rangle = 0\}$ is $O(|F|^2)$.

Fix v as above. There are two cases. If $v^\perp \not\equiv (\lambda - x_0)$, then there are $O(|F|)$ choices of $x \in \lambda$ such that $\langle x, v \rangle = 0$ and $\langle x, x \rangle = 1$. But if $v^\perp \equiv (\lambda - x_0)$, then the number of choices for x could be as large as $O(|F|^2)$. But $(\lambda - x_0)^\perp$ only has cardinality $O(|F|)$, hence the number of v in the second category is at most $O(|F|)$. Thus the number of pairs (x, v) which can generate a line in λ is at most $O(|F|^3)$. But each line is generated by $\sim |F|^2$ pairs (x, v) . Thus the number of lines in λ is at most $O(|F|)$, which clearly implies the Wolff axiom for both $k = 2$ and $k = 3$. This completes the proof of Proposition 1.3. \square

4. Some basic algebraic geometry

Here we review some basic facts from algebraic geometry (for proofs see [Harris 1992], for example), and apply them to our Kakeya problem. The material we will need is not very advanced; basically, we need the concept of the dimension of an algebraic variety, and we need to know that this dimension behaves in the expected way with respect to intersections, projections, cardinality, etc. We shall also rely heavily on the basic fact that the dimension of an algebraic variety is always an *integer* (in contrast to, say, the “half-dimensional” field G mentioned in the introduction).

Let \bar{F} denote the algebraic closure of F and $n \geq 1$. An *algebraic variety* in \bar{F}^n is defined to be the zero locus of a collection Q_1, \dots, Q_k of \bar{F} -valued polynomials on the affine space \bar{F}^n . In this paper we shall always assume that our algebraic varieties have bounded degree, thus $k = O(1)$ and all the polynomials Q_1, \dots, Q_k have degree $O(1)$.

An algebraic variety V in \bar{F}^n has a well-defined *integer-valued* dimension $0 \leq d \leq n$; there are several equivalent definitions of this dimension, for instance d is

the smallest nonnegative integer such that generic affine spaces in \bar{F}^n of codimension greater than d are disjoint from V . (See [Harris 1992] for more equivalent definitions of dimension). If V has dimension n then it must be all of \bar{F}^n , while if V has dimension 0 then it can only consist of at most $O(1)$ points. Of course, the algebraic geometry notion of dimension is consistent with the linear algebra notion of dimension, thus for instance 3-spaces have dimension 3.

An algebraic variety is *irreducible* if it does not contain any proper sub-variety of the same dimension. Every algebraic variety of dimension k can be decomposed as a union of $O(1)$ irreducible varieties of dimension at most k .

We define an *algebraic variety* in F^n of dimension d to be a restriction to F^n of an algebraic variety in \bar{F}^n of dimension d . Observe that if V is a variety in F^n of dimension d then $|V| \lesssim |F|^d$ (this can be shown, for instance, by taking generic intersections with affine spaces of codimension d).

Let $L \subseteq \text{Gr}(F^n, 1)$ be a collection of lines which point in different directions. In the introduction we observed that this implies the Wolff axiom, that not too many lines in L can lie inside a k -space. In fact we can generalize this to k -dimensional varieties; see [Mockenhaupt and Tao 2004, Proposition 8.1]:

Lemma 4.1 (Generalized Wolff property). *Let $V \subseteq F^n$ be an algebraic variety in F^n of dimension k , and let $L \subseteq \text{Gr}(F^n, 1)$ be a collection of lines in F^n which point in different directions. Then*

$$|\{l \in L : l \subseteq V\}| \lesssim |F|^{k-1}.$$

Remark. The lines in Proposition 1.3 violate this property, but of course they do not point in different directions. (On the other hand, one can show that the lines arising from the Heisenberg example do obey this generalized Wolff property. It may be that in the three-dimensional theory, one needs to extend this lemma further, to cover not only varieties over F , but also over subfields of F such as G .)

Proof. We may of course assume that $|F| \gg 1$, since the claim is obvious for $|F|$ bounded.

We can embed F^n in the projective space PF^{n+1} , which we think of as the union of F^n with the hyperplane at infinity. By replacing the defining polynomials of V with their homogeneous counterparts, we can thus extend V to a k -dimensional variety \bar{V} in PF^{n+1} (see e.g. [Harris 1992]).

We break up \bar{V} into irreducible components, each of dimension at most k . We can assume that none of the irreducible components are contained inside the hyperplane at infinity, since we could simply remove those components and still have an extension of V . In particular we see that the intersection of \bar{V} with the hyperplane at infinity is at most $k - 1$ -dimensional.

Let l be a line in L , which we can extend to be a projective line \bar{l} in PF^{n+1} by adding a single point at infinity (the direction of l). Observe that the restriction of \bar{V} to \bar{l} is an algebraic variety of dimension either 0 or 1; in other words, either the projective line \bar{l} lies inside \bar{V} , or else \bar{l} intersects \bar{V} in at most $O(1)$ points. Thus in order for l to be contained in V , the direction of l must lie inside \bar{V} (assuming that $|F|$ is sufficiently large). But by the previous paragraph the number of such directions is at most $O(|F|^{k-1})$. Since the lines in L point in different directions, we are done. \square

As a consequence of this lemma we see that a Besicovitch set cannot have high intersection with an algebraic variety:

Corollary 4.2. *Let $V \subseteq F^n$ be an algebraic variety in F^n of dimension at most $n - 1$, and let $L \subseteq \text{Gr}(F^n, 1)$ be a collection of lines in F^n which point in different directions. Then*

$$|\{(p, l) \in V \times L : p \in l\}| \lesssim |F|^{n-1}.$$

The trivial upper bound for the left-hand side is $|F||L| \lesssim |F|^n$. Thus this lemma gains a power of $|F|$ over the trivial bound.

Proof. As in the proof of Lemma 4.1, we observe that every line l in F^n is either contained in V , or else intersects V in at most $O(1)$ points. The lines of the second type contribute at most $O(|L|) = O(|F|^{n-1})$ incidences, while by Lemma 4.1 the lines of the first type contribute at most $O(|F||F|^{\dim(V)-1}) = O(|F|^{n-1})$ incidences, and we are done. \square

A further consequence is that the *lines* of a Besicovitch set cannot have large intersection with an algebraic variety:

Corollary 4.3. *Let $L \subseteq \text{Gr}(F^n, 1)$ be a collection of lines in F^n which point in different directions, and let $P \subseteq F^n$ be a set of points containing all the lines in L . Let $W \subseteq \text{Gr}(F^n, 1)$ be an algebraic variety of lines of dimension at most $n - 1$. Then*

$$|L \cap W| \lesssim |F|^{n-2} + |F|^{-1}|P|.$$

Again, this lemma gains a power of $|F|$ over the trivial bound of $|F|^{n-1}$ (assuming P is not too huge).

Proof. Consider the set

$$X := \{(p, l) \in F^n \times W : p \in l\}.$$

This is an algebraic variety in $F^n \times \text{Gr}(F^n, 1)$ of dimension at most n . Now consider the map $\phi : X \rightarrow F^n$ given by $\phi(p, l) := p$. Observe that for any p in the image of ϕ , the fibers $\phi^{-1}(p)$ are either 0-dimensional (i.e. have cardinality $O(1)$), or at least 1-dimensional. This implies (see e.g. [Harris 1992]) that we have a decomposition

$\phi(X) := P_1 \cup P_2$, where the fibers $\phi^{-1}(p)$ are 0-dimensional for all $p \in P_1$, and P_2 is contained in an algebraic variety of dimension at most $n - 1$.

By the construction of P_1 we have

$$\begin{aligned} |\{(p, l) \in P_1 \times (L \cap W) : p \in l\}| &\lesssim |\{p \in P_1 : p \in l \text{ for some } l \in L \cap W\}| \\ &\lesssim |\{p \in P_1 : p \in P\}| \\ &\leq |P|. \end{aligned}$$

Also, by Corollary 4.2 we have

$$|\{(p, l) \in P_2 \times (L \cap W) : p \in l\}| \lesssim |F|^{n-1}.$$

Adding the two estimates, we see that

$$|F||L \cap W| = |\{(p, l) \in \phi(X) \times (L \cap W) : p \in l\}| \lesssim |F|^{n-1} + |P|$$

and the claim follows. □

To apply these results to our four-dimensional problem, we need some notation for reguli.

Definition 4.4. A *frame* f is a quadruplet $f = (l_1, l_2, l_3, \lambda)$ where $\lambda \in \text{Gr}(F^4, 3)$ is a 3-space in F^4 , and $l_1, l_2, l_3 \in \text{Gr}(F^4, 1)$ are distinct, mutually skew lines in F^4 which lie inside λ . If f is a frame, we write $\lambda(f)$ for λ . If $f = (l_1, l_2, l_3, \lambda)$ is a frame, we use $L(f)$ to denote the set of lines $l \in \text{Gr}(F^4, 1)$ which intersect l_1, l_2 , and l_3 , and $r(f) \subseteq \lambda$ to denote the union of all the lines in $L(f)$.

The set $r(f)$ is called the *regulus* generated by the frame f . It is a quadric in λ , that is, the zero locus of a quadratic polynomial in λ , and hence an algebraic variety of dimension 2. (The prototypical regulus is the set $\{(x, y, xy, 0) \in F^4 : x, y \in F\}$, where the lines l_i are of the form $\{(x, y_i, xy_i, 0) : x \in F\}$ for some distinct $y_1, y_2, y_3 \in F$. All reguli can be shown to be projectively equivalent to this example.) Since the lines in a frame are mutually skew, we see that this quadratic polynomial is irreducible (so the regulus is not a (double) plane, or the union of two planes), and that the lines $L(f)$ have cardinality $\sim |F|$ and are finitely overlapping.

Corollary 4.5 (Three-regulus lemma). *Let $L \subseteq \text{Gr}(F^4, 1)$ be a collection of lines in F^4 which point in different directions, and let $P \subseteq F^4$ be a set of points containing all the lines in L . Let f_1, f_2, f_3 be three frames such that the 3-spaces $\lambda(f_1), \lambda(f_2), \lambda(f_3)$ are parallel and disjoint. Then*

$$|\{l \in L : l \cap r(f_i) \neq \emptyset \text{ for all } i = 1, 2, 3\}| \lesssim |F|^2 + |F|^{-1}|P|.$$

Again, this bound improves by roughly $|F|$ over the trivial bound of $|F|^3$, if $|P|$ is not much larger than $|F|^3$. The hypothesis that the 3-spaces $\lambda(f_i)$ are parallel can be substantially relaxed, but we will not need to do so here.

Proof. Fix f_1, f_2, f_3 , and let $W \subseteq \text{Gr}(F^4, 1)$ denote the set

$$(4-1) \quad W := \{l \in \text{Gr}(F^4, 1) : l \cap r(f_i) \neq \emptyset \text{ for all } i = 1, 2, 3\}.$$

Since the $r(f_i)$ and l are algebraic varieties, it is clear (e.g. by using resultants; see e.g. [Harris 1992]) that the relationship $l \cap r(f_i) \neq \emptyset$ is equivalent to some finite set of explicit algebraic relations between the defining parameters of l and f_i . Thus W is an algebraic variety in $\text{Gr}(F^4, 1)$. In light of Corollary 4.3, it will suffice to verify that W has dimension at most 3. (We apologize to algebraic geometry sophisticates for the appalling crudeness of the following argument.)

Let p be a point in $r(f_1)$. Let ϕ_p be the stereographic projection from $\lambda(f_2)$ to $\lambda(f_3)$, thus $\phi_p(x) = y$ if and only if p, x, y are collinear. W is isomorphic to

$$\{(p, y) \in r(f_1) \times \lambda(f_3) : y \in r(f_3) \cap \phi_p(r(f_2))\}$$

(basically because two points determine a line, and because the planes $\lambda(f_i)$ are disjoint). In other words, one can think of W as a bundle over $r(f_1)$ whose fiber at p is $r(f_3) \cap \phi_p(r(f_2))$.

Note that $\phi_p : \lambda(f_2) \rightarrow \lambda(f_3)$ is an invertible linear map, so that $\phi_p(r(f_2))$ is an irreducible quadric surface in $\lambda(f_3)$. The set $r(f_3) \cap \phi_p(r(f_2))$ thus has dimension at most 2, and in fact will have dimension at most 1 unless $\phi_p(r(f_2)) \equiv r(f_3)$ (by irreducibility). However, as p varies, the quadric surfaces $\phi_p(r(f_2))$ move by translation. Since $r(f_2)$ is not a plane, we thus see that there can be at most a one-dimensional family of points p for which $\phi_p(r(f_2)) \equiv r(f_3)$.

To summarize, as p varies over the two-dimensional variety $r(f_1)$, the fiber $r(f_3) \cap \phi_p(r(f_2))$ is at most one-dimensional, except possibly for a one-dimensional family of points p for which the fiber is two-dimensional. From this it is clear that W has dimension at most 3, and we are done. (To compute the dimension properly one should work in the algebraically closed field \bar{F} here. But this causes no difficulty, since the above geometric considerations are valid for all fields of characteristic larger than two.) \square

Corollary 4.5 is the analogue of the three lines lemma used in [Schlag 1998]. Our strategy will now be to start with a Besicovitch set and construct many frames f and many lines $l \in L$ so that $r(f)$ intersects L , in order to exploit the above Corollary. To do this we shall need some basic combinatorial tools, which we now pause to review.

5. Some basic combinatorics

We shall frequently use the following elementary observation: If B is a finite set and $\mu : B \rightarrow \mathbb{R}^+$ is a function such that

$$\sum_{b \in B} \mu(b) \geq X,$$

then

$$\sum_{b \in B: \mu(b) \geq X/2|B|} \mu(b) \geq X/2.$$

We refer to this as a *popularity argument*, since we are restricting B to the values b which are popular in the sense that μ is large. The argument will be used iteratively many times.

We shall frequently use a version of the Cauchy–Schwarz and Hölder inequalities:

Lemma 5.1. *Let A, B be finite sets, and let \sim be a relation connecting pairs $(a, b) \in A \times B$ such that*

$$|\{(a, b) \in A \times B : a \sim b\}| \gtrsim X$$

for some $X \gg |B|$. Then

$$|\{(a, a', b) \in A \times A \times B : a \neq a'; a, a' \sim b\}| \gtrsim \frac{X^2}{|B|}$$

and

$$|\{(a, a', a'', b) \in A \times A \times A \times B : a, a', a'' \text{ distinct}; a, a', a'' \sim b\}| \gtrsim \frac{X^3}{|B|^2}$$

Proof. Define for each $b \in B$, define $\mu(b) := |\{a \in A : a \sim b\}|$. By hypothesis, we have

$$\sum_{b \in B} \mu(b) \gtrsim X.$$

In particular, by the popularity argument we have

$$\sum_{b \in B: \mu(b) \gtrsim X/|B|} \mu(b) \gtrsim X.$$

By hypothesis, we have $X/|B| \gg 1$. From this and the foregoing, we obtain

$$\sum_{b \in B: \mu(b) \gtrsim X/|B|} \mu(b)(\mu(b) - 1) \gtrsim X(X/|B|)$$

and

$$\sum_{b \in B: \mu(b) \gtrsim X/|B|} \mu(b)(\mu(b) - 1)(\mu(b) - 2) \gtrsim X(X/|B|)(X/|B|).$$

The claims follow. \square

A typical application of Lemma 5.1 is the following standard incidence bound:

Corollary 5.2. *For an arbitrarily collection $P \subseteq F^n$ of points and $L \subseteq \text{Gr}(F^n, 1)$ of lines, we have*

$$(5-1) \quad |\{(p, l) \in P \times L : p \in l\}| \lesssim |P|^{1/2}|L| + |P|$$

Proof. We may of course assume that the left-hand side of (5-1) is $\gg |P|$, since the claim is trivial otherwise. From Lemma 5.1 we have

$$|\{(p, l, l') \in P \times L \times L : p \in l \cap l'; l \neq l'\}| \gtrsim |P|^{-1} |\{(p, l) \in P \times L : p \in l\}|^2.$$

On the other hand, $|l \cap l'|$ has cardinality $O(1)$ if $l \neq l'$, thus

$$|\{(p, l, l') \in P \times L \times L : p \in l \cap l'; l \neq l'\}| \lesssim |L|^2.$$

Combining the two estimates we obtain the result. \square

The preceding estimate will be most useful when $|L|$ is small — in particular if $|L| = O(|F|)$. When $|L|$ is large, we have an alternate estimate:

Proposition 5.3 [Mockenhaupt and Tao 2004]. *Let the notation be as in Corollary 5.2. If we further assume that the lines in L point in different directions, then*

$$(5-2) \quad |\{(p, l) \in P \times L : p \in l\}| \lesssim |P|^{1/2}|L|^{3/4}|F|^{1/4} + |P| + |L|.$$

Proof. A proof is given after [Mockenhaupt and Tao 2004, Proposition 8.6]; the argument there is essentially due to Nets Katz, but the original result of this type dates back to Wolff [1995; 1998b].

For the convenience of the reader we now sketch an informal “probabilistic” derivation of (5-2). Let I denote the set in (5-2). We may assume that $|I| \gg |P|, |L|$ since the claim is trivial otherwise.

Observe that a randomly chosen point $p \in P$ and a randomly chosen line $l \in L$ have a probability $|I|/|P||L|$ of being incident (so that $p \in l$). Thus, given two random lines $l_1, l_2 \in L$ and a random point $p \in P$, we expect² the chance that p is incident to both l_1 and l_2 is $(|I|/|P||L|)^2$. Since there are $|P|$ possible values for p , the chance that two random lines $l_1, l_2 \in L$ intersect at all is thus heuristically $|P|(|I|/|P||L|)^2$.

As a consequence, the probability that three random lines $l_1, l_2, l_3 \in L$ form a triangle is heuristically $(|P|(|I|/|P||L|)^2)^3$. (There is the chance that this triangle

²This of course assumes independence of various random events, which is usually not the case. To make the argument rigorous one must use such tools as Lemma 5.1, which can be viewed as a statement that certain events are positively correlated. See [Mockenhaupt and Tao 2004, Proposition 8.6] for details.

is degenerate, but the hypothesis $|I| \gg |P|, |L|$ can be used to show that the probability of this occurring is low). On the other hand, given two intersecting lines $l_1, l_2 \in L$, there are at most $O(|F|)$ lines $l_3 \in L$ which can intersect them both, since we may apply the Wolff axiom to the 2-plane spanned by l_1 and l_2 . Combining these estimates we obtain

$$\left(|P| \left(\frac{|I|}{|P||L|}\right)^2\right)^3 \lesssim |P| \left(\frac{|I|}{|P||L|}\right)^2 \frac{|F|}{|L|}$$

and (5-2) follows.³ □

Remark. One only requires the Wolff axiom on L to obtain (5-2). In particular one can easily obtain Theorem 1.2 as a consequence of (5-2). It is likely that one can generalize Theorem 1.4 to obtain a further improvement to (5-2), but we do not pursue this question here.

6. A heuristic proof of Theorem 1.4

We now give a heuristic explanation for why we can improve upon Theorem 1.2 in four dimensions, in the spirit of the probabilistic arguments in Proposition 5.3. In later sections we shall make this heuristic argument rigorous.

Suppose for contradiction we have a family $L \subseteq \text{Gr}(F^4, 1)$ of lines in different directions of cardinality $|L| \sim |F|^3$ which are contained in a set $P \subseteq F^4$, also of cardinality $|P| \sim |F|^3$. Arguing as in Proposition 5.3 we see that any two lines in L have a (heuristic) probability $\sim 1/|F|$ of intersecting.

Also, a random line $l \in L$ and a random 3-space $\lambda \in \text{Gr}(F^4, 3)$ have a probability $1/|F|^2$ of being incident (so that $l \subseteq \lambda$). Thus we expect a 3-space λ to contain $|L|/|F|^2 \sim |F|$ lines in L .

Now consider the set of all quintuples $(l_1, l_2, l^1, l^2, l^3) \in L^5$ of lines such that l_i intersects l^j for all $i = 1, 2, j = 1, 2, 3$. From the above heuristics we see that there should be about $|L|^5 (1/|F|)^6 \sim |F|^9$ such quintuples. On the other hand, for generic quintuples $(l_1, l_2, l^1, l^2, l^3)$ of the above form, the lines l^1, l^2, l^3 must lie in a 3-space λ , and l_1, l_2 must lie in the regulus generated by the frame (l^1, l^2, l^3, λ) . (For this heuristic argument we ignore the possibility that the quintuple could degenerate).

To count the number of possible reguli, observe that there are $|L|^2 \sim |F|^6$ choices for l^1, l^2 , which determines λ . From our previous heuristic we see that λ can contain at most $O(|F|)$ choices for l^3 , thus there are at most $O(|F|^7)$ reguli.

³It is an instructive exercise to obtain similar heuristic probabilistic derivations of such estimates as (5-1) (using the fact that two random lines intersect in at most one point) or (1-2) (using the fact that a random regulus contains at most $O(|F|)$ lines). See also Section 6 below.

Dividing $|F|^9$ by $|F|^7$, we thus see that a generic regulus $r(f)$ of the above type must contain at least $|F|^2$ pairs (l_1, l_2) of lines in L . But $r(f)$ only has $O(|F|)$ lines to begin with. Thus a generic regulus $r(f)$ must have extremely large intersection with P , so that $|r(f) \cap P| \sim |r(f)| \sim |F|^2$.

Since a random $p \in P$ and $l \in L$ have a probability $1/|F|^2$ of being incident, this means that a random line $l \in L$ and a random regulus $r(f)$ have a probability ~ 1 of intersecting. In particular, if we select three parallel reguli $r(f_1), r(f_2), r(f_3)$, a large fraction of lines in L must be incident to all three reguli. But this contradicts Corollary 4.5, since $|L| \sim |F|^3$ and $|P| \sim |F|^3$.

7. Preliminary refinements

We now begin the rigorous proof of Theorem 1.4, which will broadly follow the heuristic outline of the previous section.

Let $P_0 \subseteq F^4$ be a Besicovitch set. We may assume that

$$(7-1) \quad |P_0| \lesssim |F|^{3+\frac{1}{16}}.$$

since the claim is trivial otherwise. We may also assume that $|F| \gg 1$ for similar reasons.

Since P_0 is a Besicovitch set, there exists a set $L_0 \subseteq \text{Gr}(F^4, 1)$ of lines in different directions such that $|L_0| \sim |F|^3$ and P_0 contains every line in L_0 . In particular the incidence set

$$I_0 := \{(p, l) \in P_0 \times L_0 : p \in l\}$$

has cardinality $|I_0| = |F||L_0| \sim |F|^4$.

Given any line l in L_0 and a randomly selected 3-space λ in $\text{Gr}(F^4, 3)$, the probability that l lies in λ is $\sim 1/|F|^2$. Since $|L_0| \sim |F|^3$, one thus expects every 3-space λ contains about $|F|$ lines in L_0 on the average. A similar heuristic leads us to expect every 2-plane $\pi \in \text{Gr}(F^4, 2)$ to contain at most $O(1)$ lines on the average.

Although these statements need not be true for all 3-spaces λ , certain variants do hold if we refine L_0 and P_0 slightly, as stated in the next result. We write $A \approx B$ to mean $A \gtrsim B$ and $A \lesssim B$.

Proposition 7.1. *There exists a quantity*

$$(7-2) \quad 1 \lesssim \alpha \lesssim N^{\frac{1}{16}},$$

a subset⁴ P_1 of P_0 and a subset L_1 of L_0 such that the following properties hold.

⁴In the course of this argument we shall need to refine the set P_0 to a slightly smaller set P_1 , and then further to P_2 , and similarly refine L_0 to L_1 and then L_2 , while also refining some auxiliary sets H_0 to H_1 , and \mathcal{F}_0 to \mathcal{F}_1 to \mathcal{F}_2 to \mathcal{F}_3 . These refinements are largely technical and as a first

Many incidences: We have the incidence bound

$$(7-3) \quad |\{(p, l) \in P_1 \times L_1 : p \in l\}| \gtrsim |F|^4.$$

Cardinality and multiplicity bounds: We have the cardinality bound

$$(7-4) \quad |P_1| \lesssim \alpha |F|^3$$

and the multiplicity bound

$$(7-5) \quad |\{l \in L_1 : p \in l\}| \approx \alpha^{-1} |F|$$

for all $p \in P_1$.

No 3-space degeneracy: For any 3-space $\lambda \in \text{Gr}(F^4, 3)$, we have

$$(7-6) \quad |\{l \in L_1 : l \subset \lambda\}| \lesssim \alpha^2 |F|.$$

No 2-plane degeneracy: For any 2-plane $\pi \in \text{Gr}(F^4, 2)$, we have

$$(7-7) \quad |\{l \in L_1 : l \subset \pi\}| \lesssim \alpha^4.$$

The quantity α measures the improvement over Wolff’s bound $|P_0| \gtrsim |F|^3$. As one can see from (7-2), it is rather close to 1.

Proof. We follow standard “iterated refinement” arguments (see [Wolff 1998a; Łaba and Tao 2001b; Christ 1998; Tao and Wright 2003]; our argument here is particularly close to that in [Łaba and Tao 2001b]). The purpose of the iteration is mainly to obtain the property (7-7).

Define the multiplicity function μ_0 on P_0 by

$$\mu_0(p) := |\{l \in L_0 : p \in l\}|.$$

Then we have

$$\sum_{p \in P_0} \mu_0(p) = |I_0|.$$

If we divide $\mu_0(p)$ into dyadic “pigeonholes” and apply the dyadic pigeonhole principle (observing that $\log |F| \approx 1$), we conclude that there exists a multiplicity $\alpha^{-1} |F|$ such that

$$\sum_{p \in P_0: \mu_0(p) \sim \alpha^{-1} |F|} \mu_0(p) \approx |I_0| \approx |F|^4.$$

Fix this α , and define

$$P'_0 := \{p \in P_0 : \mu_0(p) \sim \alpha^{-1} |F|\}$$

approximation one can view these sets as being essentially the same (although the sets $\mathcal{F}_2, \mathcal{F}_3$ are significantly smaller than $\mathcal{F}_0, \mathcal{F}_1$).

and

$$I'_0 := \{(p, l) \in P'_0 \times L_0 : p \in l\} \subseteq I_0.$$

Then by construction, $|I'_0| \gtrsim |I_0| \sim |F|^4$, and

$$|P'_0| \approx |I_0|/(\alpha^{-1}|F|) \approx \alpha|F|^3.$$

By (7-1) we thus have $\alpha \lesssim N^{\frac{1}{16}}$. To get the other half of (7-2), we observe from Proposition 5.3 that

$$|I'_0| \lesssim |P'_0|^{1/2}|L_0|^{3/4}|F|^{1/4} + |P'_0| + |L_0|;$$

applying the above estimates, we thus obtain $\alpha \gtrsim 1$. Thus (7-2) holds.

Set $N := \log \log |F|$; the point of this choice of N is that both $|F|^{C/N}$ and C^N are ≈ 1 for any fixed choice of constant C . We shall inductively construct sets

$$(7-8) \quad P'_0 =: P^{(0)} \supset P^{(1)} \supset \dots \supset P^{(N)}$$

and

$$(7-9) \quad L_0 =: L^{(0)} \supset L^{(1)} \supset \dots \supset L^{(N)}$$

as follows.⁵

As indicated above, we set $P^{(0)} := P'_0$ and $L^{(0)} := L_0$. Now suppose inductively that $P^{(k)}$ and $L^{(k)}$ have already been constructed for some $0 \leq k < N$. We define the incidence set

$$I^{(k)} := \{(p, l) \in P^{(k)} \times L^{(k)} : p \in l\}.$$

Clearly we have

$$\sum_{l \in L^{(k)}} |l \cap P^{(k)}| = |I^{(k)}|.$$

Thus if we set

$$L^{(k+1)} := \left\{ l \in L^{(k)} : |l \cap P^{(k)}| \geq \frac{|I^{(k)}|}{2|L^{(k)}|} \right\}$$

then by the popularity argument we have

$$\sum_{l \in L^{(k+1)}} |l \cap P^{(k)}| \geq |I^{(k)}|/2.$$

We rewrite this as

$$\sum_{p \in P^{(k)}} |\{l \in L^{(k+1)} : p \in l\}| \geq |I^{(k)}|/2.$$

⁵The use of such a large number of refinements is of course overkill (one could probably get away with $N = 5$, in fact), but reducing the number of refinements used does not alter the exponent $\frac{1}{16}$, since $F^{C/N}$ and C^N were ≈ 1 anyway.

Thus if we set

$$P^{(k+1)} := \left\{ p \in P^{(k)} : |\{l \in L^{(k+1)} : p \in l\}| \geq \frac{|I^{(k)}|}{4|P^{(k)}|} \right\},$$

we get, again by the popularity argument,

$$\sum_{p \in P^{(k+1)}} |\{l \in L^{(k+1)} : p \in l\}| \geq |I^{(k)}|/4$$

or in other words

$$|I^{(k+1)}| \geq |I^{(k)}|/4.$$

We repeat this construction for $k = 0, 1, \dots, N - 1$, creating a nested sequence of sets of points (7-8) and sets of lines (7-9). By construction and the fact that $4^{-N} \approx 1$, we clearly have

$$|I^{(k)}| \approx |I^{(0)}| = |I'_0| \gtrsim |F|^4$$

for all k . Furthermore,

$$|P^{(k)}| \leq |P'_0| \lesssim \alpha |F|^3 \quad \text{and} \quad |L^{(k)}| \leq |L_0| \lesssim |F|^3.$$

Thus, setting $P_1 := P^{(N)}$ and $L_1 := L^{(N-1)}$, we see that (7-3), (7-4), and (7-5) hold. (To get the upper bound in (7-5), simply bound the left-hand side by $\mu_0(p)$.)

It remains only to verify the nondegeneracy conditions (7-6), (7-7).

We first verify (7-6). Let λ be a 3-space. Since λ is clearly an algebraic variety of dimension 3, we can invoke Corollary 4.2 and conclude that

$$|\{(p, l) \in \lambda \times L^{(N-2)} : p \in l\}| \lesssim |F|^3.$$

From the construction of $P^{(N-1)}$ we thus have

$$|P(\lambda)| \lesssim |F|^2 \alpha$$

where $P(\lambda) := \lambda \cap P^{(N-1)}$.

Let $L(\lambda)$ denote those lines in L_1 which lie in λ . By the construction of L_1 we have

$$|\{(p, l) \in P(\lambda) \times L(\lambda) : p \in l\}| \gtrsim |F| |L(\lambda)|.$$

On the other hand, from Proposition 5.3 we have

$$|\{(p, l) \in P(\lambda) \times L(\lambda) : p \in l\}| \lesssim |P(\lambda)|^{1/2} |L(\lambda)|^{3/4} |F|^{1/4} + |P(\lambda)| + |L(\lambda)|.$$

Combining all three estimates and using (7-2) we obtain

$$|L(\lambda)| \lesssim \alpha^2 |F|$$

which is (7-6).

In fact, this argument gives (7-6) if L_1 is replaced by $L^{(k)}$ for any $1 \leq k \leq N - 1$.

We now prove (7–7). Following [Wolff 1998a; Łaba and Tao 2001b], we define the *plate number* \mathbf{p}_k for $0 \leq k \leq N - 1$ to be the quantity

$$\mathbf{p}_k := \sup_{\pi \in \text{Gr}(F^4, 2)} |\{l \in L_k : l \subset \pi\}|.$$

We observe the bounds

$$(7-10) \quad 1 \leq \mathbf{p}_k \lesssim |F|;$$

the former bound comes since L_k is nonempty, while the latter bound comes since a 2-plane can contain at most $O(|F|)$ lines in different directions.

Clearly the plate numbers are nonincreasing in k . From this, (7–10), the pigeon-hole principle and the fact that $|F|^{1/N} \approx 1$, we can find $2 \leq k \leq N - 1$ such that

$$(7-11) \quad \mathbf{p}_{k-1} \approx \mathbf{p}_k.$$

Fix this k . We can find a 2-plane $\pi \in \text{Gr}(F^4, 2)$ such that the set

$$L_k(\pi) := \{l \in L_k : l \subset \pi\}$$

has cardinality \mathbf{p}_k .

Fix π , and let $P_k(\pi)$ denote the set

$$P_k(\pi) := P_k \cap \pi.$$

By the construction of L_k , every line in L_k contains $\gtrsim |F|$ points in P_k , thus every line in $L_k(\pi)$ contains $\gtrsim |F|$ points in $P_k(\pi)$. In particular we see that

$$|\{(p, l) \in P_k(\pi) \times L_k(\pi) : p \in l\}| \gtrsim |F| \mathbf{p}_k.$$

Applying Corollary 5.2 we conclude that

$$|F| \mathbf{p}_k \lesssim |P_k(\pi)|^{1/2} \mathbf{p}_k + |P_k(\pi)|;$$

from this and (7–10) we thus have

$$(7-12) \quad |P_k(\pi)| \gtrsim |F| \mathbf{p}_k.$$

Let $P'_k(\pi)$ denote the space of all points p in $P_k(\pi)$ such that at least half of all the lines in $\{l \in L_{k-1} : p \in l\}$ are contained in $L_{k-1}(\pi)$. We have two cases.

Case 1 (parallel case): $|P'_k(\pi)| \geq \frac{1}{2} |P_k(\pi)|$. In this case we have

$$\begin{aligned} |\{(p, l) \in P_k(\pi) \times L_{k-1}(\pi) : p \in l\}| &\geq |\{(p, l) \in P'_k(\pi) \times L_{k-1}(\pi) : p \in l\}| \\ &\geq \frac{1}{2} |\{(p, l) \in P'_k(\pi) \times L_{k-1} : p \in l\}| \\ &\gtrsim |P'_k(\pi)| \alpha^{-1} |F| \\ &\gtrsim |P_k(\pi)| \alpha^{-1} |F|, \end{aligned}$$

while by definition of p_{k-1} we have

$$|L_{k-1}(\pi)| \leq p_{k-1}.$$

Applying Corollary 5.2 we thus see that

$$|P_k(\pi)|\alpha^{-1}|F| \lesssim |P_k(\pi)|^{1/2}p_{k-1} + |P_k(\pi)|,$$

which by (7-2) implies that

$$|P_k(\pi)||F|^2 \lesssim \alpha^2 p_{k-1}^2.$$

But combining this with (7-11), (7-12) we obtain

$$p_k \gtrsim \alpha^{-2}|F|^3.$$

But this contradicts (7-10) by (7-2). Hence this case cannot occur.

Case 2 (transverse case): $|P'_k(\pi)| \leq \frac{1}{2}|P_k(\pi)|$. In this case we have (by a computation similar to Case 1)

$$|\{(p, l) \in P_k(\pi) \times L_k : p \in l; l \not\subset \pi\}| \gtrsim \alpha^{-1}|P_k(\pi)||F|.$$

Thus, if L_{k-1}^* denotes the lines $l \in L_{k-1}$ which are incident to a point in $P_k(\pi)$ but are not contained in π , then we have

$$(7-13) \quad |L_{k-1}^*| \gtrsim \alpha^{-1}|P_k(\pi)||F| \gtrsim \alpha^{-1}|F|^2 p_k$$

by (7-12).

We now use Wolff’s hairbrush argument [Wolff 1995], [Wolff 1998b], as modified to deal with plates in [Wolff 1998a], [Łaba and Tao 2001b]. We can foliate L_{k-1}^* as the disjoint union of

$$L_{k-1}^*(\lambda) := \{l \in L_{k-1}^* : l \in \lambda\}$$

where λ ranges over the 3-spaces containing π . For each such λ , observe from the analogue of (7-6) for $L^{(k-1)}$ that

$$(7-14) \quad |L_{k-1}^*(\lambda)| \lesssim \alpha^2|F|.$$

Also, if we define

$$P_{k-1}^*(\lambda) := \{p \in P_{k-1} : p \in \lambda \setminus \pi\}$$

then by the construction of L_{k-1} , we have

$$|\{(p, l) \in P_{k-1}^*(\lambda) \times L_{k-1}^*(\lambda) : p \in l\}| \gtrsim |L_{k-1}^*(\lambda)||F|.$$

Applying Corollary 5.2 we obtain

$$|L_{k-1}^*(\lambda)||F| \lesssim |P_{k-1}^*(\lambda)|^{1/2}|L_{k-1}^*(\lambda)| + |P_{k-1}^*(\lambda)|,$$

which by (7-14), (7-2) implies that

$$|P_{k-1}^*(\lambda)| \gtrsim \alpha^{-2} |L_{k-1}^*(\lambda)| |F|.$$

Summing in λ , we obtain

$$|P_{k-1}| \gtrsim \alpha^{-2} |L_{k-1}^*| |F| \gtrsim \alpha^{-3} |F|^3 \mathbf{p}_k.$$

Since $|P_{k-1}| \lesssim \alpha |F|^3$ by construction, we obtain $\mathbf{p}_k \lesssim \alpha^4$, and the claim follows. \square

8. Construction of reguli

We now continue the proof of Theorem 1.4. We begin by refining P_1 and L_1 a little further. By (7-3) we have

$$\sum_{l \in L_1} |l \cap P_1| \approx |F|^4.$$

Thus if we set

$$L_2 := \{l \in L_1 : |l \cap P_1| \approx |F|\}$$

then by the popularity argument we get

$$\sum_{l \in L_2} |l \cap P_1| \approx |F|^4$$

or equivalently

$$\sum_{p \in P_1} |\{l \in L_2 : p \in l\}| \approx |F|^4.$$

Thus if we set

$$P_2 := \{p \in P_1 : |\{l \in L_2 : p \in l\}| \approx \alpha^{-1} |F|\}$$

then by (7-4), (7-5), and the popularity argument we have

$$(8-1) \quad \sum_{p \in P_2} |\{l \in L_2 : p \in l\}| \approx |F|^4.$$

In particular,

$$(8-2) \quad |P_2| \approx \alpha |F|^3.$$

The next task is to generate a large number of frames, and a large number of lines in L incident to the reguli generated by these frames. As a frame is a fairly complicated combinatorial object (consisting of three lines and a 3-space), we will first begin by counting some simpler objects which eventually will be combined together to form frames.

By (8–1) we have

$$|\{(p, l) \in P_2 \times L_2 : p \in l\}| \approx |F|^4.$$

Since $|L_2| \lesssim |F|^3$, we thus see from Lemma 5.1 that

$$|\{(p_1, p_2, l) \in P_2 \times P_2 \times L_2 : p_1, p_2 \in l; p_1 \neq p_2\}| \gtrsim |F|^8/|L_2| \approx |F|^5.$$

By the definition of P_2 , we see that for each (p_1, p_2, l) as above there are $\gtrsim \alpha^{-1}|F|$ lines $l_1 \in L_2$ that contain p_1 but are distinct from l , and similarly there are $\gtrsim \alpha^{-1}|F|$ lines $l_2 \in L_2$ containing p_2 but distinct from l . Thus

$$|H_0| \gtrsim \alpha^{-2}|F|^7,$$

where

$$H_0 := \{(p_1, p_2, l, l_1, l_2) \in P_2 \times P_2 \times L_2 \times L_2 \times L_2 : \\ p_1, p_2 \in l; p_1 \neq p_2; p_1 \in l_1; p_2 \in l_2; l \neq l_1, l_2\}$$

is the space of ‘‘H-shaped’’ objects.

Let $H_1 \subseteq H_0$ be the set of elements (p_1, p_2, l, l_1, l_2) in H_0 such that l_1 and l_2 are skew. We claim that

$$|H_0 \setminus H_1| \lesssim \alpha^{-1}|F|^6.$$

Indeed, to choose an element (p_1, p_2, l, l_1, l_2) in $H_0 \setminus H_1$ (which is a degenerate H, i.e. a triangle), we first choose $p_1 \in P_2$ (of which there are $\lesssim \alpha|F|^3$ choices), and then choose the distinct lines l, l_1 incident to p_1 (of which there are $\lesssim (\alpha^{-1}|F|)^2$ choices). Since l_2 must lie in the 2-plane generated by l and l_1 , and the lines of L_1 point in different directions, there are only $O(|F|)$ choices for l_2 . Since p_2 is uniquely determined as $p_2 = l \cap l_2$, the claim follows.

From the bounds above and (7–2) we see that

$$(8–3) \quad |H_1| \gtrsim \alpha^{-2}|F|^7.$$

By construction, if $h = (p_1, p_2, l, l_1, l_2) \in H_1$, then l_1 and l_2 are skew. Thus l_1 and l_2 lie in a unique 3-space $\lambda(h)$, which then must also contain p_1, p_2, l .

Let $S_0 \subset L_2 \times L_2$ denote the pairs (l_1, l_2) of skew lines in L_2 . For each pair $(l_1, l_2) \in S_0$, we define the *connecting set* $C(l_1, l_2) \subset L_2$ to be the set of all lines $l \in L_2$ which are distinct from l_1, l_2 , but intersect both l_1, l_2 in points $p_1 \in P_2$ and $p_2 \in P_2$ respectively. Observe the identity

$$\sum_{(l_1, l_2) \in S_0} |C(l_1, l_2)| = |H_1|.$$

Since $|S_0| \leq |L_2|^2 \lesssim |F|^6$, we thus see from (8–3) that if we define

$$S_1 := \{(l_1, l_2) \in S_0 : |C(l_1, l_2)| \gtrsim \alpha^{-2}|F|\},$$

the popularity argument yields

$$(8-4) \quad \sum_{(l_1, l_2) \in S_1} |C(l_1, l_2)| \gtrsim \alpha^{-2} |F|^7.$$

If $(l_1, l_2) \in S_1$, we define the set $C^{(3)}(l_1, l_2) \subseteq C(l_1, l_2)^3$ to be the space of all triplets $(l^1, l^2, l^3) \in C(l_1, l_2)^3$ such that the six points $l^i \cap l_j$ for $i = 1, 2, 3, j = 1, 2$ are all disjoint.

We now use the nondegeneracy property (7-7) to obtain a lower bound for the size of $C^{(3)}(l_1, l_2)$.

Lemma 8.1 (Many triple connections between skew lines). *For any $(l_1, l_2) \in S_1$, we have $|C^{(3)}(l_1, l_2)| \gtrsim \alpha^{-4} |F|^2 |C(l_1, l_2)|$.*

Proof. Fix l_1, l_2 . We choose $l^1 \in C(l_1, l_2)$ arbitrarily; of course, there are $|C(l_1, l_2)|$ choices for l^1 .

Fix l^1 . From (7-7) we have

$$|\{l^2 \in C(l_1, l_2) : l^2 \cap l_1 = l^1 \cap l_1\}| \lesssim \alpha^4$$

(since such lines lie in the 2-plane spanned by $l^1 \cap l_1$ and l_2 . Similarly if the roles of l_1 and l_2 are interchanged. Since

$$|C(l_1, l_2)| \gtrsim \alpha^{-2} |F|,$$

we thus see from (7-2) that there are $\gtrsim \alpha^{-2} |F|$ choices for l^2 such that

$$l^2 \cap l_j \neq l^1 \cap l_j \quad \text{for } j = 1, 2.$$

Fix l^2 . Arguing as above we see that there are $\gtrsim \alpha^{-2} |F|$ choices for l^3 such that $l^3 \cap l_j \neq l^i \cap l_j$ for $i = 1, 2$ and $j = 1, 2$. The claim follows. \square

From this lemma and (8-4) we see that

$$\sum_{(l_1, l_2) \in S_1} |C^{(3)}(l_1, l_2)| \gtrsim \alpha^{-6} |F|^9.$$

Observe that if $(l_1, l_2) \in S_1$ and $(l^1, l^2, l^3) \in C^{(3)}(l_1, l_2)$, the various incidence assumptions in the definition of S_1 and $C^{(3)}(l_1, l_2)$ force $f := (l^1, l^2, l^3, \lambda)$ to be a frame, where λ is the unique 3-space spanned by l^1 and l^2 . Observe that l_1, l_2 both lie in $L_2 \cap L(f)$. Thus, if \mathcal{F}_0 denotes the space of all frames generated in this manner, then

$$(8-5) \quad \sum_{f \in \mathcal{F}_0} |L_2 \cap L(f)|^2 \gtrsim \alpha^{-6} |F|^9.$$

Let $f \in \mathcal{F}_0$. Since the lines in $L(f)$ are contained in a regulus, they have finite overlap. Since each line in L_2 contains $\approx |F|$ points in P_1 by construction, we thus

see that⁶

$$|P_1 \cap r(f)| \gtrsim |F| |L_2 \cap L(f)|$$

so by (8-5) we have

$$\sum_{f \in \mathcal{F}_0} |P_1 \cap r(f)|^2 \gtrsim \alpha^{-6} |F|^{11}.$$

By (7-5), each point in P_1 is incident to $\approx \alpha^{-1} |F|$ lines in L_1 . Thus we have

$$\sum_{f \in \mathcal{F}_0} |\{l \in L_1 : l \cap r(f) \cap P_1 \neq \emptyset\}|^2 \gtrsim \alpha^{-8} |F|^{13}.$$

We observe the cardinality bound

$$(8-6) \quad |\mathcal{F}_0| \lesssim \alpha^2 |F|^7.$$

Indeed, to choose a frame (l^1, l^2, l^3, λ) in \mathcal{F}_0 , we observe that there are

$$O(|L_1|^2) = O(|F|^6)$$

choices for the skew pair (l^1, l^2) . This determines λ , and then by (7-6) we thus see that there are $O(\alpha^2 |F|)$ choices for l^3 , and (8-6) follows. In particular, if we define

$$(8-7) \quad \mathcal{F}_1 := \{f \in \mathcal{F}_0 : |\{l \in L_1 : l \cap r(f) \cap P_1 \neq \emptyset\}| \gtrsim \alpha^{-5} |F|^3\}$$

then by the popularity argument

$$(8-8) \quad \sum_{f \in \mathcal{F}_1} |\{l \in L_1 : l \cap r(f) \cap P_1 \neq \emptyset\}|^2 \gtrsim \alpha^{-8} |F|^{13}.$$

Since the summand on the left-hand side can be crudely bounded by $|L_1|^2 = O(|F|^6)$, we thus have the crude bound⁷

$$(8-9) \quad |\mathcal{F}_1| \gtrsim \alpha^{-8} |F|^7$$

(compare with (8-6)).

For any frame $f \in \mathcal{F}_1$, there are only $O(|F|^3)$ possible orientations for $\lambda(f)$. By (8-9) and the pigeonhole principle, there therefore exists a 3-space $\lambda_0 \in \text{Gr}(F^4, 3)$ such that

$$(8-10) \quad |\mathcal{F}_2| \gtrsim \alpha^{-8} |F|^4$$

⁶One could also obtain this bound using Corollary 5.2 and the crude bound

$$|L_2 \cap L(f)| \leq |L(f)| \lesssim |F|.$$

⁷The bounds on $|\mathcal{F}_1|$, and later on $|\mathcal{F}_2|, |\mathcal{F}_3|$, might not be best possible, however an improvement on this part of the argument does not directly improve the gain $\frac{1}{16}$.

where

$$\mathcal{F}_2 := \{f \in \mathcal{F}_1 : \lambda(f) \text{ is a translate of } \lambda_0\}.$$

Fix this λ_0 . Let \mathcal{F}_3 be a maximal subset of \mathcal{F}_2 such that the reguli $\{r(f) : f \in \mathcal{F}_3\}$ are all distinct. Since each $r(f)$ contains at most $O(|F|)$ lines, each regulus can arise from at most $O(|F|^3)$ frames. We thus see from (8–10) that

$$(8-11) \quad |\mathcal{F}_3| \gtrsim \alpha^{-8} |F|.$$

From (8–7) we have

$$\sum_{f \in \mathcal{F}_3} |\{l \in L_1 : l \cap r(f) \cap P_1 \neq \emptyset\}| \gtrsim \alpha^{-5} |F|^3 |\mathcal{F}_3|.$$

From (7–2), (8–11) the right-hand side is $\gg |F|^3 \gtrsim |L_1|$. Thus we can use Lemma 5.1, and obtain

$$\sum_{\substack{f_1, f_2, f_3 \in \mathcal{F}_3 \\ f_1, f_2, f_3 \text{ distinct}}} |\{l \in L_1 : l \cap r(f_i) \cap P_1 \neq \emptyset \text{ for } i = 1, 2, 3\}| \gtrsim \alpha^{-15} |F|^3 |\mathcal{F}_3|^3.$$

From the pigeonhole principle, we may thus find distinct frames f_1, f_2, f_3 in \mathcal{F}_3 such that

$$(8-12) \quad |L_*| \gtrsim \alpha^{-15} |F|^3,$$

where $L_* \subseteq L_1$ is the collection of lines

$$L_* := \{l \in L_1 : l \cap r(f_i) \cap P_1 \neq \emptyset \text{ for } i = 1, 2, 3\}.$$

Now we consider the problem of obtaining upper bounds on $|L_*|$. The crude upper bound of $|L_1| \sim |F|^3$ is clearly not enough to obtain a contradiction. However, thanks to the three-regulus lemma we can improve this bound by about $|F|$:

Proposition 8.2. *We have*

$$(8-13) \quad |L_*| \lesssim |F|^{2+\frac{1}{16}}.$$

Proof. If the 3-spaces $\lambda(f_1), \lambda(f_2), \lambda(f_3)$ are disjoint, this follows directly from Corollary 4.5 and (7–1).

By symmetry it remains to consider the case when $\lambda(f_1)$ and $\lambda(f_2)$ (for instance) are equal. Then the lines in L_* must either be parallel to $\lambda(f_1)$, or else intersect $P_1 \cap r(f_1) \cap r(f_2)$. There are at most $|F|^2$ lines in the first category (in fact there are far fewer, thanks to (7–6)). In the second category, we observe that $r(f_1) \cap r(f_2)$ is at most one-dimensional (since $r(f_1), r(f_2)$ are irreducible and distinct) and hence has cardinality $O(|F|)$. On the other hand, by (7–5) every point in $r(f_1) \cap r(f_2) \cap P_1$ is incident to $\approx \alpha^{-1} |F|$ lines in L_1 . Thus we certainly have $\lesssim |F|^{2+\frac{1}{16}}$ incidences in this case as well. \square

Combining (8–13) with (8–12) we obtain

$$\alpha \gtrsim |F|^{\frac{1}{16}}$$

and hence by (8–2)

$$|P_0| \gtrsim |P_2| \approx \alpha |F|^3 \gtrsim |F|^{3+\frac{1}{16}}$$

as desired. This concludes the proof of Theorem 1.4.

9. Remarks

It seems likely that this Theorem can be generalized in several ways. The exponent $\frac{1}{16}$ is probably not sharp, and also the result should have extension to other dimensions, perhaps through more sophisticated use of algebraic geometry. In dimensions 5 and higher there are other, more “arithmetic” arguments that give slight improvements to $|F|^{(n+2)/2}$ for Besicovitch sets; see [Bourgain 1999; Katz and Tao 1999; 2002a; 2002b; Mockenhaupt and Tao 2004; Rogers 2001]. Nevertheless, if one can make an improvement of the order of $\frac{1}{16}$ in, say, five dimensions by these “geometric” techniques, this will be quite competitive with the results in, say, [Katz and Tao 2002b]. In the Euclidean setting one can improve the bound $(n+2)/2$ in all dimensions $n \geq 3$ by a small number (10^{-10}) for the *upper Minkowski dimension* problem for Besicovitch sets [Katz et al. 2000; Łaba and Tao 2001a; 2001b], but this argument seems special to the upper Minkowski problem and does not directly impact the finite field question.

Also, the argument can probably be extended to obtain an estimate on the Kakeya maximal function for finite fields; see [Mockenhaupt and Tao 2004]. In principle, the finite field results should also extend to the Euclidean setting \mathbb{R}^n , but there are unpleasant technical difficulties in the process, due, for instance, to the presence of near-degenerate reguli in \mathbb{R}^n . Also in the finite field case one is aided considerably by the fact that dimensions must be integer; for instance, the intersection of two lines is either empty, 0-dimensional (a point), or 1-dimensional (a line). In the (δ -discretized) Euclidean case there is a continuum of cases: two distinct $1 \times \delta$ tubes can intersect in a set of length $\sim \delta/\theta$, where $\delta < \theta < 1$ is the angle between the two tubes. This introduces a new dyadic parameter θ into the analysis (measuring the degeneracy of the angle), and often the cases of small θ and large θ need to be treated separately. See [Wolff 1995], for example. Here we have more complicated algebraic objects, such as the variety (4–1), and to capture the possible degeneracies of this object seems to require a large number of additional dyadic parameters. It is possible that various rescaling arguments, such as the two-ends and bilinear reductions mentioned above, may be used to reduce the number of such parameters, but the extension of this argument to the Euclidean case still appears to be quite nontrivial.

Difficulties of these kinds cause considerable complication in such papers as [Schlag 1998], although some could perhaps be alleviated using the “two-ends” reduction in [Wolff 1995] and the “bilinear reduction” in [Tao et al. 1998].

Acknowledgments

The author thanks Tony Carbery, Nets Katz, Wilhelm Schlag, and Jim Wright for helpful discussions, and is also indebted to David Gieseke and Allen Knutson for their explanation of some of the basics of algebraic geometry. The author is particularly indebted to Nets Katz for emphasizing the importance of reguli to this problem. Finally, the author thanks the anonymous referee for a careful reading and the detection of several misprints.

References

- [Bourgain 1991] J. Bourgain, “Besicovitch type maximal operators and applications to Fourier analysis”, *Geom. Funct. Anal.* **1**:2 (1991), 147–187. MR 92g:42010 Zbl 0756.42014
- [Bourgain 1999] J. Bourgain, “On the dimension of Kakeya sets and related maximal inequalities”, *Geom. Funct. Anal.* **9**:2 (1999), 256–282. MR 2000b:42013 Zbl 0930.43005
- [Christ 1998] M. Christ, “Convolution, curvature, and combinatorics: a case study”, *Internat. Math. Res. Notices* **19** (1998), 1033–1048. MR 2000a:42026 Zbl 0927.42008
- [Gowers 1998] W. T. Gowers, “A new proof of Szemerédi’s theorem for arithmetic progressions of length four”, *Geom. Funct. Anal.* **8**:3 (1998), 529–551. MR 2000d:11019 Zbl 0907.11005
- [Harris 1992] J. Harris, *Algebraic geometry*, Graduate Texts in Mathematics **133**, Springer, New York, 1992. MR 93j:14001 Zbl 0779.14001
- [Katz and Tao 1999] N. H. Katz and T. Tao, “Bounds on arithmetic projections, and applications to the Kakeya conjecture”, *Math. Res. Lett.* **6**:6 (1999), 625–630. MR 2000m:28006 Zbl 0980.42013
- [Katz and Tao 2002a] N. Katz and T. Tao, “Recent progress on the Kakeya conjecture”, *Publ. Mat.* Vol. Extra (2002), 161–179. MR 2003m:42036 Zbl 1024.42010
- [Katz and Tao 2002b] N. H. Katz and T. Tao, “New bounds for Kakeya problems”, *J. Anal. Math.* **87** (2002), 231–263. MR 2003i:28006 Zbl 1027.42014
- [Katz et al. 2000] N. H. Katz, I. Łaba, and T. Tao, “An improved bound on the Minkowski dimension of Besicovitch sets in \mathbf{R}^3 ”, *Ann. of Math. (2)* **152**:2 (2000), 383–446. MR 2002i:28006 Zbl 0980.42014
- [Łaba and Tao 2001a] I. Łaba and T. Tao, “An improved bound for the Minkowski dimension of Besicovitch sets in medium dimension”, *Geom. Funct. Anal.* **11**:4 (2001), 773–806. MR 2003b:28006 Zbl 1005.42009
- [Łaba and Tao 2001b] I. Łaba and T. Tao, “An x-ray transform estimate in \mathbb{R}^n ”, *Rev. Mat. Iberoamericana* **17**:2 (2001), 375–407. MR 2003a:44003 Zbl 1024.44002
- [Mockenhaupt and Tao 2004] G. Mockenhaupt and T. Tao, “Restriction and Kakeya phenomena for finite fields”, *Duke Math. J.* **121**:1 (2004), 35–74. MR 2004m:11200 Zbl 02103577
- [Rogers 2001] K. M. Rogers, “The finite field Kakeya problem”, *Amer. Math. Monthly* **108**:8 (2001), 756–759. MR 2002g:11175 Zbl 1028.43007
- [Schlag 1998] W. Schlag, “A geometric inequality with applications to the Kakeya problem in three dimensions”, *Geom. Funct. Anal.* **8**:3 (1998), 606–625. MR 99g:42025 Zbl 0939.42012

- [Tao and Wright 2003] T. Tao and J. Wright, “ L^p improving bounds for averages along curves”, *J. Amer. Math. Soc.* **16**:3 (2003), 605–638. MR 2004j:42005 Zbl 01896622
- [Tao et al. 1998] T. Tao, A. Vargas, and L. Vega, “A bilinear approach to the restriction and Keakeya conjectures”, *J. Amer. Math. Soc.* **11**:4 (1998), 967–1000. MR 99f:42026 Zbl 0924.42008
- [Wolff 1995] T. Wolff, “An improved bound for Keakeya type maximal functions”, *Rev. Mat. Iberoamericana* **11**:3 (1995), 651–674. MR 96m:42034 Zbl 0848.42015
- [Wolff 1998a] T. Wolff, “A mixed norm estimate for the X-ray transform”, *Rev. Mat. Iberoamericana* **14**:3 (1998), 561–600. MR 2000j:44006 Zbl 0927.44002
- [Wolff 1998b] T. Wolff, “Recent work connected with the Keakeya problem”, pp. 129–162 in *Prospects in mathematics* (Princeton, NJ, 1996), edited by H. Rossi, Amer. Math. Soc., Providence, RI, 1998. MR 2000d:42010 Zbl 0934.42014

Received April 19, 2002. Revised September 10, 2002.

TERENCE TAO
DEPARTMENT OF MATHEMATICS
UCLA
LOS ANGELES CA 90095-1555
tao@math.ucla.edu
<http://www.math.ucla.edu/~tao>

